

Vibrio fluvialis in Patients with Diarrhea, Kolkata, India

Goutam Chowdhury, Gururaja P. Pazhani,
Devarati Dutta, Sucharita Guin,
Sanjucta Dutta, Santanu Ghosh,
Hidemasa Izumiya, Masahiro Asakura,
Shinji Yamasaki, Yoshifumi Takeda, Eiji Arakawa,
Haruo Watanabe, Asish K. Mukhopadhyay, Mihir
K. Bhattacharya, K. Rajendran,
Gopinath Balakrish Nair,
and Thandavarayan Ramamurthy

We identified 131 strains of *Vibrio fluvialis* among 400 nonagglutinating *Vibrio* spp. isolated from patients with diarrhea in Kolkata, India. For 43 patients, *V. fluvialis* was the sole pathogen identified. Most strains harbored genes encoding hemolysin and metalloprotease; this finding may contribute to understanding of the pathogenicity of *V. fluvialis*.

Many members of the family *Vibrionaceae* cause diarrheal disease; among these, *Vibrio cholerae* O1/O139 and *V. parahaemolyticus* are responsible for several epidemics and pandemics (1,2). In Indonesia, >20% of diarrheal infections are caused by pathogenic *Vibrio* spp. (3). Some of these *Vibrio* spp. can grow in thiosulfate–citrate–bile salts–sucrose agar as yellow colonies and do not agglutinate with *V. cholerae* O1 antiserum. These species are broadly defined as nonagglutinating (NAG) vibrios.

The emerging etiologic agent *V. fluvialis* has caused sporadic cases and outbreaks of diarrhea in several countries (4–6). Species-specific minimal biochemical tests, e.g., lysine decarboxylase, ornithine decarboxylase, arginine dihydrolase, and L-arabinose, are used to identify

V. fluvialis; without these tests, it may be confused with NAG vibrios, *V. cholerae*, and even *Aeromonas* spp. In most resource-poor countries, these tests are not performed, which may lead to labeling of *V. fluvialis* as a NAG vibrio.

Although *V. fluvialis* is known to cause diarrhea, the mechanisms involved in its pathogenicity are not well established. To evaluate the prevalence of *V. fluvialis* in India and possible mischaracterization as an NAG vibrio, we examined cases in which isolates from hospitalized patients with diarrhea were identified as NAG vibrios and characterized the strains using phenotypic and genetic methods.

The Study

We examined 400 isolates identified as NAG vibrios that were collected during 2002–2009 from 11,904 stool specimens from patients with diarrhea admitted to the Infectious Diseases and Beliaghata General Hospital, Kolkata, India. Specimens were screened for common enteric pathogens, according to standard protocols (7). Oxidase, string test, and arginine dihydrolase–positive strains that did not agglutinate with *V. cholerae* O1 polyvalent or O139 monovalent antiserum were further confirmed as *V. fluvialis* by using a multiplex PCR targeting the *toxR* gene of *V. fluvialis* and the *ompW* gene of *V. cholerae* (8,9). Isolates were also subjected to PCRs targeting different virulence-associated genes encoding the repeat in toxin (*rtxA*, *rtxC*), heat-stable enterotoxin (*stn*), type 3 secretion system (*vcsC2*, *vcsV2*, *vcsN2* and *vspD*), cholera toxin (*ctxA*), toxin co-regulated pilus (*tcpA*), thermostable direct-hemolysin (*tdh*), TDH-related hemolysin (*trh*), *V. fluvialis* hemolysin (VFH), and metalloproteases, according to published methods (10–12).

Expression of VFH was determined in vitro by using erythrocytes from rabbit and sheep. Cytotoxin assay was performed with HeLa and Chinese hamster ovary cell lines by using sterile culture filters of the *V. fluvialis* strains that were isolated as a sole pathogen. Antimicrobial drug susceptibility testing was performed by using the disk diffusion method with commercially available disks (Becton Dickinson, Sparks Glencoe, MD, USA), according to Clinical and Laboratory Standards Institute criteria (13). Because these guidelines do not include interpretive criteria for *V. fluvialis*, breakpoints for *Enterobacteriaceae* were adopted. *Escherichia coli* ATCC 25922 was used as a quality control strain.

Pulsed-field gel electrophoresis was performed according to the PulseNet standardized protocol for *V. cholerae* (14). Gel Compare II software (Applied Maths NV, Sint-Martens-Latem, Belgium) was used for electrophoresis pattern comparison that runs on Dice similarity index and unweighted pairgroup with arithmetic mean method.

Author affiliations: National Institute of Cholera and Enteric Diseases, Kolkata, India (G. Chowdhury, G.P. Pazhani, D. Dutta, S. Guin, S. Dutta, S. Ghosh, A.K. Mukhopadhyay, M.K. Bhattacharya, K. Rajendran, G. Nair, T. Ramamurthy); National Institute of Infectious Diseases, Tokyo, Japan (H. Izumiya, E. Arakawa, H. Watanabe); Osaka Prefecture University Graduate School of Life and Environmental Sciences, Osaka, Japan (M. Asakura, S. Yamasaki); and National Institute of Cholera and Enteric Diseases Collaborative Research Center of Okayama University for Infectious Diseases in India, Kolkata (Y. Takeda).

DOI: <http://dx.doi.org/10.3201/eid1811.120520>

Table 1. Prevalence of *Vibrio fluvialis* among patients with diarrhea, Kolkata, India, 2002–2009

Year	No. samples	No. (%) <i>V. fluvialis</i> isolates	No. (%) patients	
			Sole infection	Mixed infection
2002	2,285	16 (0.7)	5 (0.2)	11 (0.5)
2003	1,673	8 (0.5)	1 (0.1)	7 (0.4)
2004	2,430	19 (0.8)	6 (0.2)	13 (0.5)
2005	1,472	17 (1.1)	7 (0.5)	10 (0.7)
2006	930	12 (1.3)	4 (0.4)	8 (0.9)
2007	842	9 (1.1)	2 (0.2)	7 (0.8)
2008	1,124	24 (2.1)	8 (0.7)	16 (1.4)
2009	1,153	26 (2.2)	10 (0.9)	16 (1.4)
Total	11,909	131 (1.1)	43 (0.4)	88 (0.7)

Among the 400 isolates presumptively identified NAG vibrios, multiplex PCR confirmed 131 and 269 strains (each strain representing a case) as *V. fluvialis* and *V. cholerae*, respectively. The overall prevalence rate of *V. fluvialis* among 11,904 hospitalized patients with diarrhea was 1.1%. Abrupt appearance of *V. fluvialis* was identified in 2002, although the surveillance of diarrheal infection was initiated at the Infectious Diseases and Beliaghata General Hospital in 1996 (www.niced.org.in/annual_reports.htm). The isolation rate of *V. fluvialis* gradually increased from 0.7% in 2002 to 2.2% in 2009 (Table 1). Of the 131 strains of *V. fluvialis*, 43 (33%) were identified as the sole pathogen; the remaining 88 (67%) were isolated as a co-pathogen with either *V. cholerae*, *V. parahaemolyticus*, *E. coli*, *Shigella* spp., parasites, or enteric viruses (data not shown). Among the mixed infections, *V. fluvialis* with *V. cholerae* was isolated most often (17%), followed by *V. fluvialis* and *V. parahaemolyticus* (8%). The presence of *Vibrio* spp. as mixed pathogens indicates that these patients likely acquired the infection from contaminated water or food. We analyzed the date of admission and place from where the patients resided and found no evidence for clusters of infection or small outbreaks caused by *V. fluvialis*.

V. fluvialis infection was much more often detected in adults (73%) than in children <5 years of age (27%). Clinical symptoms of sole infection caused by *V. fluvialis* were similar to that of cholera: watery diarrhea (86%), severe dehydration status (28%), and abdominal pain (12%) (Table 2). Several previous investigations have identified cholera-like diarrheal outbreaks caused by *V. fluvialis* (4,5).

All the *V. fluvialis* strains were negative for the virulence genes commonly reported in *V. cholerae* and *V. parahaemolyticus*, but >90% were positive for genes encoding VFH and metalloproteases. More than 80% of the strains expressed hemolysin against rabbit and sheep red blood cells. Hemolysin is a widely distributed virulence factor in most pathogenic *Vibrio* spp. Metalloprotease produced by *V. fluvialis* is related to hemagglutination proteases of *V. vulnificus*, which enhances permeability and hemorrhagic activities (12). These factors may increase the virulence of *V. fluvialis* and contribute to diarrhea.

When the culture filtrates were tested, cytotoxic effect was readily noticed in the Chinese hamster ovary and HeLa cell lines, i.e., cytoplasmic vacuolation, cell rounding, and destruction of the monolayer. In most strains isolated as a sole pathogen, the cytotoxic endpoint titer was 2–256 (online Technical Appendix Table 1, wwwnc.cdc.gov/EID/pdfs/12-0520-Techapp.pdf). The cell vacuolation phenomenon has been reported as a virulence factor in several enteric pathogens (online Technical Appendix References).

In this study, *V. fluvialis* strains were highly resistant to ampicillin (92%), streptomycin (85%), furazolidone (85%), and sulfamethoxazole/trimethoprim (70%) (online Technical Appendix Table 2). About half the number of strains were resistant to ciprofloxacin and 45% to nalidixic acid; the lower resistance rate for nalidixic acid compared with fluoroquinolones is unexpected and warrants further investigation to confirm the additional mechanisms. In a previous study, we found that some *V. fluvialis* strains carried the plasmid-mediated quinolone resistance gene allele *qnrA1* and a gene encoding the aminoglycoside acetyltransferase (*aac(6′)-Ib-cr*), which reduces ciprofloxacin activity (15). Fluoroquinolone resistance and intermediate susceptibility to erythromycin (92%) are the unique features of the *V. fluvialis* isolated in this study;

Table 2. Clinical features of *Vibrio fluvialis*-infected patients with diarrhea, Kolkata, India, 2002–2009

Clinical feature	No. (%) patients	
	Sole infection	Mixed infection
Type of diarrhea		
Watery	36 (86)	72 (81)
Bloody mucus, loose	7 (16)	16 (19)
Dehydration status		
Severe	12 (28)	14 (16)
Some or rare	31 (72)	74 (84)
Fever		
Yes	4 (9)	9 (10)
No	39 (91)	79 (90)
Abdominal pain		
Yes	5 (12)	11 (12)
No	38 (88)	77 (88)
Age		
>5 y	30 (70)	66 (75)
≤5 y	13 (30)	22 (25)
Sex		
M	23 (53)	58 (66)
F	20 (47)	30 (34)




Figure. Dendrogram of *NotI*-digested pulsed-field gel electrophoresis (PFGE) profiles with representative *Vibrio fluvialis* isolates. Clustering identified 4 clades (A–D). AM, ampicillin; S, streptomycin; G, gentamicin; NA, nalidixic acid; CIP, ciprofloxacin; C, chloramphenicol; E, erythromycin; SXT, sulfamethoxazole-trimethoprim; FZ, furazolidone; NOR, norfloxacin. Scale bar indicates degree of similarity.

this trend was not recorded in other *Vibrio* spp., e.g., *V. cholerae* and *V. parahaemolyticus*.

Although the *V. fluvialis* strains exhibited distinct *NotI* restriction profiles in the denrogram analysis, at least 4 major clades were identified (Figure). Clades A and B, with strains isolated during 2002–2007, exhibited less antimicrobial drug resistance than did clade C and D strains identified during 2008–2009; multidrug-resistant strains, especially those resistant to fluoroquinolones, were identified in higher numbers in clades C and D (Figure).

Conclusions

Our results demonstrate an emerging trend of prevalence of *V. fluvialis* among patients with acute diarrhea patients in Kolkata. The expression of cytotoxic activity and hemolysin may contribute to understanding the pathogenicity of *V. fluvialis*. Further epidemiologic studies are necessary to elucidate the public health importance of *V. fluvialis*-mediated diarrhea.

This work was supported in part by the Ministry of Health, Labor and Family Welfare of Japan (Project H17-Shinkou-3); Initiative for Global Research Network on Infectious Diseases, Ministry of Education, Culture, Sports, Science and Technology, Japan; and intramural grants of the Indian Council of Medical Research, New Delhi, India.

Mr Chowdhury is a doctoral candidate at the National Institute of Cholera and Enteric Diseases, Kolkata, India. His main research interest is the pathogenesis and molecular biology of enteric *Vibrio* spp.

References

- Kaper JB, Morris JG Jr, Levine MM. Cholera. *Clin Microbiol Rev.* 1995;8:48–86.
- Nair GB, Ramamurthy T, Bhattacharya SK, Dutta B, Takeda Y, Sack DA. Global dissemination of *Vibrio parahaemolyticus* serotype O3:K6 and its serovariants. *Clin Microbiol Rev.* 2007;20:39–48. <http://dx.doi.org/10.1128/CMR.00025-06>
- Lesmana M, Subekti DS, Tjaniadi P, Simanjuntak CH, Punjabi NH, Campbell JR, et al. Spectrum of *Vibrio* species associated with acute diarrhea in North Jakarta, Indonesia. *Diagn Microbiol Infect Dis.* 2002;43:91–7. [http://dx.doi.org/10.1016/S0732-8893\(02\)00373-5](http://dx.doi.org/10.1016/S0732-8893(02)00373-5)
- Huq MI, Alam AKMJ, Brenner DJ, Morris GK. Isolation of vibrio-like group EF-6, from patients with diarrhoea. *J Clin Microbiol.* 1980;11:621–4.
- Thekdi R, Lakhani AG, Vachha SM, Chandrakapure MR. *Vibrio fluvialis* (group F *Vibrio*) in Maharashtra. *Indian J Med Res.* 1982;76:80–5.
- Kobayashi K, Ohnaka T. Food poisoning due to newly recognized pathogens. *Asian Med J.* 1989;32:1–12.
- World Health Organization. Manual for laboratory identification of acute enteric infections. Geneva: The Organization; 1987.
- Chakraborty R, Sinha S, Mukhopadhyay AK, Asakura M, Yamasaki S, Bhattacharya SK, et al. Species-specific identification of *Vibrio fluvialis* by PCR targeted to the conserved transcriptional activation and variable membrane tether regions of the *toxR* gene. *J Med Microbiol.* 2006;55:805–8. <http://dx.doi.org/10.1099/jmm.0.46395-0>
- Nandi B, Nandy RK, Mukhopadhyay S, Nair GB, Shimada T, Ghose AC. Rapid method for species-specific identification of *Vibrio cholerae* using primers targeted to the gene of outer membrane protein *OmpW*. *J Clin Microbiol.* 2000;38:4145–51.
- Chatterjee S, Ghosh K, Raychoudhuri A, Chowdhury G, Bhattacharya MK, Mukhopadhyay AK, et al. Incidence, virulence factors, and clonality among clinical strains of non-O1, non-O139 *Vibrio cholerae* isolates from hospitalized diarrheal patients in Kolkata, India. *J Clin Microbiol.* 2009;47:1087–95. <http://dx.doi.org/10.1128/JCM.02026-08>
- Han JH, Lee JH, Choi YH, Park JH, Choi TJ, Kong IS. Purification, characterization and molecular cloning of *Vibrio fluvialis* hemolysin. *Biochim Biophys Acta.* 2002;1599:106–14. [http://dx.doi.org/10.1016/S1570-9639\(02\)00407-7](http://dx.doi.org/10.1016/S1570-9639(02)00407-7)
- Miyoshi S, Sonoda Y, Wakiyama H, Rahman MM, Tomochika K, Shinoda S, et al. An exocellular thermolysin-like metalloprotease produced by *Vibrio fluvialis*: purification, characterization, and gene cloning. *Microb Pathog.* 2002;33:127–34. <http://dx.doi.org/10.1006/mpat.2002.0520>
- Clinical and Laboratory Standards Institute. Performance standards for antimicrobial susceptibility testing. Document M100–S21. Wayne (PA): The Institute; 2011.

14. Kam KM, Luey CK, Tsang YM, Law CP, Chu MY, Cheung TL, et al. Molecular subtyping of *Vibrio cholerae* O1 and O139 by pulsed-field gel electrophoresis in Hong Kong: correlation with epidemiological events from 1994 to 2002. *J Clin Microbiol*. 2003;41:4502–11. <http://dx.doi.org/10.1128/JCM.41.10.4502-4511.2003>
15. Chowdhury G, Pazhani GP, Nair GB, Ghosh A, Ramamurthy T. Transferable plasmid-mediated quinolone resistance in association with extended-spectrum β -lactamases and fluoroquinolone-acetylating aminoglycoside-6'-N-acetyltransferase in clinical isolates of *Vibrio fluvialis*. *Int J Antimicrob Agents*. 2011;38:169–73.

Address for correspondence: Thandavarayan Ramamurthy, National Institute of Cholera and Enteric Diseases, P-33, CIT Road, Scheme XM, Beliaghata, Kolkata 700010, India; email: tramu@vsnl.net

The opinions expressed by authors contributing to this journal do not necessarily reflect the opinions of the Centers for Disease Control and Prevention or the institutions with which the authors are affiliated.

etymologia

Coxsackievirus

[kok-sak'e-vi'ræs]

Named for Coxsackie, the small town on the Hudson River where they were first isolated, human coxsackieviruses are nonenveloped, positive-sense, single-stranded RNA viruses in the family *Picornaviridae*, genus *Enterovirus*. They were first described by Gilbert Dalldorf, who was investigating suspected poliomyelitis outbreaks in upstate New York in the summer of 1947. Coxsackieviruses are divided into 2 groups, A and B. In suckling mice, group A viruses cause generalized myositis and flaccid paralysis, and group B viruses cause focal myositis and spastic paralysis. With the discovery of coxsackieviruses, Dalldorf also helped popularize the suckling mouse as an inexpensive laboratory animal model.

Sources

1. Dalldorf G. The Coxsackie viruses. *Bull N Y Acad Med*. 1950;26:329–35.
2. Dalldorf G, Sickles GM, Plager H, Gifford R. A virus recovered from the feces of "poliomyelitis" patients pathogenic for suckling mice. *J Exp Med*. 1949;89:567–82. <http://dx.doi.org/10.1084/jem.89.6.567>
3. Dorland's Illustrated Medical Dictionary. 32nd ed. Philadelphia: Elsevier Saunders; 2012.
4. Racaniello V. Coxsackie NY and the virus named after it, August 10, 2009 [cited 2012 Aug 21]. <http://www.virology.ws/2009/08/10/coxsackie-ny-and-the-virus-named-after-it/>
5. Tao Z, Song Y, Li Y, Liu Y, Jiang P, Lin X, et al. Coxsackievirus B3, Shandong Province, China, 1990–2010. *Emerg Infect Dis*. 2012;18:zzz–zzz.

Address for correspondence: Ronnie Henry, Centers for Disease Control and Prevention, 1600 Clifton Rd NE, Mailstop E03, Atlanta, GA 30333, USA; email: boq3@cdc.gov

DOI: <http://dx.doi.org/10.3201/eid1811.ET1811>

Medscape
CME[™]

Sign up to receive email announcements when
a new article is available.

Get an online subscription at wwwnc.cdc.gov/eid/subscribe.htm

Vibrio fluvialis in Patients with Diarrhea, Kolkata, India

Technical Appendix

Technical Appendix Table 1. Results of assays of clinical *Vibrio fluvialis* strains to determine ability to lyse rabbit erythrocytes and cytotoxic effect on CHO and HeLa cells.

Strain identification	<i>V. fluvialis</i> hemolysin	<i>V. fluvialis</i> metalloprotease	Hemolytic titer	Cytotoxic titer	
				Chinese hamster ovary	HeLa
IDH00612	+	+	16	64	4
IDH00629	+	+	2	4	2
IDH00653	+	+	4	32	16
IDH01036	+	+	32	32	8
IDH01577	+	+	4	16	16
H8942	+	+	8	16	8
H17768	+	+	16	8	4
L15318	+	+	2	2	2
K24681	+	+	64	128	256
J11969	+	+	32	32	128

Technical Appendix Table 2. Antimicrobial drug resistance of *Vibrio fluvialis*

Drug	Resistant, %	Intermediate, %
Ampicillin	92	8
Chloramphenicol	15	45
Ciprofloxacin	50	20
Sulfamethoxazole/ trimethoprim	70	14
Erythromycin	8	92
Furazolidone	85	15
Gentamicin	22	10
Nalidixic acid	45	15
Neomycin	10	88
Norfloxacin	40	35
Streptomycin	85	15
Tetracycline	10	15

References

1. Figueroa-Arredondo P, Heuser JE, Akopyants NS, Morisaki JH, Giono-Cerezo S, Enríquez-Rincón F, et al. Cell vacuolation caused by *Vibrio cholerae* hemolysin. *Infect Immun*. 2001; 69:1613–24.
2. Abrami L, Fivaz M, Glauser PE, Parton R, Van der Goot FG. A pore-forming toxin interacts with a GP-I-anchored protein and causes vacuolization of the endoplasmic reticulum. *J. Cell Biol*. 1998; 140:525–40.
3. Istivan TS, Coloe PJ, Fry BN, Ward P, Smith SC. Characterization of a haemolytic phospholipase A(2) activity in clinical isolates of *Campylobacter concisus*. *J Med Microbiol*. 2004;53:483–93.