

Molecular typing analysis showed that both case-patients had a monoclonal infection caused by *C. difficile* ribotype 027. Both isolates had *tcdA*, *tcdB*, and *cdtB* and had a deletion in *tcdC* (data not shown).

In summary, the described severe cases of *C. difficile* infection in Chile were caused by epidemic *C. difficile* ribotype 027. One of these case-patients required urgent colectomy. These results demonstrate that epidemic *C. difficile* 027 strains are present in South America, highlighting the need for enhanced screening for this ribotype in other regions of the continent.

Acknowledgments

We thank Nigel Minton (University of Nottingham) for kindly providing *C. difficile* strain R20291.

This work was supported by grants from the Comisión Nacional de Investigación en Ciencia y Tecnología (FONDECYT REGULAR 1100971) to M.A.-L. and S.B. and by grants from MECESUP UAB0802, Comisión Nacional de Investigación en Ciencia y Tecnología (FONDECYT REGULAR 1110569) and from the Research Office of Universidad Andrés Bello (DI-35-11/R) (to D.P.-S); and from the Medical Research Center, Facultad de Medicina of the Pontificia Universidad Católica de Chile, Resident Research Project (PG-20/11) to C. H.-R.

**Cristian Hernández-Rocha,¹
Jonathan Barra-Carrasco,¹
Marjorie Pizarro-Guajardo,
Patricio Ibáñez,
Susan M. Bueno,
Mahfuzur R. Sarker,
Ana María Guzman,
Manuel Álvarez-Lobos,
and Daniel Paredes-Sabja**

Author affiliations: Pontificia Universidad Católica de Chile, Santiago, Chile (C. Hernández-Rocha, P. Ibáñez, S.M. Bueno, A.M. Guzman, M. Álvarez-Lobos.);

¹These authors contributed equally to this article.

Universidad Andrés Bello, Santiago (J. Barra-Carrasco, M. Pizarro-Guajardo, D. Paredes-Sabja); and Oregon State University, Corvallis, Oregon, USA (M.R. Sarker, D. Paredes-Sabja)

DOI: <http://dx.doi.org/10.3201/eid1808.120211>

References

- McDonald LC, Killgore GE, Thompson A, Owens RC Jr, Kazakova SV, Sambol SP, et al. An epidemic, toxin gene-variant strain of *Clostridium difficile*. *N Engl J Med*. 2005;353:2433–41. <http://dx.doi.org/10.1056/NEJMoa051590>
- Quesada-Gómez C, Rodríguez C, Gamboa-Coronado Mdel M, Rodríguez-Cavallini E, Du T, Mulvey MR, et al. Emergence of *Clostridium difficile* NAP1 in Latin America. *J Clin Microbiol*. 2010;48:669–70. <http://dx.doi.org/10.1128/JCM.02196-09>
- Balassiano IT, Yates EA, Domingues RM, Ferreira EO. *Clostridium difficile*: a problem of concern in developed countries and still a mystery in Latin America. *J Med Microbiol*. 2012;61:169–79. <http://dx.doi.org/10.1099/jmm.0.037077-0>
- Wilson KH, Kennedy MJ, Fekety FR. Use of sodium taurocholate to enhance spore recovery on a medium selective for *Clostridium difficile*. *J Clin Microbiol*. 1982;15:443–6.
- Dawson LF, Donahue EH, Cartman ST, Barton RH, Bundy J, McNerney R, et al. The analysis of para-cresol production and tolerance in *Clostridium difficile* 027 and 012 strains. *BMC Microbiol*. 2011;11:86. <http://dx.doi.org/10.1186/1471-2180-11-86>
- Fedorko DP, Williams EC. Use of cycloserine-cefoxitin-fructose agar and L-proline-aminopeptidase (PRO Discs) in the rapid identification of *Clostridium difficile*. *J Clin Microbiol*. 1997;35:1258–9.
- Rupnik M. *Clostridium difficile* toxinotyping. *Methods Mol Biol*. 2010;646:67–76. http://dx.doi.org/10.1007/978-1-60327-365-7_5
- Bidet P, Barbut F, Lalande V, Burghoffer B, Petit JC. Development of a new PCR-ribotyping method for *Clostridium difficile* based on ribosomal RNA gene sequencing. *FEMS Microbiol Lett*. 1999;175:261–6. <http://dx.doi.org/10.1111/j.1574-6968.1999.tb13629.x>

Address for correspondence: Daniel Paredes-Sabja, Laboratorio de Mecanismos de Patogénesis Bacteriana, Departamento de Ciencias Biológicas, Facultad de Ciencias Biológicas, Universidad Andrés Bello, Santiago, Chile; email: daniel.paredes.sabja@gmail.com

Zoonotic Pathogens among White-Tailed Deer, Northern Mexico, 2004–2009

To the Editor: Intense wildlife management for hunting affects risks associated with zoonotic pathogens (1). White-tailed deer (*Odocoileus virginianus*) are increasingly managed by fencing, feeding, watering, and translocation to increase incomes from hunting in northern Mexico (2). These deer also play a major role in dissemination and reintroduction of pathogens and vectors from Mexico into the United States (3,4). White-tailed deer are suitable reservoir hosts for *Mycobacterium bovis* (1), and an *M. bovis*-positive white-tailed deer was recently found in Tamaulipas in northeastern Mexico (2). Brucellosis is widespread in many animal hosts in Latin America (5) and thus of interest in white-tailed deer. Another major zoonosis, sometimes linked to raw deer meat consumption, is hepatitis E, which is caused by genotypes of hepatitis E virus (HEV) (6). HEV is increasingly prevalent in red deer (*Cervus elaphus*) (7), but its prevalence in white-tailed deer is unknown.

The objective of this study was to determine the prevalence of zoonotic pathogens in white-tailed deer in northern Mexico. This study was conducted under a scientific collecting permit issued by the Mexican Division of Animal and Wildlife Health and on 8 ranches in 3 states in northern Mexico (≈26–28°N, 99–100°W).

Serum samples (n = 347) were collected during 2004–2009 in a cross-sectional survey for antibodies against HEV, *Brucella* spp., and mycobacteria. Deer were opportunistically sampled during live-capture operations as described by Cantú et al. (8). Bleeding was performed by using jugular venipuncture and vacuum tubes

without anticoagulant. Samples were allowed to clot and centrifuged to collect serum that was stored at -20°C .

Serum samples were tested for IgG against HEV by ELISA as described (7). Serum samples were also tested for antibodies against *Brucella* spp. by using a commercial ELISA (Ingezim Brucella Compac 2.0 Ingenasa, Madrid, Spain), according to the manufacturer's instructions. Detection of antibodies cross-reacting with 2 widely used mycobacterial antigens, bovine purified protein derivative (PPD) and paratuberculosis protoplasmatic antigen 3 (PPA3), was conducted as described (9). The sensitivity and specificity of this assay have not been established for white-tailed deer, but it has been used in seroprevalence studies of wild boar and fallow deer (9,10).

Insufficient volumes of serum samples prevented testing for antibodies against all pathogens (Table). Limited serum volume and lack of other (organ) samples also precluded additional analyses to verify presence of pathogens.

Prevalence was 62.7% (95% CI 54%–70%) for antibodies against HEV, 0.4% (95% CI 0%–2%) for antibodies against *Brucella* spp., 8.9% (95% CI 6%–13%) for antibodies against bovine PPD, and 2.6% (95% CI 1%–5%) for antibodies against PPA3 (Table). Antibody responses to bovine PPD were detected in deer from at 6 of 8 sampling sites; in deer

from 3 of these sites, antibodies were also detected against PPA3 antigen. Seroprevalence against bovine PPD was higher than that against PPA3 (χ^2 10.9, df 1, $p < 0.01$).

This cross-sectional survey of white-tailed deer in northern Mexico detected antibodies to several pathogens relevant to public and animal health. High prevalence of antibodies against HEV and frequent detection of antibodies against mycobacterial antigens are public health concerns. Prevalence of antibodies against HEV were 3× higher than that reported for red deer in Europe (7). This result suggests wide circulation of HEV in the study region and warrants further research, including detection and sequencing of virus RNA. Low *Brucella* spp. antibody prevalence confirms results of a study in this region (8).

Antibody responses to bovine PPD were detected in serum samples from deer from most sampling sites, occasionally in the absence of antibodies against PPA3. These results, and a recent report of an *M. bovis*-positive white-tailed deer from this region (2), suggest that these deer may be contracting *M. bovis* in northern Mexico. If one considers that white-tailed deer are *M. bovis* reservoirs in other parts of North America and that risk factors such as supplemental feeding are present in northern Mexico, there is a high risk for pathogen transmission to animals and humans (1). White-tailed deer are probably exposed to several

pathogens that are relevant to animal health in northern Mexico.

Although this cross-sectional survey provided only an indication of pathogen prevalence in the study populations, high antibody prevalence to HEV and mycobacterial antigens requires antigen-targeted surveillance. Risks associated with pathogen translocation by white-tailed deer are also relevant to neighboring states in Mexico and the United States.

Acknowledgments

We thank M. Villar and B. Peralta for providing ELISA antigen and A. Cantú and co-workers for providing deer serum samples.

This study was supported by the Caesar Kleberg Wildlife Research Institute, Texas A&M University–Kingsville, and European Union Framework Program grant 212414 TB-STEP.

**Citlaly Medrano,
Mariona Boadella,
Hugo Barrios, Antonio Cantú,
Zeferino García,
José de la Fuente,
and Christian Gortazar**

Author affiliations: Instituto de Investigación en Recursos Cinegéticos, Ciudad Real, Spain (C. Medrano, M. Boadella, J. de la Fuente, C. Gortazar); Universidad Autónoma de Tamaulipas, Ciudad Victoria, Mexico (C. Medrano, H. Barrios); Instituto Nacional de Investigaciones Forestales, Agrícolas, y Pecuarias, Tamaulipas, Mexico (A. Cantú); Instituto Nacional de Investigaciones Forestales, Agrícolas, y

Table. Prevalence of serum antibodies against zoonotic pathogen antigens among white-tailed deer on 8 ranches, northern Mexico, 2004–2009*

Ranch no.	State	Municipality	Sample size	No. positive/no. tested (%)			
				HEV	<i>Brucella</i> spp.	Bovine PPD	PPA3
1	Tamaulipas	Guerrero	1	0/1 (0)	0/1 (0)	0/1 (0)	0/1 (0)
2	Tamaulipas	Guerrero	106	41/71 (57.7)	0/102 (0)	9/102 (8.8)	0/102 (0)
3	Tamaulipas	Nuevo Laredo	37	21/27 (77.7)	0	0/37 (0)	0/37 (0)
4	Nuevo León	Unknown	35	0	0	3/35 (8.5)	2/35 (5.7)
5	Coahuila	Guerrero	35	1/1 (100)	0/33 (0)	2/33 (6.0)	1/33 (3.0)
6	Coahuila	Guerrero	50	6/17 (35.2)	0/40 (0)	3/40 (7.5)	0/40 (0)
7	Coahuila	Hidalgo	17	4/6 (66.6)	1/16 (6.2)	3/16 (18.7)	5/16 (31.2)
8	Coahuila	Guerrero	66	16/19 (84.2)	0/39 (0)	7/39 (18.0)	0/39 (0)
NA	Total	NA	347	89/142 (62.7)	1/231 (0.4)	27/303 (8.9)	8/303 (2.6)

*HEV, hepatitis E virus; PPD, purified protein derivative; PPA3, paratuberculosis protoplasmatic antigen 3; NA, not applicable.

Pecuarías, Morelos, Mexico (Z. García); and Oklahoma State University, Stillwater, Oklahoma, USA (J. de la Fuente)

DOI: <http://dx.doi.org/10.3201/eid1808.111902>

References

- Miller R, Kaneene JB, Fitzgerald SD, Schmitt SM. Evaluation of the influence of supplemental feeding of white-tailed deer (*Odocoileus virginianus*) on the prevalence of bovine tuberculosis in the Michigan wild deer population. *J Wildl Dis.* 2003;39:84–95.
- Barrios-García HB, Guizarnotegui-Blanco JA, Zapata-Campos CC, Almazán-García C, González-Alanis P, Villareal-Peña R, et al. Identification of *Mycobacterium tuberculosis* complex by histopathology and PCR in white-tailed deer (*Odocoileus virginianus*) in Tamaulipas, Mexico. *Journal of Animal and Veterinary Advances.* 2012;11:1036–40.
- Pérez de León AA, Strickman DA, Knowles DP, Fish D, Thacker E, De La Fuente J, et al. One Health approach to identify research needs in bovine and human babesioses: workshop report. *Parasit Vectors.* 2010;3:36.
- Pound JM, George JE, Kammlah DM, Lohmeyer KH, Davey RB. Evidence for role of white-tailed deer (*Artiodactyla: Cervidae*) in epizootiology of cattle ticks and southern cattle ticks (*Acari: Ixodidae*) in reinfestations along the Texas/Mexico border in south Texas: a review and update. *J Econ Entomol.* 2010;103:211–8. <http://dx.doi.org/10.1603/EC09359>
- Lucero NE, Ayala SM, Escobar GJ, Jacob NR. *Brucella* isolated in humans and animals in Latin America from 1968 to 2006. *Epidemiol Infect.* 2008;136:496–503. <http://dx.doi.org/10.1017/S0950268807008795>
- Cooper K, Huang FF, Batista L, Rayo CD, Bezanilla JC, Toth TE, et al. Identification of genotype 3 hepatitis E virus (HEV) in serum and fecal samples from pigs in Thailand and Mexico, where genotype 1 and 2 HEV strains are prevalent in the respective human populations. *J Clin Microbiol.* 2005;43:1684–8. <http://dx.doi.org/10.1128/JCM.43.4.1684-1688.2005>
- Boadella M, Casas M, Martín M, Vicente J, Segalés J, de la Fuente J, et al. Increasing contact with hepatitis E virus in red deer, Spain. *Emerg Infect Dis.* 2010;16:1994–6.
- Cantú A, Ortega-S JA, Mosqueda J, García-Vázquez Z, Henke SE, George JE. Prevalence of infectious agents in free-ranging white-tailed deer in northeastern Mexico. *J Wildl Dis.* 2008;44:1002–7.
- Boadella M, Lyashchenko K, Greenwald R, Esfandiari J, Jaroso R, Carta T, et al. Serologic tests for detecting antibodies against *Mycobacterium bovis* and *Mycobacterium avium* subspecies *paratuberculosis* in Eurasian wild boar (*Sus scrofa scrofa*). *J Vet Diagn Invest.* 2011;23:77–83. <http://dx.doi.org/10.1177/104063871102300111>
- Boadella M, Barasona JA, Diaz-Sanchez S, Lyashchenko KP, Greenwald R, Esfandiari J, et al. Performance of immunochromatographic and ELISA tests for detecting fallow deer infected with *Mycobacterium bovis*. *Prev Vet Med.* 2012;104:160–4. <http://dx.doi.org/10.1016/j.prevetmed.2011.10.005>

Address for correspondence: Christian Gortazar, Instituto de Investigación en Recursos Cinegéticos, Ronda de Toledo s/n, 13005 Ciudad Real, Spain; email: christian.gortazar@uclm.es

KIs Virus and Blood Donors, France

To the Editor: KIs-V is a new putative virus identified recently in the blood of persons in Japan (1). First partial sequence of KIs-V was characterized unexpectedly, when PCR primers were used that were directed primarily to the consensus domain of helicase of positive-stranded RNA viruses. Extensive physicochemical and molecular analysis suggested that KIs-V is an enveloped virus with a circular, double-stranded DNA genome of ≈9,500 bp (prototype isolate: GenBank accession no. AB550431); its genetic diversity is presumed to be extremely low because the 4 complete genomes already characterized in Japan harbor strict identical sequences. The 13 potential genes identified by in silico analysis exhibit an overall low sequence homology to other known viral proteins (1). Until now, KIs-V epidemiologic data have been related

only to the original study in which the authors analyzed plasma samples from 516 blood donors categorized into 4 groups by alanine aminotransferase (ALT) level (either ≤60 IU/L or >60 IU/L) and the presence or absence of hepatitis E virus (HEV) antibodies. As a result, KIs-V DNA was detectable at elevated prevalence in the high ALT level/HEV antibody-positive group (36%, n = 100); viral loads, checked for a few samples, ranged from 10⁶ to 10⁸ copies/mL. KIs-V DNA also was identified in HEV antibody-negative samples, with low or high ALT level (≈0.8%, n = 120, and 1%, n = 100, respectively) (1).

To gain insights about the potential presence of this virus in the blood of persons in France, we investigated KIs-V DNA in the plasma of 576 healthy blood donors (mean age 40 years; 306 men; men:women 1:1.13). Blood samples were collected in vacuum tubes (Vacutainer, SST, Becton Dickinson, Meylan, France) and centrifuged, and plasma aliquots were stored at –80°C until use. Nucleic acids were extracted from 1-mL plasma volumes (MagNA Pure LC, Roche Diagnostics, Meylan, France) (2) and tested for KIs-V DNA by using the same nested PCR system for screening Japanese blood donors (1). Briefly, one tenth (5 μL) of extracted nucleic acids were first amplified by using primers 101-C (5'-CAACACCGCA ATCACAAAGT-3') and N101-B (5'-AACATTGAAACGTCATGT CC-3') (0.8 μM each) in a 50-μL mix containing deoxynucleotide triphosphates (0.2 mM each) (Roche) and 2 U Taq DNA polymerase (Invitrogen, Cergy Pontoise, France) with its corresponding buffer. One microliter of the amplification mixture was subsequently used in a second-round PCR with primers KS-2 (5'-CTCGTCCTCGTCGTCATC GTA-3') and N101-D (5'-CA TTTGCTCCCGCTGGAGATG-3') under the same conditions as above. The amplification conditions for first-