

**Joana Abrantes, Ana M. Lopes,
Kevin P. Dalton, Pedro Melo,
Jorge J. Correia,
Margarida Ramada,
Paulo C. Alves, Francisco Parra,
and Pedro J. Esteves**

Author affiliations: Centro de Investigação em Biodiversidade e Recursos Genéticos/InBIO Laboratório Associado—Universidade do Porto (CIBIO/UP), Vairão, Portugal (J. Abrantes, A.M. Lopes, P.C. Alves, P.J. Esteves); Université de Nantes, Nantes, France (J. Abrantes, A.M. Lopes); Faculdade de Ciências da Universidade do Porto, Porto, Portugal (A.M. Lopes, P.C. Alves); Universidad de Oviedo, Asturias, Spain (K.P. Dalton, F. Parra); VetNatura, Lisbon, Portugal (P. Melo, M. Ramada); Faculdade de Medicina Veterinária—Universidade Técnica de Lisboa, Lisbon (J.J. Correia); University of Montana, Missoula Montana, USA (P.C. Alves); and Instituto de Investigação e Formação Avançada em Ciências e Tecnologias da Saúde (CESPU), Gandra, Portugal (P.J. Esteves)

DOI: <http://dx.doi.org/10.3201/eid1911.130908>

References

1. Abrantes J, van der Loo W, Le Pendu J, Esteves PJ. Rabbit haemorrhagic disease (RHD) and rabbit haemorrhagic disease virus (RHDV): a review. *Vet Res.* 2012;43:12. <http://dx.doi.org/10.1186/1297-9716-43-12>
2. Alda F, Gaitero T, Suarez M, Merchan T, Rocha G, Doadrio I. Evolutionary history and molecular epidemiology of rabbit haemorrhagic disease virus in the Iberian Peninsula and Western Europe. *BMC Evol Biol.* 2010;10:347. <http://dx.doi.org/10.1186/1471-2148-10-347>
3. Muller A, Freitas J, Silva E, Le Gall-Recule G, Zwingelstein F, Abrantes J, et al. Evolution of rabbit haemorrhagic disease virus (RHDV) in the European rabbit (*Oryctolagus cuniculus*) from the Iberian Peninsula. *Vet Microbiol.* 2009;135:368–73. <http://dx.doi.org/10.1016/j.vetmic.2008.09.057>
4. Dalton KP, Nicieza I, Balseiro A, Mugerza MA, Rosell JM, Casais R, et al. Variant rabbit hemorrhagic disease virus in young rabbits, Spain. *Emerg Infect Dis.* 2012;18:2009–12. <http://dx.doi.org/10.3201/eid1812.120341>
5. Le Gall-Reculé G, Zwingelstein F, Boucher S, Le Normand B, Plassiart G, Portejoie Y, et al. Detection of a new variant of rabbit haemorrhagic disease virus in France. *Vet Rec.* 2011;168:137–8. <http://dx.doi.org/10.1136/vr.d697>
6. Capucci L, Fallacara F, Grazioli S, Lavazza A, Pacciarini ML, Brocchi E. A further step in the evolution of rabbit hemorrhagic disease virus: the appearance of the first consistent antigenic variant. *Virus Res.* 1998;58:115–26. [http://dx.doi.org/10.1016/S0168-1702\(98\)00106-3](http://dx.doi.org/10.1016/S0168-1702(98)00106-3)
7. Abrantes J, Esteves PJ. Not-so-novel Michigan rabbit calicivirus. *Emerg Infect Dis.* 2010;16:1331–2. <http://dx.doi.org/10.3201/eid1608.091803>
8. Capucci L, Fusi P, Lavazza A, Pacciarini ML, Rossi C. Detection and preliminary characterization of a new rabbit calicivirus related to rabbit hemorrhagic disease virus but nonpathogenic. *J Virol.* 1996;70:8614–23.
9. Strive T, Wright JD, Robinson AJ. Identification and partial characterisation of a new Lagovirus in Australian wild rabbits. *Virology.* 2009;384:97–105. <http://dx.doi.org/10.1016/j.virol.2008.11.004>
10. Tamura K, Peterson D, Peterson N, Stecher G, Nei M, Kumar S. MEGA5: molecular evolutionary genetics analysis using maximum likelihood, evolutionary distance, and maximum parsimony methods. *Mol Biol Evol.* 2011;28:2731–9. <http://dx.doi.org/10.1093/molbev/msr121>

Address for correspondence: Pedro J. Esteves, CIBIO-UP, Centro de Investigação em Biodiversidade e Recursos Genéticos, Universidade do Porto, InBIO, Laboratório Associado, Campus Agrário de Vairão, Rua Padre Armando Quintas, Vairão, Portugal; email: pjesteves@cibio.up.pt

Letters

Letters commenting on recent articles as well as letters reporting cases, outbreaks, or original research are welcome. Letters commenting on articles should contain no more than 300 words and 5 references; they are more likely to be published if submitted within 4 weeks of the original article's publication. Letters reporting cases, outbreaks, or original research should contain no more than 800 words and 10 references. They may have 1 Figure or Table and should not be divided into sections. All letters should contain material not previously published and include a word count.

***Mycobacterium yongonense* in Pulmonary Disease, Italy**

To the Editor: *Mycobacterium yongonense* is a recently described species (*I*) that belongs to the *M. avium* complex (MAC) and is associated with pulmonary infection. The strain on which the description of species was based was isolated in South Korea from the sputum of a patient with unspecified pulmonary disease. We describe 2 *M. yongonense* strains isolated from patients in Italy.

Patient 1 was a 74-year-old woman who had experienced fatigue, diarrhea, and weight loss. Her medical history included liver cirrhosis resulting from hepatitis C virus infection and surgery for colon cancer; the patient also reported tuberculosis in childhood. Chest radiograph revealed a cavitory lesion, a finding confirmed by computed tomography scan (Figure). Cultures in liquid and solid media grew a nonchromogenic mycobacterium from sputum and stool samples; results were negative for urine samples.

The patient was treated with clarithromycin, rifabutin, and ethambutol and showed some improvement. A bronchoscopic investigation was performed, and microscopic examination of bronchoalveolar lavage samples revealed the presence of acid-fast bacilli that subsequently were grown in culture. The patient began improving markedly starting with the second month of treatment, which will be continued for a total of 18 months.

Patient 2 was a 74-year-old woman, living in a community of nuns, who reported cough and dyspnea. Her medical history included renal failure and surgery for breast cancer. A bronchoalveolar lavage was performed; samples yielded in culture *Pseudomonas aeruginosa* and a nonchromogenic mycobacterium.




Figure. Computed tomography scan showing a cavity (arrow) in the left lung of a 74-year-old woman (patient 1) in Italy. Laboratory testing suggests that the woman was infected with *Mycobacterium yongonense*.

The patient was treated with cefepime, to which *P. aeruginosa* was susceptible in vitro, and rapidly improved. The isolation of the nontuberculous mycobacterium was considered irrelevant, and no specific treatment was undertaken.

To determine the specific mycobacteria species isolated from these patients, we conducted a commercial line-probe assay (GenoType Mycobacterium CM; Hain Lifesciences, Nehren, Germany). Both strains were identified as *M. intracellulare*. However, the known cross-reaction of *M. intracellulare* probe with most MAC species (2) led us to determine the complete sequence of the 16S rRNA gene. Both strains showed 100% similarity with *M. yongonense* and *M. marseillense* (3) strains.

To confirm this unusual finding, we investigated other genetic regions. We detected 100% identity with *M. yongonense* in the internal transcribed spacer 1 region and in a 1,384-bp region of the *hsp65* gene and found 2 mismatches in a 420-bp fragment of the *sodA* gene (99.5%

similarity). In contrast, *M. marseillense* showed 6 mismatches (98.6% similarity) in the internal transcribed spacer 1 region and 24 (98.3% similarity) in *hsp65*; no *sodA* sequence is available in GenBank for this species. Partial sequencing of other genetic targets not available in GenBank for *M. yongonense* enabled us to confirm the close relatedness of the strains to *M. intracellulare* (100% similarity in *dnaK* gene; 99.3% identity in *gyrB* and *gyrC* genes).

The finding of the same novel *Mycobacterium* species in these 2 unrelated patients reflects variability in the significance of nontuberculous mycobacteria isolated from clinical specimens. *M. yongonense* was probably a contaminant in the second case, but in the first, its involvement as causative agent of disease seems incontrovertible. The specific criteria of the American Thoracic Society (4) were fulfilled: radiographic imaging clearly documented the presence of a cavitory pulmonary lesion, no other pathogen possibly responsible of disease was detected by bronchoscopic

investigation, and the same mycobacterium was isolated repeatedly from sputum (its presence in stool probably results from swallowed sputum) and bronchoalveolar lavage samples. Confirmation is further provided by the response to the specific therapy, according to international guidelines (4,5), for MAC pulmonary disease (MICs were 2, 1, and 8 $\mu\text{g/mL}$ for clarithromycin, rifabutin, and ethambutol, respectively).

The initial description of *M. yongonense* noted that it has a distinct *rpoB* sequence (1), identical to that of a distantly related scotochromogenic species, *M. parascrofulaceum*. In a more recent article (6), the same authors investigated 2 more strains of *M. yongonense* with similar characteristics and suggested that the recent acquisition of the *rpoB* gene resulted from a lateral gene transfer event from *M. parascrofulaceum*. The *rpoB* genes of the strains we investigated, however, were substantially different from that of *M. scrofulaceum* and were instead related to that of *M. intracellulare* (99.4% similarity) and, less closely, to that of other species belonging to the MAC, including *M. marseillense* (97.4%). Discrepancy in the *rpoB* sequence means some uncertainty remains that our strains are *M. yongonense*, but the 100% identity in major phylogenetically relevant regions strongly supports this hypothesis and suggests the possibility of a variant of the species preceding the acquisition of the *rpoB* gene from *M. parascrofulaceum*. Less evidence exists for identifying the strains as *M. marseillense* because of the clear divergence in the genes investigated, other than 16S rRNA.

The complete epidemiology of *M. yongonense* is unknown, in part because few strains have been identified. However, as in the cases we describe, use of suboptimal identification methods may mean that some isolates have been misidentified as other mycobacteria species.

GenBank accession numbers for the *M. yongonense* strains identified in this study (FI-13004 and FI-13005) are KF224989–KF224999.

**Enrico Tortoli,
Alessandro Mariottini,
Piera Pierotti,
Tullia M. Simonetti,
and Gian Maria Rossolini**

Author affiliations: San Raffaele Scientific Institute, Milan, Italy (E. Tortoli); Careggi University Hospital, Florence, Italy (A. Mariottini, M.T. Simonetti, G.M. Rossolini); SS. Maria Annunziata Hospital, Florence (P. Pierotti); University of Florence, Florence (G.M. Rossolini); and University of Siena, Siena, Italy (G.M. Rossolini)

DOI: <http://dx.doi.org/10.3201/eid1911.130911>

References

- Kim BJ, Math RK, Jeon CO, Yu HK, Park YG, Kook YH, et al. *Mycobacterium yongonense* sp. nov., a slow-growing nonchromogenic species closely related to *Mycobacterium intracellulare*. *Int J Syst Evol Microbiol*. 2013;63:192–9. <http://dx.doi.org/10.1099/ijs.0.037465-0>
- Tortoli E, Pecorari M, Fabio G, Messinò M, Fabio A. Commercial DNA probes for mycobacteria incorrectly identify a number of less frequently encountered species. *J Clin Microbiol*. 2010;48:307–10. <http://dx.doi.org/10.1128/JCM.01536-09>
- Ben Salah I, Cayrou C, Raoult D, Drancourt M. *Mycobacterium marseillense* sp. nov., *Mycobacterium timonense* sp. nov. and *Mycobacterium bouchedurhonense* sp. nov., novel species in the *Mycobacterium avium* complex. *Int J Syst Evol Microbiol*. 2009;59:2803–8. <http://dx.doi.org/10.1099/ijs.0.010637-0>
- Griffith DE, Aksamit T, Brown-Elliott BA, Catanzaro A, Daley C, Gordin F, et al. An official ATS/IDSA statement: diagnosis, treatment and prevention of nontuberculous mycobacterial disease. *Am J Respir Crit Care Med*. 2007;175:367–416. <http://dx.doi.org/10.1164/rccm.200604-571ST>
- Clinical Laboratory and Standards Institute. Susceptibility testing of mycobacteria, nocardiae and other aerobic actinomycetes; approved standard. 2nd edition, M24–A2. Wayne (PA); The Institute; 2011.
- Kim BJ, Hong SH, Kook YH, Kim BJ. Molecular evidence of lateral gene transfer in *rpoB* gene of *Mycobacterium yongonense* strains via multilocus sequence analysis. *PLoS ONE*. 2013;8:e51846. <http://dx.doi.org/10.1371/journal.pone.0051846>

Address for correspondence: Enrico Tortoli, Emerging Bacterial Pathogens Unit, San Gabriele Bldg, San Raffaele Scientific Institute, via Olgettina 58, 20132 Milan, Italy; email: tortoli.enrico@hsr.it

Subcutaneous Infection with *Dirofilaria* spp. Nematode in Human, France

To the Editor: The article by Foissac et al. titled Subcutaneous infection with *Dirofilaria immitis* nematode in human, France (1) presents an interesting and challenging diagnostic dilemma. The paper described, but did not illustrate, the worm as having a strongly ridged external surface of the cuticle—a feature known not to exist on *Dirofilaria immitis*, the dog heartworm. However,

molecular sequencing of the specimen demonstrated much closer similarity to *D. immitis* than to *D. repens*, the most common cause of zoonotic subcutaneous dirofilariasis infection in Europe.

Well-described morphologic features of parasites, including in tissue sections, have long been the standard for diagnosis. More recently, molecular diagnostics have helped in many of these difficult cases. However, in some cases, the morphology and molecular diagnosis are discordant. On the basis of the data in the article, the worm does not seem to represent *D. repens*. A more likely possibility is some other species for which no sequences are yet available for comparison. In such a worm, the regions sequenced must be similar to *D. immitis*, and distinct from *D. repens*, to achieve the observed results.

When one encounters a case such as this, where well-validated morphologic features (Figure) are contradictory to the molecular analysis, one must exercise caution in arriving at a final diagnosis. One disadvantage



Figure. Cross-section of the filarial nematode seen in the subcutaneous nodule on the thigh of a woman in France. The features, as described in the original report (1), include prominent, longitudinal ridging of the cuticle (arrows), 2 reproductive tubes, and the intestine (asterisk). Scale bar indicates 50 μ m. Image courtesy of Jean-Philippe Dales.