

in China. Although reassortment was not detected in this co-infection, a potential risk for emergence of a new pandemic strain by reassortment between these 2 viruses (with humans as mixing vessels) should not be ignored. To reduce the risk for emergence of new viral subtypes, the public health and scientific communities should enhance surveillance for co-infection with influenza (H7N9) virus and other influenza virus subtypes.

This research was supported by the National Megaprojects of China for Infectious Disease (nos. 2012ZX10004211 and 2014ZX10004002-003-004), National Natural Science Foundation of China (nos. 81341004, 81102283, and 81370131), Outstanding Academic Leader of Health System in Shanghai (no. XBR2013078), Ministry of Science and Technology (no. KJYJ-2013-01-01), Shanghai Municipal Health and Family Planning Commission (no. 2013QLG002), Key Discipline Construction Project of Pudong Health Bureau of Shanghai (no. PWZx2014-10), Academic Leader Training Project in Health System of Pudong Health Bureau of Shanghai (no. PWRd2010-01), and Key Medical Specialties of Shanghai (no. ZK2012A28).

References

1. Zhu Y, Qi X, Cui L, Zhou M, Wang H. Human co-infection with novel avian influenza A H7N9 and influenza A H3N2 viruses in Jiangsu Province, China. *Lancet*. 2013;381:2134. [http://dx.doi.org/10.1016/S0140-6736\(13\)61135-6](http://dx.doi.org/10.1016/S0140-6736(13)61135-6)
2. Li J, Kou Y, Yu X, Sun Y, Zhou Y, Pu X, et al. Human co-infection with avian influenza and seasonal influenza viruses, China. *Emerg Infect Dis*. 2014;20:1953–5. <http://dx.doi.org/10.3201/eid2011.140897>
3. Ramakrishnan MA, Tu Z, Singh S, Chockalingam A, Gramer M, Wang P, et al. The feasibility of using high resolution genome sequencing of influenza A viruses to detect mixed infections and quasispecies. *PLoS ONE*. 2009;4:e7105. <http://dx.doi.org/10.1371/journal.pone.0007105>
4. Chan KH, To K, Hung I, Zhang A, Chan J, Cheng V, et al. Differences in antibody responses of individuals with natural infection and those vaccinated against pandemic H1N1 2009 influenza. *Clin Vaccine Immunol*. 2011;18:867–73. <http://dx.doi.org/10.1128/CVI.00555-10>
5. Dong L, Bo H, Bai T, Gao R, Dong J, Zhang Y, et al. Combination of serological assays to detect human antibodies to the avian influenza A H7N9 virus. *PLoS ONE*. 2014;9:e95612. <http://dx.doi.org/10.1371/journal.pone.0095612>
6. Zhang W, He Y, Xu L, Dai F, Mei Z, Qian L, et al. Full-genome analysis of influenza A(H7N9) virus from Shanghai, China, 2014. *Genome Announc*. 2014;2:e00578–14. <http://dx.doi.org/10.1128/genomeA.00578-14>

Address for correspondence: Yunwen Hu, No. 2901 Caolang Rd, Jinshan District, Shanghai 201508, China; email: yw.hu0117@126.com; and Tao Ren, No. 150 Jimo Rd, Pudong New Area, Shanghai 200120, China; email: rentao305@163.com

Nairobi Sheep Disease Virus RNA in Ixodid Ticks, China, 2013

Shangshu Gong,¹ Biao He,¹ Zedong Wang,¹ Limin Shang, Feng Wei, Quan Liu, Changchun Tu

Author affiliations: Military Veterinary Institute, Academy of Military Medical Sciences, Key Laboratory of Jilin Province for Zoonosis Prevention and Control, Changchun, People's Republic of China (S. Gong, B. He, Z. Wang, L. Shang, Q. Liu, C. Tu); College of Life Science, Jilin Agricultural University, Changchun (F. Wei); Jiangsu Co-innovation Center for Prevention and Control of Important Animal Infectious Diseases and Zoonoses, Yangzhou, People's Republic of China (C. Tu)

DOI: <http://dx.doi.org/10.3201/eid2104.141602>

To the Editor: Nairobi sheep disease virus (NSDV; genus *Nairovirus*, family *Bunyaviridae*) causes acute hemorrhagic gastroenteritis in sheep and goats (1,2). First identified in Nairobi, Kenya, in 1910, it is considered endemic in East Africa (1,3). Ganjam virus, a variant of NSDV, is found in India and Sri Lanka (2). NSDV has a limited effect on animals bred in areas to which the virus is endemic but can cause large losses of animals during introduction of new livestock or transport of animals through these areas (4). In humans, NSDV infection can cause febrile illness, headache, nausea, and vomiting (5).

Ticks are the main transmission vectors of NSDV and many other viral pathogens and therefore pose a major threat to public health (6,7). Here, we describe a newly discovered NSDV, named NSDV (China), identified by viral metagenomic analysis of ticks collected from the northeast region of the People's Republic of China (Liaoning, Jilin, and Heilongjiang provinces) during May–July, 2013, and divided into 9 groups according to tick species and sampling sites. Four tick species were morphologically identified: *Haemaphysalis longicornis* (84.8%); *Dermacentor silvarum* (7.2%); *D. nuttalli* (5.5%); and *Ixodes persulcatus* (2.5%) (online Technical Appendix Table 1, <http://wwwnc.cdc.gov/EID/article/21/4/14-1602-Techapp1.pdf>).

Of the 6,427 ticks collected, 3,410 were divided into 9 groups (average 379 ticks/group, range 163–512); each group was homogenized in SM buffer (50 mmol/L Tris, 10 mmol/L MgSO₄, 0.1 mol/L NaCl, pH 7.5). Viral RNA extraction, Solexa sequencing, and analysis are described in the online Technical Appendix. Among the sequences annotated to mammalian viruses, 65 contigs were found to target the small (n = 15), medium (n = 27), and large (n = 23) segments of the NSDV genome (online Technical Appendix Tables 2–4).

¹These authors contributed equally to this article.

To confirm the Solexa results, a 376-nt fragment of the NSDV small gene segment was amplified by reverse transcription PCR (RT-PCR) by using primers P1 (5'-AG-CAAAGAGCACATTGACTGGGC-3') and P2 (5'-CTGT-CACACCTGCCTTCCAA-3'). Ticks in 3 *H. longicornis* groups were positive for NSDV: group 1 from sheep in Jian, Jilin Province (125°34'E, 40°52'N); group 2 from cattle in Jinxing, Jilin Province (130°38'E, 42°25'N); and group 5 from sheep in Dandong, Liaoning Province (124°23'E, 40°07'N). Ticks in the other groups were negative. The obtained sequences shared 92% identity with NSDV from *H. intermedia* in India.

The full-length sequence of NSDV was then obtained from group 2 by RT-PCR by using primers based on the Solexa sequences or the conserved sequences of nairoviruses (online Technical Appendix Table 5). The complete sequences of the small, medium, and large segments of NSDV (China) (GenBank accession nos. KM464724–KM464726) contained 1,590, 5,077, and 12,081 nt, respectively; that is, they were similar to other NSDVs. Sequence comparisons showed 75.1%–89.6% identity with other NSDVs at the nucleotide level and 81.3%–96.7% at the deduced amino acid level (online Technical Appendix Table 6). Compared with other member species within the genus *Nairovirus* (Dugbe, Kupe, Hazara, and Crimean Congo hemorrhagic fever viruses), low identities (37.5%–68.6%) were observed at both nucleotide and amino acid levels (online Technical Appendix Table 6). Phylogenetic analysis based on the amino acid sequences grouped the virus together with NSDVs from Africa and South Asia (Figure).

The remaining tick samples of the NSDV-positive groups were used to determine the infection frequency by using RT-PCR to analyze primers P1 and P2. We assayed 104 tick pools (average 15 ticks/pool, range 8–40), 13 pools of 416 ticks in Jian Province and 91 pools of 1,095 ticks in Jinxing Province; 12.5% (13/104) tested positive, 38.5% (5/13) in Jian and 8.8% (8/91) in Jinxing. The higher prevalence in Jian Province may result from more ticks in the pools. Attempts to isolate virus from the positive samples in cell lines (Vero and BHK-21) and suckling mice were unsuccessful; thus, its pathogenicity could not be determined.

In Africa, NSDV is primarily transmitted by *R. appendiculatus* ticks (5). In South Asia (India and Sri Lanka), NSDV has been isolated from ticks (*H. intermedia*, *H. wellingtoni*, and *R. haemaphysaloides*), mosquitoes, sheep and humans; *H. intermedia* ticks are considered the main vector for the virus (5,8,9). NSDV had not previously been reported from East Asia. The isolate we identified, NSDV (China), is genetically divergent from the NSDVs of South Asia and Africa and is therefore a novel strain, with *H. longicornis* likely the main vector. Nairobi sheep disease has not been reported in China and East Asia, but our results




Figure. Phylogenetic analysis of Nairobi sheep disease virus (China) and other nairoviruses. The phylogenetic trees were generated in MEGA5.2 software (<http://www.megasoftware.net>). The complete coding regions for nucleocapsid protein in the small segment (A), glycoprotein precursor in the medium segment (B), and RNA dependent RNA polymerase in the large segment (C) were analyzed by the maximum-likelihood method. An emergent severe fever thrombocytopenia syndrome virus (SFTSV; family *Bunyaviridae*, genus *Phlebovirus*) was used as the outgroup. Bootstrap testing (1,000 replicates) was performed, and the bootstrap values are indicated. Sequences are identified by their GenBank accession numbers, followed by the virus name, host, and country. Black triangles indicate novel strain NSDV (China). Scale bars indicate substitutions per site. CCHFV, Crimean-Congo hemorrhagic fever virus.

indicate the risk of its occurrence in these regions, where *H. longicornis* is widely distributed (10). More extensive investigation to clarify the natural circulation of NSDV among ticks should be conducted and surveillance of sheep improved to prevent outbreaks of Nairobi sheep disease in China and East Asia.

This work was supported by the Science and Technology Basic Work Program from the Ministry of Science and Technology of China (2013FY113600), and the Military Medical Innovation Program of Academy of Military Medical Sciences (2012CXJJ019).

References

1. Montgomery E. On a tick-borne gastro-enteritis of sheep and goats occurring in East Africa. *J Comp Pathol Ther.* 1917;30:28–57. [http://dx.doi.org/10.1016/S0368-1742\(17\)80002-3](http://dx.doi.org/10.1016/S0368-1742(17)80002-3)
2. Marczinke BI, Nichol ST. Nairobi sheep disease virus, an important tick-borne pathogen of sheep and goats in Africa, is also present in Asia. *Virology.* 2002;303:146–51. <http://dx.doi.org/10.1006/viro.2002.1514>
3. Weinbren MP, Gourlay RN, Lumsden WHR, Weinbren WM. An epizootic of Nairobi sheep disease in Uganda. *J Comp Pathol Ther.* 1958;68:174–87. [http://dx.doi.org/10.1016/S0368-1742\(58\)80018-1](http://dx.doi.org/10.1016/S0368-1742(58)80018-1)
4. Lasecka L, Baron MD. The nairovirus Nairobi sheep disease virus/Ganjam virus induces the translocation of protein disulphide isomerase-like oxidoreductases from the endoplasmic reticulum to the cell surface and the extracellular space. *PLoS ONE.* 2014;9:e94656. <http://dx.doi.org/10.1371/journal.pone.0094656>
5. Yadav PD, Vincent MJ, Khristova M, Kale C, Nichol ST, Mishra AC, et al. Genomic analysis reveals Nairobi sheep disease virus to be highly diverse and present in both Africa, and in India in the form of the Ganjam virus variant. *Infect Genet Evol.* 2011;11:1111–20. <http://dx.doi.org/10.1016/j.meegid.2011.04.001>
6. Perera LP, Peiris JSM, Weilgama DJ, Calisher CH, Shope RE. Nairobi sheep disease virus isolated from *Haemaphysalis intermedia* ticks collected in Sri Lanka. *Ann Trop Med Parasitol.* 1996;90:91–3.
7. Liu Q, He B, Huang SY, Wei F, Zhu XQ. Severe fever with thrombocytopenia syndrome, an emerging tick-borne zoonosis. *Lancet Infect Dis.* 2014;14:763–72. [http://dx.doi.org/10.1016/S1473-3099\(14\)70718-2](http://dx.doi.org/10.1016/S1473-3099(14)70718-2)
8. Rajagopalan PK, Sreenivasan MA, Paul SD. Isolation of Ganjam virus from the bird tick *Haemaphysalis wellingtoni* Nuttall and Warburton 1907. *Indian J Med Res.* 1970;58:1195–6.
9. Joshi MV, Geevarghese G, Joshi GD, Ghodke YS, Mourya DT, Mishra AC. Isolation of Ganjam virus from ticks collected of domestic animals around Pune, Maharashtra, India. *J Med Entomol.* 2005;42:204–6. <http://dx.doi.org/10.1093/jmedent/42.2.204>
10. Hoogstraal H, Roberts FH, Kohls GM, Tipton VJ. Review of *Haemaphysalis* (Kaiseriana) *longicornis* Neumann (resurrected) of Australia, New Zealand, New Caledonia, Fiji, Japan, Korea, and Northeastern China and USSR, and its parthenogenetic and bisexual populations (Ixodoidea, Ixodidae). *J Parasitol.* 1968;54:1197–213. <http://dx.doi.org/10.2307/3276992>

Address for correspondence: Quan Liu, Key Laboratory of Jilin Province for Zoonosis Prevention and Control, Military Veterinary Institute, Academy of Military Medical Sciences, 666 Liuying West Rd, Jingyue Economic Development Zone, Changchun, 130122, People's Republic of China; email: liuquan1973@hotmail.com

Avian Influenza A(H10N7) Virus–Associated Mass Deaths among Harbor Seals

Rogier Bodewes, Theo M. Bestebroer, Erhard van der Vries, Josanne H. Verhagen, Sander Herfst, Marion P. Koopmans, Ron A.M. Fouchier, Vanessa M. Pfankuche, Peter Wohlsein, Ursula Siebert, Wolfgang Baumgärtner, Albert D.M.E. Osterhaus

Author affiliations: Erasmus Medical Centre, Rotterdam, the Netherlands (R. Bodewes, T.M. Bestebroer, E. van der Vries, J.H. Verhagen, S. Herfst, M.P. Koopmans, R.A.M. Fouchier, A.D.M.E. Osterhaus); University of Veterinary Medicine, Hannover, Germany (V.M. Pfankuche, P. Wohlsein, U. Siebert, W. Baumgärtner, A.D.M.E. Osterhaus); Artemis One Health, Utrecht, the Netherlands (A.D.M.E. Osterhaus)

DOI: <http://dx.doi.org/10.3201/eid2104.141675>

To the Editor: Avian influenza A viruses occasionally cross the species barrier; influenza A(H5N1) virus and the recently emerged influenza A(H7N9) virus are prime examples of bird-to-human transmission (1,2). In addition, avian influenza A viruses can cross to various other mammalian species, including pinnipeds (e.g., seals) (3,4).

Recently, mass deaths have occurred among harbor seals (*Phoca vitulina*); hundreds of carcasses washed up the shores of Sweden (March 2014), Denmark (July 2014), and Germany (October 2014). Approximately 1,400 dead harbor seals were seen in the coastal waters of Schleswig-Holstein in Germany alone, where the population is ≈12,000 animals.

We report the investigation of the deaths of 17 seals from different age groups that were found dead on the islands of Helgoland and Sylt, Germany, during the second week of October 2014. Complete necropsies were performed on the carcasses, which were in variable nutritional conditions, ranging from very poor to good. Necropsy results showed consistently poorly retracted lungs with severe congestion, occasional diffuse consolidation, and multifocal firm nodular areas of gray-yellow discoloration with varying numbers of metazoic parasites. Histologic examinations showed acute necrotizing bronchitis and adenitis of bronchial glands with sloughing of epithelial cells (Figure, panel A). Occasionally, mild interstitial pneumonia was found. Multifocal pyogranulomatous to necrotizing pneumonia was associated with parasite infestation. A few animals had suppurative to necrotizing or nonsuppurative rhinitis and tracheitis.

Because mass deaths among seals were caused by phocine distemper virus in the same area in 1988 and 2002,

Nairobi Sheep Disease Virus RNA in Ixodid Ticks, China, 2013

Technical Appendix

Tick Collection, RNA extraction and Processing, Solexa Sequencing, and Analysis of Data

In the present study, ticks were collected respectively from sheep, cattle and by netting over the vegetation (Technical Appendix Table 1). Of the ticks collected from animals, some were fed, but the ticks collected from grass were unfed. All the ticks were adult with most of them being female. The ticks were pooled and examined according to their species and sampling site.

Tick samples were prepared for megagenomic analysis as described by He et al. (2013). The tick number of each pool was shown in Technical Appendix Table 1. A total of 9 groups were pooled and homogenized in SM buffer [1:10 (w/v); 50 mM Tris, 10 mM MgSO₄, 0.1 M NaCl, pH7.5], respectively. The homogenized samples were centrifuged at 8000×g at 4°C for 30 min to remove cell debris and foreign materials, and the supernatants were immediately filtered through 0.45-μm and 0.22-μm filters (Millipore, Billerica, MA). Host genome and other free nucleic acids were eliminated by digestion of DNase (Ambion, Austin, TX), Benzonase Nuclease (Novagen, San Diego, CA) and RNase I (Fermentas, Ontario, Canada).

The viral RNAs and DNAs were then extracted immediately using TRIzol (TaKaRa, Dalian, China) according to the manufacturer's protocol. Total viral nucleic acids were dissolved in RNase-free H₂O (TaKaRa) and used immediately for the following reverse transcription with SuperScript III reverse transcription (Invitrogen, Carlsbad, CA) using random primers according to the manufacturer's protocol. To synthesize dsDNA, a Klenow fragment (New England

Biolabs, Beijing, China) was added to the cDNA mixture, and incubated at 37°C for 60 min. After inactivation of the enzyme, phosphates and free single-stranded bases in the dscDNA reaction was removed using shrimp alkaline phosphatase and exonuclease I (TaKaRa).

To obtain sufficient viral nucleic acid, SISPA was employed to amplify the dscDNA with the Accuprime Taq DNA Polymerase System (Invitrogen) according to the manufacturer's protocol. Briefly, a 50 µl reaction system containing 10 µl of the above dscDNA mixture, random primers (20 mM), 10×Accuprime buffer I, and Taq DNA Polymerase (1 U) was denatured at 94°C for 2 min, followed by 40 cycles of 94°C denaturing for 30 s, 55°C annealing for 30 s, 68°C extending for 1 min with final 68°C extension for 8 min. The PCR products were then purified using the QIAquick PCR Purification Kit (Qiagen, Hilden, Germany) and dissolved in 50 µl TE buffer (100 mM Tris-HCl, 10 mM EDTA, pH8.0). The purified PCR products of the 9 groups were pooled together and then subjected to Solexa sequencing in one lane by the Beijing Genome Institute (BGI, Shenzhen, China).

Virus Isolation Procedure

The homogenates were prepared in cooled medium, and virus isolation was conducted according to the standard procedures. Briefly, NSDV-positive samples were ground with cold serum-free minimum essential medium (MEM; Sigma-Aldrich) and centrifuged at 12,000 × g for 10 min at 4°C. Supernatants were passed through 0.22-µm syringe filters (Sartorius), and the filtrates were added to Vero and BHK cell lines with MEM containing 2% fetal bovine serum (FBS; GIBCO). The cell cultures were observed daily during incubation with 5% CO₂ at 37°C and passaged for 5 times. The cell cultures of each passage were subjected to RT-PCR screening of NSDV.

References

1. Chen SG, Pan DM. Identification of tick species in China. In: Lu BL, ed. Identification handbook of Chinese important medical animals. China: People Health Press,1982: 752–838.
2. He B, Li Z, Yang F, Zheng J, Feng Y, Guo H, et al. Virome profiling of bats from Myanmar by metagenomic analysis of tissue samples reveals more novel mammalian viruses. PLoS ONE. 2013;8:e61950.

Technical Appendix Table 1. Tick species, origin, and number screened and analyzed for presence of Nairobi sheep disease, China, 2013

Group*	Tick species†	Sampling site	Animal host‡	No. collected ticks (%)	No. ticks for	
					metagenomic analysis	No. remaining ticks§
1	<i>Haemaphysalis longicornis</i>	Jian, Jilin (125°34' E, 40°52' N)	Sheep	920 (14.3)	504	416
2	<i>H. longicornis</i>	Jinxing, Jilin (130°38' E, 42°25' N)	Cattle	1,525 (23.7)	430	1,095
3	<i>H. longicornis</i>	Chunhua, Jilin (131°04' E, 43°11' N)	Cattle	679 (10.6)	512	167
4	<i>H. longicornis</i>	Nanshan, Jilin (130°47' E, 42°84' N)	Cattle	1,826 (28.4)	487	1,339
5	<i>H. longicornis</i>	Dandong, Liaoning (124°23' E, 40° 07' N)	Sheep	326 (5.1)	326	0
6	<i>H. longicornis</i>	Huma, Heilongjiang (125°03' E, 50°49' N)	No	174 (2.7)	174	0
Subtotal				5,450 (84.8)	2,433	3,017
7	<i>Dermacentor nuttalli</i>	Harbin, Heilongjiang (125°42' E, 44°04' N)	No	143 (2.2)	143	0
		Songlin, Jilin (130°54' E, 42°84' N)	Cattle	78 (1.2)	78	0
		Huma, Heilongjiang (125°03'20" E, 50°49' N)	No	130 (2.0)	130	0
Subtotal				351 (5.5)	351	0
8	<i>D. silvarum</i>	Harbin, Heilongjiang (125°42' E, 44°04' N)	No	56 (0.9)	56	0
		Songlin, Jilin	Cattle	182 (2.8)	182	0
		Huma, Heilongjiang (125°03' E, 50°49' N)	No	225 (3.5)	225	0
Subtotal				463 (7.2)	463	0
9	<i>Ixodes persulcatus</i>	Harbin, Heilongjiang (125°42'-130°10' E, 44°04'-46°40' N)	No	31 (0.5)	31	0
		Nanshan, Jilin (130°28' E, 42°28' N)	Cattle	62 (1.0)	62	0
		Chunhua, Jilin (131°04' E, 43°11' N)	Cattle	23 (0.3)	23	0
		Huma, Heilongjiang (125°03' E, 50°49' N)	No	47 (0.7)	47	0
Subtotal				163 (2.5)	163	0
Total				6,427	3,410	3,017

* Group 1–6: each group comprises ticks collected from a single site. Group 7–9: each group comprises pooled ticks collected from multiple sites.

† Species was identified according to tick morphology described by Cheng and Pang (1992).

Group*	Tick species†	Sampling site	Animal host‡	No. collected ticks (%)	No. ticks for	
					metagenomic analysis	No. remaining ticks§

‡ No indicates that the ticks were collected by sweep netting the vegetation.

§ This column shows the numbers of remaining ticks not subjected to metagenomic analysis, in which the ticks only in groups 1 and 2 were analyzed for prevalence testing since these 2 groups were NSDV RT-PCR positive.

Technical Appendix Table 2. Sequences of S gene contigs and their identities to NSDV strain 708, China, 2013

Contig	Location	Sequence (5'→3')	Sequence identity (%)
es.1	678–877	CAAGTTCCTTGTTGTCTTCAATCCACCATGGGGTGATATCAACAAGGCAGGAAAGTCAGGGATTGCCTTAGCAGCGAC AGGTATGGCAAATTGATAGAGCTGGATGGTCCCAAATTGCAGAGGACCTGAGGGAATCTCTGAAGGGTCTTGTGG CATGGATCAATGCCACAAGGATGAAGTGGAGAACGGTAAAGAGG	87
2	865–1109	AACGGTAAAGAGGTTGTTGATGGTTTGACCAAGCACCTGCAGAAAGCCCTTGAATTAGCCAAGCAATCAAGTGCCATG AGAGCCCAAGGGGCTCAGATTGACACTGTCTTTAGCAGCTACTACTGGCTTTGGAAGGCAGGTGTGACAGCGGAGAT GTTCCGACAGTCTCACAGTTTCTTTTTGAGCTCGGCAAGGTGCCAGGGGAAATAAAAAATGAAGAAAGCACTATC AAGTATGCCTCT	92
3	1096–1259	TCAAGTATGCCTCTGAAGTGGGAAAGAAGTTGCTAGCACTCTTGTGATGATAGCTTCACTGCTAATCGGATTTACA TGCACCCTGGTGTCTAACAGCTGGGAGAATGTCTGAGCTCGGTGTTTGTCTCGGAGCAATCCCAGTTGCCAATCCT GACGATGC	92
4	1368–1478	AGGCTTCGACATTGAGAGCATGGACATAGTTGCCTCGGAGCATCTGCTACACCAGTCACTCGTTGGAAGAGATCTCC TTTTCAGAATGCCTACAACATCCGGGGAAATGC	90
5	394–572	ACGGAAGTACCCAGCCCTGAACAACCTGCTGGGTTACCAGAGAGCAGCTCTTAAGTGGAGAAAGGATACCAAGTACGG GATCAACAGAAATACAGCTGCACTGGCCGCTGCAATTGCAACCGAGTATCGGAGCTCTGCAGATATCGTGGTGAATGT CAAGGACATGCTGTCCGACATGATCA	78
6	538–680	TCGTGGTGAATGTCAAGGACATGCTGTCCGACATGATCAGGAGAAGGAACAAGATCCTGAACAGAGACGGCAGTGAA GATGTTCCGAAAAGGGGACCCGTCAGCAAAGAGCACATTGACTGGGCTAGGGACCTTGCTCAAGGTAA	88
7	457–626	GATACCAAGTACGGGATCAACAGAAATACAGTGCCTGACTGGCCGCTGCAATCGCAACCGAGTATAGAGTCCCTGGATC AATTGTGGTGAATGTCAAGGACATGCTGTCCGACATGATCAGGAGAAGGAACAAGATCCTGAACAGAGACGGCAGTG	88

Contig	Location	Sequence (5'→3')	Sequence identity (%)
		AAGATGTTCCGAAAAG	
8	406–578	AGCCCTGAACAACACTGCTAGGTTACCAGAGAGCAGCTCTTAAGTGGAGAAAAGGATACCAAGTACGGGATCAACAGAAA TACAGCTGCACTGGCCGCTGCAATTGCAACCGAGTATAGAGTCCCTGGATCAATTGTGGTGAATGTCAAGGACATGCT GTCGGACATGATCCGGAG	85
9	1365–1500	GGCCGGCTTCGACATTGAGAGCATGGACATAGTTGCCTTGGAGCATCTGCTACACCAGTCACTCGTTGGAAAGAGAT CTCCTTTTCAGAATGCCTACAACATCCGGGGAAATGCCACCAG	89
10	204–349	GCTTGTTCATCGGAGAGTGAGAAGGACTCAGTGATGTATCTGCCCTGGTGGCCGCGACTAAATTCTGTGCTCCTAT ACTGGAGTGTGCCTGGACCAGCTGCACGGGGATGGTCGAGCGTGGTCTGGACTGGTTTGAAAAACAACA	86
11	531–657	ATCAATTGTGGTGAATGTCAAGGACATGCTGTCGGACATGATCAGGAGAAGGAACAAGATCCTGAACAGAGACGGCA GTGAAGATGTTCCGAAAAGGGGACCCGTCAGCAAAGAGCACATTGACTGG	89
12	264–437	CAACTTCTGTGCTCCTATACTGGAGTGTGCCTGGACCAGCTGCACGGGGATGGTCGAGCGTGGTCTGGACTGGTTTG AAAAACAACAAGGAAACAGTGAAGATTTGGGATGCTGAGTACGGGAAGCTAAGAACGGAAGTACCCAGCCCTGAACAA CTGCTGGGTTACCAGAGAGC	81
13	562–681	TCGGACATGATCAGGGGAAGGAACAAGATCCTGAACAGAGACGGCAGTGAAGATGTTCCGAAAAGGGGACCCGTC GCAAAGAGCACATTGACTGGGCTAGGGACCTTGCTCAAGGTAAG	88
14	1345–1459	GCCCTATTCAACATTCAGAAAGCCGGCTTCGACATTGAGAGCATGGACATAGTTGCCTCGGAGCATCTGCTACACCAG TCACTCGTTGGAAAGAGATCTCCTTTTCAGAATGCTT	89
15	1366–1457	GCCGGCTTCGACATTGAGAGCATGGACATAGTTGCCTCGGAGCATCTGCTACACCAGTCACTCGTTGGAAAGAGATC TCCTTTTCAGAATGC	90

Technical Appendix Table 3. Sequences of M gene contigs and their identities to NSDV strain 708, China, 2013

Contig	Location	Sequence (5'→3')	Sequence identity, %
1	3162–3337	CTTCTGGAAGGTCAACAGGAATCATAAACTCCAGGAAAGAACTGGTCTGATGTGGAAGATGTCATCAGAAAAGGCATC AGAAAGCAAAAACCTTCTTGTTCAGTGATGGATTTTCCAGCTCTACAATTCATATTTTCAGTATATAACGGGTGACAG GAGTCTTTCAGAATG	89
2	1743–1901	GAATGCCAGGGAAGCTGCCTCTTGGCATCCACAGGACACAGGAGACATCACAATTGACTGCTCTGGAGGAAGGCAGCA	90

Contig	Location	Sequence (5'→3')	Sequence identity, %
		TTACCTGGAAATTAATATAGTTGACATACACTGTCCAGGGAAAGACAAATGGAAAGGGTTTATGCTCTACATATGCAGG	
3	4512-4678	ATTGAGCACAAAGGGACCATCATAACAGCACTACAACAAAAGTTGTGATGAGGGATATAATTGCTGGGTTGGATCTGTTTC TGGATTCTTTGTTGGAGTGAAAGACTTTTTTAAAAGAACCCTGGAGGTGACTTATTGGTCTTGTAAGCACAATACTTC CTTTAGT	88
4	4396-4542	GGGGTGTAAGAAGTTGTGCCTGGAAGTGGTTGAGAAGGACTACTGTCCTTCCTGCACCGTAGAAGATTTGAAGATATGT GTTGATGTGTCTCTTGAGCCCCCTAAAGACATACTCATTGAGCACAAAGGGACCATCATAACAGCACT	91
5	4071-4211	TGTAAGAAGTTGTGCCTGGAAGTGGTTGAGAAGGACTACTGTCCTTCCTGCACCGTAGAAGATTTGAAGATATGTGTTG ATGTGTCTCTTGAGCCCCCTAAAGACATACTCATTGAGCACAAAGGGACCATCATAACAGCA	79
6	313-468	AGAATTTTATCAAAAATGGGCATGGCCAGAGACTTCGCAGATGAGGAGTTAGACGTCTGGTGTACGAGCGGTTTGACAA CTGTAGCTCTAATGACATTGAAAATAGAATCAGGGATTTCTTCTCATCAGTGATCAGACTGAGTGCTTTGATAAA	76
7	1573-1720	GTGTCAAGCAGGCCCTTAATTGCCTTGACATTTGGATTGTGGCTTGCAGCAGGCTACCTAATAACATGCCTTGTTTCCTT TATCATCTATAATGCTGTTTTGCTTCTCAGCATTGCCATCAAGAAAAGTAAAACAAGGGAGAGAAAA	74
8	1580-1748	TAACAAGGTGATAACAGGAAAGGCATGTAAGTGGGAAACTCCACAGGATCCTGTGAGGTGCAACAGAGCTACAGAAGT GCGAGACTGGTAAATGCATACTGGTGAAACAGAAAAGCAAAGGAGTGGTAAAAGTAAAAGAGGAAAGACTGTAATCAT	88
9	3477-3629	TGCTGTGGGATGGATGGTGAGAGGTACTTTAATAAATACTTCGGTGTGAAATGGGCCTTAGAATATGTTAGAACTGATGT AGTAGTGTGTGTGAAACTCACAATGAGGAAAGACATTGTGACCTTGTGCAGGCCGGAAGCCGCTTTTCCATA	88
10	618-763	AAAATGAAAAAAGCTTAGCCAAAGAATTTTATCAAAAATGGGCATGGCCAGAGACTTCACAGATGAGGAGTTAGACGT CTGGTGTACAGAGCGGTTTGACAACCTGTAGCTCTAATGACATTGAAAATAGAATCAGGGATTTTTT	88
11	4000-4154	TCACCAATCTCATAAACACAGAGACCGATTACACAAGCAAGTTCCACTTCCACTCAAAAAGGATTTAGCTAGAGATGAC ACTTTGCAGATGGACCTGAAAGCAAGACCTAACTCCGGAGGCGGTGAGATGACAGTTTTGGTAGAGGTTAATGGG	86
12	4602-4764	GTTGGAGTGAAAGACTTTTTTAAAAGAACCCTGGAGGTGACTTATTGGTCTTGTAAAGCACAACTTCCTTTAGTGCT GACAGTTCTGTTCTTTATTTATGGCAAGAAGCTGTTCTGCCTGTGCCGACTGTGCCACAAGAAATGTTGTAGAGGCTCAG GAA	85
13	1458-1571	ATGACAACGGCTGTGTCTGTTTCATAGTGACAAAAAGGAGGACCTGAAAAGAAGCTCACCATATGCAATGGTACCACAGT CTCAGATTCAACACTCAATGAAGGGCTCGGATGT	91
14	4830-4937	AGCAGGAACGGCGAACTACTAGGGAAGGGTAAAATGACAAAAGAAGTGTGGCAAGGATGTTTCATGGATGGGCAAAGTAC AAAGAAGGCAATCAAGGAAGTAGCTTAA	91

Contig	Location	Sequence (5'→3')	Sequence identity, %
15	3992–4159	AGACAGCTCCAATCTCATAAACACAGAGACCGATTACACAGGCAAGTTCCACTTTCACTCCAAAAGGATTTAGCTAGAG ATGACACTTTGCAAATGGATCTGAAAGCAAGACCTAACTCCGGAGGCGGTGAAATGACAGTTTTGGTAGAGGTTAATGGG TTGGA	84
16	4017–4178	ACAGAGACCGATTACACAGGCAAGTTCCACTTTCACTCCAAAAGGATTTAGCTAGAGATGACACTTTGCAAATGGATCT GAAAGCAAGACCTAACTCCGGAGGCGGTGAAATGACAGTTTTGGTAGAGGTTAATGGGTTGGAGCTGCATTCAAAAAGGA TA	84
17	4462–4590	ATTTTAAGATATGTGTTGATGTGTCTCTTGAGCCCCCTAAAGACATACTCATTGAGCACAAAGGGACCATCATACAGCAC TACAACAAAAGTTGTGATGAGGGATATAATTGCTGGGTTGGATCTGTTT	90
18	4568–4711	TAATTGCTGGGTTGGATCTGTTTCCGGATTCTTTGTTGGAGTGAAAGACTTTTTTAAAAGAACCTTGGAGGTGACTTA TTGGTCTTGAAGACAATACTTCCCTTAGTGCTGACAGTTCTGTTCTTTATTTATGGCAAGAA	86
19	1037–1182	CATGCACTGCTTAAACATAGAGAATGGGTTAGCTAAACCATCCAAAATGGTTGTTCTTAATGTATTAATGACAACAGTAG AGGTTAGACTGGAATCCTGCCGTGCGTTCATAAATGCACATCAATGCATTTACACAACATGCAG	86
20	3693–3785	ATGACAATTCAAGAAATTGCTGATAATGGCATCTTGGACCTGATGCATGTGAGCAAAGTCATCTCAGCAGAGAATGCCTG CAAACCTCCAGAGC	90
21	4309–4427	GCAGATATCGTGGTGGTTGCTGAGACCAGTGAACAGCCAGGAAGGTAGAGACAGGAGCTAGGAGCGGCTTCAGAGCCTT TGCAGTAAGGGATGTAAAGAAGTTGTGCCTGGAAGTGGTT	88
22	1248–1408	ACGATGAGCTTCAACTTGACTGATGAAAGAAATAGAGCATGCATCATCAGAACAACCTTGTGAGGTCAAAGGAAAAGAGGT AAAGAAGGGACAGAGTCAATTGAGAGGTTTCCCAACAAGTATCAGACTCTTCAAGTCTGTGACAGGAAAAGAAGGCTCA	91
23	1293–1435	ATCAGAACAACCTTGTGTGGTCAAAGGAAAAGAGGTAAGAAGGGACAGAGTCAATTGAGAGGTTTCCCAACAAGTATCAG ACTCTTCAAGTCTGTGACAGGCAAAAAGAAGGCTCATGTGACAGGAGGGAGATTGGACTGACTG	92
24	3620–3764	TTTTTCCATAGGTCCTGTGTCTGTAACCTGTCTGACCCCCAGAATGTGGTGAACCGTCTCTCGAGCAACATCATGACAA TTCAAGAAATTGCTGATAACGGCATCTTGGACTTGATGCATGTGAGCAAAGTGATATCTGCAGAG	91
25	1897–2049	GCAGGGAGGCAAGCAGGCCTTAATTGCCTTGACATTTGGATTGTGGCTTGACAGGCTACCTAATAACATGCCTTGTT TCCTTCATCATCTATAATGCTGTTTTGCTTCTCAGCATTGCCATCAAGAAAGTAAACAAGGGAGAGAATGGA	88
26	994–1151	TTATGAACGAACCTCAGGCCCACTATCCAAAACCTATGATAGCATGCACTGCTTAAACATAGAGAATGGGTTAGCTAAAC CATCCAAAATGGTTGTTCTTAATGTATTAATGACAACAGTAGAGGTTAGACTGGAATCCTGCCGTGCGTTCATAAAT	86

Contig	Location	Sequence (5'→3')	Sequence identity, %
27	2243–2494	TTTATTCTGACAATTTACCCGGTGCAGGGATTCCCATTGGAGAGTCCACCTATTGGCGCAGCAACTGACAGTAGAATCAT GTATGGGGTAGTGGGGTTCATGTTTGCCTAATAATGATGCTTAATCTTAAAAATCACACTACGGTTCCAT	86

Technical Appendix Table 4. Sequences of L gene contigs and their identities to NSDV strain 708, China, 2013

Contig	Location	Sequence (5'→3')	Sequence identity (%)
1	971–1121	TAGATGCTGGCGCCTCCTACGTTCCAGGATGGGATTAGAGAGGAACCTGCAATTCTTGCACTCAGAGTTCTG CTTGATGTGTGCACAGTGGTTGTTGCTGTCCTACTGTCTTCCTTCTGTACGGGTCAAACAACAGAAA	89
2	1122–1291	TAAGAAGACATTTATAACAACTGCTTGCTGAACACAAGCCTTTCAGGGAAGAGAGTCTTCAAGGCACTAGGAAAGCTAACAGG CACCACCTTTACAAAAGTCTAGGAATGCTTTGTCCCATGTGTGTCAAACACTCTATGGCAAGATGACAGGAGATATCTGCAG AGCT	86
3	2632–2812	AGACAGATACTACTAGTTGAGGTTGGTTATCAGACCGATGTGGACGGCAAATAACAACAGACTTTAAGAAGTGGAAGACATC TTGAGGCTCCTAGAGATGTTGAAATCAAGTGTTCATTCATAGCCTGTGCAGACTGCACATCAACACCTGCGGACAACCTGGTGG ATATCTGTGCGACA	92
4	9940–10105	GACAAGACATCTCTTCTCATAAGCTTCTCAACTGGAAAGTTTCACCTGACTCAATGGCACAAGATTGTCGGCTTTACAAACAGG AACATGCTGTCATCAGTGAATTTGCAGGACAGGTTGTTGTGAACACACTTGCCAGTGAGCTTAGCTCTGTTAGGAGAGATG	92
5	6811–6964	TGCAAGCGTCTGACTGGCAGAACAACCTGGTAAAAGGCTTCCAAGGAGTGTGAGAAGCAAAGTAATTTATGAAATGGTAAAGCTT GTTGGGGAAACTGGAATGGCAACTACAACAACCTGGCCTTTGCCAAGCTCTTAATTACGATCATAGAT	94
6	5253–5423	AAGCTTAAGTGGGAGACGTTCTTACTGTTCTAGTAGAGGGAGGATTTCAGAGCATCTTTGGCAGGGACAATTCAAACAAAAAACC TTTTGAGTTTAGACCAGGCTTAGAAGTTAGAGCAGATCCAATGAATGATTTGAGCAAGCAGTCACACACACAACGATATCTGC AGA	90
7	6853–7015	AGGAGCGTGAGAAGCAAAGTAATTTATGAAATGGTAAAGCTTGTGGGGAAACTGGAATGGCAACTACAACAACCTGGCCTTT GCCAAGCTCTTAATTACGATCATAGATTTTATGCAGTGTGGCACCAAAGGCCAAGCTTGGAGGGACTAGATATCTGT	90
8	1061–1227	TGTGCACAGTGGTTGTTGCTGTCCTACTGTCTTCCTTCTGTACGGGTCAAACAACAGAAACAAGAAGACATTTATAACAACTG CTTGCTGAACACAAGCCTTTCAGGGAAGAGAGTCTTCAAGGCACTAGGAAAGCTAACAGGCACCACCTTTACAAAAGTCTC	89
9	6262–6418	AGAATGAAGAAGGATAATCCAGCGTAAGCTTCACCAAGGAAGAAGTCTTGTGAAAAGACTTGAGAAGTCATTCCTGAAGAAA	91

Contig	Location	Sequence (5'→3')	Sequence identity (%)
		TTTAAACAAGGAAGCAATGAAGTTTGTCAATCTAGTGTTCTTTTGTTCACCTCTCTGCCCTTGGTGTGTTTCATT	
10	6289–6443	AGCTTCACCAAGGAAGAAGTTCTTGTGAAAAGACTTGAGAAGTCATTCCTGAAGAAATTTAAACAAGGAAGCAATGAAGTTTGTCA ATCTAGTGTTCTTTTGTTCACCTCTCTGCCCTTGGTGTGTTTCATTACAAGTCTTTAGAGTCTTACTTGGT	92
11	8921–9083	CTGCAGATATCTCCAGACTAAAACAGACCTTGACAGCCAGAAACGTCCTGCATGGCCTTGCTGGAGGTATCAAGGAGCTGTCA CTTCCTATCTATAACCATCTTCCTAAAAGTCGTA CTTTTTAAAGACAATGTCTTCTTAGACCTTGAGGACAGATGGTCAAC	90
12	6900–7051	GGAAACTGGAATGGCAATACTACAACA ACTGGCCTTTGCCAAGCTCTTAATTACGATCATAGATTTTATGCAGTGTTGGCACCA AAAGCCCAACTTGGAGGGGGTAGGGATCTGCTAGTCCAAGAGACCGGCACAAAAGTTATCCATGCCA	91
13	4969–5112	AACAGTGACCGACAGTTAATCTTTGACATTTACAATGTACACATCTACAACAAGGAGATGGACAATTTTATGATGAAGGATGCATAA GTGTGTTAGAAGAGACTGCTGAAAGACACATGCTGTGGGAAATGGATCTTCTTAGGTCA	91
14	8692–8859	GATGGGACAGATAAGCAAGTTAAGGCATCTCTCAATAGGGACGACAATAGAGTTATCGAAGACCCCATGATTCAACTAGTACCT GAAAAGCTGAGAAGAGAGCTAGAGAGGCTGGGAGTCTCTAGGATGGAGATCGATGAACTAATGCCTTCTATAAAGCATGACGA A	86
15	4909–5055	TTGAAAGACTTGTGCCCTGAAGTCACAATTCATGCTTCTCCGTCTATGGTGTATTTGTGAACAGTGACCGACAGTTAATCTTTG ACACTTACAATGTACACATCTACAACAAGGAGATGGACAATTTTATGATGAAGGATGCATAAGT	86
16	6786–6906	GATATCAAACAGTAACTTCAATGTCTGCAAGCGTCTGACTGGCAGAACA ACTGGTGAAGGCTTCCAAGGAGTGTGAGAAGCA AAGTAATTTATGAAATGGTAAAGCTTGTGGGGAAACT	91
17	3477–3582	TGCAGCTCTGCAGATATCGATAGGGTGGTTCGGTTGACTTTACCTGGCAAACTGAAAAGGAGAGAAGGATTAAGAGAAATGTT GAAACACTAATCCTGTTGATG	89
18	5638–5741	CCTAGAACACACATAATGCTCAAAGACTGTTTCAAGATCTTCTCGGCACTGAGAACAAGAAAATCGTCAAGATGCTAAGAGGC AAGCTAAAGAAGCTGGGTGC	88
19	5026–5099	ATGGACAATTTTATGATGAAGGATGCATAAGTGTGTTAGAAGAGACTGCTGAAAGACACATGCTATGGGAAATGGA	95
20	8680–8766	TCATTTGGTAGCGACGGGACAGATAAGCAAGTTAAGGCATCTCTCAATAGGGACGACAATAGAGTTATCGAAGACCCCAT GATTCAA	90
21	8968–9092	CTTCATGGCTTGGCTGGAGGTATAAAAAGAACTGTCCCTGCCTCTTTACACCATCTTCCTCAAATCATACTTCTTCATAGA TAAGGTATTTTATCCCATGCAGATAGGTGGAACACCAAGCACAG	74
22	7836–7969	AAGAGGCTGTCAGATGAAGGACTCTGCAAAAACGCTCATAGGAGATGTCATGTGTGAGTTTTACAGTGAGTTTATGCTAT	73

Contig	Location	Sequence (5'→3')	Sequence identity (%)
		ACCATAGGGTGACACCTGCAGTCATTAAGTTCATTATCGC	
23	7537–7639	TATATATCCAAAGGAAAGCTGGCCCTAGACTGCTACAACCACATGGGACAGGGCATACACCATGCCACCTCATCAGTAAT GACCTCTTGCATGGCTGAACTGT	77

Technical Appendix Table 5. Primer sequences used to amplify the complete genome of NSDV, China, 2013

Genome segment, interest fragment	Forward (location) (5'→3')	Reverse (location) (5'→3')
Small		
1–1012	S-F1: TCTCAAAGACAAACGTGCCGCTTTGCGC (1–28)	S-R1: CTGTACACCTGCCTTCCAA (1012–993)
637–1012	S-F2: AGCAAAGAGCACATTGACTGGGC (637–659)	S-R2: CTGTACACCTGCCTTCCAA (1012–993)
826–1335	S-F3: CTTGTGGCATGGATCAATGC (826–845)	S-R3: CTGTGCGCAGGGGTTGCCAG (1335–1316)
1165–1590	S-F4: CGGATTTACATGCACCCTGGTGTC (1165–1188)	S-R4: TCTCAAAGAGATCGTTGCCGCACAGCC (1590–1564)
Medium		
1–1451	M-F1: TCTCAAAGAGATAGTTGCGGCACTAGCAGG (1–30)	M-R1: CTGTGTGCCAGATCCGCAGTCAGT (1451–1428)
1428–1979	M-F2: ACTGACTGCGGATCTGGCACACAG (1428–1451)	M-R2: GGAAACAAGGCATGTTATTAGATAGCC (1979–1953)
1953–3187	M-F3: GGCTATCTAATAACATGCCTTGTTC (1953–1979)	M-R3: ATGATTCCTGTTGACCTTCCAGAAAG (3187–3162)
3162–4431	M-F4: CTTTCTGGAAGGTCAACAGGAATCAT (3162–3187)	M-R4: TCTCAACCACTTCCAGGCACAATT (4431–4407)
4056–5075	M-F5: TACACAGGCAAGTTCACCTTTTAC (4056–4079)	M-R5: TCTCAAAGAGATAGTGGCGGCACAGCA (5077–5051)
Large		
1–1098	L-F1: TCTCAAAGATATCAATCCCCCGTTACCCAGAGTTGC (1–38)	L-R1: ACAGGAAGGAAGACAGTAGGACAGC (1098–1078)
995–2246	L-F2: CAGCATTTCCAGGAATGGATTAG (995–1018)	L-R2: TCAGGCAGAGGAAACATCTTCTTCT (2246–2222)
1775–2875	L-F3: GCATTGAGCCAATTGCGATAATATG (1775–1799)	L-R3: CAGGAGAGTTCTTTGTGAGGCTGCT (2875–2851)
2850–3833	L-F4: TAGCAGCCTCACAAAGAACTCTCCT (2850–2874)	L-R4: GCTCTAGGCACTGAACACTTGGGA (3833–3811)
3811–5046	L-F5: TCCAAGTGTTCAAGTGCCTAGAGCATG (3811–3836)	L-R5: TCCTTCGTCAAATTGTCCATCTC (5046–5023)

Genome segment, interest fragment	Forward (location) (5'→3')	Reverse (location) (5'→3')
5023–6847	L-F6: GAGATGGACAATTTTGACGAAGGA (5023–5046)	L-R6: GCCTTTCACCAGTTGTTCTGCCAGTC (6847–6822)
6696–7767	L-F7: AGACGAGAAGCTGTTGCATCAGAC (6696–6719)	L-R7: GAGTTCCTTAGGCAGTAGCCAGT (7767–7744)
7744–8737	L-F8: ACTGGGCTACTGCCTAAGAAGCTC (7744–7767)	L-R8: TGTCGTCCTATTGAGAGATGCCTT (8737–8713)
8713–9807	L-F9: AAGGCATCTCTCAATAGGGACGACA (8713–8737)	L-R9: GACTAGCTCAGACACCGTGGGC (9807–9786)
9786–11367	L-F10: GCCCACGGTGTCTGAGCTAGTC (9786–9807)	L-R10: GGTGGCAACTGCTTCAATTTCTT (11367–11345)
11345–12081	L-F11: AAGAAATTGAAGCAGTTGCCACC (11345–11367)	L-R11: TCTCAAAGAAATCGTTCCCCCCCCACCCC (12081–12053)

Technical Appendix Table 6. Gene lengths and identity between NSDV from specimens collected in China and other nairoviruses*

Virus	Small segment					Medium segment					Large segment				
	GenBank accession No.	nt	% ID	aa	% ID	GenBank accession no.	nt	% ID	aa	% ID	GenBank accession no.	nt	% ID	aa	% ID
NSDV (China)	KM464724	1590	NA	482	NA	KM464725	5077	NA	1628	NA	KM464726	12081	NA	3992	NA
NSDV (India)	AF504294	1590	87.9	482	96.7	EU697950	5094	75.1	1624	81.3	EU697949	12081	88.7	3991	96.6
NSDV(Kenya)	AF504293	1590	88.4	482	95.6	EU697952	5077	87.6	1627	93.2	EU697951	12081	89.6	3991	96.1
Dugbe virus	NC_004157	1716	59.8	483	59.6	NC_004158	4888	51.1	1551	44.8	NC_004159	12255	66.9	4036	67.6
Kupe virus	EU257626	1694	61.8	483	61.9	EU257627	4818	52.5	1549	48.1	EU257628	12330	64.8	4050	67.5
Hazara viru	KC344857	1677	63.3	485	62.1	DQ813514	4576	54.2	1421	48.8	DQ076419	11980	64.0	3923	68.6
CCHFV	GU477494	1673	59.9	482	62.0	GU477493	5377	39.5	1697	37.5	GU477492	12158	64.9	3945	64.7

*nt, nucleotide length; % ID, percentage identities of nt and aa (amino acid) between NSDV (China) and other nairoviruses; NA, not applicable; CCHFV, Crimean Congo hemorrhagic fever virus.

