

---

# KPC and NDM-1 Genes in Related *Enterobacteriaceae* Strains and Plasmids from Pakistan and the United States

Mitchell W. Pesesky,<sup>1</sup> Tahir Hussain,<sup>1</sup>  
Meghan Wallace, Bin Wang, Saadia Andleeb,  
Carey-Ann D. Burnham, Gautam Dantas

To characterize the genomic context of New Delhi metallo- $\beta$ -lactamase-1 (NDM-1) and *Klebsiella pneumoniae* carbapenemase (KPC), we sequenced 78 *Enterobacteriaceae* isolates from Pakistan and the United States encoding KPC, NDM-1, or no carbapenemase. High similarities of the results indicate rapid spread of carbapenem resistance between strains, including globally disseminated pathogens.

Pathogenic *Enterobacteriaceae*, including *Escherichia coli* and *Klebsiella pneumoniae*, are major causes of multidrug-resistant (MDR) infections in hospitals worldwide. These pathogens have recently been shown to have acquired resistance to carbapenems, and the US Centers for Disease Control and Prevention identified carbapenem-resistant *Enterobacteriaceae* as 1 of the 3 most urgent MDR threats (1). Among the *Enterobacteriaceae*,  $\beta$ -lactam resistance, including carbapenem resistance, is primarily caused by enzymatic degradation by  $\beta$ -lactamases. Two carbapenemase subclasses are especially problematic: *Klebsiella pneumoniae* carbapenemase (KPC) and New Delhi metallo- $\beta$ -lactamase-1 (NDM-1). KPC, identified in 2001 (2), has become endemic to several noncontiguous areas of the world, including the United States, Israel, Greece, South America, and China (3). NDM-1 was first described in 2008, although retrospective studies identified NDM-1 from 2006 (4) and is abundant in New Delhi water samples (5). Most patients from whom NDM-1 is isolated have an epidemiologic link to the Indian subcontinent, but NDM-1 has also recently become endemic to the Balkans and Middle East (6).

The spread of antibiotic resistance genes such as NDM-1 and KPC is facilitated by horizontal gene transfer (HGT) between bacteria (7). Among globally disseminated pathogens, HGT facilitates combination of the most effective antibiotic resistance genes from diverse geographies into

multidrug resistance plasmids that spread between strains. Recombination and transposition have created populations of these plasmids that have related architectures but vary in their composition of antibiotic drug resistance cassettes (8). This effect has enabled both KPC and NDM-1 to rapidly expand within the *Enterobacteriaceae* and other proteobacterial pathogens, such as *Acinetobacter baumannii* (9,10). Antibiotic resistance genes can also spread through clonal expansion in successful pathogenic strains, for example, KPC in *K. pneumoniae* sequence type (ST) 258 (11), and the extended-spectrum  $\beta$ -lactamase CTX-M-15 in *E. coli* ST131 (12). Both HGT and clonal expansion have enabled KPC and NDM-1 to rapidly spread to distant locations after their emergence (6,8).

The similarities in the spread and resistance spectra of KPC and NDM-1 (both provide resistance to nearly all  $\beta$ -lactam antimicrobial drugs) leads to the hypothesis that similar mobile elements will make both genes available to similar pathogen populations. We tested this hypothesis by examining clinical *Enterobacteriaceae* isolates from Pakistan and the United States encoding NDM-1, KPC, or no carbapenemase.

## The Study

We collected 450 bacterial isolates (including 195 *Enterobacteriaceae*) in Pakistan during February 2012–March 2013 from Pakistan Railway General Hospital in Rawalpindi and the Pakistan Institute of Medical Sciences in Islamabad. From this collection, we randomly selected 55 *Enterobacteriaceae* isolates for whole-genome sequencing. We then selected 23 isolates from samples collected in the United States during January 2010–June 2013 from patients in Barnes Jewish Hospital in St. Louis, Missouri, that had similar proportions of  $\beta$ -lactam susceptibility and resistance to the isolates collected in Pakistan for sequencing. All isolates were de-identified and retrieved from existing strain banks. The combined set included 33 *E. coli*, 30 *K. pneumoniae*, 9 *Enterobacter cloacae*, and 6 *Enterobacter aerogenes* (online Technical Appendix Table 1, <http://wwwnc.cdc.gov/EID/article/21/6/14-1504-Techapp1.pdf>). We extracted plasmid DNA from 9 isolates encoding NDM-1, 11 isolates encoding KPC, and 3 isolates encoding CTX-M-15 and performed shotgun sequencing on those plasmid preparations. Detailed methods are described in the online Technical Appendix.

<sup>1</sup>These first authors contributed equally to this article.


---

Author affiliations: Washington University School of Medicine in St. Louis, St. Louis, Missouri, USA (M.W. Pesesky, T. Hussain, M. Wallace, B. Wang, C.D. Burnham, G. Dantas); National University of Sciences and Technology, Islamabad, Pakistan (T. Hussain, S. Andleeb)

DOI: <http://dx.doi.org/10.3201/eid2106.141504>

Using antibiotic resistance gene predictions from the Resfams database (13) and core genome alignment, we constructed a phylogenetic tree for each species in our set, overlaid by the  $\beta$ -lactamases encoded by each isolate (Figure 1). Isolates from both locations were found to be members of the same subspecies clades (online Technical Appendix Figure 1) and to contain similar repertoires of  $\beta$ -lactamases (Figure 1), indicating that geography is not a discriminating variable for these isolates. Many of these isolates were also MDR: resistance to ciprofloxacin, trimethoprim/sulfamethoxazole, gentamicin, doxycycline, and chloramphenicol occurred in 63%, 65%, 45%, 54%, and 56% of isolates, respectively. As expected from results of previous work (8), *E. coli* ST131 isolates had high rates of CTX-M carriage (82%; Figure 1, panel A) and ciprofloxacin resistance (100%).

The variety of strains that we discovered encoding KPC and NDM-1 is consistent with existing evidence that HGT is a major factor in their spread. All KPC genes were proximal to Tn4401 and all NDM-1 genes were carried on ISAb125, mobile elements with which each gene has respectively been previously associated (14). We observed multiple examples of NDM-1 within the *K. pneumoniae* ST11 clade (15) (Figure 1, panel B; online Technical Appendix Figure 1, panel B), a close relative of ST258. This association could be caused by clonal expansion or multiple HGT events and emphasizes that lineages known to encode KPC are now also acquiring NDM-1. We also observed high rates of NDM-1 carriage in *Enterobacter* isolates (Figure 1, panels C and D), which in general showed a high number (maximum 8) and wide variety of  $\beta$ -lactamases. These isolates were also MDR: 57% of the *Enterobacter* isolates were resistant to all or all




**Figure 1.** Distribution of antimicrobial drug resistance genotypes of *Klebsiella pneumoniae* carbapenemase (KPC) and New Delhi metallo- $\beta$ -lactamase-1 (NDM-1) genes in related *Enterobacteriaceae* strains and plasmids in Pakistan and the United States. A) *Escherichia coli*; B) *K. pneumoniae*; C) *Enterobacter cloacae*; D) *E. aerogenes*. Phylogenetic trees have been annotated with the specific  $\beta$ -lactamases encoded by those isolates. \*Denotes an unnamed single nucleotide variant of the named  $\beta$ -lactamase. Scale bars indicate nucleotide substitutions per site.

but 1 of the antimicrobial drugs tested. At best, these *Enterobacter* strains are a reservoir for resistance in Pakistan; at worst, they are the vanguard of an expansion of carbapenem-resistant *Enterobacter* infections.

Previous observations have predominantly found KPC and NDM-1 to be expressed from plasmids (6,11). To characterize the sequence similarity of plasmids within the NDM- and KPC-carrying plasmid populations, we purified and sequenced plasmid DNA from 9 isolates encoding NDM-1, 11 encoding KPC, and 3 encoding CTX-M-15. Sequencing showed that these plasmids include representatives from IncHI2, IncY, IncN, IncFIA, IncFIB, IncFIC, and IncI1 incompatibility groups. Using reciprocal BLAST (<http://blast.ncbi.nlm.nih.gov/Blast.cgi>) alignment between each pair of plasmid preparations, we calculated the percentage of each plasmid shared using a 99% identity threshold. We performed this same analysis for all sequenced plasmids containing NDM-1, KPC, or CTX-M available in the National Center for Biotechnology Information database (<http://www.ncbi.nlm.nih.gov>) together with our set (Figure 2) and separately (online Technical Appendix

Figure 2). Certain components, primarily mobile elements, were abundant within these plasmids: the average plasmid shared 500 contiguous bases with 58 of the other plasmids; however, median BLAST identity for this pairwise comparison was <12%, even when considering plasmids with the same  $\beta$ -lactamase, suggesting that both carbapenemases exist within a variety of plasmid configurations.

To visualize this comparison of carbapenemase plasmids, we generated a network diagram in which each node represented a plasmid and each line represented shared sequence between 2 plasmids (Figure 2, panel B). Node size and line width correlate to the number of nucleotides contained in the plasmid or sharing interaction. This visualization shows the abundant small, shared regions that exist between most plasmid pairs, represented as thin background lines. This visualization also highlights the larger shared regions that indicate highly similar plasmids, represented by the few wide lines. These outliers were often between pairs of plasmids encoding the same  $\beta$ -lactamase but were also observed between NDM-1 and KPC containing plasmids (maximum 79% of smaller plasmid length).



**Figure 2.** Pairwise BLAST identity (<http://blast.ncbi.nlm.nih.gov/Blast.cgi>) of all CTX-M genes, *Klebsiella pneumoniae* carbapenemase (KPC), and New Delhi metallo- $\beta$ -Lactamase-1 (NDM-1) plasmids from isolates collected in Pakistan and the United States plasmid preparations, and the National Center for Biotechnology Information database complete plasmids. An all-against-all plasmid BLAST was performed and plasmid interactions were defined by the percentage of the query plasmid conserved (at  $\geq 99\%$  identity) in the subject plasmid. A) Plasmid interactions collected based on the defining  $\beta$ -lactamase of their query and subject plasmids. Box and whisker plots represent the range of pairwise sharing values within this population of plasmids. Upper and lower boundaries of the box correspond to the first and thirds quartiles; whiskers (error bars) represent 1.5 times the interquartile range; points beyond the whiskers represent outliers. B) Network map in which nodes represent individual plasmids and lines represent regions shared between plasmids. Line width is proportional to the number of nucleotides contained in fragments >500 bp in length at >99% sequence identity. Genetic elements repeated within the same plasmid DNA are represented by lines that leave and return to the same node. Plasmid sequence origin is indicated in arcs around the network.

## Conclusions

Together, this evidence supports our hypothesis that strains and plasmids known to carry either carbapenemase also have access to the other. Given the similarity of carbapenemase-negative strains to those carrying KPC or NDM-1 and the high diversity of plasmids in which they can be found, we anticipate that global carbapenem usage will encourage HGT of both of these carbapenemases into additional strain and plasmid backgrounds. Because KPC and NDM-1 are poised to cross genetic and geographic boundaries, we recommend that hospitals routinely screen *Enterobacteriaceae* strains for both genes, even in regions where they are not yet endemic. We further advocate reduced carbapenem use to limit the selection for resistance against this vital antibiotic class.

## Acknowledgments

We thank members of the Dantas lab for thoughtful discussions of the research described herein.

Research reported in this publication was supported in part by the National Institutes of Health Director's New Innovator Award (<http://commonfund.nih.gov/newinnovator/>), the National Institute of Diabetes and Digestive and Kidney Diseases (<http://www.niddk.nih.gov/>), and the National Institute of General Medical Sciences (NIGMS: <http://www.nigms.nih.gov/>), of the National Institutes of Health under award numbers DP2DK098089 and R01GM099538 to G.D. M.W.P. is supported by the NIGMS Cell and Molecular Biology Training Grant (GM:007067). T.H. is supported by the Higher Education Commission of Pakistan's International Research Support Initiative Program. The content is solely the responsibility of the authors and does not necessarily represent the official views of the funding agencies. All genome and plasmid sequences were deposited into the National Center for Biotechnology Information database (<http://www.ncbi.nlm.nih.gov>) (BioProject ID PRJNA261540).

Dr. Pesesky received his PhD from Washington University in St. Louis, Missouri, USA, in 2015 and is currently a postdoctoral fellow at the University of Washington, Seattle, Washington, USA. His research focuses on molecular and genomic investigations of high interest functions in bacteria.

## References

- Centers for Disease Control and Prevention. Antibiotic resistance threats in the United States, 2013 [cited 2014 Sept 15]. <http://www.cdc.gov/drugresistance/threat-report-2013/index.html>
- Yigit H, Queenan AM, Anderson GJ, Domenech-Sanchez A, Biddle JW, Steward CD, et al. Novel carbapenem-hydrolyzing beta-lactamase, KPC-1, from a carbapenem-resistant strain of *Klebsiella pneumoniae*. *Antimicrob Agents Chemother*. 2001;45:1151–61. <http://dx.doi.org/10.1128/AAC.45.4.1151-1161.2001>
- Munoz-Price LS, Poirel L, Bonomo RA, Schwaber MJ, Daikos GL, Cormican M, et al. Clinical epidemiology of the global expansion of *Klebsiella pneumoniae* carbapenemases. *Lancet Infect Dis*. 2013;13:785–96. [http://dx.doi.org/10.1016/S1473-3099\(13\)70190-7](http://dx.doi.org/10.1016/S1473-3099(13)70190-7)
- Castanheira M, Deshpande LM, Mathai D, Bell JM, Jones RN, Mendes RE. Early dissemination of NDM-1- and OXA-181-producing *Enterobacteriaceae* in Indian hospitals: report from the SENTRY Antimicrobial Surveillance Program, 2006–2007. *Antimicrob Agents Chemother*. 2011;55:1274–8. <http://dx.doi.org/10.1128/AAC.01497-10>
- Walsh TR, Weeks J, Livermore DM, Toleman MA. Dissemination of NDM-1 positive bacteria in the New Delhi environment and its implications for human health: an environmental point prevalence study. *Lancet Infect Dis*. 2011;11:355–62. [http://dx.doi.org/10.1016/S1473-3099\(11\)70059-7](http://dx.doi.org/10.1016/S1473-3099(11)70059-7)
- Dortet L, Poirel L, Nordmann P. Worldwide dissemination of the NDM-type carbapenemases in gram-negative bacteria. *Biomed Res Int*. 2014;2014:249856. <http://dx.doi.org/10.1155/2014/249856>
- Thomas CM, Nielsen KM. Mechanisms of, and barriers to, horizontal gene transfer between bacteria. *Nat Rev Microbiol*. 2005;3:711–21. <http://dx.doi.org/10.1038/nrmicro1234>
- Hawkey PM, Jones AM. The changing epidemiology of resistance. *J Antimicrob Chemother*. 2009;64(Suppl 1):i3–10. <http://dx.doi.org/10.1093/jac/dkp256>
- Robledo IE, Aquino EE, Vazquez GJ. Detection of the KPC gene in *Escherichia coli*, *Klebsiella pneumoniae*, *Pseudomonas aeruginosa*, and *Acinetobacter baumannii* during a PCR-based nosocomial surveillance study in Puerto Rico. *Antimicrob Agents Chemother*. 2011;55:2968–70. <http://dx.doi.org/10.1128/AAC.01633-10>
- Sartor AL, Raza MW, Abbasi SA, Day KM, Perry JD, Paterson DL, et al. Molecular epidemiology of NDM-1 producing *Enterobacteriaceae* and *Acinetobacter baumannii* isolates from Pakistan. *Antimicrob Agents Chemother*. 2014;58:5589–93. <http://dx.doi.org/10.1128/AAC.02425-14>
- Cuzon G, Naas T, Truong H, Villegas MV, Wisell KT, Carmeli Y, et al. Worldwide diversity of *Klebsiella pneumoniae* that produce beta-lactamase blaKPC-2 gene. *Emerg Infect Dis*. 2010;16:1349–56. <http://dx.doi.org/10.3201/eid1609.091389>
- Petty NK, Ben Zakour NL, Stanton-Cook M, Skippington E, Totsika M, Forde BM, et al. Global dissemination of a multidrug resistant *Escherichia coli* clone. *Proc Natl Acad Sci U S A*. 2014;111:5694–9. <http://dx.doi.org/10.1073/pnas.1322678111>
- Gibson MKFK, Dantas G. Improved annotations of antibiotic resistance functions reveals microbial resistomes cluster by ecology. *ISME*. 2014;106. <http://dx.doi.org/10.1038/ismej.2014.106>
- Patel G, Bonomo RA. Status report on carbapenemases: challenges and prospects. *Expert Rev Anti Infect Ther*. 2011;9:555–70. <http://dx.doi.org/10.1586/eri.11.28>
- Pena I, Picazo JJ, Rodriguez-Avial C, Rodriguez-Avial I. Carbapenemase-producing *Enterobacteriaceae* in a tertiary hospital in Madrid, Spain: high percentage of colistin resistance among VIM-1-producing *Klebsiella pneumoniae* ST11 isolates. *Int J Antimicrob Agents*. 2014;43:460–4. <http://dx.doi.org/10.1016/j.ijantimicag.2014.01.021>

Address for correspondence: Gautam Dantas, Washington University in St. Louis, 4444 Forest Park Blvd, Campus Box 8510, St. Louis, MO 63108, USA; email: [dantas@wustl.edu](mailto:dantas@wustl.edu)



# KPC and NDM-1 Genes in Related *Enterobacteriaceae* Strains and Plasmids from Pakistan and the United States

## Technical Appendix

### Methods and Materials

#### Sample Selection, Processing, and Phenotyping

We collected 450 Pakistani bacterial isolates (PH) initially recovered from de-identified clinical samples from urinary, blood stream, genitourinary, and wound infections collected between February 2012 and March 2013 at Pakistan Railway General Hospital, Rawalpindi, Pakistan, and the Pakistan Institute of Medical Sciences in Islamabad, Pakistan. These included all ESKAPE pathogen isolates available in the Pakistani hospital strain banks during the indicated collection period. From these 450 isolates, we chose a random subset of 55 isolates from the *Enterobacteriaceae* family (from a total of 195 *Enterobacteriaceae* in this collection) for phenotypic and genotypic analysis. We also selected 48 US *Enterobacteriaceae* isolates (WU) from banked, de-identified frozen stocks of *Escherichia coli*, *Klebsiella pneumoniae*, and *Enterobacter cloacae* at Barnes Jewish Hospital/Washington University School of Medicine in Saint Louis, Missouri, United States to have beta-lactam resistance and susceptibility phenotypes in similar proportions to the Pakistani *Enterobacteriaceae* isolates, with a particular focus on the meropenem resistance phenotype (protocols for growth and phenotyping are described below). The WU strains were originally isolated from urine, respiratory, bone, and bile specimens between January 2010 and June 2013. Of the 48 WU isolates, 23 isolates were chosen for genome sequencing to generally match the species distribution of the 55 PH isolates as well as their beta-lactam resistance profiles, with the exception of *Enterobacter aerogenes*, for which none were available in the WU collection. In total, 33 *Escherichia coli* (24 PH, 9 WU), 30 *Klebsiella pneumoniae* (19 PH, 11 WU), 9 *Enterobacter cloacae* (6 PH, 3 WU), and 6 *Enterobacter aerogenes* (all 6 PH) isolates were included for the whole genome sequencing

analysis (Technical Appendix Tables 1 and 2). We cultivated all isolates on MacConkey and sheep's blood agar (Hardy Diagnostics). We then grew single colonies in LB broth liquid culture for DNA extraction. We assessed each isolate for susceptibility to ampicillin, cefazolin, cefotetan, ceftazidime, ceftriaxone, cefepime, meropenem, ciprofloxacin, trimethoprim-sulfamethoxazole, gentamicin, doxycycline, and chloramphenicol by Kirby-Bauer disk diffusion according to Clinical and Laboratory Standards Institute guidelines and interpretive criteria (1). Prior to whole genome sequencing, the species identity of PH and WU isolates was determined with VITEK MS MALDI-TOF MS v2.0 knowledgebase (bioMerieux) as previously described (2,3). We then extracted total DNA using the Invitrogen Charge Switch gDNA Mini Bacteria kit per the manufacturer's protocol. We also extracted plasmid DNA from 11 KPC and 9 NDM-1 encoding isolates (as well as 3 CTX-M-15 encoding isolates), as determined by PCR and the genome sequencing, using the Qiagen Large Construct kit per the manufacturer's protocol. We included one non- *Enterobacteriaceae* plasmid preparation from an *Acinetobacter baumannii* isolate (PH), which had been identified to contain NDM-1 by PCR.

### **Illumina Library Preparation**

We sheared 500ng of total DNA from each isolate to ~300 bp fragments in nine rounds of shearing of ten minutes each on the BioRupter XL. In each round the power setting was 'H' and samples were treated for 30s and allowed to rest for 30s. Each sample was concentrated using the Qiagen MinElute PCR purification kit per the manufacturer's protocol. End Repair of the sheared DNA fragments was initiated with the addition of 2.5 µl of T4 DNA ligase buffer with 10mM ATP (NEB, B0202S), 1 µl of 1 mM dNTPs (NEB), 0.5 µl T4 Polymerase (NEB, M0203S), 0.5 µl T4 PNK (NEB M0201S), and 0.5 µl Taq Polymerase (NEB, M0267S). This mixture was incubated at 25°C for 30 min, then at 75°C for 20 min. Barcoded adapters were then added to the solution along with 0.8µl of T4 DNA ligase (NEB, M0202M), for the purpose of ligating the adapters to the DNA fragments. This solution was then incubated at 16°C for 40min, then 65°C for 10min. At this point the adapter-ligated DNA was purified using the Qiagen MinElute PCR purification kit per the manufacturer's protocol.

The DNA fragments were then size selected on a 2% agarose gel in 1X TBE buffer stained with Biotium GelGreen dye (Biotium). DNA fragments were combined with 2.5uL 6X Orange loading dye before loading on to the gel. Adaptor-ligated DNA was extracted from gel

slices corresponding to DNA of 250-300bp using a QIAGEN MinElute Gel Extraction kit per the manufacturer's protocol. The purified DNA was enriched by PCR using 12.5µL 2X Phusion HF Master Mix and 1µL of 10µM Illumina PCR Primer Mix in a 25µL reaction using 1µL of purified DNA as template. DNA was amplified at 98°C for 30 seconds followed by 18 cycles of 98°C for 10 seconds, 65°C for 30 seconds, 72°C for 30 seconds with a final extension of 5min. at 72°C. Afterwards, the DNA concentration was measured using the Qubit fluorometer and 10nmol of each sample (up to 106 per lane of sequencing) were pooled. Subsequently, samples were submitted for Illumina HiSeq-2500 Pair-End (PE) 101bp sequencing at GTAC (Genome Technology Access Center, Washington University in St. Louis) at 9pmol per lane.

### Genome Sequence Assembly

All sequencing reads were de-multiplexed by barcode into separate genome bins. Reads were quality trimmed to remove adapter sequence and bases on either end with a quality score below 19. Any reads shorter than 31bp after quality trimming were not used in further analysis. The best reference sequence was chosen for each isolate or plasmid by mapping 10000 reads chosen randomly from that isolate against all reference genomes (from NCBI Genome, downloaded July 14<sup>th</sup> 2014) of the same species as the isolate (in the case of genomic DNA assembly) or against all plasmid sequences containing NDM-1, KPC, or CTX-M (in the case of plasmid DNA assembly). Reads were mapped using Bowtie 2 (4) (command: `bowtie2 -x <reference_genome_index_name> -1 <forward_read_file> -2 <reverse_read_file> -q --phred33 -very-fast -I 100 -X 600 --no-discordant --no-mixed --no-unal --no-hd --no-sq --omit-sec-strand`). The genome or plasmid against which the highest percentage of reads mapped was used as the reference sequence for that assembly. It was empirically determined that if this first mapping included fewer than 60% of the reads, then the assembly would be best done completely *de novo*. For isolates with >60% of reads matching a reference sequence, all reads were mapped to that sequence (command: `bowtie2 -x <reference_genome_index_name> -1 <forward_read_file> -2 <reverse_read_file> -q --phred33 --very-sensitive-local -I 200 -X 1000 -S <sam_output>`). Variants from the reference were called using samtools (commands: `samtools view -buS <sam_file> | samtools sort -m 4000000000 - <sample_prefix> ### samtools index <bam_file> ### samtools mpileup -uD -f <reference_genome> <bam_file> | bcftools view -bcv - > <bcf_file> ### bcftools view <bcf_file>`). The variant call format file was then filtered to

remove SNPs with a quality score lower than 70 or coverage greater than twice the average coverage expected per base. Custom scripts were then used to extract DNA sequences from the reference genome with > three independent reads, to create a fragment file of regions in the sample genome matching the reference genome modified with high-quality variant information.

*De novo* assembly of the reads from each isolate was completed using Velvet (5) (commands: `velveth <output_directory> 51 -fastq -shortPaired <interleaved_reads> ### velvetg <output_directory> -ins_length 400 -exp_cov <kmer_coverage> -cov_cutoff <coverage_cutoff>`). Kmer coverage was calculated as: total read coverage\*0.50 (because the kmer length was approximately half the read length), and the coverage cutoff was calculated as the kmer coverage divided by eight. If a complete reference mapping was performed, then contigs from the *de novo* assembly and reference mapping were put in an additional velvet assembly step as long reads with the original reads files (commands: `velveth <output_directory> 51 -fastq -shortPaired -separate <forward_reads> <reverse_reads> -fasta -long <de_novo_fragments> <reference_fragments> ### velvetg <output_directory> -ins_length 400 -clean yes -conserveLong yes -exp_cov <kmer_coverage> -cov_cutoff <coverage_cutoff> -scaffolding yes -long_mult_cutoff 0`). Finally all fragments were collapsed on nucleotide identity using cd-hit (command: `cd-hit-est -I <fragment_file> -o <collapsed_file> -d 0 -M 0`). All fragments smaller than 500bp were partitioned to a separate file by a custom script. Plasmid sequences were assembled by this same method, with the sequences of all complete plasmids encoding, NDM-1, KPC, or CTX-M used as references.

### ORF Prediction and Annotation

ORF prediction for each genome was performed separately using GeneMark (6) models based on the closest reference genome (command: `gmhmmmp -m <model_name> -o <outfile> -a <contig_name_file>`). Each ORF was compared to three databases of profile hidden Markov models using HMMR (7): Pfam (command: `hmmscan --cut_ga -o /dev/null --tblast <target_out_file> --domtblast <domain_out_file> <Pfam_database_file> <protein_input_file>`), TIGRFAMs (command: `hmmscan --cut_ga -o /dev/null --tblast <target_out_file> --domtblast <domain_out_file> <database_file> <protein_input_file>`), and Resfams ([dantaslab.wustl.edu/resfams/](http://dantaslab.wustl.edu/resfams/)) (8) (command: `hmmscan --cut_ga -o /dev/null --tblast <target_out_file> --domtblast <domain_out_file> <database_file> <protein_input_file>`). All



functional annotations were concatenated into a single file by a custom script. 756 *E. coli* and 54 *K. pneumoniae* completed and draft genomes were downloaded from the National Center for Biotechnology Information (NCBI) on April 15<sup>th</sup> 2014, for the purpose of comparing to the isolate set. GeneMark models from the completed genomes were used to predict ORFs for those genomes, while for draft genomes models created from *E. coli* K12 MG1655 (for *E. coli* draft genomes) or *K. pneumoniae* KCTC 2242 (for *K. pneumoniae* draft genomes) were used for ORF prediction. All genome and plasmid sequences were deposited into NCBI (BioProject accession number: PRJNA261540).

### **In silico MLST**

Multi-Locus Sequence Typing (MLST) profiles were downloaded from PubMLST (pubmlst.org). When an absolute MLST profile could not be identified for an organism (because of ambiguous bases or incomplete assembly of one or more loci) the remaining possible sequence types (ST) based on the incomplete information were identified. In all cases a strain could be identified as one of at most 19 ST using this methodology. MLST profiles were only applied to *Escherichia coli* and *Klebsiella pneumoniae* since the PubMLST database does not contain an *Enterobacter* table.

### **Core Genome Alignment**

Command for whole genome alignment using mugsy (9): command: mugsy --directory <output\_directory> --prefix <output\_prefix> <genome\_fasta\_1> <genome\_fasta\_2> ... <genome\_fasta\_N>. For *E. coli*, *K. pneumoniae*, and *E. cloacae* a single reference genome was included in the alignment to provide context (*E. coli* K12 MG1655, *K. pneumoniae* KCTC 2242, and *E. cloacae* ATCC 13047 respectively). Poorly aligned regions (i.e. plasmids or recombined regions, which could create noise in the phylogenetic signal) were removed using Gblocks (10) (command: Gblocks <input\_file> -t=d -b3=24) leaving only the core genome alignment. Maximum likelihood trees made by RaxML (11) (command: raxmlHPC-SSE3 -s <input\_file> -n <output\_file> -m GTRGAMMA -d -f a -N 100 -x 54321 -w <output\_directory>) and FastTree (12) (command: FastTree -gtr -nt -gamma -nome <input\_file> > <output\_file>) were compared for agreement. When trees made by both methods were in agreement, the output from FastTree was used for visualization. Files were converted between various required formats by custom scripts.

Subspecies clades were defined as groups of branches descended from a common ancestor where no individual branch within the clade could have more than 0.005 substitutions per site. This definition yielded the same groupings as the *in silico* MLST described above, in all cases where all members of a clade could be assigned to a known ST (online Technical Appendix Figure 1, panels A,B).

### **Specific $\beta$ -lactamase Identification**

A BLAST database was constructed from the amino acid sequences of all  $\beta$ -lactamases cataloged in the Bush and Jacoby (13) database at [www.lahey.org/Studies/](http://www.lahey.org/Studies/) (accessed March 25<sup>th</sup>, 2014). Genes from our genomes annotated as  $\beta$ -lactamases were extracted and compared against this database by BLAST. Exact matches were then re-annotated with their specific  $\beta$ -lactamase name, while inexact matches were recorded as their closest hit plus an asterisk.

### **Plasmid Comparisons**

We compared plasmid sequences by an all-against-all pairwise nucleotide BLAST alignment. For each pair of plasmids, we calculated the percentage of each plasmid that aligned at >99% identity. We then binned the percentages from each pairwise alignment into groups based on the defining  $\beta$ -lactamase of their query and subject plasmids. We also generated network diagrams from the pairwise BLASTs using custom Python scripts and Cytoscape (14), only including regions above 99% identity and over 500 bp.

### **Isolate Characteristics**

The sampled Enterobacteriaceae isolates are phylogenetically diverse and include multiple examples of known pathogenic sequence types. We performed WGS of each isolate, totaling 33 *Escherichia coli* isolates, 30 *Klebsiella pneumoniae* isolates, 9 *Enterobacter cloacae* complex isolates, and 6 *Enterobacter aerogenes* isolates. We then used whole genome alignment of the core genomes of each species to reconstruct the phylogenetic relationships of each isolate at high resolution (Technical Appendix Figure 2). The species trees demonstrate that we sampled genomes from a variety of evolutionary clades as well as from multiple members of specific clades. They also demonstrate that clades could include isolates from both the United States and

Pakistan, allowing us to ignore the geographic variable and group the isolates by carbapenemase carriage for subsequent analyses. We also used housekeeping gene sequence from each isolate to perform *in silico* MLST, allowing us to compare our phylogenetic analysis to previously identified sequence types. We found that the clades on our tree include globally-disseminated pathogen sequence types, such as ST131 in *E. coli* and ST11 (single locus variant of ST258) in *K. pneumoniae*.

ST131 is noted for its virulence as well as for its frequent association with the CTX-M  $\beta$ -lactamases and fluoroquinolone resistance (15–17). Previous reports have found ST258, and closely related *K. pneumoniae*, to have relatively high rates of carbapenemase carriage (18,19). Despite the utility of MLST-based classification for large-scale epidemiological purposes, binning clinical isolates into sequence types masks genotypic and phenotypic variation due to HGT or single nucleotide polymorphisms, and therefore MLST cannot be used for fine-grained epidemiology or as an accurate predictor of antibiotic susceptibility. For example, two previous studies have shown that ST131 can be subdivided into three distinct lineages with different rates of antibiotic resistance (15,16). One of these studies found that the rapid global expansion of ST131 has been driven by the success of a specific subclone of ST131 that encodes fluoroquinolone resistant *gyrA* and *parC* alleles and CTX-M-15 (16), a characterization which fits 7 of our 11 ST131 isolates. We also identified a single ST131 isolate carrying KPC-2, which was resistant to all  $\beta$ -lactams tested. We also observed *K. pneumoniae* ST11 isolates carrying KPC-3, and others carrying NDM-1, which fits with reports characterizing ST11 as being highly common worldwide and frequently encoding carbapenemases (18,19).

### **Antibiotic Resistance Phenotypes**


To establish the overall susceptibility profiles of each of our strains, we performed phenotypic tests using Kirby Bauer Disk diffusion in accordance with CLSI guidelines on all 78 clinical isolates against 12 antibiotics including 7  $\beta$ -lactams (Technical Appendix Table 1). We found that 63% of all isolates were resistant to ciprofloxacin, a fluoroquinolone commonly used to treat urinary tract infections. We also found resistance to trimethoprim-sulfamethoxazole in 65% of isolates, and gentamicin, doxycycline, and chloramphenicol exhibited *in vitro* resistance in 45%, 54%, and 56% of isolates, respectively. In the  $\beta$ -lactams, we saw near universal resistance to ampicillin (96% of isolates) and variable resistance to the cephalosporins. A high

rate of resistance to meropenem was observed (31% of isolates), but this finding was not surprising since this was the property on which many of the isolates had been selected.

## References


1. Clinical and Laboratory Standards Institute. Performance standards for antimicrobial susceptibility testing; twenty-third informational supplement. 2013.
2. Manji R, Bythrow M, Branda JA, Burnham CAD, Ferraro MJ, Garner OB, et al. Multi-center evaluation of the VITEK® MS system for mass spectrometric identification of non-*Enterobacteriaceae* Gram-negative bacilli. *Eur J Clin Microbiol Infect Dis*. 2014 2014/03/01;33(3):337-46. [PubMed http://dx.doi.org/10.1007/s10096-013-1961-2](http://dx.doi.org/10.1007/s10096-013-1961-2)
3. Richter SS, Sercia L, Branda JA, Burnham CAD, Bythrow M, Ferraro MJ, et al. Identification of *Enterobacteriaceae* by matrix-assisted laser desorption/ionization time-of-flight mass spectrometry using the VITEK MS system. *Eur J Clin Microbiol Infect Dis*. 2013 2013/12/01;32(12):1571-8. [PubMed http://dx.doi.org/10.1007/s10096-013-1912-y](http://dx.doi.org/10.1007/s10096-013-1912-y)
4. Langmead B, Salzberg SL. Fast gapped-read alignment with Bowtie 2. *Nat Methods*. 2012;9:357–9. [PubMed http://dx.doi.org/10.1038/nmeth.1923](http://dx.doi.org/10.1038/nmeth.1923)
5. Zerbino DR, Birney E. Velvet: algorithms for de novo short read assembly using de Bruijn graphs. *Genome Res*. 2008;18:821–9. [PubMed http://dx.doi.org/10.1101/gr.074492.107](http://dx.doi.org/10.1101/gr.074492.107)
6. Borodovsky M, Lomsadze A. Gene identification in prokaryotic genomes, phages, metagenomes, and EST sequences with GeneMarkS suite. *Current protocols in bioinformatics / editorial board, Andreas D Baxevanis [et al]*. 2011;Chapter 4:Unit 4.5.1-17.
7. Eddy SR. A new generation of homology search tools based on probabilistic inference. *Genome Inform*. 2009;23:205–11. [PubMed http://dx.doi.org/10.1142/9781848165632\\_0019](http://dx.doi.org/10.1142/9781848165632_0019)
8. Gibson MKFK, Dantas G. Improved annotations of antibiotic resistance functions reveals microbial resistomes cluster by ecology. *ISME*. 2014. [PubMed http://dx.doi.org/10.1038/ismej.2014.106](http://dx.doi.org/10.1038/ismej.2014.106)
9. Angiuoli SV, Salzberg SL. Mugsy: fast multiple alignment of closely related whole genomes. *Bioinformatics*. 2011;27:334–42. [PubMed http://dx.doi.org/10.1093/bioinformatics/btq665](http://dx.doi.org/10.1093/bioinformatics/btq665)
10. Talavera G, Castresana J. Improvement of phylogenies after removing divergent and ambiguously aligned blocks from protein sequence alignments. *Syst Biol*. 2007;56:564–77. [PubMed http://dx.doi.org/10.1080/10635150701472164](http://dx.doi.org/10.1080/10635150701472164)

11. Stamatakis A. RAxML Version 8: A tool for phylogenetic analysis and post-analysis of large phylogenies. *Bioinformatics*. 2014 January 21, 2014. [PubMed](#)  
<http://dx.doi.org/10.1093/bioinformatics/btu033>
12. Price MN, Dehal PS, Arkin AP. FastTree 2—approximately maximum-likelihood trees for large alignments. *PLoS ONE*. 2010;5:e9490. [PubMed](#) <http://dx.doi.org/10.1371/journal.pone.0009490>
13. Bush K, Jacoby GA. Updated functional classification of beta-lactamases. *Antimicrob Agents Chemother*. 2010;54:969–76. [PubMed](#) <http://dx.doi.org/10.1128/AAC.01009-09>
14. Smoot ME, Ono K, Ruscheinski J, Wang P-L, Ideker T. Cytoscape 2.8: New features for data integration and network visualization. *Bioinformatics*. 2011;27:431–2  
[dx.doi.org/10.1093/bioinformatics/btq675](http://dx.doi.org/10.1093/bioinformatics/btq675). [PubMed](#)  
<http://dx.doi.org/10.1093/bioinformatics/btq675>
15. Petty NK, Ben Zakour NL, Stanton-Cook M, Skippington E, Totsika M, Forde BM, et al. Global dissemination of a multidrug resistant *Escherichia coli* clone. *Proc Natl Acad Sci U S A*. 2014;111:5694–9. [PubMed](#) <http://dx.doi.org/10.1073/pnas.1322678111>
16. Price LB, Johnson JR, Aziz M, Clabots C, Johnston B, Tchesnokova V, et al. The epidemic of extended-spectrum-beta-lactamase-producing *Escherichia coli* ST131 is driven by a single highly pathogenic subclone, H30-Rx. *MBio*. 2013;4:e00377–13. [PubMed](#)  
<http://dx.doi.org/10.1128/mBio.00377-13>
17. Rogers BA, Sidjabat HE, Paterson DL. *Escherichia coli* O25b-ST131: a pandemic, multiresistant, community-associated strain. *J Antimicrob Chemother*. 2011;66:1–14. [PubMed](#)  
<http://dx.doi.org/10.1093/jac/dkq415>
18. Cuzon G, Naas T, Truong H, Villegas MV, Wisell KT, Carmeli Y, et al. Worldwide diversity of *Klebsiella pneumoniae* that produce beta-lactamase blaKPC-2 gene. *Emerg Infect Dis*. 2010;16:1349–56. [PubMed](#) <http://dx.doi.org/10.3201/eid1609.091389>
19. Pena I, Picazo JJ, Rodriguez-Avial C, Rodriguez-Avial I. Carbapenemase-producing *Enterobacteriaceae* in a tertiary hospital in Madrid, Spain: high percentage of colistin resistance among VIM-1-producing *Klebsiella pneumoniae* ST11 isolates. *Int J Antimicrob Agents*. 2014;43:460–4. [PubMed](#) <http://dx.doi.org/10.1016/j.ijantimicag.2014.01.021>



**Technical Appendix Figure 1.** Phylogenetic trees for isolates from samples collected in Pakistan and the United States. Trees are separated by species, A) *Escherichia coli*, B) *Klebsiella pneumoniae*, C) *Enterobacter cloacae*, and D) *Enterobacter aerogenes*, but not rooted. Bootstrap values are 1 for each branch unless otherwise noted. For each species  $\approx 50\%$  of the genome was determined to be core, and was used for phylogenetic inference. Scale bars indicate the nucleotide substitutions per site. In a) and b) sequence types (ST) as determined by *in silico* multilocus sequence typing are indicated by boxes grouping members of the same ST together. Unk = ST does not correspond to any reported in pubMLST, Ind = exact ST could not be determined due to sequencing error. Reference genomes included for *E. coli* (K12 MG1655), *K. pneumoniae* (KCTC 2242), and *E. cloacae* (ATCC 13047) on their respective trees.





**Technical Appendix Figure 2.** Sequence conservation between plasmids containing NDM-1, KPC, or CTX-M  $\beta$ -lactamases. All plasmids from A) NCBI and B) this study that contained a NDM-1, KPC, or CTX-M  $\beta$ -lactamase were analyzed by all-against-all BLAST. Plasmid interactions were defined by the percentage of the query plasmid conserved (at >99% identity) in the subject plasmid. Plasmid interactions were plotted based on the defining  $\beta$ -lactamase of their query and subject plasmids.

**Technical Appendix Table 1.** Antibiotic drug susceptibility profiles of clinical isolates from Pakistan and the United States\*

Species, no.	Location	Phenotype	AM	CZ	CTT	CAZ	CRO	FEP	MEM	CIP	SXT	GM	D†	C†
<i>E. coli</i> , 9	U.S.A	R	78%	67%	22%	44%	44%	44%	44%	56%	33%	22%	33%	33%
		I	0%	11%	22%	0%	0%	0%	0%	0%	0%	0%	0%	11%
		S	22%	22%	56%	56%	56%	56%	44%	67%	78%	11%	0%	
<i>E. coli</i> , 24	Pakistan	R	96%	83%	13%	38%	63%	21%	0%	67%	75%	38%	67%	33%
		I	4%	17%	0%	13%	0%	21%	0%	0%	0%	0%	17%	4%
		S	0%	0%	88%	50%	38%	58%	100%	33%	25%	63%	17%	63%
<i>K. pneumoniae</i> , 11	U.S.A	R	100%	36%	0%	36%	36%	36%	36%	27%	36%	9%	18%	27%
		I	0%	0%	36%	0%	0%	0%	0%	0%	9%	0%	0%	0%
		S	0%	64%	64%	64%	64%	64%	73%	55%	91%	18%	9%	
<i>K. pneumoniae</i> , 19	Pakistan	R	100%	68%	21%	63%	63%	42%	16%	63%	68%	58%	32%	58%
		I	0%	5%	0%	0%	0%	16%	0%	11%	5%	0%	21%	11%
		S	0%	26%	79%	37%	37%	42%	84%	26%	26%	42%	47%	32%
<i>E. cloacae</i> , 3	U.S.A	R	100%	100%	100%	100%	100%	100%	100%	67%	67%	33%	33%	33%
		I	0%	0%	0%	0%	0%	0%	0%	0%	0%	33%	33%	33%
		S	0%	0%	0%	0%	0%	0%	0%	33%	33%	33%	33%	33%
<i>E. cloacae</i> , 6	Pakistan	R	83%	100%	67%	100%	100%	83%	67%	83%	83%	100%	50%	83%
		I	0%	0%	0%	0%	0%	17%	0%	17%	0%	0%	17%	0%
		S	0%	0%	33%	0%	0%	0%	33%	0%	17%	0%	33%	17%
<i>E. aerogenes</i> , 6	Pakistan	R	100%	100%	17%	100%	100%	100%	100%	100%	100%	67%	83%	100%
		I	0%	0%	83%	0%	0%	0%	0%	0%	0%	0%	17%	0%
		S	0%	0%	0%	0%	0%	0%	0%	0%	0%	0%	33%	0%

\*AM, ampicillin; CZ, cefazolin; CTT, cefotetan; CAZ, ceftazidime; CRO, ceftriaxone; FEP, cefepime; MEM, meropenem; CIP, ciprofloxacin; SXT, trimethoprim-sulfamethoxazole; GM, gentamicin; D, doxycycline; C, chloramphenicol; R, Resistant; I, Intermediate; S, Susceptible.

†For USA *E. coli* and *K. pneumoniae* in doxycycline and chloramphenicol selections, n=4.

**Technical Appendix Table 2.** Assembly metrics for whole genome assemblies

Species	Genome	Number of contigs	N50	Largest contig	Total nucleotides
<i>Escherichia coli</i>	PH100	1288	35053	385222	6309191
<i>Escherichia coli</i>	PH101–2	404	31119	177520	4857948
<i>Escherichia coli</i>	PH105	288	55979	220618	5045926
<i>Escherichia coli</i>	PH108	754	12427	95430	4930975
<i>Escherichia coli</i>	PH114	280	68389	450011	5902296
<i>Escherichia coli</i>	PH118	1028	13550	93913	5073129
<i>Escherichia coli</i>	PH129	235	76630	188170	5470109
<i>Escherichia coli</i>	PH135	156	87914	545356	4864787
<i>Escherichia coli</i>	PH141	656	14806	87255	4469903
<i>Escherichia coli</i>	PH143	355	33581	145252	4875919
<i>Escherichia coli</i>	PH151–2	368	41647	143184	5066089
<i>Escherichia coli</i>	PH156–1	720	24256	203811	5434228
<i>Escherichia coli</i>	PH18	357	39599	126851	5026978
<i>Escherichia coli</i>	PH20	513	20802	122933	4967604
<i>Escherichia coli</i>	PH31	394	40363	219779	5077941
<i>Escherichia coli</i>	PH39	465	28063	96484	4695860
<i>Escherichia coli</i>	PH51	1108	13989	85055	4988648
<i>Escherichia coli</i>	PH5	193	185476	496282	5367128
<i>Escherichia coli</i>	PH85	331	45973	149681	5118463
<i>Escherichia coli</i>	PH90	475	31325	258054	5169559
<i>Escherichia coli</i>	PH92–1	738	19526	90560	5313215
<i>Escherichia coli</i>	PH93	487	31876	685039	5927902
<i>Escherichia coli</i>	PH94	325	40628	218502	4960477
<i>Escherichia coli</i>	PH98	401	28050	272240	4791597
<i>Escherichia coli</i>	WU31	502	21374	148889	4858305
<i>Escherichia coli</i>	WU32	329	43395	166954	5598148
<i>Escherichia coli</i>	WU33	487	25419	139243	4975544
<i>Escherichia coli</i>	WU34	278	67976	229510	5206081
<i>Escherichia coli</i>	WU35	267	51428	235055	4846364
<i>Escherichia coli</i>	WU40	145	202444	388283	5052711
<i>Escherichia coli</i>	WU43	193	91836	346403	5033909
<i>Escherichia coli</i>	WU44	196	92332	610815	5473063
<i>Escherichia coli</i>	WU45	332	40224	155439	4990710
<i>Klebsiella pneumoniae</i>	PH102	2205	3247	29455	4976237
<i>Klebsiella pneumoniae</i>	PH10	192	129071	506496	5877659
<i>Klebsiella pneumoniae</i>	PH11	224	109166	648719	5657202
<i>Klebsiella pneumoniae</i>	PH124	238	66778	202869	5515528
<i>Klebsiella pneumoniae</i>	PH12	317	46831	180775	5530414
<i>Klebsiella pneumoniae</i>	PH139	263	92250	433917	5458209
<i>Klebsiella pneumoniae</i>	PH150–2	487	40719	311295	5568393
<i>Klebsiella pneumoniae</i>	PH152	354	74725	275992	5863335
<i>Klebsiella pneumoniae</i>	PH24–1	181	209112	601209	5541053
<i>Klebsiella pneumoniae</i>	PH25	439	51744	186984	5437740
<i>Klebsiella pneumoniae</i>	PH28–1	302	72750	480318	5876774
<i>Klebsiella pneumoniae</i>	PH38–1	178	243482	614325	6135768
<i>Klebsiella pneumoniae</i>	PH40	471	43868	232550	6212797
<i>Klebsiella pneumoniae</i>	PH44	251	133296	479093	6297207
<i>Klebsiella pneumoniae</i>	PH49–2	273	84383	550379	6883299
<i>Klebsiella pneumoniae</i>	PH72	2195	13163	144014	7250407
<i>Klebsiella pneumoniae</i>	PH73	997	11652	91567	5804995
<i>Klebsiella pneumoniae</i>	PH88	150	289170	512144	5630043
<i>Klebsiella pneumoniae</i>	PH9	927	12231	72933	5207334
<i>Klebsiella pneumoniae</i>	WU10	179	117943	480943	5928719
<i>Klebsiella pneumoniae</i>	WU12	1131	12084	59345	5227362
<i>Klebsiella pneumoniae</i>	WU18	961	14636	96067	5664085
<i>Klebsiella pneumoniae</i>	WU21	372	67952	249904	5556247
<i>Klebsiella pneumoniae</i>	WU23	142	208533	460529	5958526
<i>Klebsiella pneumoniae</i>	WU2	407	55286	345910	6330941
<i>Klebsiella pneumoniae</i>	WU3	705	94321	408241	6853166
<i>Klebsiella pneumoniae</i>	WU6	297	104936	426673	5919869
<i>Klebsiella pneumoniae</i>	WU7	227	79691	298601	5818526
<i>Klebsiella pneumoniae</i>	WU8	377	64077	729491	7506385
<i>Klebsiella pneumoniae</i>	WU9	141	120728	292977	5557123
<i>Enterobacter aerogenes</i>	112–2	90	177700	566534	5153448
<i>Enterobacter aerogenes</i>	PH113	450	22178	97515	4695038
<i>Enterobacter aerogenes</i>	PH134	274	107942	399309	5126469
<i>Enterobacter aerogenes</i>	PH138	776	30613	144769	9901518

Species	Genome	Number of contigs	N50	Largest contig	Total nucleotides
<i>Enterobacter aerogenes</i>	PH63	225	47345	184904	5211429
<i>Enterobacter aerogenes</i>	PH84-2	226	61644	336177	5307861
<i>Enterobacter cloacae</i>	PH23	179	83227	473778	5536166
<i>Enterobacter cloacae</i>	PH24-2	1238	20166	178617	8756501
<i>Enterobacter cloacae</i>	PH112-1	621	175912	517857	9155709
<i>Enterobacter cloacae</i>	PH125	1221	45282	215620	6004095
<i>Enterobacter cloacae</i>	PH158	552	34922	172567	9711730
<i>Enterobacter cloacae</i>	PH82	389	29307	134776	4974173
<i>Enterobacter cloacae</i>	WU26	538	46418	208322	5369428
<i>Enterobacter cloacae</i>	WU27	315	45300	155565	5251409
<i>Enterobacter cloacae</i>	WU29	457	24686	86433	4888311
Average	NA	490	67666	285311	5664479

**Technical Appendix Table 3.** Assembly metrics for plasmid assemblies

Species	Genome	Number of contigs	N50	Largest contig	Total nucleotides
<i>Acinetobacter baumannii</i>	PH147_2	98	3072	7683	181370
<i>Escherichia coli</i>	WU31	41	16473	48183	214615
<i>Escherichia coli</i>	WU32	86	11069	31992	233185
<i>Escherichia coli</i>	WU33	87	9555	47149	168182
<i>Klebsiella pneumoniae</i>	PH11	53	19073	48703	230288
<i>Klebsiella pneumoniae</i>	PH88	39	15980	48463	218205
<i>Klebsiella pneumoniae</i>	WU13	6	12943	12943	18694
<i>Klebsiella pneumoniae</i>	WU14	4	12976	12976	18501
<i>Klebsiella pneumoniae</i>	WU17	374	4432	60538	328889
<i>Klebsiella pneumoniae</i>	WU18	40	18017	43330	221015
<i>Klebsiella pneumoniae</i>	WU19	37	22152	87768	315122
<i>Enterobacter aerogenes</i>	PH112_2	80	19156	52826	450788
<i>Enterobacter aerogenes</i>	PH113	17	5703	19905	59909
<i>Enterobacter aerogenes</i>	PH134	22	4168	9715	48836
<i>Enterobacter aerogenes</i>	PH138	171	8795	35183	466555
<i>Enterobacter aerogenes</i>	PH63	37	29896	44891	282470
<i>Enterobacter cloacae</i>	PH23	111	12334	42017	516173
<i>Enterobacter cloacae</i>	PH24_2	95	15201	41616	562271
<i>Enterobacter cloacae</i>	PH82	206	9384	37900	641720
Average	NA	84	13178	38620	272463