

---

# Population Responses during the Pandemic Phase of the Influenza A(H1N1)pdm09 Epidemic, Hong Kong, China

Nelson C.Y. Yeung, Joseph T.F. Lau,  
Kai Chow Choi, Sian Griffiths

During August 2009–July 2010, we conducted 7 longitudinal telephone surveys among 503 adults in Hong Kong, China, to explore changes in their behavioral and psychological responses to the influenza A(H1N1)pdm09 virus epidemic. Trends were examined using generalized estimating equations models. Findings showed that responses varied with the course of the pandemic.

On June 11, 2009, the World Health Organization declared the influenza A(H1N1)pdm09 (pH1N1) virus outbreak a pandemic (1). Previous studies have investigated community responses to the pandemic in different countries during early stages of the epidemic (2–5). The studies investigated persons' risk perceptions and knowledge related to the virus, perceived efficacy of preventive measures, and psychological and behavioral responses. However, because of intersample variations, these cross-sectional studies did not capture within-person changes. We conducted a longitudinal cohort study to investigate changes in responses among the general Hong Kong, China, population during the pH1N1 pandemic.

## The Study

A cohort sample of 18- to 60-year-old adults in Hong Kong participated in 7 rounds of telephone surveys during August 2009–July 2010, which covered almost the entire pH1N1 pandemic period in Hong Kong. At baseline, we invited 677 adults to participate; 503 (74.3%) consented and completed the survey (online Technical Appendix Table 1, <https://wwwnc.cdc.gov/EID/article/23/5/16-0768-Techapp1.pdf>). We measured the following variables: knowledge about the modes of pH1N1 virus transmission; risk perceptions associated with the virus (perceived susceptibility to and severity of infection); perceived efficacy

and use of preventive measures (e.g., handwashing, using a facemask); psychological responses (worry about infection and emotional distress); and evaluations of the government's performance in pandemic control. Sample sizes for surveys 2–7 ranged from 452 to 481, yielding retention rates of 89.8%–95.6%.

Most participants were women (57.9%), 40–60 years of age (55.8%), employed full time (55.9%), and married (65%). Sex and age distributions were comparable to those in the local census data (6). We aimed to determine whether there were overall linear trends in participants' perceptions, psychological responses, and behavioral responses to the pandemic. We examined linear trends for these variables across the 7 time points by using generalized estimating equations (GEE) models. GEE models not only account for intracorrelated repeated measures data but also fit various data types using appropriate link functions. The analyses were conducted using PROC GENMOD (SAS Institute, Cary, NC, USA); 2-sided  $p < 0.05$  was considered significant.

Over time, >85% of the participants used a face mask and immediately visited a doctor when experiencing influenza-like symptoms. More than 50% of the participants washed their hands >10 times every day throughout the survey period ( $p > 0.05$ ). As the pandemic progressed, a decreasing percentage of participants wore masks in public areas; avoided touching their mouth, nose, and eyes; took antiviral drugs; and avoided crowded places ( $p < 0.001$ ) (online Technical Appendix Table 2). Percentages of participants feeling worried, depressed, or emotionally disturbed about pH1N1 virus decreased over time ( $p < 0.001$ ) (online Technical Appendix Table 3).

Over time, a decreasing percentage of participants recognized that touching infected persons or contaminated objects could result in virus transmission ( $p < 0.001$ ). Throughout the study period, a consistently high percentage of participants (>92%) recognized that the virus could be transmitted via respiratory droplets. Misconceptions about possible transmission through insect bites (26.1%) and water sources (34.5%) were prevalent. The percentage of participants reporting at least 1 misconception was stable over time ( $p > 0.05$ ). A consistently high percentage (>90%) of participants believed that using face masks in public areas, washing hands frequently, and avoiding crowded places could effectively prevent the spread of pH1N1 virus ( $p > 0.05$ ).

---

Author affiliations: The Chinese University of Hong Kong, The Jockey Club School of Public Health and Primary Care, Hong Kong, China (N.C.Y. Yeung, J.T.F. Lau, S. Griffiths); The Chinese University of Hong Kong, The Nethersole School of Nursing, Hong Kong (K.C. Choi)

DOI: <https://dx.doi.org/10.3201/eid2305.160768>

The percentage of participants believing that pH1N1 virus would be more harmful than seasonal influenza in terms of fatality and bodily damage increased over time ( $p < 0.001$ ). The percentages of participants who believed the population was highly susceptible to pH1N1 virus infection and who perceived a high chance of having a large-scale local outbreak in the coming year dropped significantly ( $p < 0.001$ ), but some fluctuations were observed; for example, the percentage peaked during survey round 2 (around the September influenza season).

Throughout the study period,  $\approx 12\%$ – $21\%$  of the participants gave a failing score ( $< 5$  on a 0- to 10-point scale) for the governments overall performance in controlling the pandemic ( $p > 0.05$ ). However, during survey rounds 2–7, an increasing percentage of participants believed in the governments ability to control the pandemic ( $p < 0.001$ ) (online Technical Appendix). The percentage of participants who believed that Hong Kong would not have enough vaccine or medication to deal with the pandemic decreased over time ( $p < 0.001$ ).

## Conclusions

This study investigated changes in community perceptions over the course of the pH1N1 pandemic in Hong Kong. Findings were highly comparable to those from other local cross-sectional surveys (5,7) and a systematic review (8). Knowledge regarding preventive measures and adherence to such measures was, in general, higher among our participants than among the general population in other countries (e.g., Australia, India, and the Netherlands) (9–11). The prevalence of misconceptions about some incorrect modes of transmission (e.g., insect bites) gradually declined. However,  $\approx 50\%$  of participants still held at least 1 of the 4 misconceptions regarding transmission (i.e., airborne transmission over a long distance and transmission through insect bites, water sources, and well-cooked pork). Furthermore, over time, a lower percentage of participants avoided touching their eyes, nose, and mouth to prevent virus transmission. A 2015 systematic review suggested that health authorities should provide more updated information about the virus (8). We also recommend using health campaigns to increase public awareness about different routes of pH1N1 virus transmission.

Perceived severity of pH1N1 virus infection decreased over time, which may partially explain the decline in distress and avoidance behaviors; this pattern was also observed in a recent review (8). However, an increasing proportion of participants believed that, compared with seasonal influenza, pH1N1 resulted in more deaths and more severe body damage. Perceived susceptibility to infection declined substantially as the epidemic progressed, suggesting that the public gradually perceived fewer risks from pH1N1 virus. Avoidance behaviors and use

of facemasks in the absence of influenza-like symptoms became less prevalent over time, similar to a trend seen in Malaysia (12). Mental distress among persons in Hong Kong was lower during the pH1N1 pandemic than during the SARS (severe acute respiratory syndrome) pandemic (13), possibly due to the milder consequences of pH1N1 infection. Persons in Hong Kong seemed to remain rational during the pandemic, thereby avoiding possible pandemic-associated economic threats.

Public support for the government declined over time. During survey round 5, a total of 20.6% of the participants gave a failing score to the government's performance, and 13.5% perceived that the government would not be able to control the pandemic. The poll was split as to whether the government should use the same response for pH1N1 influenza and seasonal influenza. Our findings suggest that the public should be advised of the pros and cons of pH1N1 control policies; a watchful step-down may be better accepted if the policies are understood.

This study has limitations. First, telephone surveys may be subject to self-selection bias. However, participants' demographics were comparable to those in local census data (6). Second, Hong Kong's unique experience with the SARS outbreak may have influenced the population's response to the pH1N1 pandemic; thus, our findings may not be fully generalizable to other countries. Third, we treated time as a continuous variable in the GEE models. Ideally, polynomials should be added to the linear time variable; however, given the small number of time points and absence of theoretical shapes, that was not feasible.

Our findings provide valuable information regarding overall linear trends and changes in community responses toward the pH1N1 pandemic among a Hong Kong cohort. These findings should help inform other countries in formulating appropriate pandemic control plans for influenza and other emerging infectious diseases.

## Acknowledgments

We thank the study participants, and special thanks are extended to Mei Wah Chan, Cheri Tong, and Albert Cheung for coordinating the telephone surveys.

This study was supported by the Research Fund for the Control of Infectious Diseases (PHE-1&10), Food and Health Bureau, the Hong Kong Special Administrative Region.

Dr. Yeung is a research assistant professor at the Jockey Club School of Public Health and Primary Care, the Chinese University of Hong Kong. His research interests are health promotion and improvement, particularly in community responses and preventive behaviors in response to emerging infectious diseases, chronic illness prevention and management, and mental health promotion.

## References

1. World Health Organization. World now at the start of 2009 influenza pandemic. Statement to the press by WHO Director-General Dr. Margaret Chan. June 11, 2009 [cited 2017 Mar 9]. [http://www.who.int/mediacentre/news/statements/2009/h1n1\\_pandemic\\_phase6\\_20090611/en/](http://www.who.int/mediacentre/news/statements/2009/h1n1_pandemic_phase6_20090611/en/)
2. Lau JT, Griffiths S, Choi KC, Tsui HY. Widespread public misconception in the early phase of the H1N1 influenza epidemic. *J Infect.* 2009;59:122–7. <http://dx.doi.org/10.1016/j.jinf.2009.06.004>
3. Rubin GJ, Amlôt R, Page L, Wessely S. Public perceptions, anxiety, and behaviour change in relation to the swine flu outbreak: cross sectional telephone survey. *BMJ.* 2009;339(jul02 3):b2651. <http://dx.doi.org/10.1136/bmj.b2651>
4. Effler PV, Carcione D, Giele C, Dowse GK, Goggin L, Mak DB. Household responses to pandemic (H1N1) 2009-related school closures, Perth, Western Australia. *Emerg Infect Dis.* 2010;16:205–11. <http://dx.doi.org/10.3201/eid1602.091372>
5. Cowling BJ, Ng DMW, Ip DKM, Liao Q, Lam WWT, Wu JT, et al. Community psychological and behavioral responses through the first wave of the 2009 influenza A(H1N1) pandemic in Hong Kong. *J Infect Dis.* 2010;202:867–76. <http://dx.doi.org/10.1086/655811>
6. Census and Statistics Department, Hong Kong SAR Government. Digest of Statistics, 2009 edition [cited 2017 Mar 9]. <http://www.statistics.gov.hk/pub/B10100032009AN09B0700.pdf>
7. Liao Q, Cowling BJ, Lam WW, Ng DMW, Fielding R. Anxiety, worry and cognitive risk estimate in relation to protective behaviors during the 2009 influenza A/H1N1 pandemic in Hong Kong: ten cross-sectional surveys. *BMC Infect Dis.* 2014;14:169. <http://dx.doi.org/10.1186/1471-2334-14-169>
8. Bults M, Beaujean DJMA, Richardus JH, Voeten HACM. Perceptions and behavioral responses of the general public during the 2009 influenza A (H1N1) pandemic: a systematic review. *Disaster Med Public Health Prep.* 2015;9:207–19. <http://dx.doi.org/10.1017/dmp.2014.160>
9. Seale H, McLaws ML, Heywood AE, Ward KF, Lowbridge CP, Van D, et al. The community's attitude towards swine flu and pandemic influenza. *Med J Aust.* 2009;191:267–9.
10. Kamate SK, Agrawal A, Chaudhary H, Singh K, Mishra P, Asawa K. Public knowledge, attitude and behavioural changes in an Indian population during the Influenza A (H1N1) outbreak. *J Infect Dev Ctries.* 2009;4:7–14.
11. Bults M, Beaujean DJMA, de Zwart O, Kok G, van Empelen P, van Steenbergen JE, et al. Perceived risk, anxiety, and behavioural responses of the general public during the early phase of the Influenza A (H1N1) pandemic in the Netherlands: results of three consecutive online surveys. *BMC Public Health.* 2011;11:2. <http://dx.doi.org/10.1186/1471-2458-11-2>
12. Lau JTF, Griffiths S, Choi KC, Lin C. Prevalence of preventive behaviors and associated factors during early phase of the H1N1 influenza epidemic. *Am J Infect Control.* 2010;38:374–80. <http://dx.doi.org/10.1016/j.ajic.2010.03.002>
13. Lau JTF, Griffiths S, Choi KC, Tsui HY. Avoidance behaviors and negative psychological responses in the general population in the initial stage of the H1N1 pandemic in Hong Kong. *BMC Infect Dis.* 2010;10:139. <http://dx.doi.org/10.1186/1471-2334-10-139>

Address for correspondence: Joseph T.F. Lau, The Chinese University of Hong Kong, The Jockey Club School of Public Health and Primary Care, Room 505 Postgraduate Education Centre, Prince of Wales Hospital, Shatin, Hong Kong, China; email: [jlau@cuhk.edu.hk](mailto:jlau@cuhk.edu.hk)

## EID Podcast: Stained Glass and Flu



The work of art shown here depicts the interrelationship of human, animal, and environmental health.

Stained-glass windows have been appreciated for their utility and splendor for more than 1,000 years, and this engaging work of art by stained glass artist Jenny Hammond reminds us that influenza A viruses—which can be easily spread between animals and human, use various host species, and exist in many different environments—remain an enduring and global health concern.

**Visit our website to listen:**

**EMERGING  
INFECTIOUS DISEASES**

[https://www2c.cdc.gov/  
podcasts/player.  
asp?f=8644950](https://www2c.cdc.gov/podcasts/player.asp?f=8644950)



# Population Responses during the Pandemic Phase of the Influenza A(H1N1)pdm09 Epidemic, Hong Kong, China

## Technical Appendix

### Introduction

#### Previous findings on community responses at different phases of the H1N1 epidemic

During the pre-community outbreak phase of the H1N1 epidemic from May 6 to June 6 2009, two studies in Hong Kong reported a moderate level of perceived susceptibility of contracting the disease (around 7.5% to 12.1% of the general population felt susceptible) and perceived severity of H1N1 (over 15% of general population believed that H1N1 is highly fatal), plus a low level of H1N1-related distress in the community (less than 5% were panicking). Misconceptions about the modes of H1N1 transmissions were common. These studies found that perceptions related to bodily damages, efficacy of frequent handwashing, non-availability of effective vaccines, chance of having a large-scale local outbreak, and mental distress due to influenza A/H1N1 were associated with frequent handwashing (1,2). Another study involving a series of cross-sectional surveys was also conducted, suggesting a generally low level of anxiety and a slight improvement in knowledge on modes of transmission throughout the epidemic. The perceived susceptibility and perceived severity of getting H1N1 were high in the early phase, but started to decline in the early pandemic phase epidemic and remained stable thereafter. Preventive behaviors like handwashing and wearing facemasks did not show significant change throughout the epidemic, while avoidance behaviors like avoiding going to crowded places declined gradually at the pandemic unfolds (3). During the early phase of the global H1N1 pandemic, other international studies reported a moderate level of distress in the U.S., a lack of preventive responses such as change in frequency of handwashing in the UK (4) and in India, plus a lack of support

for governmental mitigation strategies, such as school closure in Australia (5). Other studies found that Korean college students and Italian healthcare workers washed their hands or used hand sanitizer more frequently than before the beginning of the pandemic. The Chinese general population in seven urban regions and two rural regions showed an increasing level of knowledge, but a declining level of risk perception and adoption of preventive measures, while the U.S. general population showed an increasing trend in perceived risk but a decreasing trend in prevalence of precautionary activities (6). A study in Malaysia also showed that preventive measures, fear and avoidance changed in concordance with the trend in number of reported deaths (7).

## **Methods**

### **The specific dates for the seven rounds of surveys**

The study population comprised of Chinese adults (aged 18–60) who were living in Hong Kong. Seven longitudinal telephone surveys were conducted during August 2009 through July 2010 (Round 1: Aug 19 – Sep 10; Round 2: Sep 25 – Oct 5; Round 3: Oct 27 – Nov 2; Round 4: Nov 28 – Dec 10; Round 5: Jan 28 – Feb 5; Round 6: Mar 24 – Apr 1; Round 7: Jun 24 – Jul 10). During the study period, the Hong Kong Hospital Authority lowered the influenza response level from “Emergency” to “Alert” Response Level on May 24, 2010. The numbers of H1N1 death cases at the end of the seven survey dates were respectively 4, 20, 37, 41, 64, 76 and 80.

### **Details of study procedure**

At least three other independent telephone calls were made at different hours/days to unanswered telephone numbers. Over 95% of the households in Hong Kong have a fixed-line telephone installed (5). To avoid over-representing people who were not employed, the baseline survey was conducted during 6:30–10 p.m. Verbal consent was sought from the respondents before the interview commenced. Participants provided us with their name (or nickname) for identification, at least two contact phone numbers, and time that they were usually available for phone interview. Participants were given a supermarket voucher (HKD\$50 or about USD\$6.5) by mail as compensation for the completion of each survey.

## **Results**

### **Background characteristics**

The gender/age distributions were comparable to those of the recent census data (% female = 46.0%; % 40–60 = 51.5%). Proportion of female participants was higher among the respondents than among the non-responders (57.9% versus 45.4%,  $p < 0.01$ ). No significant between-group differences were found for other socio-demographic characteristics ( $p > 0.05$ ; Table 1).

### **Preventive and avoidance behaviors**

Over time, >85% of the participants used a face mask and immediately visited a doctor when experiencing influenza-like symptoms. More than 50% of the participants washed their hands >10 times every day throughout the survey period ( $p > 0.05$ ). As the pandemic progressed, a decreasing percentage of participants wore masks in public areas; avoided touching their mouth, nose, and eyes; took anti-influenza drugs; avoided crowded places; and took Traditional Chinese medicine to prevent virus infection ( $p < 0.01$ ) (Table 2).

### **Worry and distress related to H1N1**

Fewer participants worried about themselves (from 12.3% in Round 1 to 2.4% in Round 7,  $p < 0.001$ ) or their family members (from 15.3% in Round 1 to 2.4% in Round 7,  $p < 0.001$ ) contracting H1N1, while more participants believed that “it does not matter if I contracted H1N1” (from 41.4% in Round 1 to 50.7% in Round 7,  $p < 0.001$ ). Fewer participants were in panic, feeling much depressed or felt emotionally disturbed due to H1N1, the prevalence at Round 7 became <1% for these three individual items. Percentage of participants expressing any type of mental distress (feeling panic/depressed/emotionally disturbed about H1N1) also significantly decreased from Round 1 to 7 (from 16.3% to 2.8%,  $p < 0.001$ ) (Table 3).

### **Knowledge regarding modes of transmission of H1N1**

Over time, a decreasing percentage of participants recognized that touching infected persons and touching contaminated objects could transmit H1N1 ( $p < 0.001$ ). Throughout the whole study period, a consistently high percentage of participants (>92%)

recognized that H1N1 could be transmitted through droplets. Misconceptions about possible transmissions of H1N1 via insect bites (26.1%) and water sources (34.5%) were prevalent throughout the study period (Table 4).

#### **Perceived severity of H1N1**

Over time, fewer participants had mistaken that fatality rate of H1N1 was  $\geq 1\%$  (Round 1: 31.4% to Round 7: 24.8%, such percentage peaked at Round 5: 47.6%) (Table 5).

#### **Perceived susceptibility of H1N1**

Fewer participants perceived a chance of having a large-scale local outbreak in the coming year (from 41.4% in Round 1 to 3.4% in Round 7, Table 6,  $p < 0.001$ ). Similar percentages of respondents were confident in keeping themselves and their family free from H1N1 over time (over 70% at all rounds and peaking at Round 7: 92.3%, Table 6). Nevertheless, the prevalence of perceived susceptibility in contracting H1N1 fluctuated over time, in terms of susceptibility for oneself (Round 1: 16.6% to Round 7: 2.6%), one's family (Round 1: 14.9% to Round 7: 2.6%) and the general public (Round 1: 20.9% to Round 7: 2.6%) (Table 6).

#### **Perceived efficacy of preventive measures**

A consistently high percentage (>90%) of the participants believed that using face masks in public areas, washing hands frequently, and avoiding crowded places could effectively prevent the spread of H1N1 ( $p > 0.05$ ) (Table 7).

#### **Support toward government and pandemic control policies**

Less than 10% of the participants at all rounds believed that the Hong Kong government overreacted to the H1N1 pandemic ( $p > 0.05$ ; Table 8). Less than 5% of the participants at Rounds 2–6 believed that the government should cancel all control measures for controlling the H1N1 pandemic, but the percentage peaked at Round 7 (15.6%; Table 9). During Round 2 through 7, more participants believed in government's ability to control the H1N1 pandemic (from 77.1% in Round 1 to 94.9% in Round 7,  $p < 0.001$ ; Table 9), while a sizable proportion of participants believed that the government should treat H1N1 in the same manner as seasonal flu fluctuated (R1: 53.1%, R5: 38.9%, R7: 54.4%,  $p < 0.001$ ) (Table 9).

## Discussion

### Comparison between Hong Kong community responses and those in other countries

Our participants had in general higher levels of knowledge (e.g., >92% recognizing transmission via droplets) and practice of preventive measures, as compared to general populations of other countries. For instance, only three-fourth of general population in China knew that H1N1 could be transmitted by droplets; less than 10% of the military men in Singapore were aware that influenza can be spread by touch (8); less than 25% of Indians and 50% of Australians believed that handwashing was effective in preventing H1N1; only one-fourth of general population in the Netherlands regarded wearing facemasks as an effective preventive measure and only 38% used preventive measures (9); only 6.5% of Korean college students reported washing their hands more than 10 times every day during the pandemic period.

Similarly, a study conducted in October 2009 showed that over 40% of medical students in Pakistan believed that avoiding pork consumption could effectively prevent H1N1. Another study in China showed that 30% of Chinese general population believed that food could transmit H1N1. In general, the general public in Hong Kong over-estimated the fatality rate of the H1N1 (around 40%) and believed that H1N1 is associated with higher fatality and bodily damages as compared to seasonal flu (50%–60%).

We confirmed that unlike SARS, mental distress is relatively light during the H1N1 pandemic, possible due to the milder consequences of H1N1 as compared to SARS. Except for the peak of incidence of H1N1 in September, only  $\approx 10\%$  of the respondents felt worried about contracting H1N1 and very few reported distress. About two-fifth of the general public expressed would not be bothered by H1N1 infection. No panic has been caused by the pandemic. Avoidance behaviors and use of face mask in the absence of ILI symptoms became less prevalent over time. Such a trend was also understandable as a Malaysian study showed that fear and avoidance changed according to the number of reported deaths (7). The general public in Hong Kong has apparently been becoming more rational to live with the pandemic, saving potential economic threats.



In contrast, decreasing trends in perceived severity of and anxiety associated with H1N1 were observed among Dutch population as the pandemic progressed (9). Perceived susceptibility was quite low and did not show obvious trend during the study period; 10%–15% of the general public considered himself/herself susceptible of contracting H1N1 in the coming year. Such findings are comparable to those in a study in Korea, where only 8% of the college students perceived high susceptibility to contracting H1N1.

As the pandemic progressed, we see a gradual increase in the proportions of participants believing that 1) school suspension policies is not necessary, 2) the government should suspend all pandemic control measures, and 3) the government should treat H1N1 as the same way as seasonal flu. At the same time, people were less worried about contracting H1N1, less likely to feel mental health distress, and engaging in preventive behaviors less frequently over time. It is potentially attributed to people's gaining understanding about the severity of the pandemic and getting used to the pandemic over time. These findings provide some support to the idea that people's risk perceptions of the disease and behavioral responses are consistent with their perceptions about the necessity of some pandemic control policies.

Governments in other countries seemed not to be well-supported. A study in Australia showed that people were not supporting the school closure policy as they believed that H1N1 was mild and the policy could not stop H1N1 from spreading (5). Another study in Turkey showed that the public generally perceived that its government was not managing H1N1 well and had a low level of trust to government's H1N1 information. A similar trend was also observed in the Netherlands, perceived reliability of government information decreased (9). Over half of Dutch respondents believed that the threat was exaggerated by the media and government (9).

### **Summary of findings**

The general public maintains high level of preventive behaviors and a low level of mental distress throughout the study period. Compared to other countries, the general public in Hong Kong seemed to be more compliant to preventive behaviors like frequent handwashing and be more equipped with knowledge related to H1N1. Misconceptions became less prevalent but were still noticeable. Given the prior experience with SARS, Hong Kong government kept its vigilance for H1N1 by daily reporting of infection cases and frequent TV/radio commercials about H1N1 prevention. Dissemination of knowledge and frequent updates about H1N1 pandemic

from the government may partially explain the consistently high rate of preventive behaviors and relatively better knowledge. Perceived severity and susceptibility remained relatively stable and somehow low. The public rated the government very highly at the beginning at the pre-community outbreak phase (*I*) but the rating declined over time. However, the public was still supporting the government to have treat H1N1 differently from seasonal flu, and the overall support was still very reasonable.

## References

1. Lau JT, Griffiths S, Choi KC, Tsui HY. Widespread public misconception in the early phase of the H1N1 influenza epidemic. *J Infect.* 2009;59:122–7. [PubMed http://dx.doi.org/10.1016/j.jinf.2009.06.004](http://dx.doi.org/10.1016/j.jinf.2009.06.004)
2. Lau JTF, Griffiths S, Choi KC, Lin C. Prevalence of preventive behaviors and associated factors during early phase of the H1N1 influenza epidemic. *Am J Infect Control.* 2010;38:374–80. [PubMed http://dx.doi.org/10.1016/j.ajic.2010.03.002](http://dx.doi.org/10.1016/j.ajic.2010.03.002)
3. Cowling BJ, Ng DMW, Ip DKM, Liao Q, Lam WWT, Wu JT, et al. Community psychological and behavioral responses through the first wave of the 2009 influenza A(H1N1) pandemic in Hong Kong. *J Infect Dis.* 2010;202:867–76. [PubMed http://dx.doi.org/10.1086/655811](http://dx.doi.org/10.1086/655811)
4. Rubin GJ, Amlôt R, Page L, Wessely S. Public perceptions, anxiety, and behaviour change in relation to the swine flu outbreak: cross sectional telephone survey. *BMJ.* 2009;339(jul02 3):b2651. [PubMed http://dx.doi.org/10.1136/bmj.b2651](http://dx.doi.org/10.1136/bmj.b2651)
5. Effler PV, Carcione D, Giele C, Dowse GK, Goggin L, Mak DB. Household responses to pandemic (H1N1) 2009-related school closures, Perth, Western Australia. *Emerg Infect Dis.* 2010;16:205–11. [PubMed http://dx.doi.org/10.3201/eid1602.091372](http://dx.doi.org/10.3201/eid1602.091372)
6. Ibuka Y, Chapman GB, Meyers LA, Li M, Galvani AP. The dynamics of risk perceptions and precautionary behavior in response to 2009 (H1N1) pandemic influenza. *BMC Infect Dis.* 2010;10:296. [PubMed http://dx.doi.org/10.1186/1471-2334-10-296](http://dx.doi.org/10.1186/1471-2334-10-296)
7. Wong LP, Sam IC. Temporal changes in psychobehavioral responses during the 2009 H1N1 influenza pandemic. *Prev Med.* 2010;51:92–3. [PubMed http://dx.doi.org/10.1016/j.yjpm.2010.04.010](http://dx.doi.org/10.1016/j.yjpm.2010.04.010)

8. Yap J, Lee VJ, Yau TY, Ng TP, Tor PC. Knowledge, attitudes and practices towards pandemic influenza among cases, close contacts, and healthcare workers in tropical Singapore: a cross-sectional survey. BMC Public Health. 2010;10:442. [PubMed](#)  
<http://dx.doi.org/10.1186/1471-2458-10-442>
9. Bults M, Beaujean DJ, de Zwart O, Kok G, van Empelen P, van Steenberg JE, et al. Perceived risk, anxiety, and behavioural responses of the general public during the early phase of the Influenza A (H1N1) pandemic in the Netherlands: results of three consecutive online surveys. BMC Public Health. 2011;11:2. [PubMed](#) <http://dx.doi.org/10.1186/1471-2458-11-2>

**Technical Appendix Table 1.** Background characteristics, Census & Statistics Department HKSAR, 2009\*

Characteristic	n = 503	%
<b>Gender</b>		
Male	212	42.1
Female	291	57.9
<b>Age††</b>		
<30	122	24.3
30 – 39	100	19.9
40 – 49	154	30.7
50 – 60	126	25.1
<b>Education level</b>		
Form 3 or below	79	15.8
Form 4 – matriculation	217	43.3
College or above	205	40.9
<b>Marital status</b>		
Single	168	33.4
Married /cohabited	327	65.0
Divorced /widowed	8	1.6
<b>Employment status</b>		
Full-time	280	55.9
Part-time	18	3.6
Students	46	9.2
Retired	25	5.0
Housewife	113	22.6
Unemployed	18	3.6
Others	1	0.2
<b>Current health care practitioner</b>		
No	482	97.4
Yes	13	2.6

\*Based on the census data published in Hong Kong Annual Digest of Statistics, 2009

†the proportions of male and female of age 18–60 in Hong Kong were respectively

46.02% and 53.98%

††the proportions of the 4 age groups in Hong Kong (<30, 30–39, 40–49, and 50–60) were respectively 24.75%, 23.78%, 27.33%, and 24.14%.

**Technical Appendix Table 2.** Preventive and avoidance behaviors related to H1N1 influenza A/H1N1

Preventive or avoidance behavior	Round 1 (n = 503), 19 Aug – 10 Sep 09	Round 2 (n = 481), 25 Sep – 5 Oct 09	Round 3 (n = 475), 27 Oct – 2 Nov 09	Round 4 (n = 473), 28 Nov – 10 Dec 09	Round 5 (n = 452), 28 Jan – 5 Feb 10	Round 6 (n = 457), 7 – 14 Apr 10	Round 7 (n = 467), 24 Jun – 6 Jul 10	p (trend)
<b>Preventive behaviors</b>								
<b>Wearing mask when going out in case of ILI symptoms</b>								
Most not / certainly not / unsure	37 (7.4%)	51 (10.6%)	58 (12.2%)	70 (14.8%)	45 (10.0%)	58 (12.7%)	63 (13.5%)	0.012
Mostly /certainly	466 (92.6%)	430 (89.4%)	416 (87.8%)	403 (85.2%)	407 (90.0%)	398 (87.3%)	404 (86.5%)	
<b>Consult a doctor immediately if you have fever</b>								
Most not / certainly not / unsure	39 (7.8%)	22 (4.6%)	13 (2.7%)	22 (4.7%)	38 (8.4%)	49 (10.7%)	56 (12.0%)	<0.001
Mostly /certainly	464 (92.2%)	459 (95.4%)	462 (97.3%)	451 (95.3%)	414 (91.6%)	408 (89.3%)	411 (88.0%)	

	Round 1 (n = 503), 19 Aug – 10 Sep 09	Round 2 (n = 481), 25 Sep – 5 Oct 09	Round 3 (n = 475), 27 Oct – 2 Nov 09	Round 4 (n = 473), 28 Nov – 10 Dec 09	Round 5 (n = 452), 28 Jan – 5 Feb 10	Round 6 (n = 457), 7 – 14 Apr 10	Round 7 (n = 467), 24 Jun – 6 Jul 10	p (trend)
<b>Preventive or avoidance behavior</b>								
Tell immigration control if you have the flu or cold-like symptoms								
Most not / certainly not / unsure	23 (4.6%)	17 (3.5%)	8 (1.7%)	16 (3.4%)	–	–	–	0.099
Mostly /certainly	480 (95.4%)	464 (96.5%)	466 (98.3%)	457 (96.6%)	–	–	–	
Wearing mask in public areas if you have not IL symptoms in the last week								
No / Rarely	435 (86.5%)	399 (83.5%)	436 (91.8%)	438 (92.8%)	417 (92.7%)	436 (95.4%)	452 (96.8%)	<0.001
Sometimes / Always	68 (13.5%)	79 (16.5%)	39 (8.2%)	34 (7.2%)	33 (7.3%)	21 (4.6%)	15 (3.2%)	
Frequency of washing hands per day								
1 – 10	253 (50.4%)	229 (47.6%)	203 (42.7%)	206 (43.6%)	213 (47.2%)	221 (48.4%)	197 (42.3%)	0.075
>10	249 (49.6%)	252 (52.4%)	272 (57.3%)	267 (56.4%)	238 (52.8%)	236 (51.6%)	269 (57.7%)	
Avoid touching mouth, nose and eyes so as to reduce the risk of contracting Flu A (H1N1)								
No / Rarely	326 (64.8%)	311 (64.9%)	371 (78.1%)	371 (78.6%)	318 (70.8%)	347 (76.3%)	341 (73.0%)	0.003
Sometimes / Always	177 (35.2%)	168 (35.1%)	104 (21.9%)	101 (21.4%)	131 (29.2%)	108 (23.7%)	126 (27.0%)	
Take Tamiflu or other anti-flu drugs								
Most not / certainly not / unsure	468 (93.6%)	463 (97.9%)	470 (99.2%)	456 (97.2%)	445 (98.5%)	454 (99.6%)	461 (98.7%)	<0.001
Mostly /certainly	32 (6.4%)	10 (2.1%)	4 (0.8%)	13 (2.8%)	7 (1.5%)	2 (0.4%)	6 (1.3%)	
Take Traditional Chinese Medicine to prevent human swine flu								
Most not / certainly not / unsure	455 (90.8%)	442 (93.6%)	441 (93.0%)	438 (93.4%)	416 (92.0%)	430 (94.3%)	443 (94.9%)	0.009
Mostly / certainly	46 (9.2%)	30 (6.4%)	33 (7.0%)	31 (6.6%)	36 (8.0%)	26 (5.7%)	24 (5.1%)	
<b>Avoidance behaviors</b>								
Avoid going to crowded place								
Most not / certainly not / unsure	223 (44.3%)	227 (47.3%)	282 (59.4%)	284 (60.0%)	274 (60.6%)	298 (65.2%)	355 (76.0%)	<0.001
Mostly / Certainly	280 (55.7%)	253 (52.7%)	193 (40.6%)	189 (40.0%)	178 (39.4%)	159 (34.8%)	112 (24.0%)	
Avoid going out unless it is necessary								
Most not / certainly not / unsure	296 (58.8%)	319 (66.3%)	374 (78.7%)	362 (76.5%)	354 (78.3%)	351 (76.8%)	373 (79.9%)	<0.001
Mostly / Certainly	207 (41.2%)	162 (33.7%)	101 (21.3%)	111 (23.5%)	98 (21.7%)	106 (23.2%)	94 (20.1%)	
Avoid traveling abroad								
Most not / certainly not / unsure	262 (52.2%)	286 (59.5%)	330 (69.5%)	309 (65.3%)	320 (70.8%)	337 (73.7%)	373 (79.9%)	<0.001
Mostly / Certainly	240 (47.8%)	195 (40.5%)	145 (30.5%)	164 (34.7%)	132 (29.2%)	120 (26.3%)	94 (20.1%)	
Avoid going to hospitals								
Most not / certainly not / unsure	222 (44.3%)	206 (43.0%)	266 (56.0%)	256 (54.1%)	268 (59.3%)	306 (67.0%)	317 (67.9%)	<0.001
Mostly / Certainly	279 (55.7%)	273 (57.0%)	209 (44.0%)	217 (45.9%)	184 (40.7%)	151 (33.0%)	150 (32.1%)	





	Round 1 (n = 503), 19 Aug – 10 Sep 09	Round 2 (n = 481), 25 Sep – 5 Oct 09	Round 3 (n = 475), 27 Oct – 2 Nov 09	Round 4 (n = 473), 28 Nov – 10 Dec 09	Round 5 (n = 452), 28 Jan – 5 Feb 10	Round 6 (n = 457), 7 – 14 Apr 10	Round 7 (n = 467), 24 Jun – 6 Jul 10	p (trend)
<b>Knowledge and misconceptions</b>								
Could be transmitted via airborne with a long distance (from one building to another one)	162 (32.3%)	147 (30.6%)	151 (31.8%)	108 (22.8%)	107 (23.7%)	90 (19.7%)	107 (22.9%)	<0.001
Could be transmitted via insect bites	96 (19.1%)	81 (16.9%)	69 (14.5%)	79 (16.7%)	93 (20.6%)	86 (18.9%)	122 (26.1%)	<0.001
Could be transmitted via water sources (e.g., reservoirs)	125 (24.9%)	97 (20.2%)	108 (22.7%)	93 (19.7%)	119 (26.3%)	113 (24.8%)	161 (34.5%)	<0.001
Could be transmitted via well-cooked pork	12 (2.4%)	7 (1.5%)	10 (2.1%)	17 (3.6%)	16 (3.5%)	18 (3.9%)	29 (6.2%)	<0.001
Any one of above								
No	251 (49.9%)	258 (53.6%)	239 (50.3%)	293 (61.9%)	232 (51.3%)	256 (56.0%)	224 (48.0%)	0.578
Yes	252 (50.1%)	223 (46.4%)	236 (49.7%)	180 (38.1%)	220 (48.7%)	201 (44.0%)	243 (52.0%)	

**Technical Appendix Table 5.** Perceived severity of influenza A/H1N1

	Round 1 (n = 503), 19 Aug – 10 Sep 09	Round 2 (n = 481), 25 Sep – 5 Oct 09	Round 3 (n = 475), 27 Oct – 2 Nov 09	Round 4 (n = 473), 28 Nov – 10 Dec 09	Round 5 (n = 452), 28 Jan – 5 Feb 10	Round 6 (n = 457), 7 – 14 Apr 10	Round 7 (n = 467), 24 Jun – 6 Jul 10	p (trend)
<b>Perceived severity</b>								
<b>Mortality rate for adults</b>								
<1% / Don't know	345 (68.6%)	291 (60.6%)	278 (58.5%)	254 (53.7%)	237 (52.4%)	300 (65.6%)	351 (75.2%)	<0.001
≥1%	158 (31.4%)	189 (39.4%)	197 (41.5%)	219 (46.3%)	215 (47.6%)	157 (34.4%)	116 (24.8%)	
<b>Would cause severe irreversible body damages</b>								
Disagree / Don't know	418 (83.1%)	399 (83.0%)	388 (81.7%)	364 (77.0%)	330 (73.0%)	350 (76.6%)	350 (74.9%)	<0.001
Agree	85 (16.9%)	82 (17.0%)	87 (18.3%)	109 (23.0%)	122 (27.0%)	107 (23.4%)	117 (25.1%)	
<b>What would you think in the coming year, the effects of H1N1 influenza virus on the infected person would become milder, more harmful, or similar to the current situation?</b>								
Milder / Similar to current situation / the same	410 (85.6%)	390 (84.1%)	410 (88.7%)	358 (78.3%)	401 (89.5%)	428 (95.3%)	440 (94.8%)	<0.001
More harmful	69 (14.4%)	74 (15.9%)	52 (11.3%)	99 (21.7%)	47 (10.5%)	21 (4.7%)	24 (5.2%)	
<b>Perceived severity of H1N1 compared to seasonal flu</b>								
<b>Mortality rate</b>								
Much lower / a little lower / the same	275 (55.7%)	228 (47.8%)	210 (44.5%)	203 (43.3%)	174 (38.8%)	184 (40.7%)	197 (42.7%)	<0.001
A bit higher / much higher	219 (44.3%)	249 (52.2%)	262 (55.5%)	266 (56.7%)	274 (61.2%)	268 (59.3%)	264 (57.3%)	
<b>Infectivity rate</b>								
Much lower / a little lower / the same	315 (63.3%)	316 (66.0%)	291 (61.4%)	277 (58.8%)	308 (68.3%)	274 (60.4%)	311 (66.6%)	0.374
A bit higher / much higher	183 (36.7%)	163 (34.0%)	183 (38.6%)	194 (41.2%)	143 (31.7%)	180 (39.6%)	156 (33.4%)	
<b>Severity of body damages</b>								
Much lower / a little lower / the same	264 (53.8%)	227 (48.0%)	211 (44.6%)	176 (37.4%)	147 (32.9%)	172 (38.0%)	175 (37.6%)	<0.001
A bit higher / much higher	227 (46.2%)	246 (52.0%)	262 (55.4%)	294 (62.6%)	300 (67.1%)	281 (62.0%)	290 (62.4%)	
<b>Perceived severity of H1N1 compared to SARS</b>								
<b>Mortality rate</b>								
Lower / the same / don't know	492 (97.8%)	473 (98.3%)	471 (99.2%)	464 (98.1%)	443 (98.0%)	451 (98.7%)	460 (98.5%)	0.390
Higher	11 (2.2%)	8 (1.7%)	4 (0.8%)	9 (1.9%)	9 (2.0%)	6 (1.3%)	7 (1.5%)	
<b>Infectivity rate</b>								
Lower / the same / don't know	415 (83.0%)	370 (77.2%)	389 (82.1%)	400 (84.6%)	397 (87.8%)	422 (92.3%)	444 (95.1%)	<0.001
Higher	85 (17.0%)	109 (22.8%)	85 (17.9%)	73 (15.4%)	55 (12.2%)	35 (7.7%)	23 (4.9%)	

**Technical Appendix Table 6.** Perceptions related to susceptibility and community outbreak of influenza A/H1N1

	Round 1 (n = 503), 19 Aug – 10 Sep 09	Round 2 (n = 481), 25 Sep – 5 Oct 09	Round 3 (n = 475), 27 Oct – 2 Nov 09	Round 4 (n = 473), 28 Nov – 10 Dec 09	Round 5 (n = 452), 28 Jan – 5 Feb 10	Round 6 (n = 457), 7 – 14 Apr 10	Round 7 (n = 467), 24 Jun – 6 Jul 10	p (trend)
<b>Perceived susceptibility</b>								
Perceived high or very high chance of contracting Flu A (H1N1) in the coming year								
(a) The respondent	77 (16.6%)	116 (25.4%)	78 (17.3%)	66 (14.7%)	52 (11.9%)	34 (7.7%)	12 (2.6%)	<0.001
(b) Family members	69 (14.9%)	114 (25.2%)	75 (16.7%)	68 (15.2%)	48 (11.0%)	34 (7.7%)	12 (2.6%)	<0.001
(c) The general public	90 (20.9%)	133 (31.4%)	79 (20.3%)	73 (17.0%)	58 (14.0%)	34 (8.0%)	12 (2.6%)	<0.001
Feel confident that you and your family are and will be free from infection								
Unlikely / certainly not / unsure	126 (25.0%)	137 (28.6%)	117 (24.7%)	97 (20.6%)	111 (24.6%)	68 (14.9%)	36 (7.7%)	<0.001
Certainly / likely	377 (75.0%)	342 (71.4%)	357 (75.3%)	373 (79.4%)	341 (75.4%)	388 (85.1%)	430 (92.3%)	
<b>Perception related to community outbreak of H1N1</b>								
Perceived chance of having a large-scale local A/H1N1 outbreak in Hong Kong in the coming year								
Unlikely / most unlikely / certainly not / unsure	295 (58.6%)	247 (51.4%)	311 (65.6%)	404 (85.4%)	401 (88.7%)	429 (94.3%)	451 (96.6%)	<0.001
Certainly / most likely / likely	208 (41.4%)	234 (48.6%)	163 (34.4%)	69 (14.6%)	51 (11.3%)	26 (5.7%)	16 (3.4%)	
Perceived duration of H1N1 flu A epidemic								
≤12 mo	321 (67.6%)	296 (63.2%)	293 (63.7%)	266 (57.8%)	270 (61.8%)	265 (59.6%)	347 (76.9%)	0.002
>12 mo	154 (32.4%)	172 (36.8%)	167 (36.3%)	194 (42.2%)	167 (38.2%)	180 (40.4%)	104 (23.1%)	
Perceived number of death cases in the coming year								
≤50	432 (90.2%)	379 (82.8%)	363 (78.9%)	401 (86.1%)	381 (86.8%)	428 (95.1%)	458 (99.1%)	<0.001
>50	47 (9.8%)	79 (17.2%)	97 (21.1%)	65 (13.9%)	58 (13.2%)	22 (4.9%)	4 (0.9%)	
Perceived number of hidden cases in the community								
Very few / a few	217 (44.7%)	150 (32.3%)	202 (43.8%)	226 (49.1%)	205 (47.1%)	215 (49.4%)	337 (73.6%)	<0.001
Quite a lot / many	268 (55.3%)	315 (67.7%)	259 (56.2%)	234 (50.9%)	230 (52.9%)	220 (50.6%)	121 (26.4%)	

**Technical Appendix Table 7.** Perceived efficacy of measures for preventing influenza A/H1N1

	Round 1 (n = 503), 19 Aug – 10 Sep 09	Round 2 (n = 481), 25 Sep – 5 Oct 09	Round 3 (n = 475), 27 Oct – 2 Nov 09	Round 4 (n = 473), 28 Nov – 10 Dec 09	Round 5 (n = 452), 28 Jan – 5 Feb 10	Round 6 (n = 457), 7 – 14 Apr 10	Round 7 (n = 467), 24 Jun – 6 Jul 10	p (trend)
<b>Perceived efficacy</b>								
Wearing face masks in public areas								
Not effective at all / Not very effective	47 (9.4%)	31 (6.4%)	49 (10.3%)	41 (8.7%)	39 (8.6%)	39 (8.5%)	41 (8.8%)	0.940
Quite effective / Very effective	454 (90.6%)	450 (93.6%)	426 (89.7%)	432 (91.3%)	413 (91.4%)	418 (91.5%)	426 (91.2%)	
Washing hands frequently								
Not effective at all / Not very effective	3 (0.6%)	4 (0.8%)	3 (0.6%)	6 (1.3%)	7 (1.5%)	6 (1.3%)	7 (1.5%)	0.071
Quite effective / Very effective	499 (99.4%)	477 (99.2%)	472 (99.4%)	467 (98.7%)	445 (98.5%)	451 (98.7%)	460 (98.5%)	
Injection of seasonal flu vaccine								
Not effective at all / Not very effective	268 (55.0%)	266 (57.2%)	264 (57.5%)	267 (58.4%)	286 (64.9%)	286 (63.8%)	290 (63.5%)	<0.001
Quite effective / Very effective	219 (45.0%)	199 (42.8%)	195 (42.5%)	190 (41.6%)	155 (35.1%)	162 (36.2%)	167 (36.5%)	
Injection of flu A (H1N1) vaccine								
Not effective at all / Not very effective	50 (10.7%)	44 (9.9%)	32 (7.4%)	49 (11.3%)	72 (16.9%)	68 (15.7%)	58 (12.8%)	0.007
Quite effective / Very effective	416 (89.3%)	401 (90.1%)	399 (92.6%)	385 (88.7%)	354 (83.1%)	365 (84.3%)	395 (87.2%)	
Taking traditional Chinese medicine								
Not effective at all / Not very effective	186 (41.9%)	172 (42.8%)	166 (42.6%)	178 (44.0%)	146 (37.5%)	175 (42.2%)	157 (38.1%)	0.253
Quite effective / Very effective	258 (58.1%)	230 (57.2%)	224 (57.4%)	227 (56.0%)	243 (62.5%)	240 (57.8%)	255 (61.9%)	
Avoid going to crowded places								
Not effective at all / Not very effective	24 (4.8%)	18 (3.7%)	20 (4.2%)	30 (6.3%)	29 (6.4%)	26 (5.7%)	22 (4.7%)	0.523
Quite effective / Very effective	479 (95.2%)	463 (96.3%)	455 (95.8%)	443 (93.7%)	423 (93.6%)	431 (94.3%)	445 (95.3%)	

**Technical Appendix Table 8.** Evaluation of government preparedness and performance toward influenza A/H1N1

Evaluation	Round 1 (n = 503), 19 Aug – 10 Sep 09	Round 2 (n = 481), 25 Sep – 5 Oct 09	Round 3 (n = 475), 27 Oct – 2 Nov 09	Round 4 (n = 473), 28 Nov – 10 Dec 09	Round 5 (n = 452), 28 Jan – 5 Feb 10	Round 6 (n = 457), 7 – 14 Apr 10	Round 7 (n = 467), 24 Jun – 6 Jul 10	p (trend)
Evaluation on government performance								
Timeliness of prevention measures	6.1 (1.6)	6.2 (1.4)	6.2 (1.4)	6.2 (1.4)	6.0 (1.4)	6.3 (1.4)	6.2 (1.5)	
Effectiveness of prevention measures	6.0 (1.7)	6.1 (1.4)	6.2 (1.3)	6.2 (1.4)	5.9 (1.5)	6.1 (1.5)	6.2 (1.6)	
Explaining clearly to general public	6.2 (1.7)	6.3 (1.5)	6.2 (1.4)	6.2 (1.5)	5.9 (1.6)	6.1 (1.6)	6.1 (1.6)	
Adequacy of quarantine and disinfection procedure	6.2 (1.7)	6.3 (1.4)	6.2 (1.3)	6.2 (1.4)	6.1 (1.5)	6.2 (1.5)	6.1 (1.7)	
Collaboration between governmental departments	5.6 (1.8)	5.8 (1.6)	5.7 (1.4)	5.8 (1.5)	5.6 (1.5)	5.7 (1.7)	5.7 (1.6)	
General evaluation	6.0 (1.6)	6.2 (1.3)	6.2 (1.3)	6.2 (1.3)	6.0 (1.4)	6.2 (1.5)	6.3 (1.5)	
Average score	6.0 (1.5)	6.2 (1.3)	6.1 (1.2)	6.1 (1.3)	5.9 (1.4)	6.1 (1.4)	6.1 (1.5)	
Average score <5	100 (20.7%)	62 (13.1%)	55 (11.9%)	65 (14.0%)	91 (20.6%)	75 (16.7%)	88 (19.0%)	0.101
The HK Government is overreacting with its current preventive measures								
Not overreacting at all / Not overreacting / Just right	469 (93.4%)	451 (94.2%)	460 (97.0%)	462 (98.1%)	409 (90.7%)	425 (93.0%)	430 (92.1%)	0.143
Overreacting somewhat / extremely overreacting	33 (6.6%)	28 (5.8%)	14 (3.0%)	9 (1.9%)	42 (9.3%)	32 (7.0%)	37 (7.9%)	
Evaluation on government preparedness								
Hong Kong will not have enough vaccine for human swine flu								
Disagree / Don't know	322 (64.0%)	333 (69.2%)	351 (73.9%)	378 (79.9%)	377 (83.4%)	381 (83.4%)	445 (95.3%)	<0.001
Agree	181 (36.0%)	148 (30.8%)	124 (26.1%)	95 (20.1%)	75 (16.6%)	76 (16.6%)	22 (4.7%)	
Hong Kong will not have enough medication for human swine flu								
Disagree / Don't know	384 (76.5%)	397 (82.5%)	384 (80.8%)	395 (83.5%)	371 (82.1%)	377 (82.5%)	450 (96.4%)	<0.001
Agree	118 (23.5%)	84 (17.5%)	91 (19.2%)	78 (16.5%)	81 (17.9%)	80 (17.5%)	17 (3.6%)	
HK will be able to control the Flu A (H1N1) epidemic								
Disagree / Don't know	115 (22.9%)	127 (26.4%)	85 (17.9%)	76 (16.1%)	61 (13.5%)	45 (9.8%)	24 (5.1%)	<0.001
Agree	388 (77.1%)	354 (73.6%)	390 (82.1%)	397 (83.9%)	391 (86.5%)	412 (90.2%)	443 (94.9%)	

**Technical Appendix Table 9.** Perceptions on policy-related issues for influenza A/H1N1

Perception	Round 1 (n = 503), 19 Aug – 10 Sep 09	Round 2 (n = 481), 25 Sep – 5 Oct 09	Round 3 (n = 475), 27 Oct – 2 Nov 09	Round 4 (n = 473), 28 Nov – 10 Dec 09	Round 5 (n = 452), 28 Jan – 5 Feb 10	Round 6 (n = 457), 7 – 14 Apr 10	Round 7 (n = 467), 24 Jun – 6 Jul 10	p (trend)
It is not necessary to have class suspension for kindergartens/ primary schools, even there are infected cases found in the schools								
Disagree / Don't know	–	399 (83.0%)	338 (71.2%)	332 (70.2%)	282 (62.4%)	312 (68.3%)	245 (52.5%)	<0.001
Agree	–	82 (17.0%)	137 (28.8%)	141 (29.8%)	170 (37.6%)	145 (31.7%)	222 (47.5%)	
It is not necessary to have class suspension for secondary schools, even there are infected cases found in the schools								
Disagree / Don't know	–	301 (62.6%)	205 (43.2%)	217 (45.9%)	166 (36.7%)	166 (36.3%)	108 (23.1%)	<0.001
Agree	–	180 (37.4%)	270 (56.8%)	256 (54.1%)	286 (63.3%)	291 (63.7%)	359 (76.9%)	
The Hong Kong government should cancel all pandemic control measures for H1N1 influenza A								
Disagree / Don't know	–	464 (96.7%)	461 (97.1%)	453 (95.8%)	437 (96.7%)	436 (95.4%)	394 (84.4%)	<0.001
Agree	–	16 (3.3%)	14 (2.9%)	20 (4.2%)	15 (3.3%)	21 (4.6%)	73 (15.6%)	
The Hong Kong government should suspend all pandemic control measures for H1N1 influenza A								
Disagree / Don't know	–	–	–	–	330 (73.0%)	376 (82.3%)	263 (56.3%)	<0.001
Agree	–	–	–	–	122 (27.0%)	81 (17.7%)	204 (43.7%)	
The government should treat human swine flu in the same manner as the seasonal flu								
Disagree / Unsure	236 (46.9%)	249 (52.0%)	242 (50.9%)	265 (56.0%)	276 (61.1%)	220 (48.1%)	213 (45.6%)	0.225
Agree	267 (53.1%)	230 (48.0%)	233 (49.1%)	208 (44.0%)	176 (38.9%)	237 (51.9%)	254 (54.4%)	
Do you think you understand clearly the preventive measures and treatment strategies/policies advocated by the HK Government								
Not clear at all / not quite clear	205 (40.8%)	190 (39.6%)	205 (43.2%)	215 (45.5%)	159 (35.2%)	214 (47.0%)	242 (51.8%)	<0.001
Quite clear / very clear	298 (59.2%)	290 (60.4%)	270 (56.8%)	258 (54.5%)	293 (64.8%)	241 (53.0%)	225 (48.2%)	