

Carbapenemase-Producing *Enterobacteriaceae* and Nonfermentative Bacteria, the Philippines, 2013–2016

John Mark Velasco, Maria Theresa Valderama, Trent Peacock,¹ Nirdnoy Warawadee, Kathyleen Nogrado, Fatima Claire Navarro, Domingo Chua, Jr., Srijan Apichai, Ruekit Sirigade, Louis R. Macareo, Brett Swierczewski

Author affiliations: US Army Medical Directorate—Armed Forces Research Institute of Medical Sciences, Bangkok, Thailand (J.M. Velasco, M.T. Valderama, T. Peacock, N. Warawadee, K. Nogrado, S. Apichai, R. Sirigade, L.R. Macareo, B. Swierczewski); V. Luna Medical Center, Quezon City, the Philippines (F.C. Navarro, D. Chua, Jr.)

DOI: <https://doi.org/10.3201/eid2309.161237>

During 2013–2016, we isolated *bla*_{NDM}- and *bla*_{VIM}-harboring *Enterobacteriaceae* and nonfermentative bacteria from patients in the Philippines. Of 130 carbapenem-resistant isolates tested, 45 were Carba NP–positive; 43 harbored *bla*_{NDM}, and 2 harbored *bla*_{VIM}. Multidrug-resistant microbial pathogen surveillance and antimicrobial drug stewardship are needed to prevent further spread of New Delhi metallo-β-lactamase variants.

Carbapenemase-producing *Enterobacteriaceae* can efficiently hydrolyze carbapenems and most β-lactam drugs. Since the identification of New Delhi metallo-β-lactamase-1 (NDM-1) in 2008 (1), there has been great concern regarding the spread of the Ambler class B metallo-β-lactamases (MBLs). Confirmed infections with MBL-positive bacteria are rarely identified in the Philippines, but *bla*_{IMP}-harboring *Enterobacteriaceae* were reported in 2014 (2), an *Escherichia coli* (sequence type [ST] 131) isolate harboring *bla*_{NDM-1} was reported in 2014 (3), and 2 *Klebsiella pneumoniae* (ST626 and ST903) isolates harboring *bla*_{NDM-1} and *bla*_{NDM-7} genes were reported in 2016 (4).

We performed isolate identification and antimicrobial drug susceptibility testing by using the MicroScan Walk-Away 40 plus System (Beckman Coulter, Brea, CA, USA) on 1,516 gram-positive and gram-negative isolates from patients admitted to various wards in the V. Luna Medical Center, a tertiary-care military hospital in Manila, the Philippines, during August 2013–April 2016. To better

assess the distribution of carbapenem resistance and the underlying molecular mechanisms of resistance, we selected gram-negative isolates with imipenem or meropenem (or both) MICs of ≥8 μg/mL. We used microbroth dilution susceptibility testing (5) to select and verify 130 gram-negative nonrepeat isolates (i.e., each isolate was tested once) and then tested the isolates for carbapenemase production by using the Carba NP test as previously described (6). We tested all isolates with a Carba NP–positive result for *bla*_{NDM} and *bla*_{KPC} by using a multiplex real-time PCR assay as previously described (7,8); isolates with PCR-negative results were further tested, using the Xpert Carba-R PCR test with the GeneXpert IV System (both from Cepheid, Sunnyvale, CA, USA), for the presence of *bla*_{NDM}, *bla*_{KPC}, *bla*_{VIM}, *bla*_{IMP-1}, and *bla*_{OXA-48}.

Of the 130 bacterial isolates tested, 45 (35%) had positive Carba NP test results and 43 (33%) harbored *bla*_{NDM}; 25 (58%) of the *bla*_{NDM}-carrying isolates were identified as *K. pneumoniae* (online Technical Appendix Table, <https://wwwnc.cdc.gov/EID/article/23/9/16-1237-Techapp1.pdf>). None of the isolates was positive for *bla*_{KPC}. Two *Pseudomonas aeruginosa* isolates that had positive Carba NP test results were negative for *bla*_{NDM} and *bla*_{KPC} but positive for *bla*_{VIM}. During the collection period, we also tested 8 environmental samples collected from the hospital's neonatal intensive care unit and obstetrics and gynecology wards; 3 (38%) of the 8 isolates were positive for *bla*_{NDM} and identified as *K. pneumoniae* (online Technical Appendix Table).

We report the identification of *bla*_{NDM}-positive bacterial isolates in several genera of *Enterobacteriaceae* and nonfermentative bacteria in the Philippines. This finding is particularly significant because NDM-like enzymes have a broad range of activity against most β-lactam antimicrobial drugs and are often associated with serious clinical infections (9). A higher risk for plasmid-mediated transfer of NDM-1 exists through conjugation between different gram-negative bacterial strains (10), and NDM-1 can spread rapidly via nosocomial transmission or community-acquired infection. Furthermore, although limited in number, the environmental samples in this study were also positive for *bla*_{NDM}, which suggests the possibility of nosocomial transmission and local circulation.

We conducted multiplex real-time PCR testing only for *bla*_{NDM}, *bla*_{KPC}, *bla*_{VIM}, *bla*_{IMP-1}, and *bla*_{OXA-48} and did not investigate clonality; thus, further investigation into other carbapenemase genes should be conducted. In addition, further experiments should be performed to characterize the plasmids carrying the carbapenemase genes. Strengthening of multidrug-resistant microbial pathogen surveillance and antimicrobial drug stewardship is urgently needed to better characterize drug-resistance patterns and improve early detection and containment strategies in developing countries.

¹Current affiliation: US Army Institute of Surgical Research, Fort Sam Houston, Texas, USA.

Acknowledgments

We thank Bryony Soltis and the Bacteriology Section and Department of Research and Training of V. Luna Medical Center, Armed Forces of the Philippines Health Service Command, for their support.

This study was funded by a grant from the Armed Forces Health Surveillance Center–Global Emerging Infections Surveillance and Response System.

The views expressed in this article are those of the authors and do not reflect the official policy of the Department of the Army, the Department of Defense, or the US Government.

Dr. Velasco is the regional clinical investigator of the Department of Virology, Armed Forces Research Institute of Medical Sciences. His primary research interests include detection and control of emerging pathogens, epidemiology of respiratory and arthropodborne viruses, and microbial typing of multidrug-resistant bacteria.

References

1. Yong D, Toleman MA, Giske CG, Cho HS, Sundman K, Lee K, et al. Characterization of a new metallo- β -lactamase gene, *bla*_{NDM-17} and a novel erythromycin esterase gene carried on a unique genetic structure in *Klebsiella pneumoniae* sequence type 14 from India. *Antimicrob Agents Chemother*. 2009;53:5046–54. <http://dx.doi.org/10.1128/AAC.00774-09>
2. Peirano G, Lascols C, Hackel M, Hoban DJ, Pitout JD. Molecular epidemiology of *Enterobacteriaceae* that produce VIMs and IMPs from the SMART surveillance program. *Diagn Microbiol Infect Dis*. 2014;78:277–81. <http://dx.doi.org/10.1016/j.diagmicrobio.2013.11.024>
3. Peirano G, Bradford PA, Kazmierczak KM, Badal RE, Hackel M, Hoban DJ, et al. Global incidence of carbapenemase-producing *Escherichia coli* ST131. *Emerg Infect Dis*. 2014;20:1928–31. <http://dx.doi.org/10.3201/eid2011.141388>
4. Chou A, Roa M, Evangelista MA, Sulit AK, Lagamayo E, Torres BC, et al. Emergence of *Klebsiella pneumoniae* ST273 carrying *bla*_{NDM-7} and ST656 carrying *bla*_{NDM-1} in Manila, Philippines. *Microb Drug Resist*. 2016;22:585–8. <http://dx.doi.org/10.1089/mdr.2015.0205>
5. Clinical and Laboratory Standards Institute. Performance standards for antimicrobial susceptibility testing: twenty-fifth informational supplement (M100–S25). Wayne (PA): The Institute, 2015.
6. Nordmann P, Poirel L, Dortet L. Rapid detection of carbapenemase-producing *Enterobacteriaceae*. *Emerg Infect Dis*. 2012;18:1503–7. <http://dx.doi.org/10.3201/eid1809.120355>
7. Milillo M, Kwak YI, Snesrud E, Waterman PE, Lesho E, McGann P. Rapid and simultaneous detection of *bla*_{KPC} and *bla*_{NDM} by use of multiplex real-time PCR. *J Clin Microbiol*. 2013;51:1247–9. <http://dx.doi.org/10.1128/JCM.03316-12>
8. Milillo M, Kwak YI, Snesrud E, Waterman PE, Lesho E, McGann P. Correction for Milillo et al., Rapid and simultaneous detection of blaKPC and blaNDM by use of multiplex real-time PCR. *J Clin Microbiol*. 2015;53:1460. <http://dx.doi.org/10.1128/JCM.00373-15>
9. Schwaber MJ, Klarfeld-Lidji S, Navon-Venezia S, Schwartz D, Leavitt A, Carmeli Y. Predictors of carbapenem-resistant *Klebsiella pneumoniae* acquisition among hospitalized adults and effect of acquisition on mortality. *Antimicrob Agents Chemother*. 2008;52:1028–33. <http://dx.doi.org/10.1128/AAC.01020-07>
10. Kulkova N, Babalova M, Sokolova J, Kremery V. First report of New Delhi metallo- β -lactamase-1–producing strains in Slovakia. *Microb Drug Resist*. 2015;21:117–20. <http://dx.doi.org/10.1089/mdr.2013.0162>

Address for correspondence: John Mark Velasco, Department of Virology, Armed Forces Research Institute of Medical Sciences, 315/6 Rajvithi Rd, Bangkok 10400, Thailand; email: John.Velasco.ca@afirms.org

Chronic Wasting Disease Prion Strain Emergence and Host Range Expansion

Allen Herbst,¹ Camilo Duque Velásquez,¹ Elizabeth Triscott, Judd M. Aiken, Debbie McKenzie

Author affiliation: University of Alberta, Edmonton, Alberta, Canada

DOI: <https://doi.org/10.3201/eid2309.161474>

Human and mouse prion proteins share a structural motif that regulates resistance to common chronic wasting disease (CWD) prion strains. Successful transmission of an emergent strain of CWD prion, H95⁺, into mice resulted in infection. Thus, emergent CWD prion strains may have higher zoonotic potential than common strains.

Chronic wasting disease (CWD) is a contagious prion disease of cervids that is spreading globally. CWD is enzootic in multiple cervid species, including deer and elk; the major foci of disease are Colorado/Wyoming (USA), Wisconsin/Illinois (USA), and Alberta/Saskatchewan (Canada). CWD is also present in captive cervids in South Korea and wild reindeer and moose in Norway (https://www.nwhc.usgs.gov/images/cwd/cwd_map.jpg). CWD results from the conformational transformation of the host-encoded cellular prion protein (PrP^C) into protease-resistant, detergent-insoluble, β -sheet rich, amyloidogenic conformers, termed prions (PrP^{CWD}). Within their conformation, prion strains encipher the information that directs the templated misfolding and aggregation of PrP^C molecules into additional prions (1).

Although the sequence homology of PrP among mammals is high, the ability of particular prion strains to cause disease in different species is determined by the conformational compatibility between a given strain and the host PrP^C (2). We

¹These authors contributed equally to this article.

Carbapenemase-Producing Enterobacteriaceae and Nonfermentative Bacteria, the Philippines, 2013–2016

Technical Appendix

Technical Appendix Table. Molecular resistance mechanisms of carbapenem-resistant clinical (n = 45) and environmental isolates (n = 3) from a tertiary-care military hospital in Manila, the Philippines, August 2013–April 2016*

Identification	Source	Organism	Carba NP	<i>bla_{NDM}</i>	<i>bla_{KPC}</i>	<i>bla_{VIM}</i>	Month	Year	Imipenem MIC (µg/mL)	Meropenem MIC (µg/mL)	Sex	Hospital ward
Patient isolates												
PH-0138–14	Blood	<i>K. pneumoniae</i>	Pos	Pos	Neg	ND	Nov	2013	>8	>8	M	Neonatal ICU
PH-0542–14	Soft tissue	<i>E. coli</i>	Pos	Pos	Neg	ND	Jun	2014	>8	>8	M	Pediatric
PH-0630–14	Wound	<i>K. pneumoniae</i>	Pos	Pos	Neg	ND	Jul	2014	>8	>8	F	Female Medical
PH-0631–14	Blood	<i>C. freundii</i>	Pos	Pos	Neg	ND	Jul	2014	>8	>8	M	Neurosurgery
PH-0756–14	Catheter	<i>K. pneumoniae</i>	Pos	Pos	Neg	ND	Aug	2014	>8	>8	F	Medical ICU
PH-0787–14	Endotracheal tip	<i>K. pneumoniae</i>	Pos	Pos	Neg	ND	Aug	2014	>8	>8	M	Medical ICU
PH-0837–14	Catheter	<i>K. pneumoniae</i>	Pos	Pos	Neg	ND	Sep	2014	>8	>8	F	Female Surgical
PH-0846–14	Catheter	<i>E. cloacae</i>	Pos	Pos	Neg	ND	Sep	2014	>8	>8	M	Male Medical Oncology
PH-0850–14	Urine	<i>K. pneumoniae</i>	Pos	Pos	Neg	ND	Sep	2014	>8	>8	F	Female Medical
PH-0873–14	Urine	<i>K. pneumoniae</i>	Pos	Pos	Neg	ND	Oct	2014	>8	>8	M	Surgical ICU
PH-0874–14	Wound	<i>Klebsiella sp.</i>	Pos	Pos	Neg	ND	Oct	2014	>8	>8	M	Female Medical
PH-0901–14	Wound	<i>C. freundii</i>	Pos	Pos	Neg	ND	Oct	2014	>8	>8	F	Female Surgical
PH-1037–14	Catheter	<i>K. pneumoniae</i>	Pos	Pos	Neg	ND	Dec	2014	>8	>8	F	Female Medical
PH-1076–14	Urine	<i>K. pneumoniae</i>	Pos	Pos	Neg	ND	Dec	2014	>8	>8	M	Medical ICU
PH-1078–14	Blood	<i>K. pneumoniae</i>	Pos	Pos	Neg	ND	Dec	2014	>8	>8	F	Female Medical
PH-1088–14	Endotracheal tip	<i>K. pneumoniae</i>	Pos	Pos	Neg	ND	Jan	2015	>8	>8	F	Medical ICU
PH-1093–14	Tracheal aspirate	<i>K. pneumoniae</i>	Pos	Pos	Neg	ND	Dec	2014	>8	>8	F	Female Medical
PH-1099–14	Endotracheal tip	<i>Acinetobacter sp.</i>	Pos	Pos	Neg	ND	Dec	2014	*	>8	M	Surgical ICU
PH-1115–14	Blood	<i>K. pneumoniae</i>	Pos	Pos	Neg	ND	Dec	2014	>8	>8	F	Female Medical
PH-1142–15	Blood	<i>K. pneumoniae</i>	Pos	Pos	Neg	ND	Jan	2015	>8	>8	F	Female Medical
PH-1143–15	Wound	<i>K. pneumoniae</i>	Pos	Pos	Neg	ND	Jan	2015	>8	>8	M	Neurology
PH-1150–15	Catheter	<i>E. cloacae</i>	Pos	Pos	Neg	ND	Jan	2015	>8	>8	M	Neurosurgery
PH-1159–15	Catheter	<i>K. pneumoniae</i>	Pos	Pos	Neg	ND	Jan	2015	>8	>8	M	Neurology
PH-1165–15	Urine	<i>K. pneumoniae</i>	Pos	Pos	Neg	ND	Jan	2015	>8	>8	M	Pulmonary Disease
PH-1166–15	Urine	<i>K. pneumoniae</i>	Pos	Pos	Neg	ND	Feb	2015	>8	>8	M	Surgical ICU
PH-1261–15	Blood	<i>C. freundii</i>	Pos	Pos	Neg	ND	Jul	2015	8	>8	M	Nephrology
PH-1263–15	Blood	<i>E. cloacae</i>	Pos	Pos	Neg	ND	Jul	2015	>8	>8	F	Neonatal ICU
PH-1265–15	Blood	<i>E. cloacae</i>	Pos	Pos	Neg	ND	Jul	2015	8	8	F	Neonatal ICU
PH-1266–15	Blood	<i>E. cloacae</i>	Pos	Pos	Neg	ND	Jul	2015	>8	>8	F	Neonatal ICU
PH-1270–15	Blood	<i>K. pneumoniae</i>	Pos	Pos	Neg	ND	Jan	2015	>8	>8	F	Female Medical

Identification	Source	Organism	Carba				Month	Year	Imipenem		Meropenem		Sex	Hospital ward
			NP	<i>bla_{NDM}</i>	<i>bla_{KPC}</i>	<i>bla_{VIM}</i>			MIC (µg/mL)	MIC (µg/mL)				
PH-1279-15	Wound	<i>K. oxytoca</i>	Pos	Pos	Neg	ND	Jan	2015	8	>8		M	Neurology	
PH-1280-15	Urine	<i>C. freundii</i>	Pos	Pos	Neg	ND	Jan	2015	>8	>8		M	Neurology	
PH-1363-15	Wound	<i>P. aeruginosa</i>	Pos	Pos	Neg	ND	Sep	2015	>8	>8		M	Surgical ICU	
PH-1379-15	Blood	<i>K. pneumoniae</i>	Pos	Pos	Neg	ND	Sep	2015	>8	>8		M	Medical ICU	
PH-1384-15	Blood	<i>K. pneumoniae</i>	Pos	Pos	Neg	ND	Oct	2015	>8	>8		F	Medical ICU	
PH-1394-15	Wound	<i>K. pneumoniae</i>	Pos	Pos	Neg	ND	Oct	2015	>8	>8		M	Surgical ICU	
PH-1419-15	Urine	<i>K. pneumoniae</i>	Pos	Pos	Neg	ND	Oct	2015	>8	>8		F	Female Medical	
PH-1477-15	Wound	<i>E. cloacae</i>	Pos	Pos	Neg	ND	Oct	2015	>8	>8		F	Medical ICU	
PH-1478-15	Wound	<i>E. cloacae</i>	Pos	Pos	Neg	ND	Oct	2015	>8	>8		F	Female Medical	
PH-1482-15	Wound	<i>K. pneumoniae</i>	Pos	Pos	Neg	ND	Oct	2015	>8	>8		F	Medical ICU	
PH-1499-15	Wound	<i>E. cloacae</i>	Pos	Pos	Neg	ND	Oct	2015	>8	>8		F	Medical ICU	
PH-1595-16	Blood	<i>E. cloacae</i>	Pos	Pos	Neg	ND	Jan	2016	>8	>8		F	Female Medical	
PH-1641-16	Blood	<i>K. pneumoniae</i>	Pos	Pos	Neg	ND	Feb	2016	>8	>8		M	Neurology	
PH-0745-14	Wound	<i>P. aeruginosa</i>	Pos	Neg	Neg	Pos	Aug	2014	8	>8		M	Male Surgical	
PH-0905-14	Soft tissue	<i>P. aeruginosa</i>	Pos	Neg	Neg	Pos	Oct	2014	>8	>8		M	Post-anesthesia Care Unit	
ATCC strains														
ATCC 2473		<i>K. pneumoniae</i>	Pos	Pos	Neg	ND	Apr	2016						
ATCC 1705		<i>K. pneumoniae</i>	Pos	Neg	Pos	ND	Apr	2016						
ATCC 1706		<i>K. pneumoniae</i>	Neg	Neg	Neg	ND	Apr	2016						
Environmental isolates <i>bla_{NDM}</i> positive (n = 3)														
E1	Laryngoscope	<i>K. pneumoniae</i>	Neg	Pos	Neg	ND	Nov	2014	>8	>8			Neonatal ICU	
E3	Incubator	<i>K. pneumoniae</i>	Neg	Pos	Neg	ND	Nov	2014	>8	>8			Neonatal ICU	
E7	Suction 1	<i>K. pneumoniae</i>	Neg	Pos	Neg	ND	Nov	2014	>8	>8			Neonatal ICU	

*ICU, intensive care unit; ND, not done; Neg, negative; Pos, positive.