

Multidrug-Resistant *Salmonella enterica* 4,[5],12:i:- Sequence Type 34, New South Wales, Australia, 2016–2017

Alicia Arnott, Qinning Wang, Nathan Bachmann, Rosemarie Sadsad, Chayanika Biswas, Cristina Sotomayor, Peter Howard, Rebecca Rockett, Agnieszka Wiklendt, Jon R. Iredell, Vitali Sintchenko

Multidrug- and colistin-resistant *Salmonella enterica* serotype 4,[5],12:i:- sequence type 34 is present in Europe and Asia. Using genomic surveillance, we determined that this sequence type is also endemic to Australia. Our findings highlight the public health benefits of genome sequencing-guided surveillance for monitoring the spread of multidrug-resistant mobile genes and isolates.

Since the 1990s, the global incidence of infection with *Salmonella enterica* serotype 4,[5],12:i:- has increased sharply among humans, livestock, and poultry (1). This monophasic variant of *S. enterica* serovar Typhimurium ranges from pansusceptible to multidrug resistant. In 2015, an *S. enterica* strain displaying the plasmid-mediated colistin resistance *mcr-1* gene was discovered (2). In 2016, human and food isolates with *mcr-1* were identified in Portugal (3), China (4), and the United Kingdom (5). All *mcr-1*-harboring isolates were predominantly *Salmonella* 4,[5],12:i:- multilocus sequence typing (MLST) sequence type (ST) 34. Before this study, the ST34 clone, already emerged in Europe and Asia, was yet to be detected in Australia as a drug-resistant pathogen of humans. We therefore investigated the circulation of drug-resistant *Salmonella* 4,[5],12:i:- ST34 in New South Wales (NSW), Australia.

The Study

Since October 2016, all *Salmonella* isolates referred to the NSW Enteric Reference Laboratory (Centre for Infectious Diseases and Microbiology Laboratory Services, Pathology

Author affiliations: Westmead Hospital, Sydney, New South Wales, Australia (A. Arnott, R. Sadsad, C. Biswas, C. Sotomayor, R. Rockett, J.R. Iredell, V. Sintchenko); The University of Sydney, Sydney (A. Arnott, R. Sadsad, C. Sotomayor, R. Rockett, J.R. Iredell, V. Sintchenko); NSW Health Pathology, Sydney (Q. Wang, P. Howard, A. Wiklendt, V. Sintchenko); Centenary Institute, Sydney (N. Bachmann)

DOI: <https://doi.org/10.3201/eid2404.171619>

West, Sydney, NSW, Australia) have undergone whole-genome sequencing in addition to serotyping and multi-locus variable-number tandem-repeat analysis (MLVA) performed as described (6). Of the 971 isolates (96% from humans, 4% from food and animals) received from October 1, 2016, through March 17, 2017, a total of 80 (8.2%) were identified as *Salmonella* 4,[5],12:i:-, and 61 (76%) of these underwent whole-genome sequencing. Five duplicate isolates were excluded. In our retrospective study, we included 54 isolates from humans and 2 isolates from pork meat obtained from independent butchers during a routine survey conducted by the NSW Food Authority in 2016.

We extracted genomic DNA by using the chemagic Prepito-D (Perkin Elmer, Seer Green, UK) and prepared libraries by using Nextera XT kits and sequenced them on a NextSeq-500 (both by Illumina, San Diego, CA, USA) with at least 30-fold coverage. We assessed genomic similarity and STs by using the Nullarbor pipeline (7). We identified antimicrobial resistance (AMR) genes by screening contigs through ResFinder (8) and CARD (<https://card.mcmaster.ca>) by using ABRicate version 0.5 (<https://github.com/tseemann/abricate>). Markers of colistin resistance were examined by using CLC Genomics Workbench (QIAGEN, Valencia, CA, USA). We identified *Salmonella* 4,[5],12:i:- genomes recovered in Europe and Asia by using Enterobase (<https://enterobase.warwick.ac.uk/>). We confirmed phenotypic resistance on a randomly selected subset of isolates by using the BD Phoenix system (Becton Dickinson, Franklin Lakes, NJ, USA) or Etest (bioMérieux, Marcy L'Étoile, France).

We obtained 54 isolates from 53 case-patients who had a median age of 25 years (range <1 to 90 years). We detected 20 MLVA profiles; however, 2 profiles predominated: 3-13-10-NA-0211 (45%) and 3-13-11-NA-0211 (14%). All but 2 case-patients resided in areas of distinct postal codes distributed throughout NSW; we found no apparent temporal or geographic clustering. Recent overseas travel was reported by 5 case-patients: 2 to Cambodia and 1 each to Thailand, Vietnam, and Indonesia.

All 56 *Salmonella* 4,[5],12:i:- isolates were classified as ST34. The diversity between isolates was higher than that suggested by MLVA; we detected up to 112 single-nucleotide polymorphism (SNP) differences between isolates. The isolates from Australia clustered with each other and with isolates from the United Kingdom (Figure).

Combined with the steady monthly incidence of infections, these findings suggest that local circulation of *Salmonella* 4,[5],12:i:- might play a larger role as the source of infection than independent importations from overseas. Of note, 1 isolate from pork differed from 1 isolate from a human by only 10 SNPs, indicating that pork may be a source of human infection (Figure, panel A).

We detected AMR genes in 95% of ST34 isolates from NSW. The number of AMR genes (up to 13) was equivalent to that reported for ST34 isolates from the United States and United Kingdom (Figure, panel B). Of the 53 AMR isolates from NSW, 48 (90%) were classified as multidrug resistant on the basis of containing >4 AMR genes conferring resistance to different classes of antimicrobial drugs. Among the AMR isolates, 39 (73.5%) displayed multidrug resistance

patterns, all of which are associated with resistance to aminoglycosides, β -lactams, and sulfonamides. A total of 21 (40%) isolates, including 1 from pork, had the core resistance-type (R-type) ASSuT (resistant to ampicillin, streptomycin, sulfonamides, and tetracycline) conferred by the *strA-strB*, *blaTEM-1b*, *sul2*, and *tet(B)* genes (Figure, panel B). This multidrug resistance pattern is characteristic of the European clone (9), which has been reported in Europe and North America and is strongly associated with pork (10,11).

R-type ASSuTmK was found for 12 (23%) isolates from humans: genes *strA-strB*, *aph(3')-Ia*, *blaTEM-1b*, *tet(A)-tet(B)*, *sul2*, and *dfrA5* (which confers resistance against trimethoprim). Six isolates collected from case-patients who resided in the Sydney region over a 3-week period in 2017 shared R-type ASSuTmGK: genes *aac*




Figure. Maximum-likelihood phylogeny of whole-genome single-nucleotide polymorphisms (SNPs) of 153 *Salmonella enterica* 4,[5],12:i:- sequence type (ST) 34 isolates and acquired drug-resistance genes. A) SNP analysis was conducted by performing whole-genome alignment of ST34 isolates from New South Wales (NSW), Australia, and a selection of published ST34 isolates collected in the United Kingdom, United States, and Denmark by using Snippy Core (<https://github.com/tseemann/snippy>) (online Technical Appendix, <https://wwwnc.cdc.gov/EID/article/24/4/17-1619-Techapp1.pdf>). Regions of recombination were identified by using BratNextGen (www.helsinki.fi/bsg/software/BRAT-NextGen/) and removed. SNPs were identified by using SNP-sites (<https://github.com/sanger-pathogens/snp-sites>), and the phylogeny was generated by using FastTree (www.microbesonline.org/fasttree/). Phylogeny and antimicrobial resistance metadata were combined by using Microreact (<https://microreact.org/showcase>). The colistin-resistant ST34 isolate from NSW is denoted by an orange star, fluoroquinolone-resistant isolates from NSW by orange squares, and pork isolates from NSW by orange triangles. Scale bar indicates 10 SNPs. B) Year of isolation and acquisition of drug resistance. Acquired drug-resistance genes were identified by screening all isolate contigs through the ResFinder (8) and CARD (<https://card.mcmaster.ca/>) databases by using ABRicate version 0.5 (<https://github.com/tseemann/abricate>). Only genes with a 100% homology match in ≥ 1 isolate are shown. Columns depict the results for individual isolates; rows represent acquired drug-resistance genes. The antibiotic class that genes confer resistance against is indicated at right. White indicates that the specified gene was not detected, gray indicates that the specified gene was detected but sequence homology against the reference was <100%, black indicates a perfect match between the isolate and reference gene sequence. MLS, macrolide, lincosamide, and streptogramin B.

(3)-IV, *aph* (4)-Ia, *aph*(3')-Ic, *bla*TEM-1B, *sul1*, and *dfr*A5 (which also confers resistance against trimethoprim) (Figure, panel B). These 6 isolates differed by 1–18 SNPs (most by <10 SNPs), and associated cases were clustered in time and occurred in neighboring suburbs, suggesting a possible cluster with a common source.

Fluoroquinolone resistance-conferring genes *qnr*S1 (from 3 case-patients) and *aac*(6')*lb-cr* (from 1 case-patient) were detected (Figure, panel B). As reported previously (12), the *aac*(6')*lb-cr* (*aac*A4-*cr*) gene was plasmid borne (IncHI2 plasmid) and was typically a class 1 integron-associated gene cassette (13). Of these 4 case-patients, 2 reported recent travel to Indonesia and Vietnam and the other 2 had no record of recent overseas travel; hence, we could not exclude the possibility of local acquisition. The isolate from the case-patient who traveled to Vietnam also displayed resistance to colistin (MIC 4 µg/mL). Neither the *mcr-1* or *mcr-2* genes nor mutations in the *pmrAB*, *phoPQ*, and *mgrB* genes were present (14). Rather, resistance was conferred by a recently identified third mobile colistin resistance gene, *mcr-3*, carried on a plasmid (15).

Conclusions

Using genomic surveillance, we identified the presence of novel colistin resistance gene *mcr-3* and indications that multidrug-resistant *Salmonella* 4,[5],12:i:- ST34 has established endemicity in Australia. Our findings highlight the public health benefits of genome sequencing-guided surveillance for monitoring the spread of multidrug-resistant mobile genes and isolates.

Acknowledgments

We thank the NSW Food Authority and the Health Protection Branch of the NSW Ministry of Health for expert advice.

This study was funded by the NSW Department of Health through the Translational Research Grants scheme. V.S. was funded by the Australian National Health and Medical Research Council Career Development Fellowship, and A.A. was funded by the National Health and Medical Research Council Centre of Research Excellence in Emerging Infectious Diseases.

About the Author

Dr. Arnott is a postdoctoral scientist with the Centre for Infectious Diseases and Microbiology–Public Health and Marie Bashir Institute for Infectious Diseases and Biosecurity, The University of Sydney. Her primary research interests are the molecular epidemiology and genomics of emerging and novel bacterial pathogens.

References

- Switt AI, Soyer Y, Warnick LD, Wiedmann M. Emergence, distribution, and molecular and phenotypic characteristics of *Salmonella enterica* serotype 4,5,12:i:-. *Foodborne Pathog Dis*. 2009;6:407–15. <http://dx.doi.org/10.1089/fpd.2008.0213>
- Hu Y, Liu F, Lin IY, Gao GF, Zhu B. Dissemination of the *mcr-1* colistin resistance gene. *Lancet Infect Dis*. 2016;16:146–7. [http://dx.doi.org/10.1016/S1473-3099\(15\)00533-2](http://dx.doi.org/10.1016/S1473-3099(15)00533-2)
- Campos J, Cristino L, Peixe L, Antunes P. MCR-1 in multidrug-resistant and copper-tolerant clinically relevant *Salmonella* 1,4,[5],12:i:- and S. Rissen clones in Portugal, 2011 to 2015. *Euro Surveill*. 2016;21:30270. <http://dx.doi.org/10.2807/1560-7917.ES.2016.21.26.30270>
- Li XP, Fang LX, Song JQ, Xia J, Huo W, Fang JT, et al. Clonal spread of *mcr-1* in PMQR-carrying ST34 *Salmonella* isolates from animals in China. *Sci Rep*. 2016;6:38511. <http://dx.doi.org/10.1038/srep38511>
- Doumith M, Godbole G, Ashton P, Larkin L, Dallman T, Day M, et al. Detection of the plasmid-mediated *mcr-1* gene conferring colistin resistance in human and food isolates of *Salmonella enterica* and *Escherichia coli* in England and Wales. *J Antimicrob Chemother*. 2016;71:2300–5. <http://dx.doi.org/10.1093/jac/dkw093>
- Lindstedt BA, Vardund T, Aas L, Kapperud G. Multiple-locus variable-number tandem-repeats analysis of *Salmonella enterica* subsp. *enterica* serovar Typhimurium using PCR multiplexing and multicolor capillary electrophoresis. *J Microbiol Methods*. 2004;59:163–72. <http://dx.doi.org/10.1016/j.mimet.2004.06.014>
- Seemann T, Goncalves da Silva A, Bulach DM, Schultz MB, Kwong JC, Howden BP. Nullarbor [cited 2018 Feb 12]. <https://github.com/tseemann/nullarbor>
- Carattoli A, Zankari E, Garcia-Fernández A, Voldby Larsen M, Lund O, Villa L, et al. In silico detection and typing of plasmids using PlasmidFinder and plasmid multilocus sequence typing. *Antimicrob Agents Chemother*. 2014;58:3895–903. <http://dx.doi.org/10.1128/AAC.02412-14>
- Hopkins KL, Kirchner M, Guerra B, Granier SA, Lucarelli C, Porrero MC, et al. Multiresistant *Salmonella enterica* serovar 4,[5],12:i:- in Europe: a new pandemic strain? *Euro Surveill*. 2010;15:19580.
- Mulvey MR, Finley R, Allen V, Ang L, Bekal S, El Bailey S, et al. Emergence of multidrug-resistant *Salmonella enterica* serotype 4,[5],12:i:- involving human cases in Canada: results from the Canadian Integrated Program on Antimicrobial Resistance Surveillance (CIPARS), 2003–10. *J Antimicrob Chemother*. 2013;68:1982–6. <http://dx.doi.org/10.1093/jac/dkt149>
- García P, Guerra B, Bances M, Mendoza MC, Rodicio MR. IncA/C plasmids mediate antimicrobial resistance linked to virulence genes in the Spanish clone of the emerging *Salmonella enterica* serotype 4,[5],12:i:-. *J Antimicrob Chemother*. 2011;66:543–9. <http://dx.doi.org/10.1093/jac/dkq481>
- Robicsek A, Strahilevitz J, Jacoby GA, Macielag M, Abbanat D, Park CH, et al. Fluoroquinolone-modifying enzyme: a new adaptation of a common aminoglycoside acetyltransferase. *Nat Med*. 2006;12:83–8. <http://dx.doi.org/10.1038/nm1347>
- Partridge SR, Tsafnat G, Coiera E, Iredell JR. Gene cassettes and cassette arrays in mobile resistance integrons. *FEMS Microbiol Rev*. 2009;33:757–84. <http://dx.doi.org/10.1111/j.1574-6976.2009.00175.x>
- Webb HE, Granier SA, Marault M, Millemann Y, den Bakker HC, Nightingale KK, et al. Dissemination of the *mcr-1* colistin resistance gene. *Lancet Infect Dis*. 2016;16:144–5. [http://dx.doi.org/10.1016/S1473-3099\(15\)00538-1](http://dx.doi.org/10.1016/S1473-3099(15)00538-1)
- Yin W, Li H, Shen Y, Liu Z, Wang S, Shen Z, et al. Novel plasmid-mediated colistin resistance gene *mcr-3* in *Escherichia coli*. *MBio*. 2017;8:e00543-17. <http://dx.doi.org/10.1128/mBio.00543-17>

Address for correspondence: Vitali Sintchenko, Centre for Infectious Diseases and Microbiology, Level 3, Institute of Clinical Pathology and Medical Research, Westmead Hospital, Westmead, Sydney, NSW, 2145, Australia; e-mail: vitali.sintchenko@sydney.edu.au

Multidrug-Resistant *Salmonella* Sequence Type 34, New South Wales, Australia, 2016–2017

Technical Appendix

Technical Appendix Table. Publically available *Salmonella* ST34 genomes used in study of multidrug-resistant *Salmonella* sequence type 34 in New South Wales, Australia, 2016–2017. Genomes were identified using Enterobase (<https://enterobase.warwick.ac.uk/>) and downloaded from the European Nucleotide Archive (ENA: <http://www.ebi.ac.uk/ena>). The isolate ID, year and country of collection are shown.

Isolate ID	Year of collection	Country of collection
ERR1544913	2014	Denmark
ERR24391	2009	United Kingdom
ERR24396	2009	United Kingdom
ERR24404	2009	United Kingdom
ERR24406	2008	United Kingdom
SRR1503240	2011	United States
SRR1503242	2011	United States
SRR1503243	2012	United States
SRR1503245	2013	United States
SRR1635093	2013	United Kingdom
SRR1645735	2013	United Kingdom
SRR1645745	2013	United Kingdom
SRR1646268	2013	United Kingdom
SRR1646274	2012	United Kingdom
SRR1646347	2012	United Kingdom
SRR1646348	2012	United Kingdom
SRR1646355	2012	United Kingdom
SRR1646359	2012	United Kingdom
SRR1646361	2012	United Kingdom
SRR1646365	2012	United Kingdom
SRR1646369	2012	United Kingdom
SRR1646375	2012	United Kingdom
SRR1822482	2014	United States
SRR1915431	2014	United States
SRR1915433	2014	United States
SRR1915434	2015	United States
SRR1931738	2013	United States
SRR1960281	2012	United States
SRR1966431	2014	United Kingdom
SRR1966998	2014	United Kingdom
SRR1969480	2014	United Kingdom
SRR2015055	2013	United States
SRR2015056	2013	United States
SRR2015057	2013	United States
SRR2015059	2014	United States
SRR2087755	2015	United States
SRR2152996	2015	United States
SRR2153006	2015	United States
SRR2192136	2015	United States
SRR2243429	2015	United States
SRR2243431	2015	United States
SRR2243437	2015	United States
SRR2538287	2010	Australia
SRR2538288	2010	Australia
SRR2577299	2010	United States
SRR2969505	2014	United States
SRR3098637	2015	United States
SRR3098638	2015	United States

Isolate ID	Year of collection	Country of collection
SRR3175446	2016	United States
SRR3175448	2016	United States
SRR3284675	2015	United Kingdom
SRR3284691	2015	United Kingdom
SRR3284715	2015	United Kingdom
SRR3284730	2015	United Kingdom
SRR3284731	2015	United Kingdom
SRR3284765	2015	United Kingdom
SRR3284842	2014	United Kingdom
SRR3285055	2015	United Kingdom
SRR3285406	2015	United Kingdom
SRR3322734	2015	United Kingdom
SRR3322960	2016	United Kingdom
SRR3499747	2016	United States
SRR3930234	2016	United States
SRR4032968	2016	United States
SRR4176760	2015	United States
SRR4190244	2016	United States
SRR4190251	2016	United States
SRR4450025	2011	United States
SRR4450027	2011	United States
SRR4841831	2009	United States
SRR4841833	2010	United States
SRR5193083	2016	United Kingdom
SRR5193127	2016	United Kingdom
SRR5193627	2016	United Kingdom
SRR5193644	2016	United Kingdom
SRR5193673	2016	United Kingdom
SRR5194203	2016	United Kingdom
SRR5194210	2016	United Kingdom
SRR5379273	2013	United States
SRR5631558	2017	United Kingdom
SRR5631627	2017	United Kingdom
SRR5631935	2017	United Kingdom
SRR5632032	2017	United Kingdom
SRR5632036	2017	United Kingdom
SRR5632041	2017	United Kingdom
SRR5632048	2017	United Kingdom
SRR5632062	2017	United Kingdom
SRR5632073	2017	United Kingdom
SRR5632117	2017	United Kingdom
SRR5632160	2017	United Kingdom
SRR5632173	2017	United Kingdom
SRR5632200	2017	United Kingdom
SRR5632268	2017	United Kingdom
SRR5632733	2017	United Kingdom
SRR5632860	2017	United Kingdom
SRR5632863	2017	United Kingdom
SRR5632868	2017	United Kingdom