

A Mental Models Approach to Assessing Public Understanding of Zika Virus, Guatemala

Brian G. Southwell, Sarah E. Ray,
Natasha N. Vazquez, Tere Ligorria, Bridget J. Kelly

Author affiliations: RTI International, Research Triangle Park, North Carolina, USA (B.G. Southwell, N.N. Vazquez); RTI International, Atlanta, Georgia, USA (S.E. Ray); RTI International, Guatemala City, Guatemala (T. Ligorria); RTI International, Washington, DC, USA (B.J. Kelly)

DOI: <https://doi.org/10.3201/eid2405.171570>

Mental models are cognitive representations of phenomena that can constrain efforts to reduce infectious disease. In a study of Zika virus awareness in Guatemala, many participants referred to experiences with other mosquitoborne diseases during discussions of Zika virus. These results highlight the importance of past experiences for Zika virus understanding.

Current risk communication literature includes guidelines regarding crises (1,2). Creating and distributing risk-mitigation information amid a nexus of emotion, public health threats, a journalistic tendency toward sensationalism, and misinformation can be daunting (3,4), which can make general guidelines appealing. We argue here, though, that tension between widely held, preexisting mental models of disease and the circumstances of each new emergent infectious disease offers an underappreciated challenge. By better acknowledging how existing audience mindsets reflect past experiences sometimes at odds with new circumstances, we can move beyond set guidelines to call for formative research, psychologically oriented literature reviews, and social discourse monitoring as crucial steps to address emerging infectious diseases. To support our argument, we offer case evidence regarding public understanding of Zika virus in Guatemala in early 2016.

Any assessment of public health intervention potential can begin with understanding existing mental models among a population. Mental models involve how persons imagine and conceptualize phenomena (5,6; online Technical Appendix, <https://wwwnc.cdc.gov/EID/article/24/5/17-1570-Techapp1.pdf>). In the case of infectious disease, mental models can include conceptualization of disease transmission routes and processes and constraints, just as mental models of mechanical tools can include a person's understanding of engineering principles.

To clarify how Guatemalans perceived Zika virus in spring 2016, we conducted 8 focus groups (with separate

groups for men and women) and 10 in-depth individual interviews (coordinated in country by T.L.). Participants were adults 18–49 years of age in both urban and rural regions affected by the virus. We recruited participants from the database of a market research firm in Guatemala (ConsuMer). Staff conducted all focus groups and interviews in Spanish in 2 departments (Zacapa and Suchitepéquez) (7); focus groups took place in central locations (e.g., restaurants) and individual interviews occurred in participants' homes.

The importance of personal experience with mosquitoborne viral disease in informing Zika virus understanding was striking. Virtually all participants were aware of Zika as a disease affecting Guatemala at the time of the interviews. At the same time, much of the discussion with participants clearly referred to other mosquitoborne diseases, rather than their conceiving of Zika virus as a new pathogen.

This pattern is understandable given the disease context in Guatemala, where dengue fever has been endemic for years, and given the substantial outbreak of chikungunya disease in 2014. Zika virus emerged and began to spread rapidly in Guatemala in late 2015. Participants apparently drew on experience with dengue and chikungunya as a baseline in understanding Zika virus. Mosquitoes were most commonly mentioned as a vector for the Zika virus, and participants often pointed out that the type of mosquito that transmitted Zika also was responsible for chikungunya and dengue.

Some participants made distinctions between mosquitoborne diseases. Many participants (≥ 1 in each of all 8 focus groups and 6 in individual interviews), for example, noted that the presence of conjunctivitis (red eyes) distinguishes Zika from other mosquitoborne diseases. Even in such cases, however, dengue and chikungunya served as an anchoring reference against which Zika virus was compared. Anchoring bias is a human tendency to rely on an initial piece of information even when new information comes to light (8).

Such predominant understanding of mosquitoborne transmission appears to have overshadowed other possibilities for transmission routes in popular imagination. Fewer than half of the in-depth interview participants reported knowing that Zika virus could be sexually transmitted. That gap in understanding poses challenges to preventive efforts to change social interaction (as opposed to emphasizing prevention of interaction with mosquitoes). Moreover, many participants (≥ 1 in each of 6 focus groups and half in individual interviews) expressed some sense of inevitability regarding mosquitoborne disease, likely because avoiding mosquito bites in Guatemala at certain times of the year can be difficult.

Anecdotal experience with symptoms loomed large in discussion. Participants (≥ 1 in each of 4 focus groups) noted, after learning about sexual transmission possibilities, that they would look for symptoms in a partner and base sexual behavior on their assessment (insofar as they had agency to

decide). We know, however, that a person infected with Zika virus might not have easily observable symptoms. Even if persons accept the possibility of sexual transmission, they might not engage in safe sex practices with asymptomatic infected partners.

If persons understand Zika virus through a mental model informed by dengue or chikungunya, public health officials should address potential confusion, especially in light of differences (e.g., sexual transmission) that might be misunderstood or ignored. Even when not confusing the illnesses, participants clearly conceptualized Zika in comparison with relatively more familiar illnesses. In this way, they operated in similar fashion as consumers encountering novel products do (9,10). Public health messaging might leverage this tendency. If it is easiest to understand a new outbreak in comparison to a previous one, using analogy or direct comparison might be effective but will also require careful emphasis on what is new.

Acknowledgments

We thank staff of ConSUMer of Guatemala, who provided important study support in addition to the contributions described in this letter.

RTI International sponsored the study that produced the information reported here as part of an in-kind donation to the Guatemala Ministry of Public Health and Social Assistance.

About the Author

Dr. Southwell directs the Science in the Public Sphere Program in the Center for Communication Science at RTI International. He also teaches at Duke University and the University of North Carolina at Chapel Hill. His primary research interests include public understanding of science and health and the ways in which the information environment shapes human behavior.

References

- Centers for Disease Control and Prevention. Crisis and emergency risk communication. 2014 edition. Atlanta: US Department of Health and Human Services, Centers for Disease Control and Prevention; 2014.
- Sandman P, Lazard J. Crisis communication: a guide for action. Planning what to say when terrorists, epidemics, and other emergencies strike [cited 2018 Feb 20]. http://www.psandman.com/handouts/AIHA/AIHA_book.pdf
- Glik DC. Risk communication for public health emergencies. *Annu Rev Public Health*. 2007;28:33–54. <http://dx.doi.org/10.1146/annurev.publhealth.28.021406.144123>
- Southwell BG, Reynolds BJ, Fowlie K. Communication, media relations and infectious disease surveillance. In: M'ikanatha NM, Lynfield R, Van Beneden CA, de Valk H, editors. *Infectious disease surveillance*, 2nd ed. Oxford: John Wiley & Sons; 2013. p. 607–17.
- Johnson-Laird P. *Mental models*. Cambridge (MA): Harvard University Press; 1983.
- Jones NA, Ross T, Lynam PP, Leitch A. Mental models: an interdisciplinary synthesis of theory and methods. *Ecol Soc*. 2011;16:art46. <http://dx.doi.org/10.5751/ES-03802-160146>
- Ray SE, Dolina S, Jimenez K, Vazquez N, Ligorria T, Southwell B. Insights regarding Zika virus perceptions in Guatemala: key informant interviews and information environment monitoring: report prepared for Guatemala Ministry of Public Health and Social Assistance. Research Triangle Park (NC): RTI International; 2016.
- Tversky A, Kahneman D. Judgment under uncertainty: heuristics and biases. *Science*. 1974;185:1124–31. <http://dx.doi.org/10.1126/science.185.4157.1124>
- Jhang JH, Grant SJ, Campbell MC. Get it? Got it. Good! Enhancing new product acceptance by facilitating resolution of extreme incongruity. *J Mark Res*. 2013;49:247–59. <http://dx.doi.org/10.1509/jmr.10.0428>
- Moreau CP, Lehmann DR, Markman AB. Entrenched knowledge structures and consumer response to new products. *J Mark Res*. 2001;38:14–29. <http://dx.doi.org/10.1509/jmkr.38.1.14.18836>

Address for correspondence: Brian G. Southwell, RTI International, Science in the Public Sphere Program, 3040 E Cornwallis Rd, Research Triangle Park, NC 27709, USA; email: bsouthwell@rti.org

Cerebrospinal Fluid Immunoglobulins as Potential Biomarkers of Chikungunya Encephalitis

Marzia Puccioni-Sohler, Luiz Claudio Farias, Mauro Jorge Cabral-Castro, Mariano G. Zalis, Rosangela S. Kalil, Maria Cecília F. Salgado

Author affiliations: Universidade Federal do Rio de Janeiro, Rio de Janeiro, Brazil (M. Puccioni-Sohler, L.C. Farias, M.J. Cabral-Castro, M.G. Zalis); Universidade Federal do Estado do Rio de Janeiro, Rio de Janeiro (M. Puccioni-Sohler, R.S. Kalil, M.C.F. Salgado)

DOI: <https://doi.org/10.3201/eid2405.171763>

Chikungunya virus causes fever and severe polyarthritides or arthralgia and is associated with neurologic manifestations that are sometimes challenging to diagnose. We demonstrate intrathecal synthesis of chikungunya antibodies in a patient with a history of acute infection complicated by encephalitis. The specificity of the intracerebral immune response supports early chikungunya-associated encephalitis diagnosis.

Chikungunya virus (CHIKV) is an alphavirus transmitted by infected *Aedes* mosquitoes (*Ae. aegypti* and *Ae. albopictus*) (1). Global expansion epidemics have been

A Mental Models Approach to Assessing Public Understanding of Zika Virus, Guatemala

Technical Appendix

Mental Models as a Foundation for Public Communication

Cognitive psychology and brain sciences researchers developed the concept of a mental model to explain how people make sense of, or mentally represent, processes and physical entities when considering them. Because we apply the mental model concept to public understanding of Zika virus in Guatemala in our letter, some readers might find a brief summary of thinking about mental models as a tool for risk communication to be helpful.

Mental models are cognitive representations that comprise how people imagine and conceptualize a phenomenon (1–4). For example, people might understand a house to be a structure with a roof and doors and windows in which people live; they also might have ideas about appropriate building materials or the ways in which a roof physically operates to protect and comfort occupants. The complexity and dimensions of a person’s mental model of a phenomenon can constrain engagement with information about that phenomenon. Atman and colleagues (5) cite an example of how mental models can constrain popular understanding of the environment: they found evidence for some misunderstanding of the effects of radon gas in homes that stemmed from participants’ models of radon. Some people mistakenly viewed radon as a long-lasting (as opposed to relatively short-lived) gas, which undermined their sense of the utility of efforts to clear radon from a house as a way of mitigating problems associated with radon. Viewing radon as a long-lasting or permanent contaminant could discourage people from taking action because of fatalism or perceived inability to combat the contamination.

In the case of infectious disease, mental models can include conceptualization of disease transmission as well as perceptions of the mechanisms through which the disease causes effects, as we note in our research letter. Importantly, conceptualizations of disease that people develop

based on their own observations, experiences, and background knowledge can differ substantially from models described in scientific, peer-reviewed literature (6–10): for example, some people view the common cold as being a function of cold weather rather than a viral infection (9) or some middle school students hold a mental model of vaccination as a process in which the vaccine attacks a virus that is already present within a body (10). Acknowledging audience members' foundational understanding of how processes like disease transmission work, in turn, can bolster prospects for risk communication efforts.

References

1. Craik K. The nature of explanation. New York: Cambridge University Press; 1967.
2. Johnson-Laird P. Mental models. Cambridge (MA): Harvard University Press; 1983.
3. Jones NA, Ross T, Lynam PP, Leitch A. Mental models: an interdisciplinary synthesis of theory and methods. *Ecol Soc*. 2011;16:art46. <http://dx.doi.org/10.5751/ES-03802-160146>
4. Morgan MG, Fischhoff B, Bostrom A, Atman C. Risk communication: a mental models approach. New York: Cambridge University Press; 2001.
5. Atman CJ, Bostrom A, Fischhoff B, Morgan MG. Designing risk communications: completing and correcting mental models of hazardous processes, part I. *Risk Anal*. 1994;14:779–88. [PubMed http://dx.doi.org/10.1111/j.1539-6924.1994.tb00289.x](http://dx.doi.org/10.1111/j.1539-6924.1994.tb00289.x)
6. Chi MTH, Slotta JD, de Leeuw N. From things to processes: a theory of conceptual change for learning science concepts. *Learn Instr*. 1994;4:27–43. [http://dx.doi.org/10.1016/0959-4752\(94\)90017-5](http://dx.doi.org/10.1016/0959-4752(94)90017-5)
7. Duschl RA, Schweingruber HA, Shouse AW, editors. Taking science to school: learning and teaching science in grades K–8. Washington: National Academies Press; 2007.
8. Vosniadou S, Brewer W. Mental models of the earth: a study of conceptual change in childhood. *Cognit Psychol*. 1992;24:535–86. [http://dx.doi.org/10.1016/0010-0285\(92\)90018-W](http://dx.doi.org/10.1016/0010-0285(92)90018-W)
9. Sigelman CK. Age and ethnic differences in cold weather and contagion theories of colds and flu. *Health Educ Behav*. 2012;39:67–76. [PubMed http://dx.doi.org/10.1177/1090198111407187](http://dx.doi.org/10.1177/1090198111407187)
10. Jee BD, Uttal DH, Spiegel A, Diamond J. Expert-novice differences in mental models of viruses, vaccines, and the causes of infectious disease. *Public Underst Sci*. 2015;24:241–56. [PubMed http://dx.doi.org/10.1177/0963662513496954](http://dx.doi.org/10.1177/0963662513496954)