

---

# Avian Influenza A Viruses among Occupationally Exposed Populations, China, 2014–2016

Chuansong Quan,<sup>1</sup> Qianli Wang,<sup>1</sup> Jie Zhang, Min Zhao, Qigang Dai, Ting Huang, Zewu Zhang, Shenghua Mao, Yifei Nie, Jun Liu, Yun Xie, Baorong Zhang, Yuhai Bi, Weifeng Shi, Peipei Liu, Dayan Wang, Luzhao Feng, Hongjie Yu, William J. Liu, George F. Gao

To determine the seroprevalence and seroconversion of avian influenza virus (AIV) antibodies in poultry workers, we conducted a seroepidemiologic study in 7 areas of China during December 2014–April 2016. We used viral isolation and reverse transcription PCR to detect AIVs in specimens from live poultry markets. We analyzed 2,124 serum samples obtained from 1,407 poultry workers by using hemagglutination inhibition and microneutralization assays. We noted seroprevalence of AIV antibodies for subtypes H9N2, H7N9, H6N1, H5N1-SC29, H5N6, H5N1-SH199, and H6N6. In serum from participants with longitudinal samples, we noted seroconversion, with  $\geq 4$ -fold rise in titers, for H9N2, H7N9, H6N1, H5N1-SC29, H6N6, H5N6, and H5N1-SH199 subtypes. We found no evidence of H10N8 subtype. The distribution of AIV antibodies provided evidence of asymptomatic infection. We found that AIV antibody prevalence in live poultry markets correlated with increased risk for H7N9 and H9N2 infection among poultry workers.

Author affiliations: National Institute for Viral Disease Control and Prevention, Chinese Center for Disease Control and Prevention, Beijing, China (C. Quan, J. Zhang, P. Liu, D. Wang, W.J. Liu, G.F. Gao); Shandong First Medical University & Shandong Academy of Medical Sciences, Jinan, China (C. Quan, W. Shi); Fudan University School of Public Health, Shanghai, China (Q. Wang, H. Yu); Institute of Microbiology, Chinese Academy of Sciences, Beijing (M. Zhao, Y. Bi, G.F. Gao); Jiangsu Provincial Center for Disease Control and Prevention, Nanjing, China (Q. Dai); Sichuan Provincial Center for Disease Control and Prevention, Chengdu, China (T. Huang); Dongguan Municipal Center for Disease Control and Prevention, Dongguan, China (Z. Zhang); Shanghai Municipal Center for Disease Control and Prevention, Shanghai (S. Mao); Henan Provincial Center for Disease Control and Prevention, Zhengzhou, China (Y. Nie); Zaozhuang Center for Disease Control and Prevention, Zaozhuang, China (J. Liu); Jiangxi Provincial Center for Disease Control and Prevention, Nanchang, China (Y. Xie); Aviation General Hospital, Beijing (B. Zhang); Chinese Center for Disease Control and Prevention, Beijing (L. Feng, G.F. Gao)

DOI: <https://doi.org/10.3201/eid2512.190261>

Human infection with avian influenza viruses (AIVs) has been reported in China since the late 1990s. Since then, human infections with subtypes H5N1, H5N6, H6N1, H7N4, H7N9, H9N2, and H10N8 have been reported continuously and are a substantial threat to public health in the country (1–5). Birds at wholesale and retail live poultry markets are recognized incubators for novel influenza virus subtypes (6–9). Because of special occupational characteristics, poultry workers are at a high risk for repeated exposure to AIV-infected poultry. Most case-patients with H7N9 infection have had a history of contact with live poultry, and poultry workers represent a substantial proportion of cases (10). Several studies on AIV seroprevalence in occupationally exposed populations suggest that asymptomatic or clinically mild AIV infections are extensively prevalent among poultry workers (11–14). A serologic study of AIV distribution among poultry workers could directly evaluate the potential for AIVs to cross the species barrier to infect humans and might illuminate the current understanding of AIV prevalence in live poultry markets (15).

Low pathogenicity avian influenza A(H9N2) virus is distributed widely in domestic poultry around the world. A systematic review reports H9N2 virus seroprevalence in avian-exposed populations ranges from 1% to 43% by hemagglutination inhibition (HI) assays (16). Since a 2013 H7N9 infection outbreak in China, caused by a novel reassortant influenza A(H7N9) virus and associated with severe human infections, seroprevalence of the H7N9 subtype has been reported to range from 6% to 14.9% in southern China (17,18). In a previous study, the seroprevalence of H5 subtype AIVs in poultry workers was relatively low, whereas a cross-sectional study conducted in Zhejiang Province reported a seroprevalence of 4.7% for H5N1 virus antibodies (19).

Few large-scale longitudinal seroepidemiologic studies have included multiple AIV subtypes in diverse epidemic regions, especially after emergence of novel

---

<sup>1</sup>These authors contributed equally to this article.

subtypes. We conducted a prospective seroepidemiologic study in 7 representative areas across China to address gaps in the research. We characterized the seroprevalence profiles of 7 dominant human-infecting AIV subtypes among occupationally exposed workers in live poultry markets. Our aim was to further analyze human AIV infection risks for serotypes common in occupational exposure, including H5N1, H5N6, H6N1, H6N6, H7N9, H9N2, and H10N8 virus subtypes.

## Methods

### Ethics Approval

This study was approved by the Ethics Review Committee of the National Institute for Viral Disease Control and Prevention, Chinese Center for Disease Control and Prevention. The study was conducted in accordance with the principles of the Declaration of Helsinki and the standards of Good Clinical Practice as defined by the International Conference on Harmonization (<https://www.ich.org>).

### Study Design and Participants

During December 2014–April 2016, we conducted a longitudinal seroepidemiologic study to assess asymptomatic AIV infection levels among poultry workers in China. We defined poultry workers as persons who repeatedly are exposed to poultry and work in wholesale or retail live poultry markets or in backyard farms, including wholesale sellers, retail sellers, transporters, processors, or feeders. The study included 1 municipality, Shanghai, and 6 provinces, Guangdong, Henan, Jiangsu, Jiangxi, Shandong, and Sichuan (Figure 1, panels A and B; Appendix, <https://wwwnc.cdc.gov/EID/article/25/12/19-0261-App1.pdf>). The study design included 4 serologic surveys. We collected whole blood samples from participating poultry workers at an initial visit in December 2014 and again during 3 consecutive follow-up visits in April 2015, December 2015, and April 2016 (Figure 1, panel C).

We used a standardized questionnaire to collect information at initial participant enrollment and updated participant information at subsequent visits. Participant information collected was demographic data, exposure variables, whether the worker experienced influenza-like illness within the previous month, and whether they received a seasonal influenza vaccination within the previous 12 months (Appendix).

Some poultry workers in China are short-term employees with high population mobility. We attempted to conduct follow-up studies with these employees through assistance from the market managers. To ensure the sample size, we enrolled new participants at each visit to the poultry markets (Figure 2).

We also recruited a control group of 216 outpatients with noninfectious diseases on physical examination at a

general hospital in Beijing in October 2015. We collected 216 serum samples from the control group.

### Collection of Human Samples

We collected a single venous whole blood sample from each study participant at each visit by using a Vacutainer blood collection tube (Becton Dickinson, <https://www.bd.com>). We divided serum into 3 aliquots and froze at  $-80^{\circ}\text{C}$  until testing.

### Serologic Assays


We tested participant serum samples for antibodies against H5N1, H5N6, H6N1, H6N6, H7N9, H9N2, and H10N8 virus subtypes, as well as for antibodies against seasonal influenza A(H1N1)pdm09 (pH1N1) and H3N2 viruses, to exclude cross-reactivity. We selected available representative antigens on the basis of their antigenic characteristics (Table 1) and analyzed the relevant phylogenetic relationship of hemagglutinin (HA) genes (Appendix Figures 1–5).

We performed all serologic assays in a Biosafety Level 2 or 3 laboratory. First, we screened samples by using an HI assay for antibodies, as described previously (20). We tested serum samples at a starting dilution of 1:10, followed by a 2-fold dilution to the endpoint (Appendix). To confirm HI assay results, we performed a microneutralization (MN) assay on serum samples with an HI titer  $\geq 1:20$  to H5N1, H5N6, H6N1, H6N6, H7N9, or H10N8 subtypes and those with an HI titer  $\geq 1:40$  to H9N2, pH1N1, or H3N2 subtypes, as previously described (20).

We used HI and MN cutoff values in accordance with previously published data (Appendix Table 1). We considered  $\geq 1:20$  as the cutoff value for HI and MN titers for positive tests for H5N1, H5N6, H6N1, H6N6, H7N9, and H10N8 virus subtypes (11,12,21) and considered  $\geq 1:40$  as the cutoff value for HI titer and  $\geq 1:80$  as the cutoff value for MN titer for positive tests for H9N2, pH1N1, and H3N2 virus subtypes (22,23). We set a stricter dilution cutoff value for the H9N2 virus subtype. An HI titer of 1:40 commonly is used and generally is an accepted value for influenza serologic assays used in detection of seasonal influenza and avian influenza H9 infection (24). We considered participants to have seroconversion when they had a  $\geq 4$ -fold rise in antibody titer measured by HI assay between collection of  $\geq 1$  serum samples, plus an MN titer value of the later specimen being  $\geq 1:20$  or  $\geq 1:80$  for H9N2 subtype only.

### Isolation of AIVs from Environmental and Poultry Samples

For environmental and poultry samples, we used previously described sampling and detection methods (25). In brief, we randomly selected environmental sites and poultry to sample by using a multistage sampling strategy. We collected environmental samples by swabbing water




**Figure 1.** Temporal and spatial distribution of human infections with avian influenza A virus subtypes before and during serosurveillance, China. A) Geographic distribution of avian influenza A(H7N9) virus infection among humans in China during May 1997–October 2016. The number of case-patients in each province is based on data published by the World Health Organization ([https://www.who.int/influenza/human\\_animal\\_interface/avian\\_influenza/archive/en/](https://www.who.int/influenza/human_animal_interface/avian_influenza/archive/en/)) and the National Health and Family Planning Commission of the Republic of China ([http://www.nhc.gov.cn/jkj/s2907/new\\_list.shtml?tdsourcetag=s\\_pcqq\\_aiomsg](http://www.nhc.gov.cn/jkj/s2907/new_list.shtml?tdsourcetag=s_pcqq_aiomsg)). Density of shading represents the number of reported avian influenza H7N9 cases in humans in each province. Cases of other AIV subtype infections are represented by other symbols. B) Density of live poultry markets per 10,000 persons in each province included in the study, from data collected during 2013–2014. Red flags indicate locations of poultry markets selected for the serosurveillance study. C) Distribution of biweekly cases of human H7N9 infection before and during serosurveillance study. Orange bars indicate the number of biweekly cases of human H7N9 infection. Dashed lines indicate initial survey and follow-up dates for serosurveys, which were conducted before and after the third and fourth wave H7N9 epidemics. Reported cases of H5N1, H5N6, H6N1, H9N2, and H10N8 infection are noted with symbols as in panel A. AIV, avian influenza virus.

troughs, floors, and drains in poultry enclosures and collected oropharyngeal and cloacal swabs from apparently healthy poultry. We isolated avian influenza viruses in 9- to 10-day-old specific pathogen-free chicken embryos by using viral isolation procedures and following World Health Organization guidelines (20). We further analyzed hemagglutinin-positive samples by using reverse transcription PCR (RT-PCR) to identify hemagglutinin (HA) and

neuraminidase (NA) genetic subtypes (20). Except for Shandong Province, we detected AIVs from domestic poultry and live poultry market environments in all study areas.

**Data Analysis**

Our analyses were based on seroepidemiologic studies for influenza published by Horby et al. (26). We assigned each participant a unique identifier and used all data collected



**Figure 2.** Flowchart of initial participant enrollment and follow-up distribution in 7 areas of China in a study of avian influenza virus seroprevalence during December 2014–April 2016.

with the questionnaire to establish a database. We performed a multivariate logistic regression model to evaluate independent risk factors associated with seroprevalence of antibodies in poultry workers. Risk factors evaluated were age; sex; occupational exposure factors, including processing, selling, transporting, and feeding poultry; and seropositivity to human influenza pH1N1 or H3N2 viruses. For logistic regression analysis, we estimated the maximum likelihood for the odds ratio (OR) and calculated 95% CIs by using the Wald  $\chi^2$  test. We used binomial distribution to calculate 95% CIs of rate. We used Spearman correlation analysis to estimate the association between seroprevalence and local epidemic intensity of AIVs in live poultry markets by region. We used 2-tailed p values for all calculations and considered values  $<0.05$  statistically significant. We performed statistical analyses by using SAS 9.4 (SAS Institute, Inc., <https://www.sas.com>).

## Results

### Participant Characteristics

We collected 2,124 serum samples from 1,407 participants from 1 municipality, Shanghai, and 6 provinces, Guangdong, Henan, Jiangsu, Jiangxi, Shandong, and Sichuan, in China. We had paired or serial serum samples from 652

participants who had  $\geq 2$  visits during the study period. The median age of participants with completed questionnaire information was 46 years (interquartile range [IQR] 36–52 years); 54.0% (1,147/2,124) of samples were from men. The most common category of poultry exposure was poultry seller. We did not see statistically significant differences in the distribution of demographic characteristics of participants, including sex and age, over the 4-period survey. In addition, 2.8% (59/2,124) of samples came from poultry workers who reported receiving a seasonal influenza vaccine within the previous 12 months (Table 2).

Of the 216 participants in control group, the median age was 48 years (IQR 34–59 years); 45.8% were male. We saw no significant differences in their data compared with poultry workers (data not shown).

### Seroprevalence of Antibodies against AIVs

In the 2,124 samples, the overall seroprevalence of antibodies was 11.2% for H9N2 subtype and 3.9% for H7N9 subtype. Seroprevalence for H5Nx and H6Nx subtypes was lower, ranging from 1.3% to 2.1% for H5Nx and from 0.4% to 2.5% for H6Nx. We did not observe evidence of H10N8 infection (Table 3).

The seroprevalence profile was geographically distinct (Figure 3). For example, in Shandong Province, H9N2

**Table 1.** Avian influenza A antigens used in serologic hemagglutinin inhibition and microneutralization assays, China\*

Subtype	Virus strain	GISAID number
Avian influenza		
H5N1 clade 2.3.2.1c	A/chicken/Shanghai/02.12 HZ199-P/2015 (SH199)	EPI1544294
H5N1 clade 2.3.4.4	A/pigeon/Sichuan/NCXN29/2014 (SC29)	EPI590898
H5N6 clade 2.3.4.4	A/duck/Guangdong/04.22 DGCP069-O/2015	EPI660071
H6N1	A/Taiwan/2/2013	EPI459855
H6N6	A/duck/Guangxi/04.10 JX031/2015	EPI661887
H7N9	A/chicken/Guangdong/04.22 DGCP098-O/2015	EPI666285
H9N2	A/chicken/Guangdong/04.15 SZBAXQ005/2015	EPI661935
H10N8	A/chicken/Jiangxi /B18/2014	EPI1544302
Seasonal influenza		
H1N1(pdm09)	A/California/04/2009	EPI176470
H3N2	A/Beijing/CAS0001/2007	EPI1544286

\*GISAID, <https://www.gisaid.org>.

**Table 2.** Characteristics of study participants in serosurveys for avian influenza viruses, China, 2014–2016\*

Variables	2014 Dec, n = 700	2015 Apr, n = 506	2015 Dec, n = 481	2016 Apr, n = 437	Total, n = 2,124	$\chi^2$ †	p value
Sex, no. (%)							
M	369 (52.7)	264 (52.2)	278 (51.8)	236 (54.0)	1,147 (54.0)	3.94	0.27
F	331 (47.3)	242 (47.8)	203 (42.2)	201 (46.0)	977 (46.0)		
Age, y, no. (%)‡							
<21	10 (1.4)	6 (1.2)	4 (0.8)	11 (2.5)	31 (1.5)	1.43	0.23
21–40	212 (30.3)	144 (28.5)	164 (34.1)	144 (33)	664 (31.3)		
41–60	394 (56.3)	308 (60.9)	254 (52.8)	232 (53.1)	1,188 (55.9)		
>60	78 (11.1)	47 (9.3)	55 (11.4)	50 (11.4)	230 (10.8)		
Missing data	6 (0.9)	1 (0.2)	4 (0.8)	0	11 (0.5)		
Median age (range)§	46 (38–52)	47 (38–52)	45 (35–52)	45 (35–52)	46 (36–52)	6.62	0.08
Type of poultry exposure, no. (%)¶							
Processing	155 (22.1)	107 (21.1)	118 (24.5)	94 (21.5)	474 (22.3)	27.88	0.006
Selling	423 (60.4)	332 (65.6)	299 (62.2)	191 (43.7)	1,243 (58.5)		
Transportation	39 (5.6)	31 (6.1)	24 (5)	21 (4.8)	115 (5.4)		
Feeding	191 (27.3)	125 (24.7)	124 (25.8)	93 (21.3)	533 (25.1)		
Others	59 (8.4)	35 (6.9)	25 (5.2)	48 (11)	167 (7.9)		
Missing data	0	0	2 (0.4)	0	2 (0.1)		
Length of poultry exposure, y (range)§	8 (3–15)	8 (3–15)	5 (2–10)	5 (3–10)	6 (3–13)	61.63	<0.001
Vaccinated against seasonal influenza, no. (%)	23 (3.3)	8 (1.6)	20 (4.2)	8 (1.8)	59 (2.8)	8.20	0.04

\*Some participants participated in >1 survey.

†By  $\chi^2$  test, unless otherwise indicated.

‡By  $\chi^2_{CMH}$  test. Missing data were not calculated.

§By Kruskal-Wallis test.

¶Most participants had multiple exposure types. Sums of percentages exceed 2,124. Missing data were not calculated.

virus antibody seroprevalence was 23%, which was higher than in other provinces, especially Sichuan Province, which had only a 4.2% seroprevalence for this subtype. Provinces in the Yangtze River Delta, which were the first to report H7N9 infections in patients during the 2013 outbreak, exhibited higher seroprevalence rates compared with the other provinces. Shanghai had a rate of 10.3% and Jiangsu Province had a rate of 6.9%. In Sichuan Province, where a non-laboratory-confirmed H7N9-infected patient was reported before 2017, no participant tested positive for the H7N9 subtype.

Seroprevalence of H5 and H6 subtypes among poultry workers also were different by region. Detected H5 subtypes included H5N1-SH199 clade 2.3.2.1c in 5.3% of samples from Shandong Province; H5N1-SC29 clade 2.3.4.4 in 3.0% of samples from Jiangsu Province and in 3.3% of samples from Sichuan Province; and H5N6 in 4.9% of samples from Shanghai. We detected H6N1 in 5.2% of samples from Jiangsu Province and in 3.8% from Shanghai and H6N6 in 3.3% of samples from Shanghai (Figure 3).


Among the 216 participants in the control group, we found no evidence of antibodies against H7N9 virus and a lower prevalence (3.7%) of antibodies against H9N2 virus than in the poultry workers. We observed no statistically significant differences in the prevalence of antibodies against other AIV subtypes between the control group and poultry workers (Appendix Table 2).

### Seroconversion of Antibodies against AIVs among Poultry Workers

We observed seroconversion in all AIV antigens during the study period, except the H10 subtype, which might represent a new asymptomatic AIV infection among poultry workers (Figure 4, panel A). Among 652 poultry workers with paired or serial serum samples during the study, 3.5% demonstrated seroconversion for H9N2 virus, 1.4% demonstrated seroconversion for H7N9 virus, and <1% demonstrated seroconversion for H5 or H6 viruses (Figure 4, panels B and C; Appendix Tables 3–9). Because we saw no evidence of H10N8 virus, we also

**Table 3.** Seroprevalence among poultry workers surveyed for avian influenza viruses, China, 2014–2016\*

Antigen	No. (%; 95% CI) seropositive participants				
	2014 Dec, n = 700	2015 Apr, n = 506	2015 Dec, n = 481	2016 Apr, n = 437	Total, n = 2,124
Avian influenza serotype					
H5N1-SH199	6 (0.9, 0.2–1.5)	6 (1.2, 0.2–2.1)	10 (2.1, 0.8–3.4)	6 (1.4, 0.3–2.5)	28 (1.3, 0.8–1.8)
H5N1-SC29	22 (3.1, 1.8–4.4)	17 (3.4, 1.8–4.9)	2 (0.4, 0.1–1.5)	3 (0.7, 0.1–2.0)	44 (2.1, 1.5–2.7)
H5N6	28 (4, 2.5–5.5)	11 (2.2, 0.9–3.4)	2 (0.4, 0.1–1.5)	1 (0.2, 0–1.3)	42 (2.0, 1.4–2.6)
H6N1	22 (3.1, 1.8–4.4)	21 (4.1, 2.4–5.9)	5 (1, 0.1–1.9)	5 (1.1, 0.4–2.6)	53 (2.5, 1.8–3.2)
H6N6	0 (0, 0–0.5)	0 (0, 0–0.7)	7 (1.5, 0.4–2.5)	1 (0.2, 0–1.3)	8 (0.4, 0.1–0.6)
H7N9	33 (4.7, 3.1–6.3)	36 (7.1, 4.9–9.4)	6 (1.3, 0.3–2.2)	7 (1.6, 0.4–2.8)	82 (3.9, 3.0–4.7)
H9N2	48 (6.9, 5.0–8.7)	59 (11.7, 8.9–14.5)	64 (13.3, 10.3–16.3)	66 (15.1, 11.7–18.5)	237 (11.2, 9.8–12.5)
Seasonal influenza serotype					
H1N1(pdm09)	94 (13.4, 10.9–16.0)	85 (16.8, 13.5–20.1)	90 (18.7, 15.2–22.2)	79 (18.1, 14.5–21.7)	348 (16.4, 14.8–18.0)
H3N2	237 (33.9, 30.4–37.4)	165 (32.6, 28.5–36.7)	199 (41.4, 37.0–45.8)	171 (39.1, 34.6–43.7)	772 (36.3, 34.3–38.4)



**Figure 3.** Avian influenza virus seroprevalence in the studied regions of China during December 2014–April 2016. A) Geographic areas included for serosurveillance: 1 municipality, Shanghai, and 6 provinces, Guangdong, Henan, Jiangsu, Jiangxi, Shandong, and Sichuan. B) Seroprevalence against avian influenza A virus subtypes in 4 cross-sectional surveys. Colors on map correspond to colors in bar graphs. \*Seasonal influenza virus subtype.

saw no seroconversion for the subtype (Table 4; Figure 4, panel A).

Some participants showed consistently seropositive results, 15 for H7N9 subtype and 41 for H9N2 subtype and a few each for H5N1, H5N6, and H6N1 subtypes (Figure 4, panel B). One participant (no. 14.12GD72) showed HI titers at 1:20 and MN titers at 1:160 to H5N1-SH199 subtype in 4 consecutive surveys (Figure 4, panel C).

#### Risk Analysis for Asymptomatic AIV Infections

In the multivariable analysis, we identified demographic and occupational risk factors for poultry workers with asymptomatic infections. For instance, the demographic classification female (adjusted OR [aOR] 2.2, 95% CI 1.4–3.6), and occupational classification poultry seller (aOR 4.1, 95% CI 2.2–7.7) appear to be risk factors for H7N9 infection. For H9N2 subtype, female (aOR 1.6, 95% CI 1.2–2.1) and poultry seller (aOR 1.9, 95% CI 1.4–2.6) appear to be risk factors for infection. In addition, the number of years working in poultry-related occupations was associated with seroprevalence. In particular, samples from workers reporting  $\geq 3$  years of exposure were associated with seroprevalence of H9N2 subtype. Factors associated with increased risk for H5 infections included being  $>55$  years of age, being exposed to ducks, or being exposed to ill or dead poultry (Table 5).

Our study revealed a correlation between the presence of antibodies and seasonal influenza virus infection. We saw


an association between the presence of pH1N1 virus antibodies and increased seropositivity for H5N1 or H5N6 subtypes, and between occurrence of seasonal H3N2 virus antibodies in humans and positive antibody titers for H7N9 virus subtype. We also saw a positive association between elevated H6N1 seropositivity and the presence of antibodies against pH1N1 (aOR 3.0, 95% CI 1.7–5.4) and H9N2 (aOR 2.6, 95% CI 1.4–5.0) subtypes (Table 5). Seasonal influenza vaccination history was not a significant risk factor for elevated AIV antibody titers, perhaps because of low vaccination rates.

#### AIV Circulation in Poultry and Markets

We collected 6,207 samples from poultry and the environment for AIV screening and detection in this study. In Shanghai, 4.1% (20/493) of samples were positive for H7N9 subtype, as were 8.6% (41/476) of samples from Jiangsu Province. However, only 0.6% (15/2,308) of samples from Jiangxi Province, 0.6% (12/2,158) of samples from Guangdong Province, and 0.2% (1/516) of samples from Sichuan Province were positive for H7N9 subtype (Appendix Table 10).

For H9N2 subtype, 14.4% (71/493) of samples from Shanghai, 9.5% (45/476) from Jiangsu Province, and 8.3% (180/2,158) of samples from Guangdong Province were positive. However, only 4.4% (102/2,308) of samples from Jiangxi Province and 5.5% (14/256) from Henan Province were positive for H9N2 (Appendix Table 10).

Exploring the correlation between AIV circulation in poultry and seroprevalence in workers in live poultry



**Figure 4.** Seroconversion and persistent positivity for avian influenza virus (AIV) A subtypes based on HI titers in a cohort study in China during December 2014–April 2016. Each dot and line connection represents 1 participant. Red dashed lines represent positive cutoff for the HI titers; HI–positive samples were confirmed by a microneutralization assay. A) Comparison of paired samples from participants during 2 surveillance periods showing seroconversion for 6 AIV subtypes. Weighted lines and dots represent participants with seroconversion. B) Number of participants with >2 positive sample who were persistently seropositive for 6 AIV subtypes. Weighted lines and dots represent number of participants with seropositivity. C) Antibody titers of representative participants with >1 positive sample in the 4 serosurveys. HI, hemagglutinin inhibition.

markets revealed a correlation coefficient of 0.8 ( $p = 0.04$ ) for H7N9 virus and 0.5 ( $p = 0.28$ ) for H9N2 virus, indicating that prevalence of local AIVs was statistically correlated with H7N9 subtype seroprevalence. Our results also revealed that AIV prevalence in the different provinces was a key determinant of seroprevalence in the corresponding poultry workers. However, we did not observe a similar trend with other seroepidemic subtypes.

**Discussion**

We conducted a longitudinal seroepidemiologic study of occupationally exposed poultry workers in China during December 2014–April 2016. We investigated antibody

profiles of 7 AIV subtypes that have crossed the species barrier to infect humans, H5N1, H5N6, H6N1, H7N9, H9N2 and H10N8 subtypes, and H6N6 subtype, which is a potential risk to humans. We assessed seroconversion by analyzing paired serum samples from poultry workers and detecting AIV in poultry and the environment in live poultry markets.

H9N2 virus, which plays a role at the animal–human interface, serves as gene donor for H7N9 and H10N8 viruses that infect humans (27). We used a Y280/G9 lineage antigen isolated in samples from Guangdong Province in 2015 as a reference, and its seroprevalence was higher than all other AIV subtypes in our study (Appendix Figure 4). Previous serologic studies also have reported that this strain’s seroprevalence consistently is higher than other AIV subtypes in most provinces surveyed in China, reflecting the association between prevalent asymptomatic infections and frequent poultry exposure (12,16,28).

Overall, seroprevalence of antibodies against H9N2 subtype in this study was higher than reported in previous serologic studies in China and the seroprevalence was highest in Shandong Province compared with other provinces. Li et al. reported a 3.04% seroprevalence between 2009 and 2011 in occupationally exposed populations (29), and Yu

**Table 4.** Seroconversion and persistently positive findings for avian influenza virus among 652 participants with paired or serial serum samples, China, 2014–2016\*

Subtype	No. (%; 95% CI) participants	
	Seroconversion	Persistently positive
H5N1-SH199	1 (0.2, 0–0.9)	6 (0.9, 0.3–2.0)
H5N1-SC29	5 (0.8, 0.2–1.8)	4 (0.6, 0.2–1.6)
H5N6	3 (0.5, 0.0–1.3)	3 (0.5, 0.0–1.3)
H6N1	6 (0.9, 0.3–2.0)	12 (1.8, 0.8–2.9)
H6N6	4 (0.6, 0.2–1.7)	0 (0.0, 0.0–0.6)
H7N9	9 (1.4, 0.5–2.3)	15 (2.3, 1.1–3.5)
H9N2	23 (3.5, 2.1–4.9)	41 (6.3, 4.4–8.2)
H10N8	0 (0–0.6)	0 (0–0.6)

**Table 5.** Risk analysis for seropositive participants in surveys for avian influenza subtypes among poultry workers, China, 2014–2016\*

Subtypes and variables	Seropositive, no. (%)	Seronegative, no. (%)	p value†	OR (95% CI)	Adjusted OR (95% CI)
<b>H5Nx‡</b>					
Age, y					
<35	8 (10.8)	442 (21.7)	<0.001	Referent	Referent
35–55	40 (54.1)	1,231 (60.4)		1.8 (0.8–3.9)	2.3 (1.0–4.9)
>55	26 (35.1)	366 (18.0)		3.9 (1.8–8.8)	4.7 (2.1–10.7)
Exposed to ducks					
Yes	34 (45.3)	651 (31.8)	0.014	1.8 (1.1–2.8)	1.6 (1.0–2.5)
No	41 (54.7)	1,398 (68.2)		Referent	Referent
Exposed to ill or dead poultry					
Yes	15 (20.0)	221 (10.8)	0.013	2.1 (1.2–3.7)	2.3 (1.3–4.2)
No	60 (80.0)	1826 (89.2)		Referent	Referent
Seropositivity for H1N1(pdm09) virus					
Positive	24 (32.0)	316 (16.4)	<0.001	2.6 (1.6–4.3)	3.1 (1.8–4.5)
Negative	51 (68.0)	1,733 (84.6)		Referent	Referent
<b>H7N9</b>					
Sex					
F	53 (64.6)	924 (45.2)	<0.001	2.2 (1.4–3.5)	2.2 (1.4–3.6)
M	29 (35.4)	1,118 (54.8)		Referent	Referent
Poultry seller§					
Yes	70 (85.4)	1,173 (57.5)	<0.001	4.3 (2.3–8.0)	4.1 (2.2–7.7)
No	12 (14.6)	867 (42.5)		Referent	Referent
No. years of work at live poultry market					
<3	11 (13.4)	561 (27.5)	0.017	Referent	Referent
3–10	46 (56.1)	924 (45.3)		2.0 (1.1–3.5)	1.8 (1.0–3.2)
>10	25 (30.5)	557 (27.3)		1.7 (0.9–3.2)	1.3 (0.7–2.5)
Seropositivity for seasonal H3N2 virus					
Positive	44 (53.7)	743 (36.4)	0.002	2.0 (1.3–3.2)	1.9 (1.2–2.9)
Negative	38 (46.4)	1,299 (63.6)		Referent	Referent
<b>H9N2</b>					
Age, y§					
<35	56 (23.6)	394 (21.0)	0.004	2.1 (1.3–3.4)	1.9 (1.1–3.3)
35–55	156 (65.8)	1,115 (59.4)		2.1 (1.3–3.2)	1.6 (1.0–2.5)
>55	25 (10.6)	367 (19.6)		Referent	Referent
Sex					
F	134 (56.5)	843 (44.7)	<0.001	1.6 (1.2–2.1)	1.6 (1.2–2.1)
M	103 (43.5)	1,044 (55.3)		Referent	Referent
Poultry seller§					
Yes	175 (73.8)	1,068 (56.7)	<0.001	2.2 (1.6–2.9)	1.9 (1.4–2.6)
No	62 (26.2)	817 (43.3)		Referent	Referent
Poultry processor§					
Yes	67 (28.3)	407 (21.6)	0.02	1.4 (1.1–1.9)	1.3 (1.0–1.7)
No	170 (71.7)	1,478 (78.4)		Referent	Referent
No. years of work at live poultry market					
<3	37 (15.6)	535 (28.4)	<0.001	Referent	Referent
3–10	126 (53.2)	844 (44.7)		2.6 (1.8–3.7)	2.4 (1.6–3.5)
>10	74 (31.2)	508 (26.9)		2.2 (1.5–3.2)	3.0 (1.3–3.1)
<b>H6N1</b>					
Seropositivity for H1N1(pdm09) virus					
Positive	19 (35.9)	321 (15.5)	<0.001	3.0 (1.7–5.4)	3.0 (1.7–5.4)
Negative	34 (64.1)	1,750 (84.5)		Referent	Referent
Seropositivity for H9N2 virus					
Positive	13 (24.5)	224 (10.8)	0.002	2.7 (1.4–5.1)	2.6 (1.4–5.0)
Negative	40 (75.5)	1,847 (89.2)		Referent	Referent

\*Results represent only statistically significant factors from analysis of questionnaire data.

†By  $\chi^2$  test.

‡Combined the H5N1-SC29 and H5N6 data.

§Missing data.

et al. reported 4.6% of poultry workers in their study had antibodies against H9N2 virus in 2013 (30). Another previous serologic study in Tai'an, Shandong Province, reported the prevalence of antibodies against H9 subtypes among poultry workers was  $\leq 8.5\%$  during January 2011–December 2013 (31). Because no uniform standard antibody titer cutoff is available for H9N2 seropositivity, we used a

stricter cutoff value for HI titers,  $\geq 1:40$ , and for MN titers,  $\geq 1:80$ , for seroprevalence to avoid overestimation and reduce cross-reactivity with seasonal influenza viruses (32).

The higher seroprevalence in Shandong Province could be explained by 2 possibilities. Participants in this province were all poultry sellers in live poultry markets, an occupation that we noted as a statistically high risk factor



for seroprevalence. Shandong is ranked as the one of the largest egg-producing provinces in China, and it has a high prevalence of H9N2 in local chicken flocks (33,34), which could indicate that more people are exposed to AIV from the poultry industry in general.

Logistic regression analysis of risk factors showed that occupational characteristics might increase risk for infection. Seropositive participant characteristics and related AIV information provided pivotal seroevidence for subclinical AIV infection risk factors. We noted that the participant characteristics female and poultry seller were risk factors for H7N9 and H9N2 infection, which coincides with results of previous studies (18,35). Further risk factor analysis indicated that seropositivity for pH1N1 virus was a risk factor for H5 infections with H5N1 and H5N6 subtypes and for H6N1 infection and that seropositivity for H3N2 subtype was a risk factor for H7N9 infection. In addition, seroprevalence for H6N1 infection also was affected by seropositivity for H9N2 subtypes. Our results might be explained partially by cross-reactivity between HA antigen from different AIV subtypes (36,37). We noted that the prevalence of H7N9 and H9N2 viruses in poultry from local markets was associated closely with seroprevalence for these subtypes in poultry workers. We also noted that the low seasonal influenza vaccination rate (2.8%) in poultry workers might have a limited effect on potential cross-reactions between pH1N1 and H5 subtypes and between H3N2 and H7N9 subtypes.

We observed higher prevalence for certain AIV subtypes and seroprevalence for certain AIV antibodies in live poultry markets, providing further evidence of cross-species transmission from birds to humans. Since the H7N9 outbreaks of 2013, consensus that AIV was transmitted from birds to humans led the government of China to implement epidemic control measures. The measures, such as closing live poultry markets during influenza season, cleaning and disinfecting live poultry markets daily, and vaccinating poultry, have effectively reduced the chances for human exposure to AIV-contaminated environments and ill poultry (38,39). Our results demonstrate that epidemic control measures aimed at live poultry markets, including their closure, can be highly effective in human AIV infection control (9,38).

Many participants with seropositivity were residents of southern and eastern provinces. Several determinants could account for this observation. First, the high density of live poultry markets, high population density, and expansive live poultry transportation network in these regions could favor large-scale and transboundary AIV spread in poultry, thereby increasing the risk for human infection (40). Second, these regions are rich in water resources, including the Yangtze and Pearl Rivers, as well as many lakes, which are natural habitats for waterfowl and wild birds that serve

as natural hosts for various AIV subtypes, including H5Nx and H9N2 viruses, and that continually generate biological threats to public health (41,42). Studies suggest that migratory birds play a role in the evolution and spread of various zoonotic agents, and southeast China is located along the East Asian-Australian flyway, a migratory route for many bird species (43,44).

Our study had several limitations. Despite serious efforts to collect samples from the same participants during follow-up sampling, movement of vendors and poultry workers from target poultry markets reduced the possibility of obtaining paired samples. In addition, the relatively small sample size and use of only 1 location for the control group, Beijing in 2015, could be potential sources of bias.

In conclusion, our study provides serologic evidence of subclinical human AIV infection in an occupationally exposed population of poultry workers and corresponding AIV infection risk factors. Because novel influenza viruses continue to emerge, our results show the need for enhanced etiologic surveillance of AIVs in live poultry markets and humans. Implementing poultry vaccination would also reduce human infection risk. Finally, our results demonstrate the need for active surveillance to foresee dynamic AIV epidemics and inform influenza vaccine development.

### Acknowledgments

We thank the participant poultry workers and staff of the Chinese Center for Disease Control and Prevention at country, prefecture, and provincial levels for making this study possible.

This study was funded by grants from the National Natural Science Foundation of China (NSFC; no. 81502857), the National Grand Project on Prevention and Control of Major Infectious Diseases (no. 2016ZX10004222-003), and the intramural special grant for influenza virus research from the Chinese Academy of Sciences (no. KJZD-EW-L15). W.J.L. is supported by the Excellent Young Scientist Program of the NSFC (grant no. 81822040). G.F.G. is a leading principal investigator of the NSFC Innovative Research Group (grant no. 81621091). The funding bodies played no role in the study design, data collection and analysis, manuscript preparation, or in the decision to publish. The findings and conclusions in this article are those of the authors and do not necessarily represent the official position of the funding agencies.

### About the Author


Dr. Quan was a medical student at the National Institute for Viral Disease Control and Prevention, Chinese Center for Disease Control and Prevention, Beijing, China, during the study. His primary research interests include etiology and serology of influenza viruses and other emerging and reemerging viruses.

## References

- Su S, Bi Y, Wong G, Gray GC, Gao GF, Li S. Epidemiology, evolution, and recent outbreaks of avian influenza virus in China. *J Virol*. 2015;89:8671–6. <https://doi.org/10.1128/JVI.01034-15>
- Gao GF. From “A”IV to “Z”IKV: Attacks from emerging and re-emerging pathogens. *Cell*. 2018;172:1157–9. <https://doi.org/10.1016/j.cell.2018.02.025>
- Huo X, Cui L, Chen C, Wang D, Qi X, Zhou M, et al. Severe human infection with a novel avian-origin influenza A(H7N4) virus. *Sci Bull*. 2018;63:1043–50. <https://doi.org/10.1016/j.scib.2018.07.003>
- Wei SH, Yang JR, Wu HS, Chang MC, Lin JS, Lin CY, et al. Human infection with avian influenza A H6N1 virus: an epidemiological analysis. *Lancet Respir Med*. 2013;1:771–8. [https://doi.org/10.1016/S2213-2600\(13\)70221-2](https://doi.org/10.1016/S2213-2600(13)70221-2)
- Han M, Gu J, Gao GF, Liu WJ. China in action: national strategies to combat against emerging infectious diseases. *Sci China Life Sci*. 2017;60:1383–5. <https://doi.org/10.1007/s11427-017-9141-3>
- Liu D, Shi W, Shi Y, Wang D, Xiao H, Li W, et al. Origin and diversity of novel avian influenza A H7N9 viruses causing human infection: phylogenetic, structural, and coalescent analyses. *Lancet*. 2013;381:1926–32. [https://doi.org/10.1016/S0140-6736\(13\)60938-1](https://doi.org/10.1016/S0140-6736(13)60938-1)
- Liu J, Xiao H, Wu Y, Liu D, Qi X, Shi Y, et al. H7N9: a low pathogenic avian influenza A virus infecting humans. *Curr Opin Virol*. 2014;5:91–7. <https://doi.org/10.1016/j.coviro.2014.03.001>
- Quan C, Huang T, Chen X, Zhang J, Wang Q, Zhang C, et al. Genomic characterizations of H4 subtype avian influenza viruses from live poultry markets in Sichuan Province of China, 2014–2015. *Sci China Life Sci*. 2018;61:1123–6. <https://doi.org/10.1007/s11427-018-9327-4>
- Gao GF. Influenza and the live poultry trade. *Science*. 2014;344:235. <https://doi.org/10.1126/science.1254664>
- Cui B, Liao Q, Lam WWT, Liu ZP, Fielding R. Avian influenza A/H7N9 risk perception, information trust and adoption of protective behaviours among poultry farmers in Jiangsu Province, China. *BMC Public Health*. 2017;17:463–76. <https://doi.org/10.1186/s12889-017-4364-y>
- Bai T, Zhou J, Shu Y. Serologic study for influenza A (H7N9) among high-risk groups in China. *N Engl J Med*. 2013;368:2339–40. <https://doi.org/10.1056/NEJMc1305865>
- Wang M, Fu C-X, Zheng B-J. Antibodies against H5 and H9 avian influenza among poultry workers in China. *N Engl J Med*. 2009;360:2583–4. <https://doi.org/10.1056/NEJMc0900358>
- Ma C, Cui S, Sun Y, Zhao J, Zhang D, Zhang L, et al. Avian influenza A (H9N2) virus infections among poultry workers, swine workers, and the general population in Beijing, China, 2013–2016: A serological cohort study. *Influenza Other Respir Viruses*. 2019;13:415–25. <https://doi.org/10.1111/irv.12641>
- Xin L, Bai T, Zhou JF, Chen YK, Li XD, Zhu WF, et al. Seropositivity for avian influenza H6 virus among humans, China. *Emerg Infect Dis*. 2015;21:1267–9. <https://doi.org/10.3201/eid2107.150135>
- Sikkema RS, Freidl GS, de Bruin E, Koopmans M. Weighing serological evidence of human exposure to animal influenza viruses—a literature review. *Euro Surveill*. 2016;21:30388. <https://doi.org/10.2807/1560-7917.ES.2016.21.44.30388>
- Khan SU, Anderson BD, Heil GL, Liang S, Gray GC. A systematic review and meta-analysis of the seroprevalence of influenza A(H9N2) infection among humans. *J Infect Dis*. 2015;212:562–9. <https://doi.org/10.1093/infdis/jiv109>
- Yang S, Chen Y, Cui D, Yao H, Lou J, Huo Z, et al. Avian-origin influenza A(H7N9) infection in influenza A(H7N9)-affected areas of China: a serological study. *J Infect Dis*. 2014;209:265–9. <https://doi.org/10.1093/infdis/jit430>
- Wang X, Fang S, Lu X, Xu C, Cowling BJ, Tang X, et al. Seroprevalence to avian influenza A(H7N9) virus among poultry workers and the general population in southern China: a longitudinal study. *Clin Infect Dis*. 2014;59:e76–83. <https://doi.org/10.1093/cid/ciu399>
- Li LH, Yu Z, Chen WS, Liu SL, Lu Y, Zhang YJ, et al. Evidence for H5 avian influenza infection in Zhejiang province, China, 2010–2012: a cross-sectional study. *J Thorac Dis*. 2013;5:790–6. <https://doi.org/10.3978/j.issn.2072-1439.2013.12.45>
- World Health Organization. Manual for the laboratory diagnosis and virological surveillance of influenza. Geneva: The Organization; 2011 [cited 2019 Feb 10]. [https://www.who.int/influenza/gisrs\\_laboratory/manual\\_diagnosis\\_surveillance\\_influenza/en](https://www.who.int/influenza/gisrs_laboratory/manual_diagnosis_surveillance_influenza/en)
- Qi W, Su S, Xiao C, Zhou P, Li H, Ke C, et al. Antibodies against H10N8 avian influenza virus among animal workers in Guangdong Province before November 30, 2013, when the first human H10N8 case was recognized. *BMC Med*. 2014;12:205. <https://doi.org/10.1186/s12916-014-0205-3>
- Gomaa MR, Kayed AS, Elabd MA, Zeid DA, Zaki SA, El Rifay AS, et al. Avian influenza A(H5N1) and A(H9N2) seroprevalence and risk factors for infection among Egyptians: a prospective, controlled seroepidemiological study. *J Infect Dis*. 2015;211:1399–407. <https://doi.org/10.1093/infdis/jiu529>
- Liu WJ, Tan S, Zhao M, Quan C, Bi Y, Wu Y, et al. Cross-immunity against avian influenza A(H7N9) virus in the healthy population is affected by antigenicity-dependent substitutions. *J Infect Dis*. 2016;214:1937–46. <https://doi.org/10.1093/infdis/jiw471>
- Pawar SD, Tandale BV, Raut CG, Parkhi SS, Barde TD, Gurav YK, et al. Avian influenza H9N2 seroprevalence among poultry workers in Pune, India, 2010. *PLoS One*. 2012;7:e36374. <https://doi.org/10.1371/journal.pone.0036374>
- Bi Y, Chen Q, Wang Q, Chen J, Jin T, Wong G, et al. Genesis, evolution and prevalence of H5N6 avian influenza viruses in China. *Cell Host Microbe*. 2016;20:810–21. <https://doi.org/10.1016/j.chom.2016.10.022>
- Horby PW, Laurie KL, Cowling BJ, Engelhardt OG, Sturm-Ramirez K, Sanchez JL, et al.; CONSISE Steering Committee. CONSISE statement on the reporting of seroepidemiologic studies for influenza (ROSES-I statement): an extension of the STROBE statement. *Influenza Other Respir Viruses*. 2017;11:2–14. <https://doi.org/10.1111/irv.12411>
- Sun Y, Liu J. H9N2 influenza virus in China: a cause of concern. *Protein Cell*. 2015;6:18–25. <https://doi.org/10.1007/s13238-014-0111-7>
- Wang Q, Ju L, Liu P, Zhou J, Lv X, Li L, et al. Serological and virological surveillance of avian influenza A virus H9N2 subtype in humans and poultry in Shanghai, China, between 2008 and 2010. *Zoonoses Public Health*. 2015;62:131–40. <https://doi.org/10.1111/zph.12133>
- Li X, Tian B, Jianfang Z, Yongkun C, Xiaodan L, Wenfei Z, et al. A comprehensive retrospective study of the seroprevalence of H9N2 avian influenza viruses in occupationally exposed populations in China. *PLoS One*. 2017;12:e0178328. <https://doi.org/10.1371/journal.pone.0178328>
- Yu Q, Liu L, Pu J, Zhao J, Sun Y, Shen G, et al. Risk perceptions for avian influenza virus infection among poultry workers, China. *Emerg Infect Dis*. 2013;19:313–6. <https://doi.org/10.3201/eid1901.120251>
- Li S, Zhou Y, Song W, Pang Q, Miao Z. Avian influenza virus H9N2 seroprevalence and risk factors for infection in occupational poultry-exposed workers in Tai'an of China. *J Med Virol*. 2016;88:1453–6. <https://doi.org/10.1002/jmv.24483>
- World Health Organization. Recommendations and laboratory procedures for detection of avian influenza A(H5N1) virus in specimens from suspected human cases. Geneva: The Organization; 2007. [cited 2019 Aug 12]. [https://www.who.int/influenza/resources/documents/h5n1\\_laboratory\\_procedures](https://www.who.int/influenza/resources/documents/h5n1_laboratory_procedures)

33. National Bureau of Statistics of China. Statistical Yearbook of China [in Chinese]. Beijing: The Bureau; 2014. [cited 2019 Aug 12]. <http://www.stats.gov.cn/tjsj/ndsj/2014/indexch.htm>
34. Li Y, Liu M, Sun Q, Zhang H, Zhang H, Jiang S, et al. Genotypic evolution and epidemiological characteristics of H9N2 influenza virus in Shandong Province, China. *Poult Sci*. 2019;98(9):3488-95. PubMed <https://doi.org/10.3382/ps/pez151>
35. Ma MJ, Zhao T, Chen SH, Xia X, Yang XX, Wang GL, et al. Avian influenza A virus infection among workers at live poultry markets, China, 2013–2016. *Emerg Infect Dis*. 2018;24:1246–56. <https://doi.org/10.3201/eid2407.172059>
36. Khuntirat BP, Yoon IK, Blair PJ, Krueger WS, Chittaganpitch M, Putnam SD, et al. Evidence for subclinical avian influenza virus infections among rural Thai villagers. *Clin Infect Dis*. 2011; 53:e107–16. <https://doi.org/10.1093/cid/cir525>
37. Kallewaard NL, Corti D, Collins PJ, Neu U, McAuliffe JM, Benjamin E, et al. Structure and function analysis of an antibody recognizing all influenza A subtypes. *Cell*. 2016;166:596–608. <https://doi.org/10.1016/j.cell.2016.05.073>
38. Yuan J, Lau EH, Li K, Leung YH, Yang Z, Xie C, et al. Effect of live poultry market closure on avian influenza A(H7N9) virus activity in Guangzhou, China, 2014. *Emerg Infect Dis*. 2015;21:1784–93. <https://doi.org/10.3201/eid2110.150623>
39. Peiris JSM, Cowling BJ, Wu JT, Feng L, Guan Y, Yu H, et al. Interventions to reduce zoonotic and pandemic risks from avian influenza in Asia. *Lancet Infect Dis*. 2016;16:252–8. [https://doi.org/10.1016/S1473-3099\(15\)00502-2](https://doi.org/10.1016/S1473-3099(15)00502-2)
40. Fournié G, Guitian J, Desvaux S, Cuong VC, Dung H, Pfeiffer DU, et al. Interventions for avian influenza A (H5N1) risk management in live bird market networks. *Proc Natl Acad Sci U S A*. 2013;110:9177–82. <https://doi.org/10.1073/pnas.1220815110>
41. Wang D, Yang L, Zhu W, Zhang Y, Zou S, Bo H, et al. Two outbreak sources of influenza A (H7N9) viruses have been established in China. *J Virol*. 2016;90:5561–73. <https://doi.org/10.1128/JVI.03173-15>
42. Gao GF. For a better world: Biosafety strategies to protect global health. *Biosaf Health*. 2019;1(1):1-3. <https://doi.org/10.1016/j.bsheal.2019.03.001>
43. Tian H, Zhou S, Dong L, Van Boeckel TP, Cui Y, Newman SH, et al. Avian influenza H5N1 viral and bird migration networks in Asia. *Proc Natl Acad Sci U S A*. 2015;112:172–7. <https://doi.org/10.1073/pnas.1405216112>
44. Bi Y, Shi W, Chen J, Chen Q, Ma Z, Wong G, et al. CASCIRE surveillance network and work on avian influenza viruses. *Sci China Life Sci*. 2017;60:1386–91. <https://doi.org/10.1007/s11427-017-9251-2>

Address for correspondence: George F. Gao or William J. Liu, Chinese Center for Disease Control and Prevention, 155 Changbai Rd, Changping District, Beijing 102206, China; email: gaofu@chinacdc.cn or liujun@ivdc.chinacdc.cn; Hongjie Yu, Fudan University School of Public Health, No. 138 Yixueyuan Rd, Xuhui District, Shanghai, 200032, China; email: yhj@fudan.edu.cn



**EMERGING  
INFECTIOUS DISEASES**

**December 2018**

# Zoonotic Infections

- Outbreak of HIV Infection Linked to Nosocomial Transmission, China, 2016–2017
- Autochthonous Human Case of Seoul Virus Infection, the Netherlands.
- Reemergence of St. Louis Encephalitis Virus in the Americas
- Spatial Analysis of Wildlife Tuberculosis Based on a Serologic Survey Using Dried Blood Spots, Portugal
- Comparison of Highly Pathogenic Avian Influenza H5 Guangdong Lineage Epizootic in Europe (2016–17) with Previous HPAI H5 Epizootics
- *Capnocytophaga canimorsus* Capsular Serovar and Disease Severity, Helsinki, Finland, 2000–2017
- Rat Lungworm Infection in Rodents Across Post-Katrina New Orleans, Louisiana
- Crimean-Congo Hemorrhagic Fever Virus, Mongolia, 2013–2014
- Emerging Multidrug-Resistant Hybrid Pathotype Shiga toxin–producing *Escherichia coli* O80 and Related Strains of Clonal Complex 165, Europe.
- Terrestrial Bird Migration and West Nile Virus Circulation, United States
- Substance Use and Adherence to HIV Preexposure Prophylaxis for Men Who Have Sex with Men
- Genomic Characterization of  $\beta$ -Glucuronidase–Positive *Escherichia coli* O157:H7 Producing Stx2a
- Highly Pathogenic Clone of Shiga Toxin–Producing *Escherichia coli* O157:H7, England and Wales
- CTX-M-65 Extended-Spectrum  $\beta$ -Lactamase–Producing *Salmonella enterica* Serotype Infantis, United States
- Novel Type of Chronic Wasting Disease Detected in Moose (*Alces alces*), Norway
- Survey of Ebola Viruses in Frugivorous and Insectivorous Bats in Guinea, Cameroon, and the Democratic Republic of the Congo, 2015–2017
- Prevalence of Avian Influenza A(H5) and A(H9) Viruses in Live Bird Markets, Bangladesh
- Rat Hepatitis E Virus as a Cause of Persistent Hepatitis after Liver Transplant
- Influences of Community Interventions on Zika Prevention Behaviors of Pregnant Women, Puerto Rico, July 2016–June 2017
- Emergent Sand Fly–Borne Phleboviruses in the Balkan Region

To revisit the December 2018 issue, go to:

<https://wwwnc.cdc.gov/eid/articles/issue/24/12/table-of-contents>

# Avian Influenza A Viruses among Occupationally Exposed Populations, China, 2014–2016

## Appendix

### Supplementary Methods

#### Selection of Study Regions

Shanghai and the 6 provinces selected for study all had previously reported human infections with avian influenza virus (AIV) or outbreaks of AIVs in poultry; a high density of poultry and population; were located in different key regions of China, Jiangsu and Shanghai in the east, Sichuan in the west, Jiangxi and Guangdong in the south, and Henan and Shandong in the north; and had a local Chinese Center for Disease Control and Prevention willing to take part in this study.

#### Questionnaire

We developed a structured questionnaire and administered it to each participant during a face-to-face interview after obtaining their consent to participate in the study. Questionnaire information included demographic characteristics (name, sex, age, work address); telephone number; occupational exposure history variables, including type of exposure (processing, selling, transportation, feeding and others), duration of work exposure to poultry, and which species of poultry they were exposed to; status of seasonal influenza vaccination within last 1 year; whether participant had influenza-like illness within 1 month.

### **Analysis of Antigenic Characteristic**

Based on the antigenic and genetic characteristics of avian influenza A(H5) reported by the World Health Organization (WHO) and previous studies (1–3), viruses circulating and characterized during September 2014–September 2016 in mainland China belonged to clades 2.3.2.1c and 2.3.4.4 of HA genes. The 2.3.2.1c clade contained 2 candidate virus strains A/chicken/Ghana/20/2015 and A/chicken/Guiyang/1153/2016. Both viruses are located in different groups and have different antigenic characteristics. We selected an A/chicken/Shanghai/02.12 HZ199-P/2015 (H5N1-SH199) strain based on the genetic similarity. Although A(H5) 2.3.4.4 clades are the more prevalent strains and A/Sichuan/26221/2014 is the recommended reference strain, this clade also could be divided into 2 groups. Based on the prevalence of viruses, we selected A/pigeon/Sichuan/NCXN29/2014 (H5N1-SC29) and A/duck/Guangdong/04.22DGCP069-O/2015 (H5N6) for testing antigens, which were slightly different (Appendix, Figures 1 and 2).

According to data published by WHO, the available A(H7N9) candidate influenza vaccine virus is A/Anhui/1/2013 or A/Shanghai/2/2013. Additionally, prevalent viruses of H7N9 in China have similar antigenicity, according to a report by Wang, et al. (4). Therefore, the antigens used in our study could represent the prevalent H7N9 virus (Appendix, Figure 3).

According to data published by WHO, the available influenza A(H9N2) candidate vaccine virus is the A/chicken/Hong Kong/G9/1997 or derivative virus A/Hong Kong/308/2014-like (1). Additionally, the G9 clade could be divided into subgroups depending on antigenicity as reported by Li, et al. (5). Therefore, the antigens we used could represent the prevalent H9N2 virus (Appendix, Figure 4). On the basis of previous studies (6,7), antigens of H6Nx were similar, so we believed viruses used in this study could cover prevalent virus subtypes (Appendix, Figure 5).

## **Molecular and Phylogenetic Analyses**

Through the antigenic analysis of the different AIV subtypes based on the reports and phylogenetic methods described previously (1), we performed a detailed phylogenetic analysis for HA gene segments with other prevailing AIVs. We concluded that selected strains were representative (Appendix, Figures 1–5).

## **Hemagglutination Inhibition (HI) Assay**

We inactivated antigens by using 0.05%  $\beta$ -propiolactone for HI assay. We treated 100  $\mu$ L serum samples with 300  $\mu$ L of receptor-destroying enzyme (RDE) and incubated in a water bath at 37°C for 16–18h. Then, we inactivated RDE(II) (Denka Seiken Co. Ltd, <https://denka-seiken.com>) at 56°C for 30 minutes before adding 100  $\mu$ L phosphate buffer saline. To remove nonspecific inhibitors, we added 25  $\mu$ L of packed chicken red blood cells (RBCs) to 500  $\mu$ L of the mixtures. We incubated solutions in a 37°C water bath for 1h.

We performed serial 2-fold dilutions of RDE-treated serum from 1:10–1:160 in 25  $\mu$ L phosphate buffer saline a 96-well microtitration plate. We added 25  $\mu$ L of 4 haemagglutinin unit antigens and incubated at room temperature for 45 min. Then we added 50  $\mu$ L of the RBC solution and incubated again for 45 minutes at room temperature. We considered titers accurate when hemagglutination was completely inhibited. We used negative and positive serum control samples (rabbit antibodies against the specific virus antigen) for each assay. We took titers as the reciprocal of the dilution levels of the wells. We assigned final titers of <1:10 a value of 1:5.

## **Microneutralization (MN) Assay**

We heat inactivated serum samples at 56°C for 30 minutes and then conducted serial 2-fold dilutions from 1:20–1:640 in triplicate. We diluted equal volumes of heat-inactivate serum and virus to the 50% tissue culture infective dose (TCID<sub>50</sub>) at 200, added these together, and incubated at 37°C for 1 h. We used the Reed-Muench method to determine the TCID<sub>50</sub>/100  $\mu$ L. We transferred the mixture to a confluent layer of Madin-Darby canine kidney cells and

incubated for 60 h at 37°C with 5% CO<sub>2</sub>. Then we tested virus hemagglutination activity in 0.5% RBCs and considered absence of hemagglutination as a positive result for antibodies to the antigen (8). We used rabbit antibodies raised against homologous viruses as positive controls for assays. In each assay, we used negative, positive, and cell control serum, and virus back titration. We used the same negative and positive controls used in the MN assay that we used in HI assays. We defined the microneutralization titer as the highest dilution of serum that completely inhibited absence of hemagglutination in 50% of the wells. We assigned final titers <1:20 a value of 1:10.

## References

1. World Health Organization. Antigenic and genetic characteristics of zoonotic influenza viruses and candidate vaccine viruses developed for potential use in human vaccines. Geneva: The Organization; 2019 Feb 21. [cited 2019 Aug 13].  
[http://www.who.int/influenza/vaccines/virus/characteristics\\_virus\\_vaccines/en](http://www.who.int/influenza/vaccines/virus/characteristics_virus_vaccines/en)
2. Bi Y, Chen Q, Wang Q, Chen J, Jin T, Wong G, et al. Genesis, evolution and prevalence of H5N6 avian influenza viruses in China. *Cell Host Microbe*. 2016;20:810–21. [PubMed](#)  
<https://doi.org/10.1016/j.chom.2016.10.022>
3. Yang L, Zhu W, Li X, Bo H, Zhang Y, Zou S, et al. Genesis and dissemination of highly pathogenic H5N6 avian influenza viruses. *J Virol*. 2017;91:e02199-16. [PubMed](#)  
<https://doi.org/10.1128/JVI.02199-16>
4. Wang D, Yang L, Zhu W, Zhang Y, Zou S, Bo H, et al. Two outbreak sources of influenza A (H7N9) viruses have been established in China. *J Virol*. 2016;90:5561–73. [PubMed](#)  
<https://doi.org/10.1128/JVI.03173-15>

5. Li X, Shi J, Guo J, Deng G, Zhang Q, Wang J, et al. Genetics, receptor binding property, and transmissibility in mammals of naturally isolated H9N2 Avian Influenza viruses. *PLoS Pathog.* 2014;10:e1004508. [PubMed https://doi.org/10.1371/journal.ppat.1004508](https://doi.org/10.1371/journal.ppat.1004508)
6. Xin L, Bai T, Zhou JF, Chen YK, Li XD, Zhu WF, et al. Seropositivity for avian influenza H6 virus among humans, China. *Emerg Infect Dis.* 2015;21:1267–9. [PubMed https://doi.org/10.3201/eid2107.150135](https://doi.org/10.3201/eid2107.150135)
7. Wang G, Deng G, Shi J, Luo W, Zhang G, Zhang Q, et al. H6 influenza viruses pose a potential threat to human health. *J Virol.* 2014;88:3953–64. [PubMed https://doi.org/10.1128/JVI.03292-13](https://doi.org/10.1128/JVI.03292-13)
8. Liu WJ, Tan S, Zhao M, Quan C, Bi Y, Wu Y, et al. Cross-immunity against avian influenza A(H7N9) virus in the healthy population is affected by antigenicity-dependent substitutions. *J Infect Dis.* 2016;214:1937–46. [PubMed https://doi.org/10.1093/infdis/jiw471](https://doi.org/10.1093/infdis/jiw471)
9. Wang M, Fu CX, Zheng BJ. Antibodies against H5 and H9 avian influenza among poultry workers in China. *N Engl J Med.* 2009;360:2583–4. [PubMed https://doi.org/10.1056/NEJMc0900358](https://doi.org/10.1056/NEJMc0900358)
10. Bai T, Zhou J, Shu Y. Serologic study for influenza A (H7N9) among high-risk groups in China. *N Engl J Med.* 2013;368:2339–40. [PubMed https://doi.org/10.1056/NEJMc1305865](https://doi.org/10.1056/NEJMc1305865)
11. Gomaa MR, Kayed AS, Elabd MA, Zeid DA, Zaki SA, El Rifay AS, et al. Avian influenza A(H5N1) and A(H9N2) seroprevalence and risk factors for infection among Egyptians: a prospective, controlled seroepidemiological study. *J Infect Dis.* 2015;211:1399–407. [PubMed https://doi.org/10.1093/infdis/jiu529](https://doi.org/10.1093/infdis/jiu529)
12. Chen Y, Zheng Q, Yang K, Zeng F, Lau SY, Wu WL, et al. Serological survey of antibodies to influenza A viruses in a group of people without a history of influenza vaccination. *Clin Microbiol Infect.* 2011;17:1347–9. [PubMed https://doi.org/10.1111/j.1469-0691.2011.03538.x](https://doi.org/10.1111/j.1469-0691.2011.03538.x)



13. Qi W, Su S, Xiao C, Zhou P, Li H, Ke C, et al. Antibodies against H10N8 avian influenza virus among animal workers in Guangdong Province before November 30, 2013, when the first human H10N8 case was recognized. *BMC Med.* 2014;12:205. [PubMed https://doi.org/10.1186/s12916-014-0205-3](https://doi.org/10.1186/s12916-014-0205-3)

**Appendix Table 1.** Avian influenza A virus subtypes, reference strains, cutoff values for hemagglutinin and microneutralization assays, and reference studies for an investigation of avian influenza viruses among poultry workers, China, 2014–2016\*

Subtypes	Reference strains	HI cutoff	MN cutoff	Reference no., study location
A(H1N1)pdm09	A/California/04/2009	40	80	8, China
H3N2	A/Beijing/CAS0001/2007	40	80	8, China
H5N1	A/chicken/Shanghai/02.12 HZ199-P/2015 (SH199)	20	NA	9, China
H5N1	A/pigeon/Sichuan/NCXN29/2014 (SC29)	20	NA	9, China
H5N6	A/duck/Guangdong/04.22 DGCP069-O/2015	20	NA	9, China
H6N1	A/Taiwan/2/2013	20	20	6, China
H6N6	A/duck/Guangxi/04.10 JX031/2015	20	20	6, China
H7N9	A/chicken/Guangdong/04.22 DGCP098-O/2015	20	40	10, China
H9N2	A/chicken/Guangdong/04.15 SZBAXQ005/2015	40	80	11, Egypt; 12, China
H10N8	A/chicken/Jiangxi/B18/2014	20	20	13, China

\*HI, hemagglutinin assay; MN, microneutralization assay; NA, not applicable.

**Appendix Table 2.** Seroprevalence of avian influenza A virus subtypes in the control group and study population of occupationally-exposed poultry workers\*

Subtypes	Participants	Seroprevalence, no.(%)	p value
H5N1-SH199	Control group	3 (1.4)	0.2
	Poultry workers	28 (1.3)	
H5N1-SC29/H5N6	Control group	3 (1.4)	0.04
	Poultry workers	75 (3.5)	
H6N1	Control group	2 (0.9)	0.07
	Poultry workers	53 (2.5)	
H6N6	Control group	0 (0)	0.5
	Poultry workers	8 (0.4)	
H7N9	Control group	0 (0)	<0.01
	Poultry workers	82 (3.9)	
H9N2	Control group	8 (3.7)	<0.001
	Poultry workers	237 (11.2)	
A(H1N1)pdm09	Control group	47 (23.6)	0.03
	Poultry workers	340 (16.0)	
H3N2	Control group	75 (34.7)	0.6
	Poultry workers	773 (36.4)	

\*No. of general population is 216; No. of poultry workers is 2124.

**Appendix Table 3.** Participants with seroconversion and persistently positive titers for avian influenza H5N1-SH199, China, 2014–2016\*

Subject ID	Age, y/sex	Occupation	Length of exposure, y	Dec 2014		Apr 2015		Dec 2015		Apr 2016	
				HI	MN	HI	MN	HI	MN	HI	MN
<b>Seroconversion</b>											
15.8SH14	47, F	Seller	16	NA <sup>†</sup>	NA	<b>5</b>	<b>ND</b>	<b>20</b>	<b>40</b>	NA	NA
<b>Persistently positive</b>											
14.11JS124	57, M	Seller	30	20	20	20	20	NA	NA	NA	NA
14.11JS31	30, F	Processor, seller	3	20	40	40	40	NA	NA	NA	NA
14.12GD72	34, M	Seller	5	20	160	20	160	20	160	20	160
16.1SD031	33, F	Processor, seller	10	NA	NA	NA	NA	20	40	20	40
16.1SD042	58, M	Processor, seller	26	NA	NA	NA	NA	20	80	20	80
16.1JS24	65, F	Feeder	2	NA	NA	NA	NA	20	40	20	40

\*Bold text represents titers exhibiting seroconversion. HI, hemagglutinin inhibition assay; ID, identification; MN, microneutralization assay; NA, not available; ND, not detected.

**Appendix Table 4.** Participants with seroconversion and persistently positive titers for avian influenza H5N1-SC29, China, 2014–2016\*

Subject ID	Age, y/sex	Occupation	Length of exposure, y	Dec 2014		Apr 2015		Dec 2015		Apr 2016	
				HI	MN	HI	MN	HI	MN	HI	MN
<b>Seroconversion</b>											
14.11SC60	61/M	Seller	5	<b>5</b>	<b>ND</b>	<b>20</b>	<b>40</b>	NA <sup>†</sup>	NA	NA	NA
14.11SC64	62/M	Seller	2	<b>5</b>	<b>ND</b>	<b>20</b>	<b>40</b>	NA	NA	NA	NA
14.11SC55	52/F	Seller	6	<b>10</b>	<b>40</b>	<b>40</b>	<b>80</b>	NA	NA	NA	NA
14.11JSWX54	48/F	Seller	24	<b>5</b>	<b>ND</b>	<b>20</b>	<b>80</b>	NA	NA	NA	NA
16.1SD76	39/M	Processor, seller	17	NA	NA	NA	NA	<b>5</b>	<b>ND</b>	<b>20</b>	<b>40</b>
<b>Persistently positive</b>											
14.11JS73	45/F	Seller	10	20	20	20	40	NA	NA	NA	NA
14.11JSWX39	43/F	Others	20	20	80	40	40	NA	NA	NA	NA
14.11SC86	50/M	Seller, transporter	5	20	40	20	40	NA	NA	NA	NA
14.11JSWX40	49/M	Other	30	20	40	20	40	NA	NA	NA	NA

\*Bold text represents titers exhibiting seroconversion. HI, hemagglutinin inhibition assay; ID, identification; MN, microneutralization assay; NA, not available; ND, not detected.

**Appendix Table 5.** Participants with seroconversion and persistently positive titers for avian influenza H5N6, China, 2014–2016\*

Participant ID	Age,y/sex	Occupation	Length of exposure, y	Dec 2014		Apr 2015		Dec 2015		Apr 2016	
				HI	MN	HI	MN	HI	MN	HI	MN
<b>Seroconversion</b>											
14.11JS67	50, F	Seller	10	<b>5</b>	<b>ND</b>	<b>20</b>	<b>20</b>	NA	NA	NA	NA
14.11SC85	57, F	Seller, transporter	30	<b>5</b>	<b>ND</b>	<b>20</b>	<b>20</b>	NA	NA	NA	NA
14.12GD72	34, M	Processor, seller	5	10	ND	10	ND	<b>5</b>	<b>ND</b>	<b>80</b>	<b>40</b>
<b>Persistently positive</b>											
14.11SC98	61, M	Processor	14	20	40	20	40	NA	NA	NA	NA
14.11JS123	58, F	Processor	4	20	80	20	160	NA	NA	NA	NA
14.11JS122	43, F	Other	2	20	20	20	20	NA	NA	NA	NA

\*Bold text represents titers exhibiting seroconversion. HI, hemagglutinin inhibition assay; ID, identification; MN, microneutralization assay; NA, not available; ND, not detected.

**Appendix Table 6.** Participants with seroconversion and persistently positive titers for avian influenza H6N1, China, 2014–2016\*

Participant ID	Age, y/sex	Occupation	Length of exposure, y	Dec 2014		Apr 2015		Dec 2015		Apr 2016	
				HI	MN	HI	MN	HI	MN	HI	MN
<b>Seroconversion</b>											
14.11JS82	47/F	Feeder	10	<b>5</b>	<b>ND</b>	<b>20</b>	<b>40</b>	NA	NA	NA	NA
14.12HN91	40/M	Other	5	<b>5</b>	<b>ND</b>	<b>40</b>	<b>80</b>	NA	NA	NA	NA
14.12 GD7	61/M	Other	2	5	ND	<b>5</b>	<b>ND</b>	<b>20</b>	<b>160</b>	NA	NA
15.5JS1	59/F	Feeder	23	<b>5</b>	<b>ND</b>	<b>20</b>	<b>40</b>	NA	NA	NA	NA
16.1JS112	59/M	Seller	13	NA	NA	NA	NA	<b>5</b>	<b>ND</b>	<b>20</b>	<b>40</b>
16.1JS54	28/M	Feeder	3	NA	NA	NA	NA	<b>5</b>	<b>ND</b>	<b>20</b>	<b>80</b>
<b>Persistently positive</b>											
14.11JS39	43/F	Other	20	20	40	20	40	NA	NA	NA	NA
14.11JS6	40/F	Processor, seller	12	40	320	20	80	NA	NA	NA	NA
14.11JSWX06	27/M	Seller	5	40	80	40	80	NA	NA	NA	NA
14.11JS99	59/M	Feeder	2	40	40	40	40	NA	NA	NA	NA
14.11JS102	33/M	Feeder	3	40	80	40	40	NA	NA	NA	NA
14.11JS12	60/M	Processor, seller	13	40	40	40	40	NA	NA	NA	NA
14.11JS85	50/M	Feeder	6	20	80	20	80	NA	NA	NA	NA
14.11JSWX49	43/F	Selling	24	20	40	20	40	NA	NA	NA	NA
16.1JS24	65/F	Feeding	2	NA	NA	NA	NA	20	160	20	160
14.12 GD 09	54/F	Seller	4	40	20	NA	NA	10	ND	20	20
14.12GD72	34/M	Seller	5	160	40	160	40	80	20	5	ND
14.12GD115	28/F	Seller	3	40	40	40	40	20	80	20	80

\*Bold text represents titers exhibiting seroconversion. HI, hemagglutinin inhibition assay; ID, identification; MN, microneutralization assay; NA, not available; ND, not detected.

**Appendix Table 7.** Participants with seroconversion for avian influenza H6N6, China, 2014–2016\*

Participant ID	Age,		Length of exposure, y	Dec 2014		Apr 2015		Dec 2015		Apr 2016	
	y/sex	Occupation		HI	MN	HI	MN	HI	MN	HI	MN
15.8SH2	62/M	Seller	32	NA	NA	<b>5</b>	<b>ND</b>	<b>20</b>	<b>40</b>	NA	NA
15.8SH46	35/M	Seller	3	NA	NA	<b>5</b>	<b>ND</b>	<b>20</b>	<b>40</b>	NA	NA
15.8SH46	66/F	Seller	8	NA	NA	<b>5</b>	<b>ND</b>	<b>20</b>	<b>40</b>	NA	NA
16.1JS24	65/M	Seller	2	NA	NA	NA	NA	<b>5</b>	<b>ND</b>	<b>20</b>	<b>40</b>

\*Bold text represents titers exhibiting seroconversion. HI, hemagglutinin inhibition assay; ID, identification; MN, microneutralization assay; NA, not available; ND, not detected.

**Appendix Table 8.** Participants with seroconversion and persistently positive titers for avian influenza H7N9, China, 2014–2016\*

Participant ID	Age, y/sex	Occupation	Length of exposure, y	Dec 2014		Apr 2015		Dec 2015		Apr 2016	
				HI	MN	HI	MN	HI	MN	HI	MN
<b>Seroconversion</b>											
14.11JS79	46/F	Feeder	10	<b>5</b>	<b>ND</b>	<b>40</b>	<b>20</b>	NA	NA	NA	NA
14.11JS77	59/F	Feeder	23	<b>5</b>	<b>ND</b>	<b>80</b>	<b>20</b>	NA	NA	NA	NA
14.11JS108	54/M	Transporter	10	<b>5</b>	<b>ND</b>	<b>40</b>	<b>40</b>	NA	NA	NA	NA
14.11GD63	23/M	Seller	5	<b>5</b>	<b>ND</b>	<b>20</b>	<b>20</b>	5	ND	NA	NA
14.12SH6	29/M	Seller	5	<b>5</b>	<b>ND</b>	NA	NA	NA	NA	<b>40</b>	<b>20</b>
15.8SH48	31/M	Seller	2	NA	NA	<b>5</b>	<b>ND</b>	<b>20</b>	<b>20</b>	NA	NA
15.8SH25	23/M	Seller	3	<b>5</b>	<b>ND</b>	<b>20</b>	<b>40</b>	NA	NA	NA	NA
16.1SD32	46/F	Processor, seller	20	NA	NA	NA	NA	<b>10</b>	<b>ND</b>	<b>40</b>	<b>320</b>
16.1JS110	27/M	Seller	13	NA	NA	NA	NA	<b>5</b>	<b>ND</b>	<b>20</b>	<b>80</b>
<b>Persistently positive</b>											
14.11JS73	45/M	Seller	10	20	160	20	160	NA	NA	NA	NA
14.11JS22	44/M	Seller	8	40	80	40	80	NA	NA	NA	NA
14.11JS11	29/M	Seller	10	20	40	20	80	NA	NA	NA	NA
14.11JX100	46/F	Seller	10	20	80	20	80	NA	NA	NA	NA
14.12HN48	47/F	Seller	18	20	160	20	80	NA	NA	NA	NA
14.11JX60	42/F	Seller	3	80	40	80	40	NA	NA	NA	NA
14.11JS27	34/F	Processor, seller	7	20	40	20	40	NA	NA	NA	NA
14.11JSWX25	47/F	Seller	8	20	40	40	40	NA	NA	NA	NA
14.11JSWX7	35/F	Seller	5	80	80	80	160	NA	NA	NA	NA
14.11JSWX11	47/F	Seller	10	20	80	20	80	NA	NA	NA	NA
14.11JS76	56/F	Feeder	22	80	40	40	40	NA	NA	NA	NA
14.12HN52	46/F	Seller	6	40	320	80	160	NA	NA	NA	NA
16.1JS24	46/F	Feeder	2	NA	NA	NA	NA	20	160	40	160
15.1SH23	37/F	Seller	8	20	20	NA	NA	NA	NA	40	40
14.12GD1	47/F	Seller	18	20	40	20	40	5	ND	5	ND

\*Bold text represents titers exhibiting seroconversion. HI, hemagglutinin inhibition assay; ID, identification; MN, microneutralization assay; NA, not available; ND, not detected.

**Appendix Table 9.** Participants with seroconversion and persistently positive titers for avian influenza H9N2, China, 2014–2016\*

Participant ID	Age, y/sex	Occupation	Length of exposure, y	Dec 2014		Apr 2015		Dec 2015		Apr 2016	
				HI	MN	HI	MN	HI	MN	HI	MN
<b>Seroconversion</b>											
14.11JS102	33/F	Feeder	3	<b>10</b>	<b>20</b>	<b>40</b>	<b>80</b>	NA	NA	NA	NA
14.11JS114	43/F	Transporter	16	NA	NA	NA	NA	<b>10</b>	<b>ND</b>	<b>40</b>	<b>160</b>
14.11JS8	57/M	Processor, seller	25	<b>10</b>	<b>40</b>	<b>40</b>	<b>80</b>	NA	NA	NA	NA
14.11JS95	61/M	Feeder	15	NA	NA	NA	NA	<b>10</b>	<b>160</b>	<b>40</b>	<b>80</b>
14.11JSWX45	50/F	Feeder	25	NA	NA	NA	NA	<b>5</b>	<b>ND</b>	<b>160</b>	<b>320</b>
14.12GD4	40/F	Seller	10	20	40	<b>10</b>	<b>ND</b>	<b>80</b>	<b>160</b>	<b>20</b>	<b>80</b>
14.12GD11	38/F	Seller	11	5	ND	<b>5</b>	<b>ND</b>	<b>40</b>	<b>160</b>	NA	NA
14.12GD53	49/M	Processor	11	40	40	<b>10</b>	<b>ND</b>	NA	NA	<b>40</b>	<b>80</b>
14.12GD72	34/M	Seller	5	20	80	<b>20</b>	<b>80</b>	<b>80</b>	<b>320</b>	40	160
14.12SC39	55/F	Processor	1	5	ND	<b>10</b>	<b>ND</b>	<b>80</b>	<b>80</b>	20	80
14.12SC7	42/M	Seller	5	10	ND	NA	NA	<b>5</b>	<b>ND</b>	<b>80</b>	<b>80</b>
14.12GD17	31/M	Seller	2	10	ND	<b>10</b>	<b>ND</b>	<b>40</b>	<b>80</b>	5	ND
15.5GD91	33/M	Seller	3	NA	NA	<b>10</b>	<b>ND</b>	<b>40</b>	<b>80</b>	80	640
15.8SH64	54/M	Processor	6	NA	NA	<b>10</b>	<b>ND</b>	<b>40</b>	<b>80</b>	NA	NA
16.1GD1	40/F	Seller	6	NA	NA	NA	NA	<b>5</b>	<b>ND</b>	<b>80</b>	<b>320</b>
16.1GD21	33/F	Seller	1	NA	NA	NA	NA	<b>5</b>	<b>ND</b>	<b>40</b>	<b>160</b>
16.1GD40	42/F	Seller	8	NA	NA	NA	NA	<b>5</b>	<b>ND</b>	<b>40</b>	<b>160</b>
16.1GD74	28/M	Seller	5	NA	NA	NA	NA	<b>5</b>	<b>ND</b>	<b>40</b>	<b>160</b>
16.1JS30	62/M	Feeder	1	NA	NA	NA	NA	<b>5</b>	<b>ND</b>	<b>40</b>	<b>80</b>
16.1JS51	47/M	Feeder	1	NA	NA	NA	NA	<b>5</b>	<b>ND</b>	<b>40</b>	<b>160</b>
16.1JS57	44/M	Feeder	2	NA	NA	NA	NA	<b>5</b>	<b>ND</b>	<b>40</b>	<b>160</b>
16.1SD76	39/M	Processor, seller	17	NA	NA	NA	NA	<b>10</b>	<b>ND</b>	<b>80</b>	<b>160</b>
16.5GD99	30/F	Seller	3	NA	NA	NA	NA	<b>5</b>	<b>ND</b>	<b>40</b>	<b>160</b>
<b>Persistently positive</b>											
14.11JS22	44/M	Seller	8	40	160	40	320	NA	NA	NA	NA
14.11JS3	53/M	Seller	20	40	160	40	160	NA	NA	NA	NA
14.11JS31	30/F	Processor, seller	3	40	640	40	320	NA	NA	NA	NA
14.11JS5	47/F	Processor, seller	15	NA	NA	NA	NA	40	160	40	640
14.11JS85	50/M	Feeder	6	40	160	40	80	NA	NA	NA	NA
14.11JS89	52/M	Feeder	9	40	160	40	80	NA	NA	NA	NA
14.11JSWX06	28/M	Seller	5	160	160	160	320	NA	NA	NA	NA
14.11JSWX7	35/F	Seller	5	40	160	80	160	NA	NA	NA	NA
14.11JSWX8	47/F	Seller	10	40	320	80	320	NA	NA	NA	NA
14.11JX100	46/F	Seller	10	40	320	40	160	NA	NA	NA	NA
14.11JX66	48/F	Seller	8	80	320	80	640	NA	NA	NA	NA


Participant ID	Age, y/sex	Occupation	Length of exposure, y	Dec 2014		Apr 2015		Dec 2015		Apr 2016	
				HI	MN	HI	MN	HI	MN	HI	MN
14.11SC28	59/F	Seller, feeder	8	40	320	80	160	NA	NA	NA	NA
14.12GD1	47/F	Seller	8	40	80	40	80	10	ND	20	160
14.12GD23	28/F	Seller	6	40	160	80	160	NA	NA	NA	NA
14.12GD24	33/F	Seller	10	80	160	80	80	5	ND	NA	NA
14.12GD3	40/F	Seller	9	80	ND	80	320	NA	NA	NA	NA
14.12HN102	34/M	Other	6	40	160	40	160	NA	NA	NA	NA
14.12HN52	46/F	Processor, seller	6	40	320	40	80	NA	NA	NA	NA
15.5GD113	35, M	Seller	0.5	NA	NA	40	80	NA	NA	40	160
15.5SC143	31/F	Seller	6	NA	NA	80	80	80	160	NA	NA
15.8SH12	50/F	Seller	30	NA	NA	80	320	40	640	NA	NA
15.8SH62	48/F	Seller, feeder	17	NA	NA	40	80	40	160	NA	NA
16.1JS101	49/M	Seller	7	NA	NA	NA	NA	80	320	80	320
16.1JS102	46/M	Seller	9	NA	NA	NA	NA	40	160	40	160
16.1JS107	42/F	Seller	15	NA	NA	NA	NA	40	160	40	160
16.1JS109	55/F	Seller	13	NA	NA	NA	NA	160	160	160	640
16.1JS110	27/M	Seller	13	NA	NA	NA	NA	40	160	40	160
16.1JS111	49/F	Seller	13	NA	NA	NA	NA	160	640	160	640
16.1JS112	24/M	Seller	13	NA	NA	NA	NA	40	160	40	160
16.1JS120	45/F	Seller	6	NA	NA	NA	NA	40	80	40	160
16.1JS15	46/F	Feeder	4	NA	NA	NA	NA	40	160	40	160
16.1JS24	65/F	Feeder	2	NA	NA	NA	NA	40	160	40	160
16.1JS96	52/F	Seller	20	NA	NA	NA	NA	80	320	80	320
16.1SC19	45/M	Seller	1	NA	NA	NA	NA	80	160	80	80
16.1SD1	30/M	Processor, seller	1	NA	NA	NA	NA	40	160	80	640
16.1SD18	35/F	Processor, seller	15	NA	NA	NA	NA	40	160	40	80
16.1SD2	39/M	Processor, seller	10	NA	NA	NA	NA	40	80	40	80
16.1SD21	46/M	Processor, seller	6	NA	NA	NA	NA	40	160	40	80
16.1SD33	48/M	Processor, seller	20	NA	NA	NA	NA	80	160	40	160
16.1SD40	37/F	Processor, seller	13	NA	NA	NA	NA	40	320	80	160
16.1SD58	30/M	Processor, seller	5	NA	NA	NA	NA	40	160	40	80

\*Bold text represents titers exhibiting seroconversion. HI, hemagglutinin inhibition assay; ID, identification; MN, microneutralization assay; NA, not available; ND, not detected.


**Appendix Table 10.** The number of avian influenza A isolates from live poultry markets during December 2014–December 2015, China

Province	Sampling date	Sample		No.	H5N1, no. (%)	H5N6, no. (%)	H7N9, no. (%)	H9N2, no. (%)
		source						
Guangdong	2015	Chickens		490	0	24 (4.9)	4 (0.8)	74 (15.1)
	2015	Ducks		1,326	1 (0.08)	293 (22.1)	6 (0.5)	60 (4.5)
	2015	Environment		342	0	35 (10.2)	2 (0.6)	46 (13.5)
	Total			2,158	1 (0.05)	352 (16.3)	12 (0.6)	180 (8.3)
Jiangxi	2015	Chickens		1,098	0	115 (10.5)	8 (0.7)	38 (3.5)
	2015	Ducks		735	1 (0.1)	90 (12.2)	4 (0.5)	8 (1.1)
	2015	Environment		475	1 (0.2)	99 (20.8)	3 (0.6)	56 (11.8)
	Total			2,308	2 (0.09)	304 (13.2)	15 (0.7)	102 (4.4)
Shanghai	2015	Chickens		481	13 (2.7)	0	19 (3.9)	69 (14.3)
	2015	Environment		12	2 (16.7)	0	1 (8.3)	2 (16.7)
	Total			493	15 (3.0)	0	20 (4.1)	71 (14.4)
Jiangsu	Dec 2014–Mar 2015	Chickens		361	10 (2.8)	5 (1.4)	18 (5.0)	42 (11.6)
	Dec 2014–Jan 2015	Ducks		28	4 (14.3)	0	2 (7.1)	0
	Dec 2014–Mar 2015	Environment		87	11 (12.6)	5 (5.7)	21 (24.1)	3 (3.4)
	Total			476	25 (5.3)	10 (2.1)	41 (8.6)	45 (9.5)
	Dec 2014–Apr 2015	Chickens		190	1 (0.5)	1 (0.5)	0	0
Sichuan	Dec 2014–Apr 2015	Ducks		241	2 (0.8)	17 (7.1)	1 (0.4)	6 (2.5)
	Dec 2014–Apr 2015	Environment		85	18 (21.2)	2 (2.4)	0	0
	Total			516	21 (4.1)	20 (3.9)	1 (0.2)	7 (1.4)
Henan	Apr 2015	Chickens		245	0	0	0	9 (3.7)
	Apr 2015	Ducks		11	0	0	0	5 (45.5)
	Total			256	0	0	0	14 (5.5)
Total			6,207	64 (1.0)	686 (11.1)	48 (0.8)	419 (6.8)	






**Appendix Figure 1.** Phylogenetic relationship of avian influenza A(H5) clade 2.3.2.1 HA genes in a study of avian influenza viruses among occupationally-exposed poultry workers, China, 2014–2016. Green dots indicate sequences similar to recommended vaccine strains. Pink triangles indicate virus strains isolated by our laboratory. Bold text indicates human virus strains. Red text indicates reference strain used in this study. Scale bar indicates nucleotide substitutions per site.




**Appendix Figure 2.** Phylogenetic relationship of avian influenza A(H5) clade 2.3.4 HA genes in a study of avian influenza viruses among occupationally-exposed poultry workers, China, 2014–2016. Green dots indicate available candidate influenza vaccine strains. Pink triangles indicate virus strains isolated by our laboratory. Bold text indicates human virus strains. Red text indicates reference strain used in this study. Scale bar indicates nucleotide substitutions per site.



**Appendix Figure 3.** Phylogenetic relationship of avian influenza A(H7) HA genes in a study of avian influenza viruses among occupationally-exposed poultry workers, China, 2014–2016. Green dots indicate available candidate influenza vaccine strains. Blue diamonds indicated reported sequences from this study. Pink triangles indicate virus strains isolated by our laboratory. Bold text indicates human virus strains. Red text indicates reference strain used in this study. Scale bar indicates nucleotide substitutions per site.



**Appendix Figure 4.** Phylogenetic relationship of avian influenza A(H9) HA genes in a study of avian influenza viruses among occupationally-exposed poultry workers, China, 2014–2016. Green dots indicate available candidate influenza vaccine strains. Blue diamonds indicated reported sequences from this study. Pink triangles indicate virus strains isolated by our laboratory. Bold text indicates human virus strains. Red text indicates reference strain used in this study. Scale bar indicates nucleotide substitutions per site.



**Appendix Figure 5.** Phylogenetic relationship of avian influenza A(H6) HA genes in a study of avian influenza viruses among occupationally-exposed poultry workers, China, 2014–2016. Diamonds indicate antigenic characteristics reported from this study. Pink triangles indicate virus strains isolated by our laboratory. Red text indicates reference strain used in this study. Scale bar indicates nucleotide substitutions per site.