

Case Investigations of Infectious Diseases Occurring in Workplaces, United States, 2006–2015

Chia-ping Su,¹ Marie A. de Perio, Kristin J. Cummings, Anna-Binney McCague,²
Sara E. Luckhaupt, Marie Haring Sweeney

Workers in specific settings and activities are at increased risk for certain infectious diseases. When an infectious disease case occurs in a worker, investigators need to understand the mechanisms of disease propagation in the workplace. Few publications have explored these factors in the United States; a literature search yielded 66 investigations of infectious disease occurring in US workplaces during 2006–2015. Reported cases appear to be concentrated in specific industries and occupations, especially the health-care industry, laboratory workers, animal workers, and public service workers. A hierarchy-of-controls approach can help determine how to implement effective preventive measures in workplaces. Consideration of occupational risk factors and control of occupational exposures will help prevent disease transmission in the workplace and protect workers' health.

Despite ongoing efforts to reduce, eliminate, and eradicate infectious diseases, infections continue to pose a global threat to human health. More than 2,000 years ago, Hippocrates noted that “there are many handicrafts and arts which cause those who exercise them certain... plagues” (1). Recent experiences with anthrax, severe acute respiratory syndrome (SARS), influenza A(H1N1), and Ebola virus disease have highlighted the importance of focusing on workplaces not only to identify at-risk populations but also to understand mechanisms of disease propagation and to implement successful control and prevention measures (2–5).

In the United States, work-related infectious diseases are identified in multiple ways. Workers, employers, or workplace health and safety offices may note an unusual case or cluster of disease. Local and state public health de-

partments receive case reports of infectious disease from clinicians and laboratories, and they may conduct investigations, sometimes with assistance from the Centers for Disease Control and Prevention (CDC). CDC's National Institute for Occupational Safety and Health (NIOSH) is the federal public health agency responsible for conducting research and making recommendations to prevent occupational safety and health risks, including work-related infectious diseases. The NIOSH Health Hazard Evaluation (HHE) program responds to requests from workers, employers, and public health agencies and conducts investigations of hazards including infectious diseases that occur in workplaces. Nongovernmental researchers also carry out investigations.

Investigators of work-related infectious disease must consider multiple factors related to the disease, workplace, and workers. Few publications have explored these factors. To illustrate the range of work-related infectious diseases that have been identified in the United States during 2006–2015 and to benefit future investigations, we examined the peer-reviewed literature and HHE reports on infectious diseases occurring in US workplaces. We describe occupational factors to consider and a systematic approach to control and prevent infectious disease in the workplace. We also note ways that specialized resources may be useful during the course of an investigation.

Literature Review

Methods

As defined by the World Health Organization, work-related diseases may have multiple causes, in which the work environment and other risk factors can play a role (https://www.who.int/occupational_health/activities/

Author affiliations: National Institute for Occupational Safety and Health, Cincinnati, Ohio, USA (C.-P. Su, M.A. de Perio, S.E. Luckhaupt, M.H. Sweeney); Centers for Disease Control and Prevention, Atlanta, Georgia, USA (C.-P. Su, A.-B. McCague); National Institute for Occupational Safety and Health, Morgantown, West Virginia, USA (K.J. Cummings, A.-B. McCague)

DOI: <https://doi.org/10.3201/eid2503.180708>

¹Current affiliation: Taiwan Centers for Disease Control, Taipei, Taiwan, and Institute of Epidemiology and Preventive Medicine, National Taiwan University, Taipei, Taiwan.

²Current affiliation: Washington Family Medicine Residency Program, Washington Hospital System, Washington, Pennsylvania, USA.

occupational_work_diseases/en). Work-related infectious diseases can be defined as those primarily caused by occupational exposure to biologic agents. These biologic agents, such as bacteria, fungi, viruses, and parasites, can be transmitted from human to human, from animal to human, or through environmental contact (6). We focused on studies of infectious disease occurring among workers in specific occupational groups or workplaces.

For the literature review, we followed the methods of Haagsma et al. (6). In March 2016, we searched PubMed for articles published since 2006. Our search strategy combined 3 groups of relevant keywords, including case investigation, workers/workplace/occupational, and infectious diseases. We also searched the NIOSH HHE reports database (<https://www2a.cdc.gov/hhe/search.asp>), using a similar strategy. We classified infectious diseases occurring in workplaces by industry category (work setting), using 2-digit codes of the North American Industry Classification System (NAICS) (<https://www.census.gov/eos/www/naics>), and by occupation (type of job). We describe detailed methods of our search of the literature and HHE reports in the Appendix (<https://wwwnc.cdc.gov/EID/article/25/3/18-0708-App1.pdf>).

Results

The literature search yielded 67 articles and 7 HHE reports from 66 investigations of infectious diseases occurring in US workplaces during 2006–2015 (Table 1; Appendix). Cases were concentrated in specific industries and occupations, especially healthcare and work involving contact with animals, reflecting the potential for disease transmission from patients and animals. Complementing previous findings from a systematic review by Haagsma et al. (6), our review of worksite case investigations found that several work-related infectious disease outbreaks or individual cases have been reported among laboratory and public service workers.

Work-related cases were associated with a variety of infectious pathogens. Bacteria were responsible for most reported cases, followed by viruses, fungi, and parasites or protozoa. As noted previously (6), respiratory viruses and zoonotic pathogens still threaten workers' health, especially for healthcare personnel and animal-contact workers. However, we also found reports of some emerging or reemerging pathogens, such as Ebola virus, lymphocytic choriomeningitis virus, norovirus, *Bacillus anthracis*, and *Yersinia pestis*, that caused several workplace disease clusters.

Specific Considerations

Many factors may combine to increase the risk for infection among workers during pathogen transmission. Categories of risk factors for work-related infections include disease

factors (such as transmission mode), workplace factors, and worker factors. Considering each of these categories during case investigations is useful in planning and implementing prevention and control strategies.

Disease Factors

Infectious disease can be transmitted via direct contact (including percutaneous), droplet, airborne (aerosol), vehicles (such as food, water, and fomites), and vectors (Table 2). The studies included in our review suggest that occupations involving interaction with the general population, particularly ill persons, pose an increased risk for infection.

As one example, teachers and public service workers may acquire respiratory virus infections, including influenza and measles, because their work may bring them in contact with persons who are ill (7–10). Workers in the healthcare industry are also at risk for influenza as well as airborne (such as tuberculosis [TB]) and percutaneously transmitted (such as HIV) infection from patients (11–13). In 2014, work-related Ebola virus infections among healthcare personnel were a substantial component of the Ebola epidemic worldwide; 2 healthcare personnel acquired Ebola virus disease within the United States (14,15). Occupational contact with human corpses can also result in infectious disease. In 2007, an embalmer in New York, New York, USA, contracted *Mycobacterium tuberculosis* from a cadaver (16).

Disease transmission patterns are also relevant to those whose work brings them in contact with animals, putting them at risk for zoonotic infections. Occupational exposure to livestock and poultry contributed substantially to work-related infectious diseases. Twenty-nine cases of *Campylobacter* infection occurred over a period of several years among workers at a poultry processing plant (17), and sealpox virus infections were reported among animal rescue workers in a marine mammal rehabilitation center (18).

Transmission of pathogens in the workplace may occur in 2 directions: workers can acquire infections in the workplace and then also may serve as vectors that spread the disease to others, such as clients and co-workers. We found that workers in food preparation and serving-related occupations have been identified as sources of transmission in foodborne outbreaks. Two delicatessen workers infected with *Salmonella* from occupational contact with chicken became the source of disease transmission in a 2007 salmonellosis cluster in Minnesota (19). Transmission of norovirus gastroenteritis among workers and customers at a restaurant has also been reported (20). These examples show that preventing workers from acquiring infections in workplaces may also prevent disease transmission among the general public.

Pathogens in the environment can also serve as a source of worker infections through the respiratory route.

Table 1. Reported case investigations of infectious disease occurring in workplaces, by industry categories, occupations, and diseases, United States, 2006–2015*

Industry category (NAICS code)	Occupations	Infectious diseases	References†
Agriculture, forestry, fishing, and hunting (11)	Hunter	Brucellosis	(61)
	Farmer	Variant influenza A(H3N2); <i>Escherichia coli</i> infection	(83); (71)
	Rodent breeder	LCMV infection	(82)
Construction (23)	Laborer	Coccidioidomycosis	(23,25)
Manufacturing (31–33)	Drum maker	Anthrax	(2)
	Poultry vaccine production worker	Salmonellosis	(29)
	Poultry-processing worker	Campylobacteriosis	(17)
	Furniture company worker	Tuberculosis	(54)
	Slaughterhouse inspector	Q fever	(65)
	Automobile manufacturing worker	Legionnaires' disease	(81)
Transportation (48)	Truck driver	<i>Streptococcus suis</i> infection; cryptosporidiosis	(79); (88)
	Pilot, flight attendant	Malaria	(37)
	Laboratory worker	Vaccinia virus infection, HIV infection, plague, cowpox, meningococcal disease, brucellosis	(13,30–35,86)
Administrative support and waste management and remediation services (56)	Landscaper	Tularemia	(21)
Education services (61)	School employee, teacher	Influenza	(8)
Healthcare and social assistance (62)	Healthcare worker (security guard, nurse, nursing aide, physician, volunteer, environmental services)	Mumps; MRSA skin infection; norovirus gastroenteritis; adenovirus 14 infection; RSV infection; <i>Trichophyton tonsurans</i> skin infection; meningococcal disease; influenza; salmonellosis; Ebola virus disease; measles; TB;	(51); (52); (56); (57); (62); (64); (66); (11,68); (87); (14); (92); (12,77)
		<i>E. coli</i> infection	(72)
		Plague	(59)
		MRSA skin infection	(63)
	Childcare worker	HIV infection	(36)
		MAC infection	(22)
		Coccidioidomycosis	(24)
		Histoplasmosis	(26)
	Wildlife biologist	Norovirus gastroenteritis; salmonellosis; <i>E. coli</i> infection	(20); (19); (89)
		TB	(16)
Arts, entertainment, and recreation (71)	Animal caretaker	Tuberculosis; sealpox virus infection	(28); (18)
	Adult film performer	Salmonellosis	(74)
	Spa maintenance worker	Melioidosis; dengue fever	(75); (70)
	Filmmaker	Measles	(9,10)
Day camp counselor	Police officer	Meningococcal disease	(66)
	Firefighter	Cryptosporidiosis	(88)
Food services (72)	Cook, food server	Cryptosporidiosis; Shiga toxin-producing <i>E. coli</i> infection; TB; coccidioidomycosis;	(78); (71); (12); (90)
	Embalmers	Legionellosis; TB	(73); (53)
	Animal refugee worker		
Other services except public administration (81)	Pet store worker		
	Missionary worker		
	Missionary worker		
Public administration (92)	US Customs officer		
	Police officer		
	Firefighter		
	Correctional officer		
	Military		

*An expanded version of this table showing complete details on all cases is available online (<https://wwwnc.cdc.gov/EID/article/25/3/18-0708-T1.htm>). HIV, human immunodeficiency virus; LCMV, lymphocytic choriomeningitis virus; MAC, *Mycobacterium avium* complex; MRSA, methicillin-resistant *Staphylococcus aureus*; RSV, respiratory syncytial virus; TB, tuberculosis. NAICS, 2012 North American Industry Classification System (<https://www.census.gov/eos/www/naics/>).

†Reference numbers >50 and additional details on the literature search are available in the Appendix (<https://wwwnc.cdc.gov/EID/article/25/3/18-0708-App1.pdf>).

A 2006 report described a 21-year-old healthy landscaper diagnosed with tularemia. Traditionally thought of as a zoonotic pathogen, *Francisella tularensis* can also be acquired via aerosolized bacteria during occupational activities such as lawn mowing and leaf blowing (21). Another report described 2 spa maintenance workers infected with *Mycobacterium avium* complex organisms, which live in water and are highly resistant to disinfectants,

such as chlorine. Occupational exposure to aerosolized bacteria during routine cleaning and maintenance of spa filters and tubs was the likely cause of this outbreak (22).

Workplace Factors

Regardless of transmission mode, workplace factors can contribute to the propagation of infection. It is crucial to identify aspects of the workplace that pose biologic haz-

Table 2. Reported case investigations of infectious disease pathogens occurring in workplaces, by mode of transmission and source of disease, United States, 2006–2015*

Mode of transmission	Disease (reference no.)†		
	Human	Animal	Environment
Direct contact (including percutaneous)	MRSA (52,63) <i>Trichophyton tonsurans</i> (64) Ebola virus (14, 15) HIV (13,36)	MRSA (63) <i>Bacillus anthracis</i> (2) <i>Brucella</i> spp. (61) <i>Coxiella burnetii</i> (65) <i>Yersinia pestis</i> (59) Sealpox virus (18) <i>Streptococcus suis</i> (79) LCMV (82) <i>Burkholderia pseudomallei</i> (75)	
Droplet	Adenovirus 14 (57) influenza virus (7,8,11,68) <i>Neisseria meningitidis</i> (66) RSV (62) Mumps virus (51)	Influenza virus (83)	
Airborne	<i>Mycobacterium tuberculosis</i> (12,40,53,54,77,85) Measles virus (9,10,92)	<i>Mycobacterium tuberculosis</i> (28)	<i>Coccidioides</i> spp. (23–25,90) MAC (22) <i>Legionella pneumophila</i> (73,81) <i>Francisella tularensis</i> (21) <i>Histoplasma capsulatum</i> (26)
Vehicles (fecal–oral)	Norovirus (20,56) <i>E. coli</i> (72)	<i>Salmonella</i> spp. (19,74) <i>E. coli</i> (71,89) <i>Cryptosporidium</i> (78,88) <i>Campylobacter jejuni</i> (17)	
Vectors		<i>Plasmodium</i> spp. (37) Dengue virus (70)	

*LCMV, lymphocytic choriomeningitis virus; MAC, *Mycobacterium avium* complex; MRSA, methicillin-resistant *Staphylococcus aureus*; RSV, respiratory syncytial virus.

†Reference numbers >50 and additional details on the literature search are available in the Appendix (<https://wwwnc.cdc.gov/EID/article/25/3/18-0708-App1.pdf>).

ards, because those aspects may be amenable to controls. Such factors can include workplace characteristics, work practices and processes, and engineering and administrative issues. For example, work-related fungal respiratory infection is a concern in some areas. Outbreaks of coccidioidomycosis have occurred among construction workers and outdoor film production workers in California (23,24); during 2011–2014, a total of 44 cases of coccidioidomycosis were identified among workers constructing solar power farms (25). Outdoor workplaces with hot, dry conditions in areas of endemicity pose a risk for coccidioidomycosis because soil-disruptive activities and high winds expose workers to dust harboring *Coccidioides* spores. Other factors may also contribute. In 2012, Nebraska officials reported an investigation of a cluster of histoplasmosis among 32 day camp counselors (26). The probable infection source was campsite contamination of soil and picnic tables by bat guano, which probably became aerosolized during camp activities or cleanup.

Engineering factors can also promote disease transmission in the workplace. Previous studies have shown that workplace environmental characteristics contributed to *M. tuberculosis* transmission among workers (27). During the outbreak investigation of TB among workers at an elephant refuge in 2009, investigators found that shared air between the administrative area and the

barn, along with pressure-washing of the barn by workers, contributed to transmission (28). Laboratories are another example of unique workplaces where engineering controls (such as biosafety cabinets and local exhaust ventilation) are essential for reducing or eliminating potential biologic hazards to workers. Laboratory workers have acquired *Salmonella* infection, vaccinia virus infection, plague, cowpox, brucellosis, meningococcal disease, and HIV infection through accidental direct contact with pathogens in the workplace over the past decade (13,29–35).

Administrative issues, including workplace policies and practices, can also play an important role in disease transmission in the workplace. In 2014, the California Department of Public Health reported an occupational HIV outbreak among adult film performers (36). Some adult film production companies rely on HIV testing results as a control and require performers to engage in penetrative sex without a condom. This approach is problematic, because during acute infection, a performer can transmit the infection even when HIV test results are negative. This example highlights that the lack of a protective administrative policy in the workplace could lead to infectious disease transmission among workers.

For workers in high-risk environments, employers have a responsibility to provide adequate prophylaxis for prevention of infectious disease. However, sometimes

employers or workers are unfamiliar with or unable to comply with the relevant recommendations. For example, CDC staff found 4 malaria cases among employees of a commercial airline who had all traveled to Ghana and stayed at the same hotel before disease onset. None had used antimalarial chemoprophylaxis provided by the company (37). It is unclear why they did not use the chemoprophylaxis, but this cluster underscores the importance of a comprehensive malaria prevention program that includes education and counseling.

Worker Factors

Individual characteristics, such as impaired immunity, inadequate prophylaxis, and socioeconomic and language factors, may increase the risk for acquisition and transmission of infectious diseases. For example, a case of fatal laboratory-acquired infection caused by *Y. pestis* occurred in a laboratorian in 2009 (32). No additional cases or major deficiencies in engineering controls were identified in this laboratory. A postmortem examination revealed that the affected worker had hereditary hemochromatosis, a condition that increases susceptibility to infection with certain bacterial pathogens. In another situation, *Campylobacter* infection among poultry-processing workers was found to occur most frequently during the first weeks of work, after which the workers develop immunity that may be protective against future infection (17). Therefore, investigators should consider that individual host susceptibility to certain diseases may play a role in disease transmission in workplaces.

Documented nosocomial transmission puts healthcare personnel at substantial risk for acquiring or transmitting several vaccine-preventable diseases, including hepatitis B, influenza, measles, mumps, rubella, pertussis, and varicella. The Advisory Committee on Immunization Practices and the Hospital Infection Control Practices Advisory Committee recommend that healthcare personnel be vaccinated or have documentation of immunity for all of these diseases; employers must formulate a comprehensive vaccination policy for all healthcare personnel (38,39). Workers in other settings, such as public services, may also be at risk for exposure to vaccine-preventable diseases, such as measles. In 2007, an airport officer contracted measles from an international traveler and may have transmitted it to a second airport worker (9). In 2011, a US Customs and Border Protection officer contracted measles after processing an arriving refugee with measles (10). All of these situations underscore that immunizations should be kept current for workers whose jobs involve frequent contact with the public.

In addition to individual susceptibility and immunity, a worker's socioeconomic status could contribute to risk. Texas officials reported a TB outbreak investigation among

workers in a meatpacking plant in 2011. The index case was in a foreign-born patient who had cavitary TB with acid-fast-positive smears. Most of the patient's work contacts were foreign-born. Investigators found that low economic status, limited access to healthcare, and communication and language barriers caused delays in diagnosis and played significant roles in TB transmission among immigrant workers (40).

Approach to Controlling Exposure to Protect Workers

Investigators of an infectious disease outbreak that is potentially related to a workplace must consider the "who, what, where, when, and why" questions of field epidemiology in the context of a unique and sometimes contentious environment. Stakeholders, including workers, employers, labor unions, trade associations, regulatory agencies, and members of the public, may have distinct and, at times, competing priorities. Barriers to implementing solutions may include those common to all epidemics, such as expense and time constraints, and others more specific to the occupational setting, such as a poorly developed safety culture.

To reduce worker exposure to potential occupational hazards, including biologic agents, we need feasible, effective measures that can be implemented in the workplace. Occupational health and safety specialists have long used the hierarchy of controls (Figure 1) to eliminate or minimize exposure to any occupational hazard. The control methods at the top of the graphic are potentially more effective and protective than those at the bottom. Here are examples of control recommendations made for identified occupational hazards using the hierarchy.

Elimination and Substitution

Elimination and substitution are the most effective ways to reduce occupational hazards but sometimes can be difficult to implement in an existing work process. For infectious diseases, one should first assess the feasibility of not working in an area in which biohazards are present. If avoidance is not feasible, then decontaminating surfaces, items, and areas in a hazardous workplace can eliminate the possibility of transmission of infectious agents to workers. For environmentally associated laboratory infections, CDC has published guidance on methods for sterilization and disinfection in laboratories and on the levels of antimicrobial activity associated with liquid chemical germicides (41). However, employers and workers should also be aware of potential health risks when using such disinfectants (42).

Training on early recognition of the key signs and symptoms of common communicable diseases and the




Figure. The hierarchy of controls for controlling exposures to occupational hazards. Source: NIOSH, <https://www.cdc.gov/niosh/topics/hierarchy/default.html>.

use of sick leave by workers may help to reduce disease transmission in the workplace. For example, CDC guidance for responding to influenza in schools during the 2009 influenza A(H1N1) pandemic included directing students and staff with influenza-like illness to stay home when ill (43). A recent study showed that offering paid sick leave to workers is likely to reduce the spread of disease in workplaces by increasing the rate at which sick workers stay home (44). However, the effectiveness of those measures still needs to be closely evaluated.

Engineering Controls

Engineering controls in the field of occupational health and safety are physical changes to work processes or equipment to remove hazardous conditions or to place a barrier between workers and hazards. Engineering controls effectively protect workers without placing the primary responsibility of implementation on the worker. Ventilation is by far the most common engineering control, especially for airborne pathogens. Previous modeling data showed that the risk for TB infection in healthcare settings decreases exponentially as room ventilation rates increase (45). Thus, guidelines for preventing transmission of TB in healthcare settings recommend engineering controls such as local exhaust and general ventilation to prevent the spread and reduce the concentration of infectious droplet nuclei in the air. Supplementing ventilation systems with high-efficiency particulate air filtration, ultraviolet germicidal irradiation, or both can further enhance system performance and reduce the spread of airborne disease (46).

Nonventilation engineering controls also can reduce or eliminate pathogen exposure in the workplace. For example,

dust suppression during construction, including continuous soil-wetting and proper covering, could decrease the risk for coccidioidomycosis among outdoor workers (23). In addition, evidence shows that engineering controls can reduce healthcare personnel's exposure to bloodborne pathogens. Safety needle devices with built-in engineering controls reduce the risk for needlestick injury among healthcare personnel (47).

Administrative Controls

Administrative controls are methods such as standard operating procedures that change the way work is performed. Their effectiveness depends on the availability of the control, employer commitment, and worker acceptance. Regular monitoring and reinforcement are necessary to ensure that workers follow policies and procedures consistently. The CDC publication *Biosafety in Microbiological and Biomedical Laboratories* addresses recommended practices for working safely from a biosafety perspective (41).

Developing and implementing a mandated workplace health regulation can be an effective administrative control. In December 1991, mandatory hepatitis B vaccination of all healthcare personnel (at employers' expense) became a federal standard under the Occupational Safety and Health Act (48). The regulation accelerated the use of hepatitis B vaccine for healthcare personnel. As the result of routine vaccination and improved infection control precautions, the number of hepatitis B virus infections among healthcare personnel decreased from $\approx 10,000$ in 1982 to ≈ 304 in 2004 (39). In addition, administrative measures of TB control, such as developing and implementing a written TB infection-control plan, prevent disease transmission in healthcare settings (46).

Personal Protective Equipment

Various types of personal protective equipment (PPE) are available to minimize exposure to hazards in workplaces. PPE, such as respirators, gloves, goggles, and coveralls, provides a physical barrier between the worker and the infectious agent. Compared with other methods of controlling exposure, PPE is the least effective but probably best-known method used for infectious disease prevention. Proper use of PPE requires a comprehensive program and a high level of worker involvement and commitment. When engineering, work practice, and administrative controls are not feasible or do not provide sufficient protection, PPE may be the only reliable method of disease prevention for workers. Therefore, employers must provide their workers with PPE that is appropriate to the task and the correct size for the user, along with proper training on use and on donning and doffing methods.

In the early stages of epidemics of emerging infectious diseases, such as Ebola virus disease, lack or misuse of PPE can lead to infections in healthcare personnel. In 2014, the transmission of Ebola virus to 2 nurses who provided care to an Ebola-infected patient at a US hospital revealed the importance of directive PPE recommendations and standardized training efforts for healthcare personnel. Interventions included a system of trained observers supervising the donning and doffing of PPE (5,49). Even though the quality of evidence is low, the risk for contamination may be reduced by double-gloving, following directives for donning and doffing procedures, and instituting more active training (50).

Additional Resources

The NIOSH Surveillance Program works with partners at CDC and the Council of State and Territorial Epidemiologists to promote inclusion of standard occupational information in the National Notifiable Disease Surveillance system (in which most conditions included are infectious) and in electronic medical records, to facilitate detection of possible disease transmission in workplaces (<https://www.cdc.gov/niosh/topics/ehr/default.html>). Once a disease cluster or outbreak is suspected, the NIOSH HHE program can be a resource for technical assistance and consultation during the case investigation in a workplace. Workers, employers, or public health professionals can request an evaluation of health hazards in the workplace (<https://www.cdc.gov/niosh/hhe>). NIOSH makes recommendations aimed at controlling the hazard; these are voluntary but can reduce risk and improve the health and safety of the workforce.

Conclusions

We found that cases of work-related infectious diseases in the United States during 2006–2015 appeared to be concentrated

in specific industries and occupations, especially in healthcare and among laboratory, animal, and public service workers. The biosafety programs in these industries could be strengthened. Case investigations of infectious disease occurring in a workplace can be challenging with regard to linking the symptoms of a disease to a specific pathogen or exposure source and identifying effective preventive strategies. A multidisciplinary approach that includes epidemiologists, physicians, industrial hygienists, and engineers may be beneficial.

Emerging and reemerging work-related infectious diseases will continue to threaten workers' health. Previously published literature has identified several high-risk occupations, but other occupations may also be at risk. Indeed, we cannot know with certainty which industry or which workers will be at risk in the future. Considering occupational risk factors and controlling exposures among workers when investigating infectious disease may help prevent disease transmission in the workplace. In addition, because person-to-person transmission between workers and members of the public can propagate some disease outbreaks, assessment of worker infection may contribute to control of disease outbreaks in communities.

Acknowledgments

We thank Onnalee A. Gomez, Seleen S. Collins, and Stephen Martin for their technical assistance and consultation. We also highly appreciate the comments and suggestions of Derry Stover and Kenneth Rosenman on an earlier version of this paper.

About the Author

Dr. Su was an Epidemic Intelligence Service officer at the CDC's National Institute for Occupational Safety and Health at the time of the study. He is now a medical officer and infectious diseases physician at the Field Epidemiology Training Program at the Taiwan Centers for Disease Control in Taipei. His primary research interests include outbreak investigation and all infectious diseases of public health importance, especially influenza, emerging infectious diseases, and infectious diseases that occur in the workplace.

References

1. Price GM. Medicine and the industries. *Bulletin of the American Academy of Medicine*. 1917;18:134–9.
2. Centers for Disease Control and Prevention (CDC). Cutaneous anthrax associated with drum making using goat hides from West Africa—Connecticut, 2007. *MMWR Morb Mortal Wkly Rep*. 2008;57:628–31.
3. Fowler RA, Lapinsky SE, Hallett D, Detsky AS, Sibbald WJ, Slutsky AS, et al.; Toronto SARS Critical Care Group. Critically ill patients with severe acute respiratory syndrome. *JAMA*. 2003;290:367–73. <http://dx.doi.org/10.1001/jama.290.3.367>
4. Suarathana E, McFadden JD, Laney AS, Kreiss K, Anderson HA, Hunt DC, et al. Occupational distribution of persons with confirmed 2009 H1N1 influenza. *J Occup Environ Med*. 2010;52:1212–6. <http://dx.doi.org/10.1097/JOM.0b013e3181fd32e4>


5. Cummings KJ, Choi MJ, Esswein EJ, de Perio MA, Harney JM, Chung WM, et al. Addressing infection prevention and control in the first U.S. community hospital to care for patients with Ebola virus disease: context for national recommendations and future strategies. *Ann Intern Med*. 2016;165:41–9. <http://dx.doi.org/10.7326/M15-2944>
6. Haagsma JA, Tariq L, Heederik DJ, Havelaar AH. Infectious disease risks associated with occupational exposure: a systematic review of the literature. *Occup Environ Med*. 2012;69:140–6. <http://dx.doi.org/10.1136/oemed-2011-100068>
7. Lessler J, Reich NG, Cummings DA. New York City Department of Health and Mental Hygiene Swine Influenza Investigation Team. Outbreak of 2009 pandemic influenza A(H1N1) at a New York City school. *N Engl J Med*. 2009;361:2628–36.
8. Centers for Disease Control and Prevention (CDC). Outbreak of 2009 pandemic influenza A (H1N1) at a school—Hawaii, May 2009. *MMWR Morb Mortal Wkly Rep*. 2010;58:1440–4.
9. Chen TH, Kutty P, Lowe LE, Hunt EA, Blostein J, Espinoza R, et al. Measles outbreak associated with an international youth sporting event in the United States, 2007. *Pediatr Infect Dis J*. 2010;29:794–800. <http://dx.doi.org/10.1097/INF.0b013e3181dbaacf>
10. Centers for Disease Control and Prevention (CDC). Measles outbreak associated with an arriving refugee—Los Angeles County, California, August–September 2011. *MMWR Morb Mortal Wkly Rep*. 2012;61:385–9.
11. Magill SS, Black SR, Wise ME, Kallen AJ, Lee SJ, Gardner T, et al. Investigation of an outbreak of 2009 pandemic influenza A virus (H1N1) infections among healthcare personnel in a Chicago hospital. *Infect Control Hosp Epidemiol*. 2011;32:611–5. <http://dx.doi.org/10.1086/660097>
12. Medrano BA, Salinas G, Sanchez C, Miramontes R, Restrepo BI, Haddad MB, et al. A missed tuberculosis diagnosis resulting in hospital transmission. *Infect Control Hosp Epidemiol*. 2014;35:534–7. <http://dx.doi.org/10.1086/675833>
13. Joyce MP, Kuhar D, Brooks JT. Notes from the field: occupationally acquired HIV infection among health care workers—United States, 1985–2013. *MMWR Morb Mortal Wkly Rep*. 2015;63:1245–6.
14. Chevalier MS, Chung W, Smith J, Weil LM, Hughes SM, Joyner SN, et al.; Centers for Disease Control and Prevention (CDC). Ebola virus disease cluster in the United States—Dallas County, Texas, 2014. *MMWR Morb Mortal Wkly Rep*. 2014;63:1087–8.
15. Yacisin K, Balter S, Fine A, Weiss D, Ackelsberg J, Prezant D, et al.; Centers for Disease Control and Prevention (CDC). Ebola virus disease in a humanitarian aid worker—New York City, October 2014. *MMWR Morb Mortal Wkly Rep*. 2015;64:321–3.
16. Anderson JA, Meissner JS, Ahuja SD, Shashkina E, O'Flaherty T, Proops DC. Confirming *Mycobacterium tuberculosis* transmission from a cadaver to an embalmer using molecular epidemiology. *Am J Infect Control*. 2015;43:543–5. <http://dx.doi.org/10.1016/j.ajic.2015.01.027>
17. de Perio MA, Niemeier RT, Levine SJ, Gruszynski K, Gibbins JD. *Campylobacter* infection in poultry-processing workers, Virginia, USA, 2008–2011. *Emerg Infect Dis*. 2013;19:286–8. <http://dx.doi.org/10.3201/eid1902.121147>
18. Roess AA, Levine RS, Barth L, Monroe BP, Carroll DS, Damon IK, et al. Sealpox virus in marine mammal rehabilitation facilities, North America, 2007–2009. *Emerg Infect Dis*. 2011;17:2203–8. <http://dx.doi.org/10.3201/eid1712.101945>
19. Hedican E, Miller B, Ziemer B, LeMaster P, Jawahir S, Leano F, et al. Salmonellosis outbreak due to chicken contact leading to a foodborne outbreak associated with infected delicatessen workers. *Foodborne Pathog Dis*. 2010;7:995–7. <http://dx.doi.org/10.1089/fpd.2009.0495>
20. Centers for Disease Control and Prevention (CDC). Norovirus outbreak associated with ill food-service workers—Michigan, January–February 2006. *MMWR Morb Mortal Wkly Rep*. 2007;56:1212–6.
21. Hofinger DM, Cardona L, Mertz GJ, Davis LE. Tularemic meningitis in the United States. *Arch Neurol*. 2009;66:523–7. <http://dx.doi.org/10.1001/archneurol.2009.14>
22. Moraga-McHaley SA, Landen M, Krapfl H, Sewell CM. Hypersensitivity pneumonitis with *Mycobacterium avium* complex among spa workers. *Int J Occup Environ Health*. 2013;19:55–61. <http://dx.doi.org/10.1179/2049396712Y.0000000015>
23. Cummings KC, McDowell A, Wheeler C, McNary J, Das R, Vugia DJ, et al. Point-source outbreak of coccidioidomycosis in construction workers. *Epidemiol Infect*. 2010;138:507–11. <http://dx.doi.org/10.1017/S0950268809990999>
24. Wilken JA, Marquez P, Terashita D, McNary J, Windham G, Materna B; Centers for Disease Control and Prevention. Coccidioidomycosis among cast and crew members at an outdoor television filming event—California, 2012. *MMWR Morb Mortal Wkly Rep*. 2014;63:321–4.
25. Wilken JA, Sondermeyer G, Shusterman D, McNary J, Vugia DJ, McDowell A, et al. Coccidioidomycosis among workers constructing solar power farms, California, USA, 2011–2014. *Emerg Infect Dis*. 2015;21:1997–2005. <http://dx.doi.org/10.3201/eid2111.150129>
26. Centers for Disease Control and Prevention. Notes from the field: histoplasmosis outbreak among day camp attendees—Nebraska, June 2012. *MMWR Morb Mortal Wkly Rep*. 2012;21:747–8.
27. Davidow AL, Mangura BT, Wolman MS, Bur S, Reeves R, Thompson V, et al. Workplace contact investigations in the United States. *Int J Tuberc Lung Dis*. 2003;7(Suppl 3):S446–52.
28. Murphree R, Warkentin JV, Dunn JR, Schaffner W, Jones TF. Elephant-to-human transmission of tuberculosis, 2009. *Emerg Infect Dis*. 2011;17:366–71. <http://dx.doi.org/10.3201/eid1703.101668>
29. Centers for Disease Control and Prevention. *Salmonella* serotype enteritidis infections among workers producing poultry vaccine—Maine, November–December 2006. *MMWR Morb Mortal Wkly Rep*. 2007;56:877–9.
30. Centers for Disease Control and Prevention. Laboratory-acquired vaccinia exposures and infections—United States, 2005–2007. *MMWR Morb Mortal Wkly Rep*. 2008;57:401–4.
31. Centers for Disease Control and Prevention. Laboratory-acquired vaccinia virus infection—Virginia, 2008. *MMWR Morb Mortal Wkly Rep*. 2009;58:797–800.
32. Centers for Disease Control and Prevention. Fatal laboratory-acquired infection with an attenuated *Yersinia pestis* strain—Chicago, Illinois, 2009. *MMWR Morb Mortal Wkly Rep*. 2011;60:201–5.
33. McCollum AM, Austin C, Nawrocki J, Howland J, Pryde J, Vaid A, et al. Investigation of the first laboratory-acquired human cowpox virus infection in the United States. *J Infect Dis*. 2012;206:63–8. <http://dx.doi.org/10.1093/infdis/jis302>
34. Traxler RM, Guerra MA, Morrow MG, Haupt T, Morrison J, Saah JR, et al. Review of brucellosis cases from laboratory exposures in the United States in 2008 to 2011 and improved strategies for disease prevention. *J Clin Microbiol*. 2013;51:3132–6. <http://dx.doi.org/10.1128/JCM.00813-13>
35. Sheets CD, Harriman K, Zipprich J, Louie JK, Probert WS, Horowitz M, et al. Fatal meningococcal disease in a laboratory worker—California, 2012. *MMWR Morb Mortal Wkly Rep*. 2014;63:770–2.
36. Wilken JA, Ried C, Rickett P, Arno JN, Mendez Y, Harrison RJ, et al. Occupational HIV transmission among male adult film performers—multiple states, 2014. *MMWR Morb Mortal Wkly Rep*. 2016;65:110–4. <http://dx.doi.org/10.15585/mmwr.mm6505a3>
37. Centers for Disease Control and Prevention. Notes from the field: malaria imported from West Africa by flight crews—Florida

- and Pennsylvania, 2010. MMWR Morb Mortal Wkly Rep. 2010;59:1412.
38. Bolyard EA, Tablan OC, Williams WW, Pearson ML, Shapiro CN, Deitchman SD; Hospital Infection Control Practices Advisory Committee. Guideline for infection control in healthcare personnel, 1998. Infect Control Hosp Epidemiol. 1998;19:407–63. <http://dx.doi.org/10.2307/30142429>
 39. Advisory Committee on Immunization Practices; Centers for Disease Control and Prevention. Immunization of health-care personnel: recommendations of the Advisory Committee on Immunization Practices (ACIP). MMWR Recomm Rep. 2011;60(RR-7):1–45.
 40. Kambali S, Nantsupawat N, Lee M, Nugent K. A workplace tuberculosis case investigation in the presence of immigrant contacts from high prevalence countries. J Community Health. 2015;40:576–80. <http://dx.doi.org/10.1007/s10900-014-9946-3>
 41. CDC. Biosafety in microbiological and biomedical laboratories (BMBL), 5th edition [cited 2017 May 4]. <https://www.cdc.gov/biosafety/publications/bmbl5>
 42. Quinn MM, Henneberger PK, Braun B, Delclos GL, Fagan K, Huang V, et al.; National Institute for Occupational Safety and Health (NIOSH), National Occupational Research Agenda (NORA) Cleaning and Disinfecting in Healthcare Working Group. Cleaning and disinfecting environmental surfaces in health care: Toward an integrated framework for infection and occupational illness prevention. Am J Infect Control. 2015; 43:424–34. <http://dx.doi.org/10.1016/j.ajic.2015.01.029>
 43. Centers for Disease Control and Prevention. CDC guidance for state and local public health officials and school administrators for school (K–12) responses to influenza during the 2009–2010 school year [cited 2017 May 4]. <https://www.cdc.gov/h1n1flu/schools/schoolguidance.htm>
 44. Piper K, Youk A, James AE III, Kumar S. Paid sick days and stay-at-home behavior for influenza. PLoS One. 2017;12:e0170698. <http://dx.doi.org/10.1371/journal.pone.0170698>
 45. Fennelly KP, Nardell EA. The relative efficacy of respirators and room ventilation in preventing occupational tuberculosis. Infect Control Hosp Epidemiol. 1998;19:754–9. <http://dx.doi.org/10.2307/30141420>
 46. Jensen PA, Lambert LA, Iademarco MF, Ridzon R; CDC. Guidelines for preventing the transmission of *Mycobacterium tuberculosis* in health-care settings, 2005. MMWR Recomm Rep. 2005;54(RR-17):1–141.
 47. Ballout RA, Diab B, Harb AC, Tarabay R, Khamassi S, Akl EA. Use of safety-engineered devices by healthcare workers for intravenous and/or phlebotomy procedures in healthcare settings: a systematic review and meta-analysis. BMC Health Serv Res. 2016;16:458. <http://dx.doi.org/10.1186/s12913-016-1705-y>
 48. Bloodborne pathogens: the standard, 29 C.F.R. Sect. 1910.1030 (1991).
 49. Hageman JC, Hazim C, Wilson K, Malpiedi P, Gupta N, Bennett S, et al. Infection prevention and control for Ebola in health care settings—West Africa and United States. MMWR Suppl. 2016;65:50–6. <http://dx.doi.org/10.15585/mmwr.su6503a8>
 50. Verbeek JH, Ijaz S, Mischke C, Ruotsalainen JH, Mäkelä E, Neuvonen K, et al. Personal protective equipment for preventing highly infectious diseases due to exposure to contaminated body fluids in healthcare staff. Cochrane Database Syst Rev. 2016;4:CD011621.

Address for correspondence: Sara E. Luckhaupt, Centers for Disease Control and Prevention, 1090 Tusculum Ave, Mailstop R-17, Cincinnati, OH 45226, USA; email: sluckhaupt@cdc.gov

EID Podcast: A Tale of Two Kitchens, Meals, and Microbes

Byron Breedlove
and Martin I. Meltzer



Visit our website to listen:
[https://wwwnc.cdc.gov/eid/
article/24/6/ac-2406_article](https://wwwnc.cdc.gov/eid/article/24/6/ac-2406_article)

EMERGING INFECTIOUS DISEASES

Case Investigations of Infectious Diseases Occurring in Workplaces, United States, 2006–2015

Appendix

Methods

We conducted a literature review by referring to the methods of Haagsma et al. (6).

In March 2016, with the assistance of a CDC librarian, we searched articles published since 2006 in the PubMed database (<http://www.ncbi.nlm.nih.gov/PubMed>). Our search strategy was to combine 3 groups of relevant keywords, including case investigation, workers/workplace, and infectious diseases. Below is the detailed search strategy:

("outbreak" OR "cluster" OR "investigation" OR "epidemiology" OR "cases" OR "case report" OR "Notes") AND ("Occupational diseases" OR "Environmental Exposure" OR "laborers" OR "occupational exposure" OR "workers" OR "employees" OR "workplace" OR "industry") AND ("infectious diseases" OR "infection" OR "communicable disease" OR "Bacterial Infections" OR "viral infection" OR "pathogen" OR "zoonoses" OR "animals" OR Parasitic Diseases(Mesh)) AND eng(LA) AND United states(PL) NOT review(PT)

OR

("occupational diseases"(mesh) and "infectious"(title) AND ("2006/01/01"(PDat): "2016/12/31"(PDat)) AND Humans(Mesh) AND English(lang))) OR ((((((("Occupational Exposure"(Mesh) AND Humans(Mesh) AND English(lang))) AND (((((infectious or infection* or fungal or mycotic or mycoses or mycosis or viral or virus*)) OR (((("Bacterial Infections and Mycoses"(Mesh)) OR "Viruses"(Mesh)) OR "Bacteria"(Mesh)) OR "Fungi"(Mesh) OR pathogen*))) AND ("occupational disease" OR "occupational diseases" OR laborer* OR "occupational exposure" OR "occupational exposures" OR worker* OR employee* OR workplace* OR industry OR industries) or (pathogen* and occupation*))) AND Humans(Mesh)

AND English(lang))) AND Humans(Mesh) AND English(lang))) AND (((("Disease Outbreaks"(Mesh)) OR "Case Reports" (Publication Type)) OR ("Epidemiologic Studies"(Mesh) OR "Epidemiologic Study Characteristics as Topic"(Mesh))) OR ("Epidemiology"(Mesh) OR "epidemiology" (Subheading) OR "Molecular Epidemiology"(Mesh) OR "Bias (Epidemiology)"(Mesh) OR "Confounding Factors (Epidemiology)"(Mesh))) AND Humans(Mesh) AND English(lang))) AND ("2006/01/01"(PDat): "2016/12/31"(PDat)) AND Humans(Mesh) AND English(lang)) Filters: Publication date from 2006/01/01 to 2016/12/31; Humans; English

In addition, we searched NIOSH Health Hazard Evaluation (HHE) reports in the database (<https://www2a.cdc.gov/hhe/search.asp>) by using 2 filters:

1. Health effect: Viral, bacterial, parasitic, and fungal diseases.
2. Year published: 2006 to 2015.

We identified 2,402 articles that met the search criteria. After scanning the titles and/or abstracts, we excluded 2,045 for the following reasons:

1. Did not occur in the United States.
2. Not relevant to work-related infectious disease.

We also identified 40 HHE reports in the database and reviewed these along with the remaining articles. A total of 67 articles and 7 HHE reports from 66 infectious disease case investigations among workers were determined to be relevant to our topic, because each publication satisfied the following criteria:

1. It described a work-related infectious disease case investigation.
2. It mentioned one specific pathogen or disease.
3. It mentioned at least one specific occupational group or workplace.
4. It reported at least one symptomatic patient or seroconversion among workers.
5. The event occurred in the United States from 2006 to 2015.

The pathogens/diseases, industries (types of activity at the places of work), occupations (kinds of work), numbers of ill workers, transmission routes, and sources of exposure identified in this literature review are listed in Appendix Table 1.

Appendix Table 1. Data from reported case investigations of infectious diseases occurring in workplaces, United States, 2006–2015*

Year	State	Pathogen or disease	Industry (work setting)	Occupation	No. ill workers	Transmission mode†	Source of exposure‡	References
2006	MI	Norovirus	Food services	Server, cook	15	Direct contact	Co-workers and/or patrons	(20)
2006	ME	Salmonellosis	Manufacturing (poultry vaccine production)	Unspecified employees in the same room	5 confirmed	Vehicles	Environment (lab)	(29)
2006	IL	Mumps	Healthcare (hospital)	Employees, not specified	7	Airborne	Patient	(51)
2006	RI	Methicillin-resistant Staphylococcus aureus (MRSA) soft-tissue infections	Healthcare (hospital)	Protective service (security guards)	5	Direct contact	Patient	(52)
2006	Unk	Tuberculosis	Public administration (Navy)	Sailors	139 latent TB	Airborne	Co-worker	(53)
2006	Mult	Tuberculosis (TB)	Manufacturing (furniture installation company)	Unspecified employees in the same building	42 latent TB	Airborne	Co-worker	(54)
2006	NM	Tularemia	Landscaping services	Landscaper	1	Aerosol or vectors	Animal (rabbits?) or environment	(21)
2007	NY	Tuberculosis	Funeral service	Embalmer	1	Airborne	Human corpse	(16)
2007	MN	Salmonellosis	Food services (grocery store delicatessen)	Food preparation	2	Vehicles	Animal (chicken)	(19)
2007	CA	Coccidioidomycosis	Construction (pipeline installation contractor)	Construction laborers	12	Airborne	Environment	(23, 55)
2007	MI	Measles	U.S. Customs	Federal airport officer	2	Airborne	International traveler	(9)
2007	OR	Norovirus	Healthcare (long-term residential treatment facility)	Healthcare (nurse, doctor, nursing aid, others)	242	Vehicles	Co-workers, patients, or visitors	(56)
2007	TX	Adenovirus 14	Healthcare (hospital)	Not specified healthcare workers	6	Droplet	Patient	(57)
2007	CT	Anthrax	Manufacturing	Drum maker	1	Direct contact	Animal (goat hide)	(2, 58)
2007	AZ	Plague	National Park Service	Wildlife biologist	1	Droplet	Animal (mountain lion carcass)	(59)
2005–2007	Mult	Vaccinia virus	Academic institution/government facility	Laboratory workers	5	Direct contact	Environment (lab)	(30)
2008	MI	Norovirus	Food services	Dining service staff	3	Vehicles	Contaminated food?	(60)
2008	VA	Vaccinia virus	Academic institution	Laboratory worker	1	Direct contact	Environment (lab)	(31)
2008	Unk	Human immunodeficiency virus (HIV)	Laboratory	Technician	1	Needle puncture	Environment (lab)	(13)

Year	State	Pathogen or disease	Industry (work setting)	Occupation	No. ill workers	Transmission mode†	Source of exposure†	References
2007 – 2008	Mult	Brucellosis	Agriculture	Hunters	3	Direct contact	Animal (feral swine)	(61)
2007 – 2008	WA	Respiratory syncytial virus (RSV)	Healthcare (inpatient and outpatient cancer care)	Not specified healthcare workers	4	Droplet	Patients	(62)
2008	CA	MRSA skin infection	Zoo	Animal caretaker	20	Direct contact	Animal (elephant)	(63)
2009	IL	Plague	University	Laboratory worker	1	Direct contact	Environment (lab)	(32)
2009	NY	H1N1 influenza	High school	School employees	4	Droplet	Human	(7)
2009	MO	<i>Trichophyton tonsurans</i> Q fever	Healthcare	Staff, volunteer	21	Direct contact	Human	(64)
2009	Unk		Manufacturing (slaughterhouse)	Inspector	1	Unknown	Animal (sheep and goat)	(65)
2009	CA	Meningococcal disease	Emergency response and healthcare	Police officer and respiratory therapist	2	Droplet	Patient	(66)
2009	HI	H1N1 influenza	School	Teacher	1	Droplet	Patient	(8)
2009	IL	H1N1 influenza	Hospital	Healthcare personnel (physician, nurse, other)	20	Droplet	Patient	(11)
2009	UT	H1N1 influenza	Hospital	Physicians	5	Droplet	Patient	(67, 68)
2009	TN	Tuberculosis	Elephant refuge	Employees (caregiver and administrative)	9 latent TB	Airborne	Animal (elephant)	(28, 69)
2009	NM	<i>Mycobacterium avium</i> complex	Spa	Maintenance worker	2	Airborne	Environment	(22)
2007 – 2009	Unk	Sealpox virus	Marine mammal rehabilitation facility	Marine mammal workers	2	Direct contact	Animal (seals or sea lions)	(18)
2010	Unk	Cowpox	Research lab	Lab worker	1	Direct contact	Environment (lab)	(33)
2010	FL, PA	Malaria	Commercial airline	Flight attendants and pilots	4	Vectors	?	(37)
2010	GA, NE	Dengue	Religious organizations	Missionary workers	7	Vectors	?	(70)
2010	CO	Shiga toxin–producing <i>E. coli</i>	Dairy farm	Employees at the dairy	10	Vehicles	Animal (cows)	(71)
2010	CO	Shiga toxin–producing <i>E. coli</i>	Child care center	Not specified staff	1 confirmed 11 suspect	Vehicles	Human	(72)
2010	Unk	Legionnaires' disease	Military base	Not specified base workers	7 confirmed 22 probable	Droplet	Environment	(73)
2010	Mult	Salmonellosis	Pet store	Employee	3	Vehicles	Animal (guinea pig)	(74)
2010	TX	Tuberculosis	Jail, hospital	Hospital and correctional staff	6 active TB	Airborne	Patient	(12)
2010	Unk	Melioidosis	Reptile importation and distribution center	Not specified	1	Direct contact	Animal (reptile)	(75)
2011	AZ	Tuberculosis	Hospital	Nursing assistant, nurse, medical support assistant, physician, Officer	1 active TB 18 latent TB	Airborne	Patient	(76, 77)
2011	CA	Measles	U.S. Customs and Border Protection		1	Airborne	Patient (arriving refugee)	(10)
2011	IN, MI	Cryptosporidiosis	Fire department	Firefighter (response to a fire in a barn)	3	Vehicles	Animal (cattle)	(78)

Year	State	Pathogen or disease	Industry (work setting)	Occupation	No. ill workers	Transmission mode†	Source of exposure†	References
2011	MN	<i>Streptococcus suis</i>	Trucking company	Driver	1	Vehicles	Animal (swine)	(79)
2011	TX	Tuberculosis	Meat-packing plant	Not specified workers	20 latent TB	Airborne	Co-workers	(40)
2008 – 2011	VA	Campylobacteriosis	Poultry-processing plant	Poultry workers	29	Vehicles	Animal (chicken)	(17, 80)
2008 – 2011	Mult	Brucellosis	University, public health lab	Laboratory workers	5	Direct contact	Environment (lab)	(34)
2009 – 2011	NY	Legionnaires' disease	Automobile and scrap metal shredding facility	Employees	4	Droplet	Environment	(81)
2012	CA	Coccidioidomycosis	Outdoor television-filming event	Cast or crew members	5 confirmed 5 probable	Airborne	Environment	(24)
2012	NE	Histoplasmosis	Day camp	Counselors	32	Airborne	Environment	(26)
2012	CA	Meningococcal disease	Research laboratory	Microbiologist	1	Direct contact	Environment (lab)	(35)
2012	IN, KY	Lymphocytic choriomeningitis virus (LCMV)	Rodent-breeding facilities	Employees	31	Direct contact	Animal	(82)
2012	OH	Variant influenza A(H3N2)	State fair	Swine exhibitor	20	Droplet	Animal (swine)	(83)
2012	AK	Tuberculosis	Long-term-care facility	Nursing assistant, environmental services	16 latent TB	Airborne	Resident/patient	(84, 85)
2013	MA	Vaccinia virus	Research laboratory	Laboratory worker	1	Direct contact	Environment (lab)	(86)
2013	MN	Salmonellosis	Hospital	Phlebotomist	1	Vehicle	Patient	(87)
2013	KS	Cryptosporidiosis	Emergency responders	Law enforcement officers, tow truck employees	15	Vehicle	Animal (calves)	(88)
2013	PA	<i>E. coli</i> O157:H7	Restaurant	Food handlers: beef grinding	9	Vehicle	Animal (beef)	(89)
2009 – 2013	CA	Coccidioidomycosis	Prison	Employees	103	Airborne	Environment	(90, 91)
2014	TX	Ebola	Hospital	Nurses	2	Direct contact	Patient	(14)
2014	NY	Ebola	International medical humanitarian organization	Humanitarian aid worker	1	Direct contact	Patient	(15)
2014	CA	HIV	Adult-film production	Performers	2	Sexual contact	Co-worker	(36)
2011 – 2014	CA	Coccidioidomycosis	Construction (solar power farms)	Construction labors	44	Airborne	Environment	(25)
2015	AZ	Measles	Outpatient healthcare facility	Nurse	1	Airborne	Patient	(92)

*Unk, unknown; Mult, multiple.

†Suspected or confirmed.

Expanded Reference List

1. Price GM. Medicine and the industries. *Bulletin of the American Academy of Medicine*. 1917;18:134–9.
2. Centers for Disease Control and Prevention (CDC). Cutaneous anthrax associated with drum making using goat hides from West Africa—Connecticut, 2007. *MMWR Morb Mortal Wkly Rep*. 2008;57:628–31. [PubMed](#)
3. Fowler RA, Lapinsky SE, Hallett D, Detsky AS, Sibbald WJ, Slutsky AS, et al.; Toronto SARS Critical Care Group. Critically ill patients with severe acute respiratory syndrome. *JAMA*. 2003;290:367–73. [PubMed](#) <http://dx.doi.org/10.1001/jama.290.3.367>
4. Suarathana E, McFadden JD, Laney AS, Kreiss K, Anderson HA, Hunt DC, et al. Occupational distribution of persons with confirmed 2009 H1N1 influenza. *J Occup Environ Med*. 2010;52:1212–6. [PubMed](#) <http://dx.doi.org/10.1097/JOM.0b013e3181fd32e4>
5. Cummings KJ, Choi MJ, Esswein EJ, de Perio MA, Harney JM, Chung WM, et al. Addressing infection prevention and control in the first U.S. community hospital to care for patients with Ebola virus disease: context for national recommendations and future strategies. *Ann Intern Med*. 2016;165:41–9. [PubMed](#) <http://dx.doi.org/10.7326/M15-2944>
6. Haagsma JA, Tariq L, Heederik DJ, Havelaar AH. Infectious disease risks associated with occupational exposure: a systematic review of the literature. *Occup Environ Med*. 2012;69:140–6. [PubMed](#) <http://dx.doi.org/10.1136/oemed-2011-100068>
7. Lessler J, Reich NG, Cummings DA. New York City Department of Health and Mental Hygiene Swine Influenza Investigation Team. Outbreak of 2009 pandemic influenza A(H1N1) at a New York City school. *N Engl J Med*. 2009;361:2628–36.
8. Centers for Disease Control and Prevention (CDC). Outbreak of 2009 pandemic influenza A (H1N1) at a school—Hawaii, May 2009. *MMWR Morb Mortal Wkly Rep*. 2010;58:1440–4. [PubMed](#)
9. Chen TH, Kutty P, Lowe LE, Hunt EA, Blostein J, Espinoza R, et al. Measles outbreak associated with an international youth sporting event in the United States, 2007. *Pediatr Infect Dis J*. 2010;29:794–800. [PubMed](#) <http://dx.doi.org/10.1097/INF.0b013e3181dbaacf>

10. Centers for Disease Control and Prevention (CDC). Measles outbreak associated with an arriving refugee— Los Angeles County, California, August–September 2011. MMWR Morb Mortal Wkly Rep. 2012;61:385–9. [PubMed](#)
11. Magill SS, Black SR, Wise ME, Kallen AJ, Lee SJ, Gardner T, et al. Investigation of an outbreak of 2009 pandemic influenza A virus (H1N1) infections among healthcare personnel in a Chicago hospital. Infect Control Hosp Epidemiol. 2011;32:611–5. [PubMed](#)
<http://dx.doi.org/10.1086/660097>
12. Medrano BA, Salinas G, Sanchez C, Miramontes R, Restrepo BI, Haddad MB, et al. A missed tuberculosis diagnosis resulting in hospital transmission. Infect Control Hosp Epidemiol. 2014;35:534–7. [PubMed](#) <http://dx.doi.org/10.1086/675833>
13. Joyce MP, Kuhar D, Brooks JT. Notes from the field: occupationally acquired HIV infection among health care workers—United States, 1985–2013. MMWR Morb Mortal Wkly Rep. 2015;63:1245–6. [PubMed](#)
14. Chevalier MS, Chung W, Smith J, Weil LM, Hughes SM, Joyner SN, et al.; Centers for Disease Control and Prevention (CDC). Ebola virus disease cluster in the United States—Dallas County, Texas, 2014. MMWR Morb Mortal Wkly Rep. 2014;63:1087–8. [PubMed](#)
15. Yacisin K, Balter S, Fine A, Weiss D, Ackelsberg J, Prezant D, et al.; Centers for Disease Control and Prevention (CDC). Ebola virus disease in a humanitarian aid worker—New York City, October 2014. MMWR Morb Mortal Wkly Rep. 2015;64:321–3. [PubMed](#)
16. Anderson JA, Meissner JS, Ahuja SD, Shashkina E, O’Flaherty T, Proops DC. Confirming *Mycobacterium tuberculosis* transmission from a cadaver to an embalmer using molecular epidemiology. Am J Infect Control. 2015;43:543–5. [PubMed](#)
<http://dx.doi.org/10.1016/j.ajic.2015.01.027>
17. de Perio MA, Niemeier RT, Levine SJ, Gruszynski K, Gibbins JD. *Campylobacter* infection in poultry-processing workers, Virginia, USA, 2008–2011. Emerg Infect Dis. 2013;19:286–8. [PubMed](#) <http://dx.doi.org/10.3201/eid1902.121147>
18. Roess AA, Levine RS, Barth L, Monroe BP, Carroll DS, Damon IK, et al. Sealpox virus in marine mammal rehabilitation facilities, North America, 2007–2009. Emerg Infect Dis. 2011;17:2203–8. [PubMed](#) <http://dx.doi.org/10.3201/eid1712.101945>

19. Hedican E, Miller B, Ziemer B, LeMaster P, Jawahir S, Leano F, et al. Salmonellosis outbreak due to chicken contact leading to a foodborne outbreak associated with infected delicatessen workers. *Foodborne Pathog Dis.* 2010;7:995–7. [PubMed http://dx.doi.org/10.1089/fpd.2009.0495](http://dx.doi.org/10.1089/fpd.2009.0495)
20. Centers for Disease Control and Prevention (CDC). Norovirus outbreak associated with ill food-service workers—Michigan, January–February 2006. *MMWR Morb Mortal Wkly Rep.* 2007;56:1212–6. [PubMed](http://dx.doi.org/10.1093/mmwr/mmwr5612a1)
21. Hofinger DM, Cardona L, Mertz GJ, Davis LE. Tularemic meningitis in the United States. *Arch Neurol.* 2009;66:523–7. [PubMed http://dx.doi.org/10.1001/archneurol.2009.14](http://dx.doi.org/10.1001/archneurol.2009.14)
22. Moraga-McHaley SA, Landen M, Krapfl H, Sewell CM. Hypersensitivity pneumonitis with *Mycobacterium avium* complex among spa workers. *Int J Occup Environ Health.* 2013;19:55–61. [PubMed http://dx.doi.org/10.1179/2049396712Y.0000000015](http://dx.doi.org/10.1179/2049396712Y.0000000015)
23. Cummings KC, McDowell A, Wheeler C, McNary J, Das R, Vugia DJ, et al. Point-source outbreak of coccidioidomycosis in construction workers. *Epidemiol Infect.* 2010;138:507–11. [PubMed http://dx.doi.org/10.1017/S0950268809990999](http://dx.doi.org/10.1017/S0950268809990999)
24. Wilken JA, Marquez P, Terashita D, McNary J, Windham G, Materna B; Centers for Disease Control and Prevention. Coccidioidomycosis among cast and crew members at an outdoor television filming event—California, 2012. *MMWR Morb Mortal Wkly Rep.* 2014;63:321–4. [PubMed](http://dx.doi.org/10.1093/mmwr/mmwr6303a1)
25. Wilken JA, Sondermeyer G, Shusterman D, McNary J, Vugia DJ, McDowell A, et al. Coccidioidomycosis among workers constructing solar power farms, California, USA, 2011–2014. *Emerg Infect Dis.* 2015;21:1997–2005. [PubMed http://dx.doi.org/10.3201/eid2111.150129](http://dx.doi.org/10.3201/eid2111.150129)
26. Centers for Disease Control and Prevention. Notes from the field: histoplasmosis outbreak among day camp attendees—Nebraska, June 2012. *MMWR Morb Mortal Wkly Rep.* 2012;21:747–8.
27. Davidow AL, Mangura BT, Wolman MS, Bur S, Reves R, Thompson V, et al. Workplace contact investigations in the United States. *Int J Tuberc Lung Dis.* 2003;7(Suppl 3):S446–52. [PubMed](http://dx.doi.org/10.1093/ajph/93.3.446)
28. Murphree R, Warkentin JV, Dunn JR, Schaffner W, Jones TF. Elephant-to-human transmission of tuberculosis, 2009. *Emerg Infect Dis.* 2011;17:366–71. [PubMed http://dx.doi.org/10.3201/eid1703.101668](http://dx.doi.org/10.3201/eid1703.101668)
29. Centers for Disease Control and Prevention. *Salmonella* serotype enteritidis infections among workers producing poultry vaccine—Maine, November–December 2006. *MMWR Morb Mortal Wkly Rep.* 2007;56:877–9. [PubMed](http://dx.doi.org/10.1093/mmwr/mmwr5612a1)

30. Centers for Disease Control and Prevention. Laboratory-acquired vaccinia exposures and infections—United States, 2005–2007. MMWR Morb Mortal Wkly Rep. 2008;57:401–4. [PubMed](#)
31. Centers for Disease Control and Prevention. Laboratory-acquired vaccinia virus infection—Virginia, 2008. MMWR Morb Mortal Wkly Rep. 2009;58:797–800. [PubMed](#)
32. Centers for Disease Control and Prevention. Fatal laboratory-acquired infection with an attenuated *Yersinia pestis* strain—Chicago, Illinois, 2009. MMWR Morb Mortal Wkly Rep. 2011;60:201–5. [PubMed](#)
33. McCollum AM, Austin C, Nawrocki J, Howland J, Pryde J, Vaid A, et al. Investigation of the first laboratory-acquired human cowpox virus infection in the United States. J Infect Dis. 2012;206:63–8. [PubMed](#) <http://dx.doi.org/10.1093/infdis/jis302>
34. Traxler RM, Guerra MA, Morrow MG, Haupt T, Morrison J, Saah JR, et al. Review of brucellosis cases from laboratory exposures in the United States in 2008 to 2011 and improved strategies for disease prevention. J Clin Microbiol. 2013;51:3132–6. [PubMed](#) <http://dx.doi.org/10.1128/JCM.00813-13>
35. Sheets CD, Harriman K, Zipprich J, Louie JK, Probert WS, Horowitz M, et al. Fatal meningococcal disease in a laboratory worker—California, 2012. MMWR Morb Mortal Wkly Rep. 2014;63:770–2. [PubMed](#)
36. Wilken JA, Ried C, Rickett P, Arno JN, Mendez Y, Harrison RJ, et al. Occupational HIV transmission among male adult film performers—multiple states, 2014. MMWR Morb Mortal Wkly Rep. 2016;65:110–4. [PubMed](#) <http://dx.doi.org/10.15585/mmwr.mm6505a3>
37. Centers for Disease Control and Prevention. Notes from the field: malaria imported from West Africa by flight crews—Florida and Pennsylvania, 2010. MMWR Morb Mortal Wkly Rep. 2010;59:1412. [PubMed](#)
38. Bolyard EA, Tablan OC, Williams WW, Pearson ML, Shapiro CN, Deitchman SD; Hospital Infection Control Practices Advisory Committee. Guideline for infection control in healthcare personnel, 1998. Infect Control Hosp Epidemiol. 1998;19:407–63. [PubMed](#) <http://dx.doi.org/10.2307/30142429>
39. Advisory Committee on Immunization Practices; Centers for Disease Control and Prevention. Immunization of health-care personnel: recommendations of the Advisory Committee on Immunization Practices (ACIP). MMWR Recomm Rep. 2011;60(RR-7):1–45. [PubMed](#)

40. Kambali S, Nantsupawat N, Lee M, Nugent K. A workplace tuberculosis case investigation in the presence of immigrant contacts from high prevalence countries. *J Community Health*. 2015;40:576–80. [PubMed](#) <http://dx.doi.org/10.1007/s10900-014-9946-3>
41. CDC. Biosafety in microbiological and biomedical laboratories (BMBL), 5th edition [cited 2017 May 4]. <https://www.cdc.gov/biosafety/publications/bmbl5>
42. Quinn MM, Henneberger PK, Braun B, Delclos GL, Fagan K, Huang V, et al.; National Institute for Occupational Safety and Health (NIOSH), National Occupational Research Agenda (NORA) Cleaning and Disinfecting in Healthcare Working Group. Cleaning and disinfecting environmental surfaces in health care: Toward an integrated framework for infection and occupational illness prevention. *Am J Infect Control*. 2015;43:424–34. [PubMed](#) <http://dx.doi.org/10.1016/j.ajic.2015.01.029>
43. Centers for Disease Control and Prevention. CDC guidance for state and local public health officials and school administrators for school (K–12) responses to influenza during the 2009–2010 school year [cited 2017 May 4]. <https://www.cdc.gov/h1n1flu/schools/schoolguidance.htm>
44. Piper K, Youk A, James AE III, Kumar S. Paid sick days and stay-at-home behavior for influenza. *PLoS One*. 2017;12:e0170698. [PubMed](#) <http://dx.doi.org/10.1371/journal.pone.0170698>
45. Fennelly KP, Nardell EA. The relative efficacy of respirators and room ventilation in preventing occupational tuberculosis. *Infect Control Hosp Epidemiol*. 1998;19:754–9. [PubMed](#) <http://dx.doi.org/10.2307/30141420>
46. Jensen PA, Lambert LA, Iademarco MF, Ridzon R; CDC. Guidelines for preventing the transmission of *Mycobacterium tuberculosis* in health-care settings, 2005. *MMWR Recomm Rep*. 2005;54(RR-17):1–141. [PubMed](#)
47. Ballout RA, Diab B, Harb AC, Tarabay R, Khamassi S, Akl EA. Use of safety-engineered devices by healthcare workers for intravenous and/or phlebotomy procedures in healthcare settings: a systematic review and meta-analysis. *BMC Health Serv Res*. 2016;16:458. [PubMed](#) <http://dx.doi.org/10.1186/s12913-016-1705-y>
48. Bloodborne pathogens: the standard, 29 C.F.R. Sect. 1910.1030 (1991).
49. Hageman JC, Hazim C, Wilson K, Malpiedi P, Gupta N, Bennett S, et al. Infection prevention and control for Ebola in health care settings—West Africa and United States. *MMWR Suppl*. 2016;65:50–6. [PubMed](#) <http://dx.doi.org/10.15585/mmwr.su6503a8>

50. Verbeek JH, Ijaz S, Mischke C, Ruotsalainen JH, Mäkelä E, Neuvonen K, et al. Personal protective equipment for preventing highly infectious diseases due to exposure to contaminated body fluids in healthcare staff. *Cochrane Database Syst Rev*. 2016;4:CD011621. [PubMed](#)
51. Bonebrake AL, Silkaitis C, Monga G, Galat A, Anderson J, Trad JT, et al. Effects of mumps outbreak in hospital, Chicago, Illinois, USA, 2006. *Emerg Infect Dis*. 2010;16:426–32. [PubMed](#)
<http://dx.doi.org/10.3201/eid1603.090198>
52. Patrozou E, Reid K, Jefferson J, Mermel LA. A cluster of community-acquired methicillin-resistant *Staphylococcus aureus* infections in hospital security guards. *Infect Control Hosp Epidemiol*. 2009;30:386–8. [PubMed](#) <http://dx.doi.org/10.1086/596611>
53. Centers for Disease Control and Prevention (CDC). Latent tuberculosis infection among sailors and civilians aboard U.S.S. Ronald Reagan—United States, January–July 2006. *MMWR Morb Mortal Wkly Rep*. 2007;55:1381–2. [PubMed](#)
54. Centers for Disease Control and Prevention (CDC). Workplace-based investigation of contacts of a patient with highly infectious tuberculosis—Maryland, District of Columbia, and Virginia, 2006. *MMWR Morb Mortal Wkly Rep*. 2008;57:94–8. [PubMed](#)
55. Das R, McNary J, Fitzsimmons K, Dobraca D, Cummings K, Mohle-Boetani J, et al. Occupational coccidioidomycosis in California: outbreak investigation, respirator recommendations, and surveillance findings. *J Occup Environ Med*. 2012;54:564–71. [PubMed](#)
<http://dx.doi.org/10.1097/JOM.0b013e3182480556>
56. Centers for Disease Control and Prevention (CDC). Recurring norovirus outbreaks in a long-term residential treatment facility—Oregon, 2007. *MMWR Morb Mortal Wkly Rep*. 2009;58:694–8. [PubMed](#)
57. Yun HC, Prakash V. Transmission of adenovirus serotype 14 in the health care setting. *Clin Infect Dis*. 2008;46:1935–6. [PubMed](#) <http://dx.doi.org/10.1086/588558>
58. Nguyen TQ, Clark N, The 2006 NYC Anthrax Working Group. Public health and environmental response to the first case of naturally acquired inhalational anthrax in the United States in 30 years: infection of a New York City resident who worked with dried animal hides. *J Public Health Manag Pract*. 2010;16:189–200. [PubMed](#)
<http://dx.doi.org/10.1097/PHH.0b013e3181ca64f2>

59. Wong D, Wild MA, Walburger MA, Higgins CL, Callahan M, Czarnecki LA, et al. Primary pneumonic plague contracted from a mountain lion carcass. *Clin Infect Dis*. 2009;49:e33–8. [PubMed http://dx.doi.org/10.1086/600818](http://dx.doi.org/10.1086/600818)
60. Centers for Disease Control and Prevention (CDC). Norovirus outbreaks on three college campuses—California, Michigan, and Wisconsin, 2008. *MMWR Morb Mortal Wkly Rep*. 2009;58:1095–100. [PubMed](http://dx.doi.org/10.1093/mmwr/mmwr5810a1)
61. Centers for Disease Control and Prevention (CDC). *Brucella suis* infection associated with feral swine hunting—three states, 2007–2008. *MMWR Morb Mortal Wkly Rep*. 2009;58:618–21. [PubMed](http://dx.doi.org/10.1093/mmwr/mmwr5810a1)
62. Chu HY, Englund JA, Podczervinski S, Kuypers J, Campbell AP, Boeckh M, et al. Nosocomial transmission of respiratory syncytial virus in an outpatient cancer center. *Biol Blood Marrow Transplant*. 2014;20:844–51. [PubMed http://dx.doi.org/10.1016/j.bbmt.2014.02.024](http://dx.doi.org/10.1016/j.bbmt.2014.02.024)
63. Centers for Disease Control and Prevention (CDC). Methicillin-resistant *Staphylococcus aureus* skin infections from an elephant calf—San Diego, California, 2008. *MMWR Morb Mortal Wkly Rep*. 2009;58:194–8. [PubMed](http://dx.doi.org/10.1093/mmwr/mmwr5810a1)
64. Shroba J, Olson-Burgess C, Preuett B, Abdel-Rahman SM. A large outbreak of *Trichophyton tonsurans* among health care workers in a pediatric hospital. *Am J Infect Control*. 2009;37:43–8. [PubMed http://dx.doi.org/10.1016/j.ajic.2007.11.008](http://dx.doi.org/10.1016/j.ajic.2007.11.008)
65. Towey A, Petitti C. OSHA compliance issues: a case of Q fever. *J Occup Environ Hyg*. 2009;6:D63–5. [PubMed http://dx.doi.org/10.1080/15459620903152956](http://dx.doi.org/10.1080/15459620903152956)
66. Centers for Disease Control and Prevention (CDC). Occupational transmission of *Neisseria meningitidis*—California, 2009. *MMWR Morb Mortal Wkly Rep*. 2010;59:1480–3. [PubMed](http://dx.doi.org/10.1093/mmwr/mmwr5910a1)
67. de Perio MA, Brueck SE, Mueller CA. Evaluation of 2009 pandemic influenza A(H1N1) virus exposure among internal medicine housestaff and fellows. *Health Hazard Evaluation Report*, HETA 2009–0206–3117. October 2010 [cited 2017 May 4], <https://www.cdc.gov/niosh/hhe/reports/pdfs/2009-0206-3117.pdf>.
68. de Perio MA, Brueck SE, Mueller CA, Milne CK, Rubin MA, Gundlapalli AV, et al. Evaluation of 2009 pandemic influenza A(H1N1) exposures and illness among physicians in training. *Am J Infect Control*. 2012;40:617–21. [PubMed http://dx.doi.org/10.1016/j.ajic.2012.01.014](http://dx.doi.org/10.1016/j.ajic.2012.01.014)
69. Niemeier RT, Mead K, de Perio MA, Martin SB Jr, Burr GA. Evaluation of potential employee exposures to *Mycobacterium tuberculosis* at an elephant refuge. *HHE Report No.* 2010–0080–

3235. April 2015 [cited 2017 May 4], <https://www.cdc.gov/niosh/hhe/reports/pdfs/2010-0080-3235.pdf>
70. Centers for Disease Control and Prevention (CDC). Dengue virus infections among travelers returning from Haiti—Georgia and Nebraska, October 2010. MMWR Morb Mortal Wkly Rep. 2011;60:914–7. [PubMed](#)
71. Centers for Disease Control and Prevention (CDC). Outbreak of Shiga toxin-producing *Escherichia coli* O111 infections associated with a correctional facility dairy—Colorado, 2010. MMWR Morb Mortal Wkly Rep. 2012;61:149–52. [PubMed](#)
72. Brown JA, Hite DS, Gillim-Ross LA, Maguire HF, Bennett JK, Patterson JJ, et al. Outbreak of Shiga toxin-producing *Escherichia coli* serotype O26: H11 infection at a child care center in Colorado. *Pediatr Infect Dis J*. 2012;31:379–83. [PubMed](#) <http://dx.doi.org/10.1097/INF.0b013e3182457122>
73. Ambrose J, Hampton LM, Fleming-Dutra KE, Marten C, McClusky C, Perry C, et al. Large outbreak of Legionnaires' disease and Pontiac fever at a military base. *Epidemiol Infect*. 2014;142:2336–46. [PubMed](#) <http://dx.doi.org/10.1017/S0950268813003440>
74. Bartholomew ML, Heffernan RT, Wright JG, Klos RF, Monson T, Khan S, et al. Multistate outbreak of *Salmonella enterica* serotype enteritidis infection associated with pet guinea pigs. *Vector Borne Zoonotic Dis*. 2014;14:414–21. [PubMed](#) <http://dx.doi.org/10.1089/vbz.2013.1506>
75. Benoit TJ, Blaney DD, Gee JE, Elrod MG, Hoffmaster AR, Doker TJ, et al.; Centers for Disease Control and Prevention (CDC). Melioidosis cases and selected reports of occupational exposures to *Burkholderia pseudomallei*—United States, 2008–2013. MMWR Surveill Summ. 2015;64:1–9. [PubMed](#)
76. de Perio MA, Niemeier RT. Evaluation of exposure to tuberculosis among employees at a medical center—Arizona. Health Hazard Evaluation Report, HETA 2011–0137–3159. April 2012 [cited 2017 May 4], <https://www.cdc.gov/niosh/hhe/reports/pdfs/2011-0137-3159.pdf>
77. de Perio MA, Niemeier RT. Evaluation of exposure to tuberculosis among employees at a medical center. *J Occup Environ Hyg*. 2014;11:D63–8. [PubMed](#) <http://dx.doi.org/10.1080/15459624.2014.888075>
78. Centers for Disease Control and Prevention (CDC). Outbreak of cryptosporidiosis associated with a firefighting response—Indiana and Michigan, June 2011. MMWR Morb Mortal Wkly Rep. 2012;61:153–6. [PubMed](#)

79. Fowler HN, Brown P, Rovira A, Shade B, Klammer K, Smith K, et al. *Streptococcus suis* meningitis in swine worker, Minnesota, USA. Emerg Infect Dis. 2013;19:330–1. [PubMed](#)
<http://dx.doi.org/10.3201/eid1902.120918>
80. de Perio MA, Gibbins JD, Niemeier RT. *Campylobacter* infection and exposures among employees at a poultry processing plant—Virginia. Health Hazard Evaluation Report, HETA 2011–0058–3157. April 2012 [cited 2017 May 4], <https://www.cdc.gov/niosh/hhe/reports/pdfs/2011-0058-3157.pdf>
81. Boylstein R, Bailey R, Piacitelli C, Schuler C, Cox-Ganser J, Kreiss K. Legionnaires’ disease at an automobile and scrap metal shredding facility, New York. Health Hazard Evaluation Report, HETA 2011–0109–3162. August 2012 [cited 2017 May 4], <https://www.cdc.gov/niosh/hhe/reports/pdfs/2011-0109-3162.pdf>
82. Knust B, Ströher U, Edison L, Albariño CG, Lovejoy J, Armeanu E, et al. Lymphocytic choriomeningitis virus in employees and mice at multipremises feeder-rodent operation, United States, 2012. Emerg Infect Dis. 2014;20:240–7. [PubMed](#)
<http://dx.doi.org/10.3201/eid2002.130860>
83. Greenbaum A, Quinn C, Bailer J, Su S, Havers F, Durand LO, et al. Investigation of an outbreak of variant influenza A(H3N2) virus infection associated with an agricultural fair—Ohio, August 2012. J Infect Dis. 2015;212:1592–9. [PubMed](#) <http://dx.doi.org/10.1093/infdis/jiv269>
84. de Perio MA, Niemeier RT. Evaluation of exposure to tuberculosis among employees at a long-term care facility. Health Hazard Evaluation Report No. 2012–0137–3178 May 2013. Revised June 2013 [cited 2017 May 4], <https://www.cdc.gov/niosh/hhe/reports/pdfs/2012-0137-3178.pdf>.
85. Jackson DA, Mailer K, Porter KA, Niemeier RT, Fearey DA, Pope L, et al. Challenges in assessing transmission of *Mycobacterium tuberculosis* in long-term-care facilities. Am J Infect Control. 2015;43:992–6. [PubMed](#) <http://dx.doi.org/10.1016/j.ajic.2015.03.035>
86. Hsu CH, Farland J, Winters T, Gunn J, Caron D, Evans J, et al.; Centers for Disease Control and Prevention (CDC). Laboratory-acquired vaccinia virus infection in a recently immunized person—Massachusetts, 2013. MMWR Morb Mortal Wkly Rep. 2015;64:435–8. [PubMed](#)
87. Centers for Disease Control and Prevention (CDC). Occupationally acquired *Salmonella* I 4,12:i:1,2 infection in a phlebotomist—Minnesota, January 2013. MMWR Morb Mortal Wkly Rep. 2013;62:525. [PubMed](#)

88. Webb LM, Tubach SA, Hunt DC; Centers for Disease Control and Prevention (CDC). Outbreak of cryptosporidiosis among responders to a rollover of a truck carrying calves—Kansas, April 2013. *MMWR Morb Mortal Wkly Rep.* 2014;63:1185–8. [PubMed](#)
89. Torso LM, Voorhees RE, Forest SA, Gordon AZ, Silvestri SA, Kissler B, et al. *Escherichia coli* O157:H7 outbreak associated with restaurant beef grinding. *J Food Prot.* 2015;78:1272–9. [PubMed](#) <http://dx.doi.org/10.4315/0362-028X.JFP-14-545>
90. de Perio MA, Niemeier RT, Burr GA. *Coccidioides* exposure and coccidioidomycosis among prison employees, California, United States. *Emerg Infect Dis.* 2015;21:1031–3. [PubMed](#) <http://dx.doi.org/10.3201/eid2106.141201>
91. de Perio MA, Burr GA. Evaluation of coccidioides exposures and coccidioidomycosis infections among prison employees. Health Hazard Evaluation Report No. 2013–0113–3198. January 2014 [cited 2017 May 4], <https://www.cdc.gov/niosh/hhe/reports/pdfs/2013-0113-3198.pdf>.
92. Jones J, Klein R, Popescu S, Rose K, Kretschmer M, Carrigan A, et al. Lack of measles transmission to susceptible contacts from a health care worker with probable secondary vaccine failure—Maricopa County, Arizona, 2015. *MMWR Morb Mortal Wkly Rep.* 2015;64:832–3. [PubMed](#) <http://dx.doi.org/10.15585/mmwr.mm6430a5>