

## References

- Godfroid J. Brucellosis in wildlife. *Rev Sci Tech*. 2002; 21:277–86. <https://doi.org/10.20506/rst.21.2.1333>
- Garin-Bastuji B, Oudar J, Richard Y, Gastellu J. Isolation of *Brucella melitensis* biovar 3 from a chamois (*Rupicapra rupicapra*) in the southern French Alps. *J Wildl Dis*. 1990;26:116–8. <https://doi.org/10.7589/0090-3558-26.1.116>
- Ferroglio E, Tolari F, Bollo E, Bassano B. Isolation of *Brucella melitensis* from alpine ibex. *J Wildl Dis*. 1998;34:400–2. <https://doi.org/10.7589/0090-3558-34.2.400>
- Muñoz PM, Boadella M, Arnal M, de Miguel MJ, Revilla M, Martínez D, et al. Spatial distribution and risk factors of brucellosis in Iberian wild ungulates. *BMC Infect Dis*. 2010;10:46. <https://doi.org/10.1186/1471-2334-10-46>
- Lindstrom E. The role of medium-sized carnivores in the Nordic boreal forest. *Finnish Game Research*. 1989;46:53–63.
- Alton GG, Jones LM, Pietz DE. Laboratory techniques in brucellosis. *Monogr Ser World Health Organ*. 1975;55:1–163.
- Wang Q, Zhao S, Wureli H, Xie S, Chen C, Wei Q, et al. *Brucella melitensis* and *B. abortus* in eggs, larvae and engorged females of *Dermacentor marginatus*. *Ticks Tick Borne Dis*. 2018;9:1045–8. <https://doi.org/10.1016/j.ttbdis.2018.03.021>
- Maquart M, Le Flèche P, Foster G, Tryland M, Ramisse F, Dønne B, et al. MLVA-16 typing of 295 marine mammal *Brucella* isolates from different animal and geographic origins identifies 7 major groups within *Brucella ceti* and *Brucella pinnipedialis*. *BMC Microbiol*. 2009;9:145. <https://doi.org/10.1186/1471-2180-9-145>
- Li S, Hu X. Serological investigation on brucellosis of cattle and sheep in Nilka County. *Chinese Livestock and Poultry Breeding*. 2017;4:41–3.
- Davis DS, Boeer WJ, Mims JP, Heck FC, Adams LG. *Brucella abortus* in coyotes. I. A serologic and bacteriologic survey in eastern Texas. *J Wildl Dis*. 1979;15:367–72. <https://doi.org/10.7589/0090-3558-15.3.367>

Addresses for correspondence: Yuanzhi Wang or Hai Jiang, School of Medicine, Shihezi University, Shihezi, Xinjiang Uygur Autonomous Region 832002, China; email: wangyuanzhi621@126.com or jianghai@icdc.com

## Multicenter Study of Azole-Resistant *Aspergillus fumigatus* Clinical Isolates, Taiwan<sup>1</sup>

Chi-Jung Wu, Wei-Lun Liu, Chih-Cheng Lai, Chien-Ming Chao, Wen-Chien Ko, Hsuan-Chen Wang, Ching-Tzu Dai, Ming-I Hsieh, Pui-Ching Choi, Jia-Ling Yang, Yee-Chun Chen

<sup>1</sup>Results from this study were presented in part at the 30th International Congress of Chemotherapy and Infection, November 24–27, 2017, Taipei, Taiwan.

Author affiliations: National Health Research Institutes, Zhunan, Taiwan (C.-J. Wu, H.-C. Wang, M.-I. Hsieh, Y.-C., Chen); National Cheng Kung University Hospital and College of Medicine, Tainan, Taiwan (C.-J. Wu, W.-C. Ko); Fu Jen Catholic University Hospital and College of Medicine, New Taipei, Taiwan (W.-L. Liu), Chi Mei Medical Center, Tainan (C.-C. Lai, C.-M. Chao); National Taiwan University Hospital and College of Medicine, Taipei, Taiwan (C.-T. Dai, P.-C. Choi, J.-L. Yang, Y.-C. Chen)

DOI: <https://doi.org/10.3201/eid2604.190840>


In a multicenter study, we determined a prevalence rate of 4% for azole-resistant *Aspergillus fumigatus* in Taiwan. Resistance emerged mainly from the environment (TR<sub>34</sub>/L98H, TR<sub>34</sub>/L98H/S297T/F495I, and TR<sub>46</sub>/Y121F/T289A mutations) but occasionally during azole treatment. A high mortality rate observed for azole-resistant aspergillosis necessitates diagnostic stewardship in healthcare and antifungal stewardship in the environment.

Worldwide emergence of azole-resistant *Aspergillus fumigatus* since the late 2000s threatens human health (1). Azole resistance in *A. fumigatus* might develop during patient therapy with medical azoles or through exposure to azole fungicides in the environment; environmental exposure predominantly involves TR<sub>34</sub>/L98H and TR<sub>46</sub>/Y121F/T289A mutations in *cyp51A* (1).

Taiwan is an island country in eastern Asia that is geographically separated from mainland Eurasia and has a long history of azole fungicide use. To delineate the influence of clinical and environmental use of azoles on resistance, we conducted a multicenter study that investigated 375 *A. fumigatus* sensu stricto isolates collected during August 2011–March 2018 from 297 patients at 11 hospitals in Taiwan (Appendix Table 1, Figure 1, <https://wwwnc.cdc.gov/EID/article/26/4/19-0840-App1.pdf>).

We confirmed the presence of azole resistance by using the Clinical Laboratory Standard Institute method (Appendix Table 1) (2). Isolates resistant to  $\geq 1$  medical azoles (itraconazole, voriconazole, posaconazole, and isavuconazole) were defined as azole-resistant *A. fumigatus* and examined for resistance mechanisms, microsatellite-based phylogenetic relatedness, and growth rates following previously described methods (3,4).

Overall, 19 isolates from 12 patients were azole-resistant *A. fumigatus*. These isolates had resistance rates of 4.0%/patient and 5.1%/isolate analyses (Appendix Tables 2, 3). Ten (83.3%) patients harbored azole-resistant *A. fumigatus* that had environmental mutations, including TR<sub>34</sub>/L98H (5 isolates, 5 patients), TR<sub>34</sub>/L98H/S297T/F495I (7 isolates, 4 patients), and TR<sub>46</sub>/Y121F/T289A (1 isolate) mutations. This observation



**Figure.** Genetic relatedness among *Aspergillus fumigatus* isolates based on microsatellite genotyping, Taiwan. Scale bar indicates percentage relatedness. AF, *A. fumigatus*; C, clinical; E, environmental; R, azole-resistant; S, azole-susceptible; STR, short tandem repeat; TW, Taiwan.

is consistent with the estimated global prevalence of azole resistance in *Aspergillus* (3%–6%) and the predominance of environmental resistance mechanisms in azole-resistant *A. fumigatus* (1,5).

Phylogenetic analysis showed that TR<sub>34</sub>/L98H/S297T/F495I isolates from 2 patients with pulmonary aspergillosis (isolates B44 and B51 in 2012, isolates E071, E073, and E074 in 2015) (Figure) belonged to a local microsatellite genotype widely distributed in the environment of Taiwan (3), indicating that this clone has locally evolved and adapted to the environment. The TR<sub>34</sub>/L98H isolates were genetically clustered with local environmental isolates or clinical isolates from China and Europe (Appendix Table 4). The TR<sub>46</sub>/Y121F/T289A isolate (S05–322) recovered in 2018, which colonized a patient without overseas travel, was genetically identical to a clone prevalent in the Netherlands and Tanzania (6), raising the concern of the intercountry transfer of resistant isolates.

All TR<sub>34</sub>/L98H/S297T/F495I, TR<sub>34</sub>/L98H, and TR<sub>46</sub>/Y121F/T289A isolates exhibited cross-resistance to difenoconazole and tebuconazole (both triazole fungicides) without fitness cost, demonstrated by normal growth rates (Appendix Figure 2). The TR<sub>34</sub>/L98H/S297T/F495I isolates and TR<sub>46</sub>/Y121F/T289A isolates were also resistant to prochloraz (an imidazole fungicide) (Appendix Table 2). The prevalence of TR<sub>34</sub>/L98H/S297T/F495I isolates in Taiwan might be attributed to widespread use of prochloraz over the past 3 decades. Studies have suggested an association between use of imidazole fungicides and emergence of azole-resistant *A. fumigatus* with TR<sub>34</sub>/L98H/S297T/F495I mutations (7,8).

In Taiwan, the annual consumption of difenoconazole and tebuconazole has exceeded that of prochloraz since 2012 (Appendix Figure 3), further creating a favorable environment for maintenance and spread of TR<sub>34</sub>/L98H, TR<sub>34</sub>/L98H/S297T/F495I, and TR<sub>46</sub>/Y121F/T289A isolates. Thus, the One Health approach to implement environmental antifungal stewardship is warranted to minimize ongoing resistance selection in the fields.

Six azole-resistant *A. fumigatus* isolates with wild-type *cyp51A* were obtained from 2 patients. Four pan-azole-resistant urinary isolates were sequentially recovered from a patient (no. 11) with *A. fumigatus* renal abscesses who was receiving voriconazole for >3 months in whom an initial urine isolate was susceptible to azole; all 5 isolates were genetically identical.

Overexpression of *cdr1B* (a drug efflux transporter) and an S269P mutation in *hmg1* (a hydroxymethylglutaryl-CoA reductase) were identified in 4 resistant isolates but not in the initial susceptible

isolate (Appendix Table 5, Figure 4), suggesting their roles involved in azole resistance (4,9). Another 2 pan-azole-resistant respiratory isolates were recovered from a patient (no. 12) who had pulmonary aspergillosis and was receiving voriconazole for 4 months. Azole-susceptible and azole-resistant isolates co-existed in this patient, which echoes the international recommendation suggesting testing multiple colonies ( $\geq 5$ ) from a single culture (1). *Cyp51A* overexpression and an F262 deletion in *hmg1* (*hmg1*<sup>F262-del</sup>) were identified in these 2 resistant isolates. Although *hmg1*<sup>F261-del</sup> was recently reported in azole-resistant *A. fumigatus* from a voriconazole-exposed patient (4), whether *cyp51A* overexpression and *hmg1*<sup>F262-del</sup> act synergistically to cause resistance warrants further studies.

Finally, reduced colony sizes were observed in all 6 azole-resistant *A. fumigatus* isolates with wild-type *cyp51A* (Appendix Figure 2). Thus, attention should be paid to select colonies of various sizes for susceptibility testing from patients with azole exposure.

Overall, 4 patients harboring azole-resistant *A. fumigatus* with environmental mutations and 2 patients harboring azole-resistant *A. fumigatus* with wild-type *cyp51A* showed development of invasive aspergillosis, and all had aspergillosis-related deaths. High mortality rates for azole-resistant aspergillosis we observed (6/6, 100%) and for those from a previous report (10) emphasize the need for a proposed integrated algorithm for management and control of azole-resistant aspergillosis (Appendix Table 6).

In conclusion, we report a health threat that arose from clinical and environmental use of azoles; environmental use contributed at a larger and global scale. These data necessitate diagnostic stewardship in the clinic and antifungal stewardship in the environment.

#### Acknowledgments

We thank Cheng-Chen Cheng, Li-Fang Chen, Li-Yu Yang, and Ya-Feng Wu for laboratory assistance; Hui-Chun Chang for collecting clinical information; and the following hospitals for sharing *Aspergillus* spp. clinical isolates: Changhua Christian Hospital, Chi Mei Medical Center-Liouying, Chung Shan Medical University Hospital, Ditmanson Medical Foundation Chia-Yi Christian Hospital, Far Eastern Memorial Hospital, Hualien Tzu Chi Hospital, Kaohsiung Chang Gung Memorial Hospital, Kaohsiung Medical University Chung-Ho Memorial Hospital, Kaohsiung Veterans General Hospital, Mennonite Christian Hospital, National Cheng Kung University Hospital, National Taiwan University Hospital, Show Chwan Memorial Hospital, Taichung Veterans General Hospital, Tainan Sin Lau Hospital, and Taipei City Hospital Heping Fuyou Branch.



This study was supported by the National Health Research Institutes of Taiwan (grants IV-105-PP-08 and IV-106-PP-08), the Ministry of Science and Technology of Taiwan (grants MOST 105-2628-B-400-004-MY2), and Gilead Sciences, Inc. (grant IN-US-131-3918).

Y.-C.C. has received honoraria for speaking and advisory board membership from Pfizer, Gilead, Merck, and Astellas, and was involved as a steering committee member of regional education programs at Gilead (Asia CARE) and Pfizer (Medical Mycology Training Network). W.-L.L. has received honoraria for speaking from Pfizer and Gilead.

### About the Author

Dr. Wu is an assistant investigator and attending physician at National Health Research Institutes, Zhunan, Taiwan. Her research interests include molecular epidemiology of infectious diseases and antimicrobial drug resistance in bacterial and fungal pathogens.

### References

1. Verweij PE, Ananda-Rajah M, Andes D, Arendrup MC, Brüggemann RJ, Chowdhary A, et al. International expert opinion on the management of infection caused by azole-resistant *Aspergillus fumigatus*. *Drug Resist Updat*. 2015;21-22:30-40. <https://doi.org/10.1016/j.drup.2015.08.001>
2. Clinical and Laboratory Standards Institute. Reference method for broth dilution antifungal susceptibility testing of filamentous fungi: approved standard, 2nd ed. M38-A2. Wayne (PA): The Institute; 2008.
3. Wang HC, Huang JC, Lin YH, Chen YH, Hsieh MI, Choi PC, et al. Prevalence, mechanisms and genetic relatedness of the human pathogenic fungus *Aspergillus fumigatus* exhibiting resistance to medical azoles in the environment of Taiwan. *Environ Microbiol*. 2018;20:270-80. <https://doi.org/10.1111/1462-2920.13988>
4. Hagiwara D, Arai T, Takahashi H, Kusuya Y, Watanabe A, Kamei K. Non-*cyp51A* azole-resistant *Aspergillus fumigatus* isolates with mutation in HMG-CoA reductase. *Emerg Infect Dis*. 2018;24:1889-97. <https://doi.org/10.3201/eid2410.180730>
5. Arendrup MC. Update on antifungal resistance in *Aspergillus* and *Candida*. *Clin Microbiol Infect*. 2014;20(Suppl 6):42-8. <https://doi.org/10.1111/1469-0691.12513>
6. Chowdhary A, Sharma C, van den Boom M, Yntema JB, Hagen F, Verweij PE, et al. Multi-azole-resistant *Aspergillus fumigatus* in the environment in Tanzania. *J Antimicrob Chemother*. 2014;69:2979-83. <https://doi.org/10.1093/jac/dku259>
7. Chen YC, Lai MH, Wu CY, Lin TC, Cheng AH, Yang CC, et al. The genetic structure, virulence, and fungicide sensitivity of *Fusarium fujikuroi* in Taiwan. *Phytopathology*. 2016; 106:624-35. <https://doi.org/10.1094/PHYTO-11-15-0285-R>
8. Chen Y, Li Z, Han X, Tian S, Zhao J, Chen F, et al. Elevated MIC values of imidazole drugs against *Aspergillus fumigatus* isolates with TR<sub>34</sub>/L98H/S297I/F495I mutation. *Antimicrob Agents Chemother*. 2018;62:e01549-17. <https://doi.org/10.1128/AAC.01549-17>
9. Fraczek MG, Bromley M, Buied A, Moore CB, Rajendran R, Rautemaa R, et al. The *cdr1B* efflux transporter is associated with non-*cyp51a*-mediated itraconazole resistance in *Aspergillus fumigatus*. *J Antimicrob Chemother*. 2013;68: 1486-96. <https://doi.org/10.1093/jac/dkt075>
10. van der Linden JW, Snelders E, Kampinga GA, Rijnders BJ, Mattsson E, Debets-Ossenkopp YJ, et al. Clinical implications of azole resistance in *Aspergillus fumigatus*, The Netherlands, 2007-2009. *Emerg Infect Dis*. 2011;17: 1846-54. <https://doi.org/10.3201/eid1710.110226>

Address for correspondence: Yee-Chun Chen, Department of Internal Medicine, National Taiwan University Hospital, No. 7 Chung-Shan South Rd, Taipei 10002, Taiwan; email: yeechunchen@gmail.com

## Knowledge of Infectious Disease Specialists Regarding Aspergillosis Complicating Influenza, United States

Mitsuru Toda, Susan E. Beekmann, Philip M. Polgreen, Tom M. Chiller, Brendan R. Jackson, Karlyn D. Beer

Author affiliations: Centers for Disease Control and Prevention, Atlanta, Georgia, USA (M. Toda, T.M. Chiller, B.R. Jackson, K.D. Beer); University of Iowa, Iowa City, Iowa, USA (S.E. Beekmann, P.M. Polgreen)

DOI: <https://doi.org/10.3201/eid2604.190953>

In an online survey, we found that nearly one fifth of physicians in the United States who responded had seen or heard about a case of invasive pulmonary aspergillosis after severe influenza at their institution. However, <10% routinely used galactomannan testing to test for this fungus in patients with severe influenza.

**I**nvasive pulmonary aspergillosis (IPA) occurs primarily among immunocompromised patients with a history of organ or stem cell transplantation, chemotherapy, or immunosuppressive medications. However, a multicenter retrospective study in the Netherlands and Belgium suggested that patients

# Multicenter Study of Azole-Resistant *Aspergillus fumigatus* Clinical Isolates, Taiwan

## Appendix

**Appendix Table 1.** Details of participating hospitals, antifungal susceptibility testing, and isolate collection for analysis of azole-resistant *Aspergillus fumigatus* clinical isolates, Taiwan\*

Hospital	Location	Period of collection	Susceptibility testing method†	
			Isolates before June 2017	Isolates during June 2017–March 2018
Chi-Mei Medical Center, Luying (CMMC)	Southern	2015 Feb–2018 Mar	CLSI M38-A2	Screening azole agar plates; confirmed by CLSI M38-A2
National Cheng-Kung University Hospital (NCKUH)	Southern	2011 Aug–2018 Mar	CLSI M38-A2	Screening azole agar plates; confirmed by CLSI M38-A2
National Taiwan University Hospital (NTUH)	Northern	2012 Feb–2018 Mar	YeastOne	Screening azole agar plates; confirmed by CLSI M38-A2
<b>TSARM hospitals</b>				
Changhua Christian Hospital	Central	2016 Jul–Sep	CLSI M38-A2	NI
Ditmanson Medical Foundation Chia-Yi Christian Hospital	Southern	2016 Jul–Sep	CLSI M38-A2	NI
Hualien Tzu Chi Hospital	Eastern	2016 Jul–Sep	CLSI M38-A2	NI
Kaohsiung Medical University Chung-Ho Memorial Hospital	Southern	2016 Jul–Sep	CLSI M38-A2	NI
Kaohsiung Veterans General Hospital	Southern	2016 Jul–Sep	CLSI M38-A2	NI
Show Chwan Memorial Hospital	Central	2016 Jul–Sep	CLSI M38-A2	NI
Taichung Veterans General Hospital	Central	2016 Jul–Sep	CLSI M38-A2	NI
Tainan Sin Lau Hospital	Southern	2016 Jul–Sep	CLSI M38-A2	NI

\*This study was approved by the Institutional Review Boards of the National Health Research Institutes (no. EC1040502-E and EC1050307) and participating hospitals: CMMC (10607-L01 and 10711-L03), NCKUH (B-ER-101–342), and NTUH (201605098RIPA). NI, no isolate; TSARM, Taiwan Surveillance of Antimicrobial Resistance of Molds.

†*A. fumigatus* sensu stricto was identified on the basis of morphologic characteristics, growth at 50°C, and sequence analyses of the internal transcribed spacer region and calmodulin gene (1). For isolates from CMMC, NCKUH, and TSARM hospitals, MICs of antifungal agents were determined by using the Clinical and Laboratory Standards Institute (CLSI) M38-A2 method. The MICs for the isolates from NTUH were determined by using the Sensititer YeastOne method (Trek Diagnostic Systems, <http://www.trekds.com>); isolates with any of the following MIC values (i.e.,  $\geq 2$ ,  $\geq 2$ , and  $\geq 0.25$   $\mu\text{g/mL}$  of itraconazole, voriconazole, and posaconazole, respectively) were reexamined by using the CLSI method. For isolates collected during June 2017–March 2018, azole resistance was screened by using azole-containing agar plates. In brief, the conidia of these isolates were directly inoculated onto 3 Sabouraud dextrose agar plates supplemented with itraconazole (2  $\mu\text{g/mL}$ ), voriconazole (1  $\mu\text{g/mL}$ ), or posaconazole (0.25  $\mu\text{g/mL}$ ), and incubated at 37°C. Colonies that grew after 2–4 d on any of the azole-containing agar plates were selected for the MIC determination by using the CLSI method.

**Appendix Table 2.** Laboratory characteristics of *Aspergillus fumigatus* clinical isolates from 12 patients with aspergillosis, Taiwan\*

Patient no.	Strain no.	Year (day)†	Cyp51A mutation	MIC or MIC <sub>50</sub> /MIC <sub>90</sub> (range), µg/mL, by CLSI M38-A2‡								
				AMB	ITR	VRC	POS	ISA	DFC	TBC	PRC	
Azole-resistant isolates§ except YL1, g054L, and g057L												
1	B44	2012 (d0)	TR <sub>34</sub> /L98H/S297T/F495I	0.5	>16	2	1	>16	>32	16	>32	
1	B51	2012 (d2)	TR <sub>34</sub> /L98H/S297T/F495I	0.5	>16	2	1	>16	>32	16	>32	
2	D007	2014	TR <sub>34</sub> /L98H/S297T/F495I	0.5	>16	2	1	>16	>32	>32	>32	
3	E071	2015 (d0)	TR <sub>34</sub> /L98H/S297T/F495I	0.5	>16	2	1	>16	>32	>32	>32	
3	E073	2015 (d3)	TR <sub>34</sub> /L98H/S297T/F495I	0.5	>16	2	1	>16	>32	>32	>32	
3	E074	2015 (d4)	TR <sub>34</sub> /L98H/S297T/F495I	0.5	>16	2	1	>16	>32	>32	>32	
4	S05–31 9	2018	TR <sub>34</sub> /L98H/S297T/F495I	0.5	>16	2	1	>16	>32	>32	>32	
5	A31	2013	TR <sub>34</sub> /L98H	0.5	>16	4	1	16	>32	32	4	
6	S05–12 2	2016	TR <sub>34</sub> /L98H	0.5	>16	8	1	8	>32	>32	2	
7	C03–0 04	2016	TR <sub>34</sub> /L98H	0.5	>16	4	1	8	32	>32	2	
8	S07–00 8	2016	TR <sub>34</sub> /L98H	0.5	>16	4	1	8	32	>32	2	
9	S05–20 5	2017	TR <sub>34</sub> /L98H	0.5	>16	4	1	8	32	>32	2	
10	S05–32 2	2018	TR <sub>46</sub> /Y121F/T289A	0.5	>16	>16	1	>16	>32	>32	>32	
11	YL1	2014 (d0)	Polymorphisms¶	0.5	0.25	0.5	0.25	1	4	4	0.25	
11	YL3	2014 (d100)	Polymorphisms¶	0.5	>16	8	1	8	32	>32	2	
11	YL4	2015 (d134)	Polymorphisms¶	0.5	>16	8	1	8	32	>32	2	
11	YL5	2015 (d165)	Polymorphisms¶	0.5	>16	2–4	0.5	2	16	16	1	
11	YL6	2015 (d185)	Polymorphisms¶	0.5	>16	8	1	8	32	>32	2	
12	g054m	2016 (d0)	Wild-type	0.12	>16	8	1	8	32	>32	2	
12	g054L	2016 (d0)	Wild-type	0.5	0.5	0.5	0.06	0.5	1	4	0.25	
12	g057m	2016 (d2)	Wild-type	0.12	>16	4	1	8	32	>32	2	
12	g057L	2016 (d2)	Wild-type	0.5	0.5	0.5	0.06	0.5	1	4	0.25	
Azole-susceptible isolates, n = 200#				0.5/1 (0.12–1)	0.5/0.5 (0.03–1)	0.5/1 (0.12–1)	0.12/0.25 (0.015–0.25)	ND (1–8)	2/4 (1–8)	4/8 (2–8)	ND	

\*AMB, amphotericin B; ATCC, American Type Culture Collection; DFC, difenoconazole; ISA, isavuconazole; ITR, itraconazole; ND, not done; POS, posaconazole; PRC, prochloraz; TBC, tebuconazole; VRC, voriconazole.

†d0 and dn indicate day of and n days after the first isolation of *A. fumigatus*, respectively.

‡*Candida parapsilosis* ATCC 22019 and *A. fumigatus* ATCC MYA 3626 were used as quality control and reference strains.

§Because of good essential agreement in the azole MIC values between the CLSI and European Committee on Antimicrobial Susceptibility Testing (EUCAST) methods (2) and the lack of established CLSI clinical breakpoints for *A. fumigatus*, the MIC interpretive criteria for resistance in this study followed the EUCAST clinical breakpoints (i.e., >2, >2, >0.25, and >1 µg/mL) for itraconazole, voriconazole, posaconazole, and isavuconazole, respectively (3). The drugs for susceptibility testing were obtained from Sigma-Aldrich (<https://www.sigmaaldrich.com>) (AMB, ITR, VRC, POS, DFC, and PRC), Toronto Research Chemicals (<https://www.trc-canada.com>) (ISA), and Chem Service (<https://www.chemservice.com>) (TBC).

¶These isolates have F46Y/G89G/M172V/N248T/D255E/L358L/E427K/C454C polymorphisms.

#The MICs of 200 azole-susceptible isolates were used. Only 62 isolates were tested for MICs for DFC and TBC.

**Appendix Table 3.** Clinical characteristics of 12 patients harboring azole-resistant *Aspergillus fumigatus* isolates, Taiwan\*

Patient no.	Strain no.	Age, y/ sex	Concurrent condition	Sample	<i>Aspergillus</i> disease†	Previous azole exposure	Sequential antifungal treatment (d)	Outcome	IA-related death
1‡	B44, B51	66/F	DM, HCV/cirrhosis, adrenal insufficiency	Sputum	IPA, unclassified	No	VRC (1), AMB (3)	Died	Yes
2	D007	49/M	AML, peritonitis	Nasal swab	Colonization	POS/VR C (2 wk)	MCF (33)	Died	No
3	E071, E074; E073	64/F	SLE, ESRD, bacterial septicemia, meningoencephalitis	Pleural effusion; sputum	Proven IPA with empyema	No	VRC (1), LAMB (3)	Died	Yes
4	S05–319	88/F	DM, influenza B	Sputum	Colonization	No	No	Alive	No
5‡	A31	59/M	Lung cancer, COPD, bronchiectasis	BAL	Proven IPA	No	AMB (3), POS (10), VRC (11)	Died	Yes
6	S05–122	90/F	COPD, steroid use, bronchiectasis	Sputum	Probable IPA	No	No	Died	Yes
7	C03–004	–	–	Sputum	–	–	–	–	–
8	S07–008	–	–	Ear	–	–	–	–	–
9	S05–205	76/F	COPD, steroid use, bronchiectasis, DM	Sputum	Colonization	No	No	Alive	No
10	S05–322	74/M	B cell lymphoma, COPD, HCV, CAD	Sputum	Colonization	No	No	Alive	No
11	YL1;§ YL3, YL4, YL5, YL6	36/M	HIV/AIDS	Urine; PCN	Proven IA, (renal abscess)	VRC (3 mo)	VRC (93), VRC/CAS (43), LAMB (44), LAMB/5FC (25), LAMB/AND (24)	Died	Yes
12	g054m, g054L,§ g057m, g057L§	39/M	MDS with RAEB-T, status post-HSCT with GVHD, bacterial septicemia	Sputum; BAL	Probable IPA	VRC (4 mo)	AMB (18), POS (21), AMB (13), AMB/AND (8), AND (5), AMB/AND (1), AND (4), AMB/AND (2)	Died	Yes

\*AMB, conventional amphotericin B; AML, acute myeloid leukemia; AND, anidulafungin; BAL, bronchoalveolar lavage; CAD, coronary artery disease; CAS, caspofungin; COPD, chronic obstructive pulmonary disease; DM, diabetes mellitus; ESRD, end-stage renal disease; 5FC, flucytosine; GVHD, graft versus host disease; HCV, hepatitis C virus infection; HSCT, allogeneic hematopoietic stem cell transplantation; IA, invasive aspergillosis; IPA, invasive pulmonary aspergillosis; LAMB, liposomal amphotericin B; MCF, micafungin; MDS with RAEB-T, myelodysplastic syndrome with refractory anemia and excess blast in transformation; PCN, percutaneous nephrostomy; POS, posaconazole; SLE, systemic lupus erythematosus; VRC, voriconazole; –, data not available.

†Clinical data for patients harboring azole-resistant *A. fumigatus* were reviewed, and IA was classified according to the EORTC/MSG definition (4).

‡Three isolates (A31, B44, and B51) from 2 patients have been described in our previous report (5).

§YL1, g054L, and g057L were azole susceptible.





**Appendix Table 4.** References for 38 oversea *Aspergillus fumigatus* strains included in microsatellite-based phylogenetic analysis during a multicenter study of azole-resistant *Aspergillus fumigatus* clinical isolates, Taiwan

Strain	Reference
C485	(6)
C96	(6)
E2619	(6)
20643.023	(7)
20684.007	(7)
E1001	(6)
20677.079	(7)
2087 m1341.17-06-2012	(8)
2091 m1428.01-07-2012	(8)
C94	(6)
094411/7/50	(8)
Case 2-90d	(9)
Japan Dec 2013	(10)
The Netherlands 7	(10, 11)
20643.017	(7)
2005-456307L	(6)
OKH50	(12)
04-202165	(13)
F2126	(6)
1042/09	(14)
14-148-2460	(6)
2107m1974.23-09-2012	(8)
A12519	(8)
R2-07-1_R	(6)
E454	(15)
Case 1-7d	(9)
Myc-2008-002 nr.42	(14)
Case 3-6d	(9)
CF/NL2992	(8)
Tanzania	(11)
CF/NL0682	(8)
VPC1651/Ei/12/2/a/3	(8)
CF/NL0645	(11)
The Netherlands 2	(10, 11)
The Netherlands 3	(10, 11)
C195	(7)
12-90032258	(13)
Case 4-36d	(9)

**Appendix Table 5.** Gene substitutions in azole-resistant *Aspergillus fumigatus* isolates and control strains, Taiwan\*

Strain no.	Azole susceptibility (mechanism)	Gene substitutions†			
		<i>hapE</i>	<i>srbA</i>	<i>hmg1</i>	<i>erg6</i>
ATCC MYA-3626	S	-	V37D	-	-
ATCC 16903	S	-	V37D, S82P	H564Y	-
F2509	S	-	V37D	-	-
F02411	S	-	A865V	P212S, H564Y	-
YL1	S	-	E957D	P212S, H564Y	-
YL3	R	-	E957D	P212S, S269P, H564Y	F186V
YL4	R	-	E957D	P212S, S269P, H564Y	F186V
YL5	R	-	E957D	P212S, S269P, H564Y	-
YL6	R	-	E957D	P212S, S269P, H564Y	F186V
g054m	R	-	A865V	F262_del, H564Y	-
g057m	R	-	A865V	F262_del, H564Y	-
g054L	S	-	V37D	-	-
g057L	S	-	V37D	-	-
B44	R (TR <sub>34</sub> /L98H/S297T/F495I)	-	V37D	H564Y	-
A31	R (TR <sub>34</sub> /L98H)	-	V37D	-	-


\*ATCC, American Type Culture Collection; del, deletion; R, resistant; S, susceptible; -, wild type.

†The *hapE*, *hmg1*, *erg6* and *srbA* genes were sequenced as described (16-18). The *hapE* and *srbA* sequences were compared with those of *A. fumigatus* f293; the *hmg1* and *erg6* sequences were compared with those of *A. fumigatus* A1163.



**Appendix Table 6.** Proposed integrated algorithm for management and control of azole-resistant aspergillosis, Taiwan\*

Category	Response
Academic society	Incorporate the issue of azole resistance into national guidelines for management of aspergillosis; hold educational programs to improve diagnosis and treatment for azole-resistant aspergillosis.
Laboratory personnel	Identify clinically relevant <i>Aspergillus</i> isolates at the species complex level and confirm <i>A. fumigatus</i> by thermotolerance test (growth at 50°C) (19); screen for azole resistance with azole agar plates for clinically relevant <i>A. fumigatus</i> isolates and screen multiple colonies ( $\leq 5$ colonies) from a single specimen (19); for colonies grown on any azole agar plate, perform azole MIC testing by using reference CLSI or EUCAST methods or an alternative Sensititer YeastOne assay (19,20); If MIC testing is not available, refer isolates to a mycology reference laboratory (19); prompt notification of the clinical team if azole-resistant <i>A. fumigatus</i> is suspected and identified.
Physicians	Be familiar with patient risk factors for invasive aspergillosis; obtain clinical specimens for fungal culture as possible; select empirical antifungal agents according to the updated local prevalence rate of azole resistance (21); antifungal susceptibility testing is recommended for <i>A. fumigatus</i> isolates from invasive diseases and should be repeated on later isolates if infection persists despite treatment (21); be aware of the possibility of azole-resistance in patients unresponsive to azole treatment; consider amphotericin B-based or azole/echinocandin combination therapy for azole-resistant aspergillosis (19,21).
Hospital environment	Segregate patients from construction or renovation, potted plants, and flowers in wards and patients' room (22); control the airborne dissemination of fungal spores, (e.g., barriers, containment, air handling, HEPA filters, sealed windows, sealing the area of construction or renovation activities if possible) (23).
Reference mycology laboratory	Identify <i>Aspergillus</i> isolates to the species level by molecular methods; confirm antifungal susceptibility of <i>Aspergillus</i> isolates with reference CLSI or EUCAST methods; perform periodic reference MIC testing of isolates of <i>A. fumigatus</i> complex ( $\geq 100$ isolates) (19); sequence <i>cyp51A</i> genes in resistant isolates to determine the nature and trends in <i>cyp51A</i> mutation distribution (19); establish molecular typing methods; collect strains.
Scientists and plant pathologists	Identify the key azole fungicides that select azole-resistant <i>Aspergillus</i> ; propose better fungicide application strategies to minimize resistance development (24).
Agricultural authority	Include azole fungicides that select azole-resistant isolates into national pesticide risk reduction programs; advise farmers to reduce culprit azole fungicide use by in rotation with alternative fungicides with different modes of action.
Governance	Update and evaluate the global situation; accredit national mycology reference laboratories; implement antifungal stewardship programs in agriculture in addition to hospitals and animal husbandry to achieve the One Health goal (24).


\*CLSI, Clinical and Laboratory Standards Institute; EUCAST, European Committee on Antimicrobial Susceptibility Testing.





**Appendix Figure 1.** Distribution of participating hospitals and isolate collection areas for *Aspergillus fumigatus*, Taiwan. Values in parentheses indicate no. *A. fumigatus* isolates/no. patients and no. azole-resistant *A. fumigatus* isolates/no. patients.



**Appendix Figure 2.** A) Radial growth of *Aspergillus fumigatus* isolates on Sabouraud dextrose agar plates at 35°C, Taiwan. The radius of the growing colony was measured after 72 hours of incubation. Values are the mean diameter of triplicate samples. Error bars indicate SD. Colonies of B) YL1, YL3, YL4, YL5, and YL6 from patient 11 and C) g054m, g054L, g057m, and g057L from patient 12 observed at 72 hours. C, clinical isolate; E, environmental isolate; R, azole-resistant; S, azole-susceptible; ↑, overexpression.



**Appendix Figure 3.** Annual sales of azole fungicides in Taiwan, 2003–2016. Annual sales of imidazole fungicides (imazalil and prochloraz) and triazole fungicides (bromuconazole, difenoconazole, epoxiconazole, propiconazole, and tebuconazole) are shown according to data derived from Domestic Manufacturers Production and Sale of Pesticides published by the Taiwan Crop Protection Industry Association (25).



**Appendix Figure 4.** mRNA expression levels of A) a drug efflux transporter gene, *cdr1B*, and B) *cyp51A* in *Aspergillus fumigatus* isolates, Taiwan. Expression levels were normalized to  $\beta$ -tubulin levels and compared with those in *A. fumigatus* ATCC MYA-3626. Error bars indicate SD. Results for the *cyp51B* gene and other transporter genes (*AfuMDR1*, *AfuMDR2*, *AfuMDR3*, *AfuMDR4*, *atrF*, and *MFS56*) were inconclusive and are not shown. ATCC, American Type Culture Collection; ITR, itraconazole; VRC, voriconazole.



## References

1. Balajee SA, Houbraeken J, Verweij PE, Hong SB, Yaghuchi T, Varga J, et al. *Aspergillus* species identification in the clinical setting. *Stud Mycol.* 2007;59:39–46. [PubMed https://doi.org/10.3114/sim.2007.59.05](https://doi.org/10.3114/sim.2007.59.05)
2. Pfaller M, Boyken L, Hollis R, Kroeger J, Messer S, Tendolkar S, et al. Comparison of the broth microdilution methods of the European Committee on Antimicrobial Susceptibility Testing and the Clinical and Laboratory Standards Institute for testing itraconazole, posaconazole, and voriconazole against *Aspergillus* isolates. *J Clin Microbiol.* 2011;49:1110–2. [PubMed https://doi.org/10.1128/JCM.02432-10](https://doi.org/10.1128/JCM.02432-10)
3. EUCAST. Subcommittee on Antimicrobial Susceptibility Testing (EUCAST-AFST). Clinical breakpoints for fungi, version 9 [cited 2020 Jan 25]. <http://www.eucast.org/astoffungi/clinicalbreakpointsforantifungals>
4. De Pauw B, Walsh TJ, Donnelly JP, Stevens DA, Edwards JE, Calandra T, et al.; European Organization for Research and Treatment of Cancer/Invasive Fungal Infections Cooperative Group; National Institute of Allergy and Infectious Diseases Mycoses Study Group (EORTC/MSG) Consensus Group. Revised definitions of invasive fungal disease from the European Organization for Research and Treatment of Cancer/Invasive Fungal Infections Cooperative Group and the National Institute of Allergy and Infectious Diseases Mycoses Study Group (EORTC/MSG) Consensus Group. *Clin Infect Dis.* 2008;46:1813–21. [PubMed https://doi.org/10.1086/588660](https://doi.org/10.1086/588660)
5. Wu CJ, Wang HC, Lee JC, Lo HJ, Dai CT, Chou PH, et al. Azole-resistant *Aspergillus fumigatus* isolates carrying TR<sub>34</sub>/L98H mutations in Taiwan. *Mycoses.* 2015;58:544–9. [PubMed https://doi.org/10.1111/myc.12354](https://doi.org/10.1111/myc.12354)
6. Chen Y, Lu Z, Zhao J, Zou Z, Gong Y, Qu F, et al. Epidemiology and molecular characterizations of azole resistance in clinical and environmental *Aspergillus fumigatus* isolates from China. *Antimicrob Agents Chemother.* 2016;60:5878–84. [PubMed https://doi.org/10.1128/AAC.01005-16](https://doi.org/10.1128/AAC.01005-16)
7. Deng S, Zhang L, Ji Y, Verweij PE, Tsui KM, Hagen F, et al. Triazole phenotypes and genotypic characterization of clinical *Aspergillus fumigatus* isolates in China. *Emerg Microbes Infect.* 2017;6:e109. [PubMed https://doi.org/10.1038/emi.2017.97](https://doi.org/10.1038/emi.2017.97)
8. Steinmann J, Hamprecht A, Vehreschild MJ, Cornely OA, Buchheidt D, Spiess B, et al. Emergence of azole-resistant invasive aspergillosis in HSCT recipients in Germany. *J Antimicrob Chemother.* 2015;70:1522–6. [PubMed https://doi.org/10.1093/jac/dku566](https://doi.org/10.1093/jac/dku566)
9. Astvad KM, Jensen RH, Hassan TM, Mathiasen EG, Thomsen GM, Pedersen UG, et al. First detection of TR<sub>46</sub>/Y121F/T289A and TR<sub>34</sub>/L98H alterations in *Aspergillus fumigatus* isolates from azole-naïve patients in Denmark despite negative findings in the environment. *Antimicrob Agents Chemother.* 2014;58:5096–101. [PubMed https://doi.org/10.1128/AAC.02855-14](https://doi.org/10.1128/AAC.02855-14)

10. Hagiwara D, Takahashi H, Fujimoto M, Sugahara M, Misawa Y, Gono T, et al. Multi-azole resistant *Aspergillus fumigatus* harboring Cyp51A TR<sub>46</sub>/Y121F/T289A isolated in Japan. J Infect Chemother. 2016;22:577–9. [PubMed https://doi.org/10.1016/j.jiac.2016.01.015](https://doi.org/10.1016/j.jiac.2016.01.015)
11. Chowdhary A, Sharma C, van den Boom M, Yntema JB, Hagen F, Verweij PE, et al. Multi-azole-resistant *Aspergillus fumigatus* in the environment in Tanzania. J Antimicrob Chemother. 2014;69:2979–83. [PubMed https://doi.org/10.1093/jac/dku259](https://doi.org/10.1093/jac/dku259)
12. Toyotome T, Hagiwara D, Kida H, Ogi T, Watanabe A, Wada T, et al. First clinical isolation report of azole-resistant *Aspergillus fumigatus* with TR<sub>34</sub>/L98H-type mutation in Japan. J Infect Chemother. 2017;23:579–81. [PubMed https://doi.org/10.1016/j.jiac.2016.12.004](https://doi.org/10.1016/j.jiac.2016.12.004)
13. Kidd SE, Goeman E, Meis JF, Slavin MA, Verweij PE. Multi-triazole-resistant *Aspergillus fumigatus* infections in Australia. Mycoses. 2015;58:350–5. [PubMed https://doi.org/10.1111/myc.12324](https://doi.org/10.1111/myc.12324)
14. Chowdhary A, Kathuria S, Xu J, Sharma C, Sundar G, Singh PK, et al. Clonal expansion and emergence of environmental multiple-triazole-resistant *Aspergillus fumigatus* strains carrying the TR<sub>34</sub>/L98H mutations in the cyp51A gene in India. PLoS One. 2012;7:e52871. [PubMed https://doi.org/10.1371/journal.pone.0052871](https://doi.org/10.1371/journal.pone.0052871)
15. Ahmad S, Khan Z, Hagen F, Meis JF. Occurrence of triazole-resistant *Aspergillus fumigatus* with TR<sub>34</sub>/L98H mutations in outdoor and hospital environment in Kuwait. Environ Res. 2014;133:20–6. [PubMed https://doi.org/10.1016/j.envres.2014.05.009](https://doi.org/10.1016/j.envres.2014.05.009)
16. Camps SM, Dutilh BE, Arendrup MC, Rijs AJ, Snelders E, Huynen MA, et al. Discovery of a *hapE* mutation that causes azole resistance in *Aspergillus fumigatus* through whole genome sequencing and sexual crossing. PLoS One. 2012;7:e50034. [PubMed https://doi.org/10.1371/journal.pone.0050034](https://doi.org/10.1371/journal.pone.0050034)
17. Hagiwara D, Arai T, Takahashi H, Kusuya Y, Watanabe A, Kamei K. Non-*cyp51A* azole-resistant *Aspergillus fumigatus* isolates with mutation in HMG-CoA reductase. Emerg Infect Dis. 2018;24:1889–97. [PubMed https://doi.org/10.3201/eid2410.180730](https://doi.org/10.3201/eid2410.180730)
18. Wang HC, Huang JC, Lin YH, Chen YH, Hsieh MI, Choi PC, et al. Prevalence, mechanisms and genetic relatedness of the human pathogenic fungus *Aspergillus fumigatus* exhibiting resistance to medical azoles in the environment of Taiwan. Environ Microbiol. 2018;20:270–80. [PubMed https://doi.org/10.1111/1462-2920.13988](https://doi.org/10.1111/1462-2920.13988)
19. Ullmann AJ, Aguado JM, Arikian-Akdagli S, Denning DW, Groll AH, Lagrou K, et al. Diagnosis and management of *Aspergillus* diseases: executive summary of the 2017 ESCMID-ECMM-ERS guideline. Clin Microbiol Infect. 2018;24(Suppl 1):e1–38. [PubMed https://doi.org/10.1016/j.cmi.2018.01.002](https://doi.org/10.1016/j.cmi.2018.01.002)

20. Wang HC, Hsieh MI, Choi PC, Wu CJ. Comparison of the Sensititre YeastOne and CLSI M38–A2 microdilution methods in determining the activity of amphotericin B, itraconazole, voriconazole, and posaconazole against *Aspergillus* species. *J Clin Microbiol*. 2018;56:e00780–18. [PubMed](#)  
<https://doi.org/10.1128/JCM.00780-18>
21. Verweij PE, Ananda-Rajah M, Andes D, Arendrup MC, Brüggemann RJ, Chowdhary A, et al. International expert opinion on the management of infection caused by azole-resistant *Aspergillus fumigatus*. *Drug Resist Updat*. 2015;21-22:30–40. [PubMed](#) <https://doi.org/10.1016/j.drug.2015.08.001>
22. Freifeld AG, Bow EJ, Sepkowitz KA, Boeckh MJ, Ito JI, Mullen CA, et al.; Infectious Diseases Society of America. Clinical practice guideline for the use of antimicrobial agents in neutropenic patients with cancer: 2010 update by the Infectious Diseases Society of America. *Clin Infect Dis*. 2011;52:e56–93. [PubMed](#)  
<https://doi.org/10.1093/cid/cir073>
23. Kanamori H, Rutala WA, Sickbert-Bennett EE, Weber DJ. Review of fungal outbreaks and infection prevention in healthcare settings during construction and renovation. *Clin Infect Dis*. 2015;61:433–44. [PubMed](#)  
<https://doi.org/10.1093/cid/civ297>
24. Chowdhary A, Meis JF. Emergence of azole resistant *Aspergillus fumigatus* and One Health: time to implement environmental stewardship. *Environ Microbiol*. 2018;20:1299–301. [PubMed](#)  
<https://doi.org/10.1111/1462-2920.14055>
25. Taiwan Crop Protection Industry Association. Domestic manufacturers production and sale of pesticides [in Chinese]. Taipei: Taiwan Crop Protection Industry Association, 1962–2016.