

9. Widjaja I, Wang C, van Haperen R, Gutiérrez-Álvarez J, van Dieren B, Okba NMA, et al. Towards a solution to MERS: protective human monoclonal antibodies targeting different domains and functions of the MERS-coronavirus spike glycoprotein. *Emerg Microbes Infect.* 2019;8:516–30. <https://doi.org/10.1080/22221751.2019.1597644>
10. Haagmans BL, van den Brand JM, Raj VS, Volz A, Wohlsein P, Smits SL, et al. An orthopoxvirus-based vaccine reduces virus excretion after MERS-CoV infection in dromedary camels. *Science.* 2016;351:77–81. <https://doi.org/10.1126/science.aad1283>

Address for correspondence: Berend-Jan Bosch, Virology Division, Department of Infectious Diseases and Immunology, Faculty of Veterinary Medicine, Utrecht University, Yalelaan 1, 3584 CL, Utrecht, the Netherlands; email: b.j.bosch@uu.nl; Bart L. Haagmans, Department of Viroscience, Erasmus Medical Center, PO Box 2040, 3000 CA, Rotterdam, the Netherlands; email: b.haagmans@erasmusmc.nl

Novel *Ehrlichia* Strain Infecting Cattle Tick *Amblyomma neumanni*, Argentina, 2018

Lucía Fargnoli, Camilo Fernandez, Lucas D. Monje

Author affiliation: Instituto de Ciencias Veterinarias del Litoral, UNL-CONICET, Esperanza, Argentina

DOI: <https://doi.org/10.3201/eid2605.190940>

In 2018, we detected a novel *Ehrlichia* strain infecting *Amblyomma neumanni* ticks in Argentina. The novel strain is phylogenetically related to the ruminant pathogen *E. ruminantium* and represents a potential risk for veterinary and public health because *A. neumanni* ticks parasitize domestic and wild ruminants and bite humans.

etymologia

Coronavirus [kə-ro'nə-vi"rus]

Ronnie Henry

The first coronavirus, avian infectious bronchitis virus, was discovered in 1937 by Fred Beaudette and Charles Hudson. In 1967, June Almeida and David Tyrrell performed electron microscopy on specimens from cultures of viruses known to cause colds in humans and identified particles that resembled avian infectious bronchitis virus. Almeida coined the term “coronavirus,” from the Latin *corona* (“crown”), because the glycoprotein spikes of these viruses created an image similar to a solar corona.

Strains that infect humans generally cause mild symptoms. However, more recently, animal coronaviruses have caused outbreaks of severe respiratory disease in humans, including severe acute respiratory syndrome (SARS), Middle East respiratory syndrome (MERS), and coronavirus disease (COVID-19).




Figure. Illustration reveals the ultrastructural morphology exhibited by coronaviruses. Note the spikes that adorn the outer surface of the virus, which impart the look of a corona surrounding the virion, when viewed electron microscopically. Photo: CDC/ Alissa Eckert, MS; Dan Higgins, MAMS

Sources

1. Almeida JD, Tyrrell DA. The morphology of three previously uncharacterized human respiratory viruses that grow in organ culture. *J Gen Virol.* 1967;1:175–8. <https://doi.org/10.1099/0022-1317-1-2-175>
2. Beaudette FR, Hudson CB. Cultivation of the virus of infectious bronchitis. *J Am Vet Med Assoc.* 1937;90:51–8.
3. Estola T. Coronaviruses, a new group of animal RNA viruses. *Avian Dis.* 1970;14:330–6. <https://doi.org/10.2307/1588476>
4. Groupe V. Demonstration of an interference phenomenon associated with infectious bronchitis virus of chickens. *J Bacteriol.* 1949;58:23–32. <https://doi.org/10.1128/JB.58.1.23-32.1949>

Address for correspondence: Ronnie Henry, Centers for Disease Control and Prevention, 1600 Clifton Rd NE, Mailstop V18-2, Atlanta, GA 30329-4027, USA; email: boq3@cdc.gov

DOI: <https://doi.org/10.3201/eid2605.ET2605>

Ehrlichia spp. are intracellular gram-negative bacteria relevant to human and animal health; they infect monocytes, neutrophils, or endothelial cells, depending on the species involved (1). The genus *Ehrlichia* (Rickettsiales: Anaplasmataceae) comprises 6 formally recognized tick-transmitted species: *E. canis*, *E. muris*, *E. chaffeensis*, *E. ewingii*, *E. minasensis*, and *E. ruminantium* (2,3). Recently, other *Ehrlichia* species have been reported and different strains of putative novel *Ehrlichia* species have been molecularly detected, but their taxonomic positions are still not clearly defined (4–6). Current knowledge about this group of pathogens suggests that a large number of *Ehrlichia* species might be not yet described.

Amblyomma neumanni ticks are relevant to human and veterinary medicine because in all stages they commonly parasitize wild and domestic ruminants and other large mammals, including humans (7). Moreover, *A. neumanni* ticks can reportedly be infected with *Rickettsia bellii* and *Rickettsia amblyomantis* and are potential vectors of the cattle pathogen *Anaplasma marginale* (7). To determine the presence of tickborne bacteria of the genus *Ehrlichia* in questing *A. neumanni* ticks in northwestern Argentina, we performed phylogenetic analyses on Anaplasmataceae-positive tick DNA samples. All procedures were approved by the Ethics and Biosafety Committee of the Facultad de Ciencias Veterinarias, Universidad Nacional del Litoral, Esperanza, Argentina.

During May 2018 (late autumn), we collected free-living ticks by dragging and by using dry ice-baited traps in Dean Funes (30°22'S, 64°21'W) and San José de la Dormida (30°21'S, 63°58'W), Córdoba Province, Argentina. Both sites are located in the Chaco Seco ecoregion. We identified all ticks by using standard taxonomic keys (7) and individually processed them for DNA extraction by using a boiling method (8). We screened DNA extracts for Anaplasmataceae by real-time PCR targeting the 16S rRNA gene, as previously described (9). We further tested samples positive for Anaplasmataceae by amplification of *Ehrlichia* genes *dsb* and *groESL*, as described elsewhere (5). We sequenced all amplicons and performed phylogenetic analyses with the maximum-likelihood method.

We collected 229 ticks from Dean Funes (70 adults, 159 nymphs) and 62 from San José de la Dormida (24 adults, 38 nymphs) and identified all ticks as *A. neumanni*. Only 1 adult tick from San José de la Dormida was positive for Anaplasmataceae by PCR. Further analysis of that sample resulted in 2 sequences of 355 bp (*dsb*) and 784 bp (*groESL*). Phylogenetic analysis of the *dsb* sequence (GenBank

accession no. MN176580) showed that the *A. neumanni* tick was infected with a species of *Ehrlichia*, which we named *Ehrlichia* sp. strain La Dormida, closely related to *E. ruminantium* (82.0%, GenBank accession no. CR925677) (Figure, panel A). Other ehrlichiae from South America included in the analysis, such as *Ehrlichia* sp. strain Córdoba from Argentina (78.0%, GenBank accession no. KY413807) and *Ehrlichia* sp. strain Natal from Brazil (79.3%, GenBank accession no. KY207546), were placed in a clade sister to the group formed by *Ehrlichia* sp. strain La Dormida and *E. ruminantium*. Furthermore, the phylogenetic analysis performed by using the *groESL* sequence (GenBank accession no. MN176581) confirmed these results (Figure, panel B).

Several recent studies conducted in South America reported finding novel ehrlichial agents infecting jaguars, horses, crab-eating foxes, opossums, sloths, and peccaries (6). Unfortunately, only short *dsb* sequences are available for those ehrlichiae from South America. Phylogenetic analysis including these sequences (210 positions included in the final dataset) placed them all together in a clade sister to the group formed by *Ehrlichia* sp. strain La Dormida and *E. ruminantium* (Appendix Figure, <https://wwwnc.cdc.gov/EID/article/26/5/19-0940-App1.pdf>).

Ehrlichia sp. strain La Dormida, associated with *A. neumanni* ticks, circulates in rural areas of northwestern Argentina. In our phylogenetic analyses, *Ehrlichia* sp. strain La Dormida genotype was unique and well separated from all other available ehrlichial sequences, suggesting that it could represent a distinct species yet to be properly characterized. In addition, these analyses positioned *Ehrlichia* sp. strain La Dormida in a separate group together with *E. ruminantium*. The species *E. ruminantium* is native to the Africa continent, where it is the etiologic agent of heartwater, a tickborne disease of major economic relevance with regard to domestic ruminants throughout sub-Saharan Africa (10). Besides *E. ruminantium*, the only other species of *Ehrlichia* known to naturally infect and cause clinical manifestations of ehrlichiosis in cattle is *E. minasensis* (3). However, *E. minasensis* is phylogenetically distant from *E. ruminantium* and closely related to *E. canis* genotypes (3).

We report *A. neumanni* ticks as a potential vector of *Ehrlichia* sp. strain La Dormida. Because it is assumed that bacteria of the genus *Ehrlichia* are not transmitted transovarially in ticks (1), infection with *Ehrlichia* must be acquired during feeding of immature ticks, which then pass the infection to adults by transstadial transmission. Regarding *E. ruminantium*, wild African ruminants are reservoirs




Figure. Maximum-likelihood trees constructed from *dsb* and *groESL* sequences of *Ehrlichia* sp. infecting *Amblyomma neumanni* ticks in Argentina compared with reference strains. A) Tree constructed by using *dsb* *Ehrlichia* sequences of approximately the same length as the sequence identified in this study (341 positions included in the final dataset). B) Tree constructed by using *groESL* *Ehrlichia* sequences of approximately the same length as the sequence identified in this study (767 positions included in the final dataset). Phylogenetic trees were constructed by using MEGA 7.0 (<https://www.megasoftware.net>), and best-fitting substitution models were determined with the Akaike Information Criterion, using the maximum-likelihood model test. Numbers represent bootstrap support generated from 1,000 replications. GenBank accession numbers are shown in parentheses. Boldface indicates the strain identified in this study. Scale bars indicate nucleotide substitutions/site.

of the bacteria (10). The novel *Ehrlichia* sp. strain La Dormida is phylogenetically related to the ruminant pathogen *E. ruminantium* and represents a potential risk for veterinary and public health because *A. neu-manni* ticks parasitize domestic and wild ruminants and bite humans.

Acknowledgments

We thank Federico Ruiz and Anibal Gomez for their support with field work and Federico Prandi.

This study was supported by Agencia Nacional de Promoción Científica y Tecnológica (PICT-2015-1084) and Agencia Santafesina de Ciencia, Tecnología e Innovación (IO-2017-00125).

About the Author

Dr. Fargnoli is a postdoctoral student at the Instituto de Ciencias Veterinarias del Litoral in Esperanza, Santa Fe, Argentina. Her research interests focus on the ecology of tickborne diseases.

References

- Brouqui P, Matsumoto K. Bacteriology and phylogeny of Anaplasmataceae. In: Raoul D, Parola P, editors. Rickettsial diseases. New York; Informa; 2007. p. 179-198.
- Dumler JS, Barbet AF, Bekker CP, Dasch GA, Palmer GH, Ray SC, et al. Reorganization of genera in the families Rickettsiaceae and Anaplasmataceae in the order Rickettsiales: unification of some species of *Ehrlichia* with *Anaplasma*, *Cowdria* with *Ehrlichia* and *Ehrlichia* with *Neorickettsia*, descriptions of six new species combinations and designation of *Ehrlichia equi* and 'HGE agent' as subjective synonyms of *Ehrlichia phagocytophila*. *Int J Syst Evol Microbiol*. 2001;51:2145-65. <https://doi.org/10.1099/00207713-51-6-2145>
- Aguiar DM, Ziliani TF, Zhang X, Melo AL, Braga IA, Witter R, et al. A novel *Ehrlichia* genotype strain distinguished by the TRP36 gene naturally infects cattle in Brazil and causes clinical manifestations associated with ehrlichiosis. *Ticks Tick Borne Dis*. 2014;5:537-44. <https://doi.org/10.1016/j.ttbdis.2014.03.010>
- Eberhardt AT, Fernandez C, Fargnoli L, Beldomenico PM, Monje LD. A putative novel strain of *Ehrlichia* infecting *Amblyomma tigrinum* associated with Pampas fox (*Lycalopex gymnocercus*) in Esteros del Iberá ecoregion, Argentina. *Ticks Tick Borne Dis*. 2020;11:101318.
- Cicuttin GL, De Salvo MN, Nava S. Two novel *Ehrlichia* strains detected in *Amblyomma tigrinum* ticks associated to dogs in peri-urban areas of Argentina. *Comp Immunol Microbiol Infect Dis*. 2017;53:40-4. <https://doi.org/10.1016/j.cimid.2017.07.001>
- Lopes MG, Muñoz-Leal S, de Lima JTR, Fournier GFDSR, Acosta IDCL, Martins TF, et al. Ticks, rickettsial and ehrlichial infection in small mammals from Atlantic forest remnants in northeastern Brazil. *Int J Parasitol Parasites Wildl*. 2018;7:380-5. <https://doi.org/10.1016/j.ijppaw.2018.10.001>
- Nava S, Venzal JM, Gonzalez-Acuña DA, Martins TF, Guglielmo AA, editors. Ticks of the Southern Cone of America: diagnosis, distribution and hosts with taxonomy, ecology and sanitary importance. London: Elsevier; 2017. p. 135-42.
- Monje LD, Costa FB, Colombo VC, Labruna MB, Antoniazzi LR, Gamieta I, et al. Dynamics of exposure to *Rickettsia parkeri* in cattle in the Paraná River delta, Argentina. *J Med Entomol*. 2016;53:660-5. <https://doi.org/10.1093/jme/tjv250>
- Monje LD, Fernandez C, Percara A. Detection of *Ehrlichia* sp. strain San Luis and *Candidatus Rickettsia andeanae* in *Amblyomma parvum* ticks. *Ticks Tick Borne Dis*. 2019;10:111-4. <https://doi.org/10.1016/j.ttbdis.2018.09.008>
- Allsopp BA. Natural history of *Ehrlichia ruminantium*. *Vet Parasitol*. 2010;167:123-35. <https://doi.org/10.1016/j.vetpar.2009.09.014>

Address for correspondence: Lucas D. Monje, Laboratorio de Ecología de Enfermedades, Instituto de Ciencias Veterinarias del Litoral, R.P. Kreder 2805, Esperanza, Santa Fe 3080, Argentina; email: lmonje@fcv.unl.edu.ar

Multidrug-Resistant *Salmonella* Serotype Anatum in Travelers and Seafood from Asia, United States

Beth E. Karp, Molly M. Leeper, Jessica C. Chen, Kaitlin A. Tagg, Louise K. Francois Watkins, Cindy R. Friedman

Author affiliations: Centers for Disease Control and Prevention, Atlanta, Georgia, USA (B.E. Karp, M.M. Leeper, J.C. Chen, L.K. Francois Watkins, C.R. Friedman); Weems Design Studio, Inc., Suwanee, Georgia, USA (K.A. Tagg)


DOI: <https://doi.org/10.3201/eid2605.190992>

A multidrug-resistant *Salmonella enterica* serotype Anatum strain reported in Taiwan was isolated in the United States from patients and from seafood imported from Asia. Isolates harbored 11 resistance determinants, including quinolone and inducible cephalosporin resistance genes. Most patients had traveled to Asia. These findings underscore the need for global One Health resistance surveillance.

A sharp increase in *Salmonella enterica* serotype Anatum infections reported in Taiwan during 2016-2017 was associated with emergence of

Novel *Ehrlichia* Strain Infecting Cattle Tick *Amblyomma neumanni*, Argentina, 2018

Appendix



Appendix Figure. Maximum Likelihood tree constructed by using shorter *dsb* sequences reported from other South American *Ehrlichia* sp. (210 positions included in the final dataset). Phylogenetic tree was constructed by using the program Mega 7.0 (<https://www.megasoftware.net/>) and best fitting substitution models were determined with the Akaike Information Criterion using the ML model test. Numbers represent bootstrap support generated from 1000 replications. GenBank accession numbers are in brackets.