

# New Sequence Types and Antimicrobial Drug-Resistant Strains of *Streptococcus suis* in Diseased Pigs, Italy, 2017–2019

Lucilla Cucco, Marta Paniccià, Francesca Romana Massacci, Alessandra Morelli, Massimo Ancora, Iolanda Mangone, Adriano Di Pasquale, Andrea Luppi, Denis Vio, Cesare Cammà, Chiara Francesca Magistrali

*Streptococcus suis* is a pathogen associated with severe diseases in pigs and humans. Human infections have a zoonotic origin in pigs. To assess circulating strains, we characterized the serotypes, sequence types, and antimicrobial susceptibility of 78 *S. suis* isolates from diseased farmed pigs in Italy during 2017–2019. Almost 60% of infections were caused by serotypes 1/2 and 9. All but 1 of the serotype 2 and 1/2 isolates were confined to a single cluster, and serotype 9 isolates were distributed along the phylogenetic tree. Besides sequence type (ST) 1, the serotype 2 cluster included ST7, which caused severe human infections in China in 1998 and 2005. A large proportion of serotype 9 isolates, assigned to ST123, were resistant to penicillin. The emergence of this clone threatens the successful treatment of *S. suis* infection. Characterizing *S. suis* isolates from pigs will promote earlier detection of emerging clones.

*Streptococcus suis* is a major pathogen in pigs and an emerging zoonotic agent (1–4). This bacterium is a natural inhabitant of the upper respiratory tract of pigs and is endemic to all pig-production countries. In pigs, *S. suis* causes meningitis, septicemia, polyserositis, arthritis, and endocarditis, mainly during the postweaning period; it is a source of concern for farmers because of potential economic losses and its effects on the welfare of infected pigs (2). Human

infection is acquired through occupational contact or ingestion of undercooked pork-derived products and is associated with meningitis, endocarditis, septicemia, deafness, and death (5).

*S. suis* is a heterogeneous species. Until 2005, *S. suis* was divided into 35 serotypes (1–34 and 1/2), based on capsular polysaccharides, but 6 serotypes were recently reclassified as belonging to other *Streptococcus* species, leaving 29 currently recognized *S. suis* serotypes (6,7). Most *S. suis* infections in humans and pigs are caused by serotype 2, but the predominant serotypes causing invasive disease in pigs vary according to time and region (8). In some countries in Europe, serotype 9 has emerged as the leading cause of invasive diseases in pigs (2,8–10); prevalence of this serotype has also recently increased in China (5).

Since 2002, the introduction of a standard multi-locus sequence typing (MLST) scheme has improved the description of the epidemiology of *S. suis* infection (8). Sequence types (STs), determined by MLST, are also better predictors of the pathogenicity of a particular isolate than are serotypes (11). Among serotype 2 isolates from pigs, ST1, a highly successful clone associated with most human infections globally, is prevalent in Europe (8). Another sequence type of serotype 2, ST7, was responsible for major *S. suis* epidemics among humans in 1998 and 2005 in China (12). Serotypes other than 2 are less frequently responsible for human infections (8). Of note, despite the increased frequency of pig infections caused by serotype 9, the first human case of serotype 9 infection was documented in Thailand in 2015 (13). That strain was assigned to ST16, an emerging sequence type known for its increased virulence potential and predominance in invasive *S. suis* infections in pigs in the Netherlands (14).

Author affiliations: Istituto Zooprofilattico Sperimentale dell'Umbria e delle Marche 'Togo Rosati,' Perugia, Italy (L. Cucco, M. Paniccià, F.R. Massacci, A. Morelli, C.F. Magistrali); Istituto Zooprofilattico Sperimentale di Abruzzo e Molise 'Giuseppe Caporale,' Teramo, Italy (M. Ancora, I. Mangone, A. Di Pasquale, C. Cammà); Istituto Zooprofilattico Sperimentale della Lombardia e della Emilia-Romagna 'Bruno Ubertini,' Brescia, Italy (A. Luppi); Istituto Zooprofilattico Sperimentale delle Venezie, Padova, Italy (D. Vio)

DOI: <https://doi.org/10.3201/eid2801.210816>

On-farm management options for controlling *S. suis* infections include improving environmental conditions (e.g., providing correct temperature, providing correct air humidity, and reducing overcrowding and pig mixing) (15). The control of viral infections, particularly porcine reproductive and respiratory virus, is also essential because they are well-known predisposing factors for the disease (16). Another tool for protecting against infection is vaccination, but available vaccines are based on bacterins and provide only nonheterologous protection (17). Thus, in many countries in Europe, including Italy, control of *S. suis* infections in pigs is based mainly on antimicrobial treatment (18). *S. suis* is generally susceptible to  $\beta$ -lactams, the main class of antimicrobials administered to control the infection on pig farms. Conversely, *S. suis* is almost always resistant to tetracycline; macrolide-lincosamide-streptogramin B; and, less frequently, aminoglycosides, chloramphenicol, vancomycin, and linezolid (15). In *S. suis*, genes encoding antimicrobial resistance (AMR) are often carried on mobile genetic elements that can be transferred to other members of the genus, including human pathogens (4,19). Thus, *S. suis* can be considered a public health concern because of its zoonotic potential (a leading cause of antimicrobial drug use in pig farming) and a reservoir of AMR genes (4,19).

Information about circulating strains is lacking in many countries, including Italy, which is one of the most prominent pig-production countries in Europe (8). We characterized the serotypes, sequence types, and antimicrobial susceptibility of 78 *S. suis* isolates from infected pigs in Italy. By providing updated epidemiologic information about *S. suis* infection, we aim to drive the use of autogenous vaccines, reduce antibiotic consumption, and protect animal health. We also assessed presence of *S. suis* clones with zoonotic potential.

## Materials and Methods

### Bacterial Isolates

We investigated isolates collected from pigs with clinical *S. suis* infection on pig farms in northern/central Italy during 2017–2019. To avoid redundancy, we included only 1 isolate per year and farm. A total of 78 *S. suis* isolates were collected from piglets with meningitis (49), pericarditis (1), arthritis (3), septicemia (17), and pneumonia (8) (Appendix 1, <https://wwwnc.cdc.gov/EID/article/28/1/21-0816-App1.xlsx>).

The samples were cultured on 5% sheep blood agar (Biolife Italiana Srl, <http://www.biolifeit.com>) at 5% CO<sub>2</sub>, 37°C, for 24–48 h. We used matrix-assist-

ed laser desorption/ionization time-of-flight mass spectrometry (Bruker Daltonics GmbH, <https://www.bruker.com>) and PCR to confirmed selected suspected  $\alpha$ -hemolytic colonies as belonging to the *S. suis* species (20).

### Serotyping and Virulence Genotyping

We identified serotype and virulence-associated genes by using PCR (Appendix 2 Table 1, <https://wwwnc.cdc.gov/EID/article/28/1/21-0816-App2.pdf>). To discriminate between different variants of *mvp*, we used whole-genome sequencing (21–23).

### Antimicrobial Susceptibility Testing

We assessed MICs by using a commercially prepared microtiter MIC panel (BOP06F, Sensititre; Trek Diagnostic Systems Inc., <https://www.thermofisher.com>) according to the manufacturer's instructions and by using *Streptococcus pneumoniae* ATCC 49619 as a quality control strain. We interpreted MIC results by using the breakpoints recommended by the Clinical Laboratory Standards Institute (24) for swine respiratory *S. suis*. The interpretative criteria for trimethoprim/sulfamethoxazole and clindamycin were those recommended for human *S. pneumoniae* (25).

### Whole-Genome Sequencing

We prepared genomic DNA from all 78 *S. suis* isolates. We extracted pure cultures from 1 mL of logarithmic-phase broth cultures by using QIAamp DNA Mini Kit (QIAGEN, <https://www.qiagen.com>) according to the manufacturer's instructions and then quantified the DNA by using the Qubit fluorometer (Thermo Fisher Scientific, <https://www.thermofisher.com>). We prepared the libraries by using the Nextera XT Library Prep kit (Illumina Inc., <https://www.illumina.com>) and then loaded them onto an Illumina NextSeq 500/550 Mid Output Reagent Cartridge version 2 kit (300 cycles) and sequenced them on an Illumina NextSeq 500 platform to generate 150-bp paired-end reads.

### Sequence Analyses

Raw data were checked for quality, trimmed by using Trimmomatic version 0.36 (26), and assembled by using SPAdes genome assembler version 3.11.1 (27). To determine distinct sequence types, we performed MLST. The allele sequences and profiles were obtained from the *S. suis* MLST database (<https://pubmlst.org/ssuis>). We uploaded sequences for new MLST allele variations to the same database for assignment of allele identification and then uploaded final allele combinations for assignment of new

MLSTs. We submitted the raw sequencing data to the National Center for Biotechnology Information Sequence Read Archive repository (BioProject PRJNA717238, Biosample SUB9357225; accession nos. SAMN18490763–SAMN18490790).

To identify potential clonal complexes and founders, we performed global optimal eBURST (<http://www.phylovis.net/goeburst> analysis). The entire *S. suis* MLST database was displayed as a single goeBURST diagram by setting the double-locus variants level and the group definition to 0 of 7 shared alleles. We conducted minimum core-genome sequence typing in silico (28).

We annotated genomes by using Prokka (<https://github.com>) and constructed a maximum-likelihood phylogenetic tree, based on the final alignment of core genome from Roary analysis, by using FastTree 2.1.11 (29). Manual annotation of the tree was performed in iTOL (v.5.7) (30). We identified AMR genes by using ABRicate (<https://github.com>) against the following databases: AMRFinderPlus, CARD, RESfinder, ARG-ANNOT (31–34).

To research putative virulence genes, we created a database containing 91 previously described genes (2,3) (Appendix 2 Table 1) and searched by using BLASTN version 2.5.0+ (35). According to O’Dea et al., (17), only genes with  $\geq 95\%$  coverage and  $\geq 99\%$  identity were considered present. We investigated the null hypothesis of a random distribution of the

number of virulence factors among the different sequence types and excluded sequence types represented by a small set of isolates ( $\leq 3$ ), resulting in 8 sequence types and 65 isolates. To show the distribution of the putative virulence genes across the sequence types, we selected the putative virulence genes that were present in  $< 90\%$  or in  $> 10\%$  of isolates. After checking the normality of the data by using the Shapiro-Wilk normality test, we performed Kruskal-Wallis rank-sum testing, followed by pairwise comparisons using the Dunn test for multiple comparisons of independent samples. To investigate the distribution of genes encoding putative virulence factors, we constructed a heat map based on the distance metric “euclidean” and complete linkage method. We performed all analyses in R (36).

## Results

### Molecular Serotyping, Virulence Genotyping, and MLST

We identified 13 serotypes: 1, 2, 1/2, 3, 4, 5, 7, 8, 9, 10, 15, 19, and 23. The most prevalent were serotypes 9, accounting for 34.6% ( $n = 27$ ) of isolates, and 1/2, accounting for 25.6% ( $n = 20$ ) of isolates. These serotypes were followed by 10 ( $n = 7$ , 9.0%), 2 ( $n = 7$ , 9.0%), and 7 ( $n = 6$ , 7.7%) (Table 1).

MLST analysis revealed that 59 (75.6%) isolates belonged to 9 sequence types (ST1, ST7, ST11, ST16, ST28, ST29, ST94, ST108, and ST123) in the *S. suis*

**Table 1.** Combination of putative virulence genes among sequence types and minimum core genome groups of *Streptococcus suis* from diseased pigs, Italy, 2017–2019\*

| Sequence type | MCG group | Serotype | Virulence profile                                  | No. isolates/total no. isolates for each sequence type (%) |
|---------------|-----------|----------|--|--|
| ST1           | 1         | 2        | <i>mrp<sup>EU</sup>/sly/epf</i>                    | 3/17 (17.6)  |
| ST1           | 1         | 1/2      | <i>mrp<sup>EU</sup>/sly/epf</i>                    | 14/17 (82.4)   |
| ST7           | 1         | 1/2      | <i>mrp<sup>EU</sup>/sly/epf</i>                    | 6/9 (66.6)   |
| ST7           | 1         | 2        | <i>mrp<sup>EU</sup>/sly/epf</i>                    | 3/9 (33.3)   |
| ST11          | N         | 1        | <i>mrp/sly/epf</i>                                 | 2/2 (100)  |
| ST16          | 1         | 9        | <i>mrp<sup>*</sup>/sly</i>                         | 3/3 (100)  |
| ST28          | 4         | 2        | <i>mrp<sup>NA1</sup>/sly</i>                       | 1/1 (100)  |
| ST29          | 4         | 7        | <i>mrp<sup>*</sup></i><br><i>mrp<sup>NA1</sup></i> | 3/6 (50)<br>3/6 (50)                                       |
| ST94          | 3         | 4        | <i>mrp<sup>NA1</sup>/sly</i>                       | 2/3 (66.7)   |
|               |           | 9        | <i>mrp<sup>NA1</sup>/sly</i>                       | 1/3 (33.3)   |
| ST108         | 3         | 23       | <i>mrp<sup>NA1</sup>/sly</i>                       | 1/1 (100)  |
| ST123         | 3         | 9        | <i>mrp<sup>NA1</sup>/sly</i>                       | 17/17 (100)  |
| ST1540        | N         | 9        | -  | 3/3 (100)  |
| ST1541        | 1         | 9        | -  | 1/1 (100)  |
| ST1542        | N         | 3        | -  | 1/1 (100)  |
| ST1543        | 3         | 4        | <i>mrp<sup>NA1</sup>/sly</i>                       | 1/1 (100)  |
| ST1544        | 3         | 4        | <i>mrp<sup>NA1</sup>/sly</i>                       | 1/2 (50)   |
|               |           | 5        | <i>mrp<sup>NA1</sup>/sly</i>                       | 1/2 (50)   |
| ST1545        | 1         | 8        | <i>mrp<sup>*</sup></i>                             | 1/1 (100)  |
| ST1546        | 1         | 8        | <i>mrp/sly</i>                                     | 1/1 (100)  |
| ST1547        | 1         | 10       | -  | 7/7 (100)  |
| ST1548        | N         | 15       | <i>sly</i>   | 1/1 (100)  |
| ST1549        | N         | 19       | -  | 1/1 (100)  |

\*Dashes indicate absence of putative virulence genes according to PCR. *mrp<sup>\*</sup>* is the *mrp* variant (22). MCG, minimum core genome; N, not groupable; ST, sequence type.

MLST database, and 10 new sequence types were identified as ST1540–1549 (ID2702–ID2711; <https://pubmlst.org/organisms/streptococcus-suis>). Most were singletons, except for ST1547 (n = 7, all serotype 10), ST1540 (n = 3, serotype 9), and ST1544 (n = 2, serotypes 4 and 5). With 17 isolates each, the predominant STs were ST1 and ST123, which accounted for 43% of all isolates. All isolates belonging to ST29 were serotype 7, and all isolates belonging to ST11 were serotype 1.

### AMR Phenotypes and Genotypes

A total of 7 (9.0%) *S. suis* isolates were resistant to antimicrobial drugs, usually tetracycline (6/7) (Table 2). Most (48/78, 61.5%) isolates were resistant to 2 antimicrobials, generally (45/48) clindamycin and tetracycline. Of the 78 isolates, 23 (29.5%) were resistant to  $\geq 3$  antimicrobials and were classified as multiresistant. Multiresistance was detected in 4/17 (23%) ST1, 1/9 (11%) ST7, and 12/17 (71%) ST123 isolates. Of 17 ST123 isolates, 14 (82%) were resistant to penicillin (Tables 1, 3; Figure 1).

### Phylogenetic Analyses

A total of 1,156 genes, corresponding to 19.88% of the pangenome (Appendix 2), comprised the core genome. Phylogenetic analysis of the collected isolates showed 4 major clusters and 2 singletons. The first cluster was composed of serotypes 1, 1/2, and 2 isolates and was characterized by low heterogeneity, even though these isolates originated from different regions and body sites. We found no relationship among the 4 isolates of the second cluster, belonging to serotypes 8 and 9. Cluster 3 recognized 2 sister groups: the first group comprised serotypes 9, 15, and 19; the second group included serotype 10 isolates. The serotype SS10 isolates were derived from 7 outbreaks of meningitis, 6 of which occurred in the

Piedmont region of northern Italy in 2018. The fourth cluster included the highest number of isolates from our collection, belonging to 6 serotypes. We found no correlations with geographic location, year, or site of origin for members of this cluster. All penicillin-resistant SS9 isolates were grouped within this cluster (Figure 1). goeBurst analysis (<http://www.phylovis.net/goeburst>) showed 5 major clusters. ST1543 and ST1544 were related to the ST94 subgroup, ST1546 to the ST1521 subgroup, and ST1545 to clonal complex (CC) 1; the other sequence types occurred as singletons (Appendix 2 Figure 1).

### Putative Virulence Genes

When we investigated the distribution of putative virulence genes in a subset of 65 isolates belonging to 8 sequence types (ST1, ST7, ST16, ST29, ST94, ST123, ST1540, and ST1547), we found 61 putative virulence genes in  $>10\%$  and  $<90\%$  of the isolates and included them in the heat map. Putative virulence genes were not randomly distributed across the 8 sequence types ( $p < 0.001$  by Kruskal-Wallis test; Figure 2). The number of putative virulence genes in ST1 and ST7 isolates did not differ. The number of putative virulence genes in ST1 and ST7 isolates differed from the number in ST123, ST29, ST1540, and ST1547 ( $p < 0.05$  by Dunn test). A block of 38 putative virulence genes was characteristic of ST1 and ST7 isolates. This block included genes encoding components of the cell wall, proteases, and molecules related to adhesion (*cps2E*, *cps2F*, *cps2C*, *neuB*, *fbps*, *sbp2*, *pgdA*, *dppIV*, *igaP*, *ssnA*, *srtF*, and *gnd*) and putative virulence genes involved in the regulation of metabolic pathways (*ccpA*, *lspA*, *ssu1889*, *revS*, *virA*, *guaB*, *sodA*, *adcR*, *purA*, *nadR*, *stp*, *stk*, *vapE*, *lysS*, *Ssads*, *fls*, *apuA*, *aroK*, *flps*, *proA*, *scrB*, *ofs*, *prtP*, *gtfA*, *perR*, and *fur*). Virulence genes harbored by the 89-kb pathogenicity island, including *SalK/SalR* and *tetM*, were not found in ST7 isolates. ST123 was

**Table 2.** Distribution of MICs among 78 *Streptococcus suis* isolates from diseased pigs, Italy, 2017–2019\*

| Antibiotic molecule | No. (%) isolates by MIC, $\mu\text{g/mL}$ |         |         |         |         |         |         |         |         |       |         |     |         |
|---------------------|---|---------|---------|---------|---------|---------|---------|---------|---------|-------|---------|-----|---------|
|                     | 0.12                                      | 0.25    | 0.5     | 1       | 2       | 4       | 8       | 16      | 32      | 64    | 128     | 256 | 512     |
| Penicillin G        | 49 (63)                                   | 7 (9)   | 5 (6)   | 14 (18) | 3 (4)   |         |         |         |         |       |         |     |         |
| Ampicillin          |   | 73 (94) | 2 (3)   | 3 (4)   |         |         |         |         |         |       |         |     |         |
| Enrofloxacin        |   | 1 (1)   | 17 (22) | 57 (73) | 2 (3)   | 1 (1)   |         |         |         |       |         |     |         |
| Tetracycline        |   |         |         | 1 (1)   | 3 (4)   | 1 (1)   |         | 73 (94) |         |       |         |     |         |
| Florfenicol         |   | 1 (1)   | 3 (4)   | 40 (51) | 32 (41) |         | 2 (3)   |         |         |       |         |     |         |
| TMP/SXT             |   |         |         |         | 67 (86) | 11 (14) |         |         |         |       |         |     |         |
| Clindamycin†        |   | 8 (10)  |         |         |         | 4 (5)   | 4 (5)   | 10 (13) | 50 (64) |       |         |     |         |
| Tylosin†            |   |         | 15 (19) |         |         |         |         |         |         |       |         |     | 61 (78) |
| Neomicin†           |   |         |         |         |         | 22 (28) | 27 (35) | 15 (19) | 7 (9)   | 5 (6) |         |     |         |
| Gentamicin†         |   |         |         |         |         | 12 (15) | 39 (50) | 17 (22) | 6 (8)   | 2 (3) |         |     |         |
| Sulfadimethoxine†   |   |         |         |         |         |         |         |         |         |       | 19 (24) |     | 57 (73) |

\*Gray shading indicates range of values actually tested for each antibiotic. Black vertical bars indicate threshold values for clinical resistance, according to the Clinical and Laboratory Standards Institute (<https://clsi.org>). TMP/SXT, trimethoprim/sulfamethoxazole.

†Two isolates were not identified.



**Table 3.** Antimicrobial resistance genes identified in the 78 *Streptococcus suis* isolates from diseased pigs, Italy, 2017–2019, by sequence type

| Antimicrobial resistance genes   | Sequence type |       |        |      |     |      |      |        |        | Total |
|--|---------------|-------|--------|------|-----|------|------|--------|--------|-------|
|  | ST1           | ST123 | ST1547 | ST29 | ST7 | ST16 | ST94 | ST1540 | Other* |       |
| <i>ermB</i> , <i>tet</i> (O)   | 11            | 15    | 3      | 3    | 8   | 3    | 1    |        | 2      | 46    |
| <i>tet</i> (O)   | 1             |       | 4      |      | 1   |      |      |        |        | 6     |
| None   |               |       |        | 1    |     |      | 1    |        | 3      | 5     |
| <i>Cv ermB</i> , <i>tet</i> (O), <i>dfr</i> (F)  | 5             |       |        |      |     |      |      |        |        | 5     |
| <i>aac6-aph2</i> , <i>ant6-ia</i> , <i>aph3-iiiA</i> , <i>spw</i> , <i>ermB</i> , <i>tet</i> (40), <i>tet</i> (W), <i>tet</i> (O), <i>tet</i> (O/W/32/O), <i>tet</i> (W/N/N) |               |       |        |      |     |      |      | 3      | 1      | 4     |
| <i>tet</i> (M)   |               |       |        | 1    |     |      |      |        |        | 1     |
| <i>ermB</i> , <i>tet</i> (M)   |               |       |        | 1    |     |      |      |        |        | 1     |
| <i>ermB</i> , <i>tet</i> (W), <i>tet</i> (O/W/32/O), <i>tet</i> (W/N/N)  |               |       |        |      |     |      |      |        | 1      | 1     |
| <i>ant6ia</i> , <i>aadE</i> , <i>ermB</i> , <i>tet</i> (O)   |               |       |        |      |     |      |      |        | 1      | 1     |
| <i>aac6-aph2</i>   |               | 1     |        |      |     |      |      |        |        | 1     |
| <i>ant6ia</i> , <i>aph3-iiiA</i> , <i>spw</i> , <i>cat</i>   |               |       |        |      |     |      | 1    |        |        | 1     |
| <i>ant6ia</i> , <i>aph3-iiiA</i> , <i>apmA</i> , <i>ermB</i> , <i>optrA</i> , <i>tet</i> (40), <i>spw</i>  |               |       |        |      |     |      |      |        | 1      | 1     |
| <i>aac6-aph2</i> , <i>ermB</i> , <i>tet</i> (O)  |               |       |        |      |     |      |      |        | 1      | 1     |
| <i>ant6-ia</i> , <i>aadE</i> , <i>ermB</i> , <i>tet</i> (W), <i>tet</i> (O), <i>tet</i> (O/W/32/O), <i>tet</i> (W/N/N)   |               |       |        |      |     |      |      |        | 1      | 1     |
| <i>ant6-ia</i> , <i>aadE</i> , <i>ermB</i> , <i>tet</i> (40), <i>tet</i> (O), <i>tet</i> (O/32/O)  |               |       |        |      |     |      |      |        | 1      | 1     |
| <i>ant6-ia</i> , <i>spw</i> , <i>lnuB</i> , <i>lsaE</i> , <i>tet</i> (O)   |               |       |        |      |     |      |      |        | 1      | 1     |
| <i>aac-aph2</i> , <i>aad</i> (6), <i>spw</i> , <i>ermB</i> , <i>erm</i> (47), <i>lnuB</i> , <i>lsaE</i> , <i>tet</i> (40), <i>tet</i> (T)                                    |               | 1     |        |      |     |      |      |        |        | 1     |
| Total  | 17            | 17    | 7      | 6    | 9   | 3    | 3    | 3      | 13     | 78    |

\*Sequence types (STs) represented by <3 isolates.

characterized by the presence of another group of 12 putative virulence genes related to adhesion (*murM*) or involved in metabolic pathways (*htpsC*, *ppc*, *troA*, *pyrF*, *nox*, *purD*, *msmK*, *gloA*, *rgg*, and *yhbU\_2*, *lysM*) (Appendix 2 Figure 3).

## Discussion

Among 78 *S. suis* isolates from diseased pigs in Italy, we identified the major serotypes associated with *S. suis* infections as serotypes 9 and SS1/2, responsible for almost 60% of cases. Previous studies in Italy have shown a predominance of serotype 2 infections, which were a minority in our study (1,37). Until 2020, PCR testing for serotyping did not differentiate serotype 2 from serotype 1/2, and many studies reported these 2 serotypes as belonging to the same category. However, such distinction is relevant because serotype 1/2 is associated with pig infections; however, different from serotype 2, its role as a zoonotic agent is still uncertain (38). With this study, we confirmed increased serotype 9 infections at pig farms, which has already been described for other countries in Europe (10). The proportion of isolates belonging to serotype 7, the third most common serotype, remained more or less stable compared with data from 2000, thus confirming the trend observed in Germany (7,37). We also detected serotypes 10 and 15 in our collection. These serotypes have not previously been detected in Italy but have been identified in Spain and the United Kingdom (37). Given the variability of serotypes and the low prevalence of serotype 2 observed in our study, complete characterization of

isolates is essential for the successful implementation of autogenous vaccines. Indeed, autogenous vaccines are well-established tools for preventing serotype 2 infections, but data about their efficacy for other serotypes, including 1/2, are still lacking (16).

All serotype 2 and 1/2 isolates, except 1, were confined to a single cluster of the phylogenetic tree. This cluster was composed of ST1 and, unexpectedly, ST7, which is a subgroup founder related to CC1. The number of putative virulence factors was higher in ST7 and ST1 isolates than in other STs in our collection, which was expected, because ST1 is the predominant sequence type associated with invasive infections in pigs in Europe. Cases of *S. suis* infection in humans in Italy are sporadic and are caused by serotype 2, ST1 (1,39). ST7, which differs from ST1 at a single locus, has not been detected in pigs in Europe (8). However, ST7 isolates are prevalent among diseased pigs in China (40–42). The epidemic strain ST7, which is characterized by the presence of an 89-kb pathogenicity island, the insertion of a 128-kb ICE (integrative and conjugative element)–phage tandem mobile genetic element, is responsible for the 2 largest outbreaks of human *S. suis* infection, which occurred in 1998 and 2005 (42). The ST7 isolates from our study lacked the virulence genes harbored by the 89-kb pathogenicity island; thus, their zoonotic potential may be lower than that of the ST7 epidemic strain in China. Moreover, they did not cluster with the newly described lineage III of ST7 (Appendix 2 Tables 2–4, Figure 2) (43). Further analysis is necessary to explain the presence of ST7 in Italy. New *S. suis* strains may


be imported by living animals or traveling humans, or they may have been derived from an early mixing of pig breeds, as previously hypothesized (3,40).

All serotype 7 isolates belonged to ST29, grouped in cluster 4, and had 2 *mrp* gene variants. The same characteristics were described for serotype 7 ST29 isolates from recent severe outbreaks among piglets in Germany and Austria (18). Thus, ST29 has been suggested as an emerging virulent sequence type in Europe (18).

In contrast to the serotype 2 and 1/2 isolates, serotype 9 isolates were distributed among different clusters in the phylogenetic tree, grouping with isolates belonging to other serotypes. High heterogeneity has been reported for serotype 9 (11,44). Three isolates from our collection were typed as serotype 9 ST16, a dominant clone in diseased pigs from the Nether-

lands (3). Although most cases in humans have been attributed to ST1 isolates, ST16 has recently been associated with cases of *S. suis* infection in humans in Thailand (13). It has been suggested that the zoonotic and virulence potential may be higher for ST16 than for other strains. In our study, the ST16 subgroup was related to CC1, harbored *mrp* and *sly* genes, and was close to ST1 and ST7 in the phylogenetic tree, in accordance with the results reported by Zheng et al. (10). The presence of ST16 in Italy suggests the need for monitoring and typing *S. suis* from diseased pigs and infected humans in a One Health scenario.

A large proportion of serotype 9 isolates were assigned to ST123 and grouped into cluster 4 in the phylogenetic tree. This sequence type was prevalent in our collection and was found in 5 regions of Italy and




**Figure 1.** Circular phylogenetic tree containing 78 *Streptococcus suis* isolates from diseased pigs, Italy, 2017–2019. The tree was inferred by using the iTOL interactive user interface (<https://itol.embl.de>). Shading over tip labels indicates sequence types. The serotypes of each isolate are also shown. The antimicrobial-resistant molecules are annotated by colors and shapes. Scale bar indicates substitutions per site. CLI, clindamycin; ENR, enrofloxacin; FFC, florfenicol; PEN, penicillin; ST, sequence type; TET, tetracycline; TMP/SXT, trimethoprim/sulfamethoxazole.

in pigs from different production companies. Most ST123 isolates were resistant to penicillin. ST123 was reported in Spain in 2009 (9). As already observed in Spain, the ST123 isolates from our study were related to the ST94 subgroup and were characterized by the presence of *sly* and *mrp<sup>NA1</sup>* genes (10).

The number of live pigs imported into Italy has increased over the past 10 years, almost doubling from 2013 to 2018 (<http://www.anas.it>). Pigs are imported from other countries in Europe, predominantly the Netherlands, Denmark, and Germany, and, to a lesser extent, from Spain and countries in eastern Europe. Imported live pigs can be carriers of new *S. suis* clones, which may then be transferred to other animals at the receiving farm (11). The differences in the *S. suis* population in our study compared with those in previous studies may result from this intensive exchange of live pigs between Italy and other countries in Europe.

We confirmed widespread resistance to tetracycline and clindamycin, as previously reported for *S. suis* isolates globally. Resistance to tetracycline was mainly associated with the presence of *tetO* and, to a lesser extent, other *tet* genes, including *tetM* and the mosaic gene *tet* (*O/W/32/O*), which was first described in *S. suis* isolates in Italy (1). Resistance to clindamycin was coupled with high MICs for tylosin and the diffuse presence of *ermB*, suggesting a macrolide/lincosamide/streptogramin B profile. Resistance to florfenicol was detected in 2 multiresistant isolates. This type of resistance is emerging in *S. suis* species (45). One of the florfenicol-resistant isolates was positive for *optrA*, an oxazolidinone/phenicol resistance determinant carried by mobile genetic elements. *optrA* in *S. suis* isolates from China has been previously described and is frequently detected in *Enterococcus* isolates from pig farms in Italy (46,47). The high levels of AMR and the detection of emerging drug-resistance determinants are a consequence of selective pressure caused by antibiotic overuse. Despite the declining trend in antibiotic consumption, the use of antibiotics in veterinary medicine is still more frequent in Italy than in other countries in Europe (48).

We observed a high level of resistance to penicillin;  $\approx 1$  in 5 isolates showed reduced susceptibility to this antimicrobial. This finding contrasts with previous observations from other countries in Europe (15,49). Resistance to ampicillin was not observed, thus confirming the hypothesis of incomplete cross-resistance between these 2 antimicrobials (49). Resistance to penicillin was mostly detected in serotype 9 isolates and was particularly frequent in ST123 isolates. Blume et al. (9) suggested that the spread of *S.*



**Figure 2.** Distribution of the putative virulence genes detected among different sequence types of *Streptococcus suis* isolates from diseased pigs, Italy, 2017–2019. Box tops and bottoms indicate interquartile ranges, horizontal lines within boxes indicate means, whiskers indicate 95% CIs, and dots indicate outliers. ST, sequence type.

*suis* serotype 9 is favored by the selective advantage conferred by the absence of heterologous immunity induced by the dominant serotype 2 clone (9). Our data suggest that penicillin resistance may also be a driver of the expansion of *S. suis* serotype 9.

The emergence of a penicillin-resistant clone among the *S. suis* population threatens the successful treatment of *S. suis* infections in pigs. Penicillin resistance in *S. suis* may favor the prescription of critical classes of antimicrobial drugs, which should be limited in veterinary medicine. Penicillin resistance in a zoonotic agent raises concerns about hampering the treatment of infections.

In conclusion, our study highlights the value of characterizing *S. suis* isolates from pigs for monitoring trends in AMR and enabling early detection of emerging clones. In addition, our data strongly suggest the need for preventive strategies to limit the spread of penicillin-resistant *S. suis* among pig populations in Italy.

#### Acknowledgments

We thank Marcelo Gottschalk for his support and advice for setting up the molecular assays.

This study was funded by the Italian Ministry of Health (RC007/2018 IZSUM). The opinions expressed by authors contributing to this journal do not necessarily reflect the opinions of the Centers for Disease Control and Prevention but do represent the institutions with which the authors are affiliated.



## About the Author

Dr. Cucco is a microbiologist at the Istituto Zooprofilattico Sperimentale dell'Umbria e delle Marche, in Perugia, Italy. Her primary research interests include epidemiology, AMR, and genomics of bacterial pathogens in veterinary medicine.

## References

- Princivalli MS, Palmieri C, Magi G, Vignaroli C, Manzin A, Camporese A, et al. Genetic diversity of *Streptococcus suis* clinical isolates from pigs and humans in Italy (2003–2007). *Euro Surveill.* 2009;14:19310. <https://doi.org/10.2807/ese.14.33.19310-en>
- Segura M, Fittipaldi N, Calzas C, Gottschalk M. Critical *Streptococcus suis* virulence factors: are they all really critical? *Trends Microbiol.* 2017;25:585–99. <https://doi.org/10.1016/j.tim.2017.02.005>
- Willemsse N, Howell KJ, Weinert LA, Heuvelink A, Pannekoek Y, Wagenaar JA, et al. An emerging zoonotic clone in the Netherlands provides clues to virulence and zoonotic potential of *Streptococcus suis*. *Sci Rep.* 2016;6:28984. <https://doi.org/10.1038/srep28984>
- Palmieri C, Varaldo PE, Facinelli B. *Streptococcus suis*, an emerging drug-resistant animal and human pathogen. *Front Microbiol.* 2011;2:235. <https://doi.org/10.3389/fmicb.2011.00235>
- Dong W, Zhu Y, Ma Y, Ma J, Zhang Y, Yuan L, et al. Multilocus sequence typing and virulence genotyping of *Streptococcus suis* serotype 9 isolates revealed high genetic and virulence diversity. *FEMS Microbiol Lett.* 2017;364:1–8. <https://doi.org/10.1093/femsle/fnx192>
- Estrada AA, Gottschalk M, Rossow S, Rendahl A, Gebhart C, Marthaler DG. Serotype and genotype (multilocus sequence type) of *Streptococcus suis* isolates from the United States serve as predictors of pathotype. *J Clin Microbiol.* 2019;57:e00377–19.
- Prüfer TL, Rohde J, Verspohl J, Rohde M, de Greeff A, Willenborg J, et al. Molecular typing of *Streptococcus suis* strains isolated from diseased and healthy pigs between 1996–2016. *PLoS One.* 2019;14:e0210801. <https://doi.org/10.1371/journal.pone.0210801>
- Goyette-Desjardins G, Auger JP, Xu J, Segura M, Gottschalk M. *Streptococcus suis*, an important pig pathogen and emerging zoonotic agent—an update on the worldwide distribution based on serotyping and sequence typing. *Emerg Microbes Infect.* 2014;3:e45. <https://doi.org/10.1038/emi.2014.45>
- Blume V, Luque I, Vela AI, Borge C, Maldonado A, Domínguez L, et al. Genetic and virulence-phenotype characterization of serotypes 2 and 9 of *Streptococcus suis* swine isolates. *Int Microbiol.* 2009;12:161–6.
- Zheng H, Du P, Qiu X, Kerdsin A, Roy D, Bai X, et al. Genomic comparisons of *Streptococcus suis* serotype 9 strains recovered from diseased pigs in Spain and Canada. *Vet Res (Faisalabad).* 2018;49:1–13. <https://doi.org/10.1186/s13567-017-0498-2>
- Estrada AA, Gottschalk M, Rossow S, Rendahl A, Gebhart C, Marthaler DG. Serotype and genotype (multilocus sequence type) of *Streptococcus suis* isolates from the United States serve as predictors of pathotype. *J Clin Microbiol.* 2019; 57:1–16. <https://doi.org/10.1128/JCM.00377-19>
- Ye C, Bai X, Zhang J, Jing H, Zheng H, Du H, et al. Spread of *Streptococcus suis* sequence type 7, China. *Emerg Infect Dis.* 2008;14:787–91. <https://doi.org/10.3201/eid1405.070437>
- Kerdsin A, Hatrongjit R, Gottschalk M, Takeuchi D, Hamada S, Akeda Y, et al. Emergence of *Streptococcus suis* serotype 9 infection in humans. *J Microbiol Immunol Infect.* 2017;50:545–6. <https://doi.org/10.1016/j.jmii.2015.06.011>
- Willemsse N, van der Ark KCH, Stockhofe-Zurwieden N, Smith H, Picavet DJ, van Solt-Smits C, et al. Clonal expansion of a virulent *Streptococcus suis* serotype 9 lineage distinguishable from carriage subpopulations. *Sci Rep.* 2019;9:15429. <https://doi.org/10.1038/s41598-019-51576-0>
- Werinder A, Aspán A, Backhans A, Sjölund M, Guss B, Jacobson M. *Streptococcus suis* in Swedish grower pigs: occurrence, serotypes, and antimicrobial susceptibility. *Acta Vet Scand.* 2020;62:36. <https://doi.org/10.1186/s13028-020-00533-3>
- Rieckmann K, Pendzialek SM, Vahlenkamp T, Baums CG. A critical review speculating on the protective efficacies of autogenous *Streptococcus suis* bacterins as used in Europe. *Porcine Health Manag.* 2020;6:12. <https://doi.org/10.1186/s40813-020-00150-6>
- O'Dea MA, Laird T, Abraham R, Jordan D, Lugsomya K, Fitt L, et al. Examination of Australian *Streptococcus suis* isolates from clinically affected pigs in a global context and the genomic characterisation of ST1 as a predictor of virulence. *Vet Microbiol.* 2018;226:31–40. <https://doi.org/10.1016/j.vetmic.2018.10.010>
- Rieckmann K, Seydel A, Szewczyk K, Klimke K, Rungelrath V, Baums CG. *Streptococcus suis* cps7: an emerging virulent sequence type (ST29) shows a distinct, IgM-determined pattern of bacterial survival in blood of piglets during the early adaptive immune response after weaning. *Vet Res (Faisalabad).* 2018;49:48. <https://doi.org/10.1186/s13567-018-0544-8>
- Du F, Lv X, Duan L, Wang L, Huang J. Characterization of a linezolid- and vancomycin-resistant *Streptococcus suis* isolate that harbors *optrA* and *vanG* operons. *Front Microbiol.* 2019;10:2026. <https://doi.org/10.3389/fmicb.2019.02026>
- Okwumabua O, O'Connor M, Shull E. A polymerase chain reaction (PCR) assay specific for *Streptococcus suis* based on the gene encoding the glutamate dehydrogenase. *FEMS Microbiol Lett.* 2003;218:79–84. <https://doi.org/10.1111/j.1574-6968.2003.tb11501.x>
- Smith HE, Reek FH, Vecht U, Gielkens ALJ, Smits MA. Repeats in an extracellular protein of weakly pathogenic strains of *Streptococcus suis* type 2 are absent in pathogenic strains. *Infect Immun.* 1993;61:3318–26. <https://doi.org/10.1128/iai.61.8.3318-3326.1993>
- Silva LMG, Baums CG, Rehm T, Wisselink HJ, Goethe R, Valentin-Weigand P. Virulence-associated gene profiling of *Streptococcus suis* isolates by PCR. *Vet Microbiol.* 2006; 115:117–27. <https://doi.org/10.1016/j.vetmic.2005.12.013>
- Fittipaldi N, Fuller TE, Teel JF, Wilson TL, Wolfram TJ, Lowery DE, et al. Serotype distribution and production of muramidase-released protein, extracellular factor and suilysin by field strains of *Streptococcus suis* isolated in the United States. *Vet Microbiol.* 2009;139:310–7. <https://doi.org/10.1016/j.vetmic.2009.06.024>
- Clinical and Laboratory Standards Institute. VET08 performance standards for antimicrobial disk. 4th ed. Wayne (PA): The Institute; 2018.
- Clinical and Laboratory Standards Institute. Performance standards for antimicrobial susceptibility testing (M100). Wayne (PA): The Institute; 2017. p. 27–39
- Bolger AM, Lohse M, Usadel B. Trimmomatic: a flexible trimmer for Illumina sequence data. *Bioinformatics.* 2014;30:2114–20. <https://doi.org/10.1093/bioinformatics/btu170>



27. Bankevich A, Nurk S, Antipov D, Gurevich AA, Dvorkin M, Kulikov AS, et al. SPAdes: a new genome assembly algorithm and its applications to single-cell sequencing. *J Comput Biol*. 2012;19:455–77. <https://doi.org/10.1089/cmb.2012.0021>
28. Wileman TM, Weinert LA, Howell KJ, Wang J, Peters SE, Williamson SM, et al. Pathotyping the zoonotic pathogen *Streptococcus suis*: novel genetic markers to differentiate invasive disease-associated isolates from non-disease-associated isolates from England and Wales. *J Clin Microbiol*. 2019;57:1–15. <https://doi.org/10.1128/JCM.01712-18>
29. Price MN, Dehal PS, Arkin AP. FastTree 2 – approximately maximum-likelihood trees for large alignments. *PLoS One*. 2010;5:e9490. <https://doi.org/10.1371/journal.pone.0009490>
30. Letunic J, Bork P. Interactive tree of life (iTOL) v3: an online tool for the display and annotation of phylogenetic and other trees. *Nucleic Acids Res*. 2016;44(W1):W242–5. <https://doi.org/10.1093/nar/gkw290>
31. Feldgarden M, Brover V, Haft DH, Prasad AB, Slotta DJ, Tolstoy I, et al. Validating the AMRFINDER tool and resistance gene database by using antimicrobial resistance genotype-phenotype correlations in a collection of isolates. *Antimicrob Agents Chemother*. 2019;63:1–20. <https://doi.org/10.1128/AAC.00483-19>
32. Jia B, Raphenya AR, Alcock B, Waglechner N, Guo P, Tsang KK, et al. CARD 2017: expansion and model-centric curation of the comprehensive antibiotic resistance database. *Nucleic Acids Res*. 2017;45(D1):D566–73. <https://doi.org/10.1093/nar/gkw1004>
33. Zankari E, Hasman H, Cosentino S, Vestergaard M, Rasmussen S, Lund O, et al. Identification of acquired antimicrobial resistance genes. *J Antimicrob Chemother*. 2012;67:2640–4. <https://doi.org/10.1093/jac/dks261>
34. Gupta SK, Padmanabhan BR, Diene SM, Lopez-Rojas R, Kempf M, Landraud L, et al. ARG-ANNOT, a new bioinformatic tool to discover antibiotic resistance genes in bacterial genomes. *Antimicrob Agents Chemother*. 2014;58:212–20. <https://doi.org/10.1128/AAC.01310-13>
35. Camacho C, Coulouris G, Avagyan V, Ma N, Papadopoulos J, Bealer K, et al. BLAST+: architecture and applications. *BMC Bioinformatics*. 2009;10:421. <https://doi.org/10.1186/1471-2105-10-421>
36. TeamCore. R: A language and environment for statistical computing [cited 2021 Dec 20]. <https://www.R-project.org>
37. Wisselink HJ, Smith HE, Stockhofe-Zurwieden N, Peperkamp K, Vecht U. Distribution of capsular types and production of muramidase-released protein (MRP) and extracellular factor (EF) of *Streptococcus suis* strains isolated from diseased pigs in seven European countries. *Vet Microbiol*. 2000;74:237–48. [https://doi.org/10.1016/S0378-1135\(00\)00188-7](https://doi.org/10.1016/S0378-1135(00)00188-7)
38. Lacouture S, Okura M, Takamatsu D, Corsaut L, Gottschalk M. Development of a mismatch amplification mutation assay to correctly serotype isolates of *Streptococcus suis* serotypes 1, 2, 1/2, and 14. *J Vet Diagn Invest*. 2020;32:490–4. <https://doi.org/10.1177/1040638720915869>
39. Mancini F, Adamo F, Creti R, Monaco M, Alfarone G, Pantosti A, et al. A fatal case of streptococcal toxic shock syndrome caused by *Streptococcus suis* carrying *tet(40)* and *tet(O)/W/32(O)*, Italy. *J Infect Chemother*. 2016;22(11):774–6. <https://doi.org/10.1016/j.jiac.2016.05.011>
40. Fittipaldi N, Xu J, Lacouture S, Tharavichitkul P, Osaki M, Sekizaki T, et al. Lineage and virulence of *Streptococcus suis* serotype 2 isolates from North America. *Emerg Infect Dis*. 2011;17:2239–44. <https://doi.org/10.3201/eid1712.110609>
41. Chen C, Zhang W, Zheng H, Lan R, Wang H, Du P, et al. Minimum core genome sequence typing of bacterial pathogens: a unified approach for clinical and public health microbiology. *J Clin Microbiol*. 2013;51:2582–91. <https://doi.org/10.1128/JCM.00535-13>
42. Guo G, Du D, Yu Y, Zhang Y, Qian Y, Zhang W. Pan-genome analysis of *Streptococcus suis* serotype 2 revealed genomic diversity among strains of different virulence. *Transbound Emerg Dis*. 2021;68:637–47.
43. Dong X, Chao Y, Zhou Y, Zhou R, Zhang W, Fischetti VA, et al. The global emergence of a novel *Streptococcus suis* clade associated with human infections. *EMBO Mol Med*. 2021;13:e13810. <https://doi.org/10.15252/emmm.202013810>
44. Dong W, Ma J, Zhu Y, Zhu J, Yuan L, Wang Y, et al. Virulence genotyping and population analysis of *Streptococcus suis* serotype 2 isolates from China. *Infect Genet Evol*. 2015;36:483–9. <https://doi.org/10.1016/j.meegid.2015.08.021>
45. Zhang C, Zhang P, Wang Y, Fu L, Liu L, Xu D, et al. Capsular serotypes, antimicrobial susceptibility, and the presence of transferable oxazolidinone resistance genes in *Streptococcus suis* isolated from healthy pigs in China. *Vet Microbiol*. 2020;247:108750. <https://doi.org/10.1016/j.vetmic.2020.108750>
46. Hong B, Ba Y, Niu L, Lou F, Zhang Z, Liu H, et al. A comprehensive research on antibiotic resistance genes in microbiota of aquatic animals. *Front Microbiol*. 2018;9:1617. <https://doi.org/10.3389/fmicb.2018.01617>
47. Fioriti S, Morroni G, Cocchitto SN, Brencciani A, Antonelli A, Di Pilato V, et al. Detection of oxazolidinone resistance genes and characterization of genetic environments in enterococci of Swine origin, Italy. *Microorganisms*. 2020;8:2021. <https://doi.org/10.3390/microorganisms8122021>
48. European Centre for Disease Prevention and Control (ECDC); European Food Safety Authority (EFSA); European Medicines Agency (EMA). ECDC/EFSA/EMA second joint report on the integrated analysis of the consumption of antimicrobial agents and occurrence of antimicrobial resistance in bacteria from humans and food-producing animals: Joint Interagency Antimicrobial Consumption and Resistance Analysis (JIACRA) Report. *EFSA J*. 2017;15:e04872.
49. van Hout J, Heuvelink A, Gonggrijp M. Monitoring of antimicrobial susceptibility of *Streptococcus suis* in the Netherlands, 2013–2015. *Vet Microbiol*. 2016;194:5–10. <https://doi.org/10.1016/j.vetmic.2016.03.014>

Address for correspondence: Chiara Francesca Magistrali, Istituto Zooprofilattico Sperimentale dell’Umbria e delle Marche ‘Togo Rosati,’ via Salvemini, 1, 06126 Perugia, Italy; email: c.magistrali@izsum.it

# New Sequence Types and Antimicrobial Drug–Resistant Strains of *Streptococcus suis* in Diseased Pigs, Italy, 2017–2019

## Appendix 2

### Read quality, number of contigs and coverage of Whole-Genome Sequencing data

The final assembly of the 78 sequences resulted in an average read quality after trimming of 34.60% (min 34.04, max 34.89) and 3,165,852 read pairs (min 1,043,985; max 5,077,416). The average number of contigs was 68 (min 25, max 177), with a mean length of 2,114,359 nucleotides (min 2,026,262; max 2,264,709). The average vertical coverage was 211 (min, 69; max, 339). The mean N50 and L50 values were 149,502 nucleotides (min 371, 66; max 293,044) and 6.95 (min 3, max 21), respectively.

### References

1. Okura M, Lachance C, Osaki M, Sekizaki T, Maruyama F, Nozawa T, et al. Development of a two-step multiplex PCR assay for typing of capsular polysaccharide synthesis gene clusters of *Streptococcus suis*. J Clin Microbiol. 2014;52:1714–9. [PubMed](#)  
<https://doi.org/10.1128/JCM.03411-13>
2. Lacouture S, Okura M, Takamatsu D, Corsaut L, Gottschalk M. Development of a mismatch amplification mutation assay to correctly serotype isolates of *Streptococcus suis* serotypes 1, 2, 1/2, and 14. J Vet Diagn Invest. 2020;32:490–4. [PubMed](#)  
<https://doi.org/10.1177/1040638720915869>
3. Silva LMG, Baums CG, Rehm T, Wisselink HJ, Goethe R, Valentin-Weigand P. Virulence-associated gene profiling of *Streptococcus suis* isolates by PCR. Vet Microbiol. 2006;115:117–27. [PubMed](#)  
<https://doi.org/10.1016/j.vetmic.2005.12.013>

4. Wisselink HJ, Reek FH, Vecht U, Stockhofe-Zurwieden N, Smits MAE, Smith HE. Detection of virulent strains of *Streptococcus suis* type 2 and highly virulent strains of *Streptococcus suis* type 1 in tonsillar specimens of pigs by PCR. *Vet Microbiol.* 1999;67:143–57. [PubMed](#) [https://doi.org/10.1016/S0378-1135\(99\)00036-X](https://doi.org/10.1016/S0378-1135(99)00036-X)
5. King SJ, Heath PJ, Luque I, Tarradas C, Dowson CG, Whatmore AM. Distribution and genetic diversity of suilysin in *Streptococcus suis* isolated from different diseases of pigs and characterization of the genetic basis of suilysin absence. *Infect Immun.* 2001;69:7572–82. [PubMed](#) <https://doi.org/10.1128/IAI.69.12.7572-7582.2001>
6. Princivalli MS, Palmieri C, Magi G, Vignaroli C, Manzin A, Camporese A, et al. Genetic diversity of *Streptococcus suis* clinical isolates from pigs and humans in Italy (2003-2007). *Euro Surveill.* 2009;14:19310. [PubMed](#) <https://doi.org/10.2807/ese.14.33.19310-en>
7. Dong X, Chao Y, Zhou Y, Zhou R, Zhang W, Fischetti VA, et al. The global emergence of a novel *Streptococcus suis* clade associated with human infections. *EMBO Mol Med.* 2021;13:e13810. [PubMed](#) <https://doi.org/10.15252/emmm.202013810>



**Appendix 2 Table 1.** Primers used to identify the *S. suis* serotypes and virulence factors by PCR\*

| Target                       | Primer sequences (5' to 3')                            |  | Reference |
|------------------------------|--|--|-----------|
|                              | Forward  | Reverse  |           |
| <b>For grouping PCR</b>      |  |  |           |
| I                            | TGGTTCAAATATCAATGCTC                                   | ATTGGTTGTGAGTGCATTG                                    | (1)       |
| II                           | TCAAAATACGCACCTAAGGC                                   | CACTCACCTGCCCAAGAC                                     |           |
| III                          | TGATTTGGGTGAGACCATG                                    | CTCATGCTGGATAACACGT                                    |           |
| IV                           | ACAGTCGGTCAAGATAATCG                                   | TCAGCTTGGGTAATATCTGG                                   |           |
| V                            | GGAAAGATGGAGGACCAGC                                    | CCAACCAGACTCATATCCCC                                   |           |
| III and VI                   | GATGCCCCAAGCGATATGCC (F1)<br>GACGCACCAAGTGATATGCC (F2) | GGACCAACAATGGCCATCTC (R1)<br>GGTCCGACAATAGCCATTTC (R2) |           |
| <b>For typing PCR</b>        |  |  |           |
| SS group I                   |  |  | (1)       |
| 3                            | GGTTTTGATTGGTCTAGTTG                                   | CTCTAAAGCTCGATATCTAC                                   |           |
| 13                           | TATGGTTAAAGGTGGAAGCTG                                  | CCTTGATATATTCCTCCA                                     |           |
| 18                           | TAATGGGATAGTTGCCATTAC                                  | ATACATAAAGTTGTCCTGCG                                   |           |
| SS group II                  |  |  | (1)       |
| 2                            | TTAGCAACGTTGCCAATAAG                                   | AATCCTCCATTAACCCCTG                                    |           |
| 6                            | GCTCACTATTTTTACATTACAC                                 | TATTAATCCGCCAAATACAG                                   |           |
| 1 and 14                     | TTAGACAGACACCTTATAGG                                   | CTAGCTTCGTTACTTGTATTC                                  |           |
| 16                           | AAGGTTATCCACGAAAGATG                                   | TCCGGCAATATTTCTTTCAAG                                  |           |
| 27                           | AGACACTGCTTGCATTATTG                                   | TCAGAATTACTTCTGTGTTGC                                  |           |
| 1 and 1/2                    | ATCGCTTTGTGGTGGCCCTT                                   | AGAAGCTTCTTTTGTGTTTGC                                  | (2)       |
| 1 and 14                     | ATCGCTTTGTGGTGGCCCTT                                   |  |           |
| SS group III                 |  |  | (1)       |
| 21                           | TATCATATTGAGAATCTTCCC                                  | TTGCGTAGCATACAAAGTTC                                   |           |
| 28                           | ATTATGTTGGTTGCAGAAGG                                   | CGACTCAATTGTTGTAGTAG                                   |           |
| 29                           | TTCTGGGATTTTAGGAATGC                                   | CATGAAATACGCACCTTGTAC                                  |           |
| 30                           | TATTGCACTAGCTTCAGAAC                                   | TGCATCCATAGTTGTATTCC                                   |           |
| SS group IV                  |  |  | (1)       |
| 4                            | GACTATCTGTATACCCAAAC                                   | TCCTTCCAAGTATTCTCTAG                                   |           |
| 5                            | ATCTTAGGAATGATTCGGAC                                   | ACCAGATATCTGAGCAAATG                                   |           |
| 7                            | AACTACCTACCTGAACCTTG                                   | AGTCTAAAAGTGATCGAGTC                                   |           |
| 17                           | TAGCATCAGTTTATACGAGG                                   | TAGTTTATCTGTGACACACC                                   |           |
| 19                           | GTGTCGCAAATCAAGTATTG                                   | AAGCTAGTACAACAAGCATG                                   |           |
| 23                           | TAATGTATGCTCTGTCACTG                                   | AACGAAACGGAATAGTTTGC                                   |           |
| SS group V                   |  |  | (1)       |
| 8                            | AAATAAGGTAGGAGCTACTC                                   | ATCCAACCTTAGCTTTCTGT                                   |           |
| 15                           | ATCGTTTTGAGATTGAGTGG                                   | TAAACGGATTCCGGTACTCA                                   |           |
| 20                           | TGTGGATTTCTGGGATAATC                                   | TGTGGACGAATTACTACTTG                                   |           |
| 22                           | GCATTATCAGGATTCTTTCC                                   | CCAATTGGGTGTTCAAAAAG                                   |           |
| 25                           | GTTTGCTCCGATCATAATAG                                   | CCAGTAAAAGGACTCAATAC                                   |           |
| SS group VI                  |  |  | (1)       |
| 9                            | GAAAGTAGGTATATCTCAGC                                   | GGGCTATTAACCTCCTATC                                    |           |
| 10                           | TTTCCCATTTGCTTATGGAC                                   | GGAATAAAAACGATTGGGAG                                   |           |
| 11                           | ATGCGATTGCAACAATTGAC                                   | AGGCATGAGTAATACATAGG                                   |           |
| 12                           | AACAGGTATTTCAAGATTGC                                   | CTCGGATAAAGATAATCAGC                                   |           |
| 24                           | TACTGAGATTTATTGGGACG                                   | AAGCGATTGGATTACATTGC                                   |           |
| 26                           | TTATACCGAAATTTGTTGCC                                   | CGTCAATCATATAAAGTGGG                                   |           |
| 33                           | GATGTTTTCAACAGGTGTAC                                   | CAAAGTACCTATTTTCAGCG                                   |           |
| SS group VII                 |  |  | (1)       |
| 31                           | ACAATCGTTTCTGCAATACG                                   | GATGAAAACATCGTTGGTAG                                   |           |
|                              | ATCAGTAGTGGGAATAGTTG                                   | TTTACTGTTTTTCGACCGTG                                   |           |
| 32                           | AACCGCTGTTGAATTAAGAG                                   | TTCGTTAGTTGAAGTGTTC                                    |           |
|                              | TAGGACTATGGTTCCTAATG                                   | TATTCTAGTTCAAGTCGCTC                                   |           |
| 34                           | AAGTTTCATTCGAGGACTTC                                   | GTATATAACACCGCAAGAAG                                   |           |
|                              | ATACAGTGATGTCTTGCAAC                                   | ATTGCTTTTTGACAATCGGC                                   |           |
| <b>For virulence factors</b> |  |  |           |
| mrp                          | GACAGATGGTGAGGAAAATGG                                  | TGAGCTTTACCTGAAGCGGT                                   | (3)       |
| epf                          | GCTACGACGGCCTCAGAAATC                                  | TGGATCAACCACTGGTGTAC                                   | (4)       |
| Sly                          | CAGCTCGTTGCCCTGTACTA                                   | ACTCTATCACCTCATCCGC                                    | (5)       |

\*The DNA for SS3 and SS14 were kindly provided by Marcelo Gottschalk (University of Montreal, Canada) and used as positive control. *S. suis* strain V20 from a previous study conducted in Italy (6) was used as control.

**Appendix 2 Table 2.** Genes lacking in Lineage III and Lineage I ST7 reference strains, but present in the ST7 isolates of this study

| Gene       | Annotation  |
|------------|---|
| group_16   | putative autolysin SsaALP                                       |
| group_2145 | IS630 family transposase ISSsu3                                 |
| elaA       | Protein ElaA  |
| group_324  | hypothetical protein  |
| group_325  | hypothetical protein  |
| group_326  | hypothetical protein  |
| group_327  | hypothetical protein  |
| pdxK       | Pyridoxine kinase   |
| hmpT       | Thiamine precursor transporter HmpT                             |
| group_330  | hypothetical protein  |
| group_331  | hypothetical protein  |
| baeS       | Signal transduction histidine-protein kinase BaeS               |
| walR_2     | Transcriptional regulatory protein WalR                         |
| group_334  | hypothetical protein  |
| mco        | Multicopper oxidase mco   |
| group_336  | IS3 family transposase ISStin7                                  |
| group_337  | hypothetical protein  |
| group_338  | hypothetical protein  |
| group_339  | hypothetical protein  |
| group_340  | hypothetical protein  |
| immR       | HTH-type transcriptional regulator ImmR                         |
| mhqD       | Putative hydrolase MhqD   |
| mhqA       | Putative ring-cleaving dioxygenase MhqA                         |
| pgl_1      | 6-phosphogluconolactonase                                       |
| group_347  | hypothetical protein  |
| lgt_1      | Phosphatidylglycerol--prolipoprotein diacylglyceryl transferase |
| group_349  | hypothetical protein  |
| group_350  | hypothetical protein  |
| copB       | Copper-exporting P-type ATPase B                                |
| group_352  | hypothetical protein  |
| actP       | Copper-transporting P-type ATPase                               |
| copY_2     | Transcriptional repressor CopY                                  |
| cadC_1     | Cadmium resistance transcriptional regulatory protein CadC      |
| group_356  | hypothetical protein  |
| group_357  | hypothetical protein  |
| group_358  | hypothetical protein  |
| group_359  | hypothetical protein  |
| group_360  | hypothetical protein  |
| cadC_2     | Cadmium resistance transcriptional regulatory protein CadC      |
| cadA       | putative cadmium-transporting ATPase                            |
| yadH       | Inner membrane transport permease YadH                          |
| yadG       | putative ABC transporter ATP-binding protein YadG               |
| cdr_1      | Coenzyme A disulfide reductase                                  |
| acr3       | Arsenical-resistance protein Acr3                               |
| arsA       | Arsenical pump-driving ATPase                                   |
| group_368  | hypothetical protein  |
| arsD       | Arsenical resistance operon trans-acting repressor ArsD         |
| group_37   | Deoxyguanosinetriphosphate triphosphohydrolase-like protein     |
| group_370  | IS6 family transposase IS1216V                                  |
| group_371  | hypothetical protein  |
| group_372  | hypothetical protein  |
| dnaG_1     | DNA primase   |
| group_374  | hypothetical protein  |
| group_375  | hypothetical protein  |
| group_376  | hypothetical protein  |
| group_377  | hypothetical protein  |
| ltrA       | Group II intron-encoded protein LtrA                            |
| group_379  | hypothetical protein  |
| group_380  | hypothetical protein  |
| group_381  | hypothetical protein  |
| group_382  | hypothetical protein  |
| group_383  | hypothetical protein  |
| group_45   | hypothetical protein  |
| yeeO_3     | putative FMN/FAD exporter YeeO                                  |
| group_82   | hypothetical protein  |
| sdpR_1     | Transcriptional repressor SdpR                                  |

**Appendix 2 Table 3.** Genes lacking in the ST7 isolates of this study, but present in Lineage I ST7 reference strains

| Gene      | Annotation  |
|-----------|---|
| group_310 | Tyrosine recombinase XerC   |
| group_311 | hypothetical protein  |
| group_312 | hypothetical protein  |
| group_313 | hypothetical protein  |
| group_314 | hypothetical protein  |
| group_315 | hypothetical protein  |
| group_316 | hypothetical protein  |
| group_317 | hypothetical protein  |
| group_318 | hypothetical protein  |
| group_319 | hypothetical protein  |
| group_320 | hypothetical protein  |
| group_321 | hypothetical protein  |
| group_22  | hypothetical protein  |
| group_395 | hypothetical protein  |
| group_396 | Adaptive-response sensory-kinase SasA                                 |
| group_397 | Response regulator ArlR   |
| group_398 | hypothetical protein  |
| group_399 | hypothetical protein  |
| yxIF_2    | putative ABC transporter ATP-binding protein YxIF                     |
| lmrA      | Multidrug resistance ABC transporter ATP-binding and permease protein |
| group_402 | hypothetical protein  |
| group_403 | hypothetical protein  |
| group_404 | hypothetical protein  |
| group_405 | hypothetical protein  |
| group_406 | hypothetical protein  |
| group_407 | hypothetical protein  |
| group_408 | hypothetical protein  |
| group_409 | hypothetical protein  |
| Int-Tn    | Transposase from transposon Tn916                                     |
| group_411 | hypothetical protein  |
| group_412 | hypothetical protein  |
| group_413 | hypothetical protein  |
| group_414 | hypothetical protein  |
| tet(M)    | tetracycline resistance ribosomal protection protein Tet(M)           |
| group_416 | hypothetical protein  |
| group_417 | hypothetical protein  |
| group_418 | hypothetical protein  |
| group_419 | hypothetical protein  |
| group_420 | hypothetical protein  |
| group_421 | hypothetical protein  |
| group_422 | hypothetical protein  |
| group_423 | hypothetical protein  |
| group_424 | hypothetical protein  |
| group_425 | hypothetical protein  |
| group_426 | hypothetical protein  |
| pcrA_2    | ATP-dependent DNA helicase PcrA                                       |
| group_428 | hypothetical protein  |
| pezT      | Toxin PezT  |
| group_430 | hypothetical protein  |
| group_431 | DNA primase   |
| degU      | Transcriptional regulatory protein DegU                               |
| group_433 | hypothetical protein  |
| group_434 | hypothetical protein  |
| group_435 | hypothetical protein  |
| bcrA_2    | Bacitracin transport ATP-binding protein BcrA                         |
| lagD      | Lactococcin-G-processing and transport ATP-binding protein LagD       |
| group_438 | hypothetical protein  |
| group_439 | hypothetical protein  |
| group_440 | hypothetical protein  |
| group_441 | hypothetical protein  |
| group_442 | hypothetical protein  |
| group_443 | hypothetical protein  |
| group_444 | 2-methoxy-6-polyprenyl-1,4-benzoquinol methylase, mitochondrial       |
| aadK      | Aminoglycoside 6-adenylyltransferase                                  |
| apt_1     | Adenine phosphoribosyltransferase                                     |
| group_447 | IS1380 family transposase ISSu5                                       |




| Gene      | Annotation              |
|-----------|-------------------------|
| group_448 | hypothetical protein    |
| group_449 | Lantibiotic macedovicin |
| group_450 | hypothetical protein    |
| group_451 | hypothetical protein    |
| group_452 | hypothetical protein    |
| group_453 | hypothetical protein    |
| glgP      | Glycogen phosphorylase  |

**Appendix 2 Table 4.** Genes lacking in the ST7 isolates of this study, but present in Lineage III ST7 reference strains

| Gene      | Annotation                                     |
|-----------|--|
| group_173 | hypothetical protein                           |
| group_127 | hypothetical protein                           |
| group_179 | hypothetical protein                           |
| group_18  | hypothetical protein                           |
| entS      | Enterobactin exporter EntS                     |
| msr(D)    | ABC-F type ribosomal protection protein Msr(D) |
| group_182 | hypothetical protein                           |
| group_183 | hypothetical protein                           |
| dinB_2    | DNA polymerase IV                              |
| group_185 | hypothetical protein                           |
| group_186 | hypothetical protein                           |
| group_187 | hypothetical protein                           |
| group_188 | hypothetical protein                           |
| group_189 | hypothetical protein                           |
| group_190 | hypothetical protein                           |
| group_191 | hypothetical protein                           |
| group_192 | hypothetical protein                           |
| group_193 | hypothetical protein                           |
| group_194 | hypothetical protein                           |
| group_195 | hypothetical protein                           |
| group_196 | hypothetical protein                           |
| group_197 | hypothetical protein                           |
| group_198 | hypothetical protein                           |
| group_199 | hypothetical protein                           |
| group_200 | hypothetical protein                           |
| metK_1    | S-adenosylmethionine synthase                  |
| group_202 | hypothetical protein                           |
| group_203 | hypothetical protein                           |
| group_204 | hypothetical protein                           |
| group_205 | hypothetical protein                           |
| group_206 | hypothetical protein                           |
| group_207 | hypothetical protein                           |
| group_208 | hypothetical protein                           |
| group_209 | hypothetical protein                           |
| group_210 | hypothetical protein                           |
| group_211 | hypothetical protein                           |
| group_212 | hypothetical protein                           |
| group_213 | hypothetical protein                           |
| group_214 | hypothetical protein                           |
| group_215 | hypothetical protein                           |
| clpP_1    | ATP-dependent Clp protease proteolytic subunit |
| group_217 | hypothetical protein                           |
| group_218 | hypothetical protein                           |
| group_219 | hypothetical protein                           |
| group_220 | hypothetical protein                           |
| group_221 | hypothetical protein                           |
| group_222 | hypothetical protein                           |
| group_223 | hypothetical protein                           |
| smc_1     | Chromosome partition protein Smc               |
| group_225 | hypothetical protein                           |
| group_226 | hypothetical protein                           |
| group_227 | hypothetical protein                           |
| group_228 | hypothetical protein                           |
| group_230 | hypothetical protein                           |
| group_231 | hypothetical protein                           |
| group_232 | hypothetical protein                           |
| group_233 | hypothetical protein                           |
| group_234 | hypothetical protein                           |

| Gene      | Annotation   |
|-----------|--|
| group_235 | hypothetical protein                                 |
| aphA      | Aminoglycoside 3'-phosphotransferase                 |
| satA      | Streptothricin acetyltransferase A                   |
| group_238 | Aminoglycoside 6-adenylyltransferase                 |
| group_239 | hypothetical protein                                 |
| group_241 | hypothetical protein                                 |
| group_242 | hypothetical protein                                 |
| group_243 | hypothetical protein                                 |
| group_244 | hypothetical protein                                 |
| group_245 | hypothetical protein                                 |
| group_250 | hypothetical protein                                 |
| group_251 | hypothetical protein                                 |
| group_253 | hypothetical protein                                 |
| group_254 | hypothetical protein                                 |
| group_255 | hypothetical protein                                 |
| group_256 | hypothetical protein                                 |
| group_257 | hypothetical protein                                 |
| group_258 | hypothetical protein                                 |
| noc       | Nucleoid occlusion protein                           |
| group_260 | hypothetical protein                                 |
| group_261 | hypothetical protein                                 |
| group_262 | hypothetical protein                                 |
| group_263 | hypothetical protein                                 |
| group_264 | hypothetical protein                                 |
| group_265 | DNA primase  |
| group_267 | Toxin PezT   |
| group_268 | hypothetical protein                                 |
| group_269 | hypothetical protein                                 |
| group_270 | hypothetical protein                                 |
| group_271 | hypothetical protein                                 |
| group_272 | hypothetical protein                                 |
| nisP      | Nisin leader peptide-processing serine protease NisP |
| regX3     | Sensory transduction protein regX3                   |
| creC      | Sensor protein CreC                                  |
| bcrA_1    | Bacitracin transport ATP-binding protein BcrA        |
| group_277 | hypothetical protein                                 |
| group_278 | hypothetical protein                                 |
| nsuA      | Lantibiotic nisin-U                                  |
| nisB      | Nisin biosynthesis protein NisB                      |
| group_281 | putative ABC transporter ATP-binding protein         |
| nisC      | Nisin biosynthesis protein NisC                      |
| nisI_1    | Nisin immunity protein                               |
| group_284 | hypothetical protein                                 |
| group_289 | hypothetical protein                                 |
| group_292 | hypothetical protein                                 |
| group_293 | hypothetical protein                                 |
| group_294 | hypothetical protein                                 |
| group_295 | PTS system mannose-specific EIID component           |
| sorC_2    | PTS system sorbose-specific EIIC component           |
| sorB_2    | PTS system sorbose-specific EIIB component           |
| group_298 | hypothetical protein                                 |
| group_299 | hypothetical protein                                 |
| xylB      | Xylulose kinase                                      |
| xylA      | Xylose isomerase                                     |
| nagC      | N-acetylglucosamine repressor                        |
| group_303 | ISL3 family transposase ISSsu13                      |
| group_304 | hypothetical protein                                 |
| group_32  | IS110 family transposase ISCARN28                    |
| group_71  | hypothetical protein                                 |
| group_78  | hypothetical protein                                 |
| gmuD_3    | 6-phospho-beta-glucosidase GmuD                      |
| group_15  | hypothetical protein                                 |
| purD_2    | Phosphoribosylamine--glycine ligase                  |
| group_290 | hypothetical protein                                 |
| group_3   | IS110 family transposase ISSsu7                      |
| group_31  | IS110 family transposase ISCARN28                    |
| group_34  | IS110 family transposase ISCARN28                    |
| group_35  | IS110 family transposase ISCARN28                    |
| group_39  | hypothetical protein                                 |

| Gene     | Annotation  |
|----------|---|
| mraY_1   | Phospho-N-acetylmuramoyl-pentapeptide-transferase |
| group_4  | IS110 family transposase ISSsu7                   |
| group_43 | hypothetical protein                              |
| glf_1    | UDP-galactopyranose mutase                        |
| group_61 | hypothetical protein                              |
| group_65 | hypothetical protein                              |
| group_67 | hypothetical protein                              |
| group_7  | IS110 family transposase ISSsu7                   |
| group_96 | IS4 family transposase ISSsu2                     |



**Appendix 2 Figure 1.** e-BURST illustration of the *Streptococcus suis* population was used together with the complete MLST database to cluster the sequence types (STs) into major clonal complexes (CCs). The representation of groups was performed with the double-locus variants (DLVs) parameters. Primary founders (green) are positioned at the center of the cluster and subgroup founders are showed in yellow. The new STs described in our study are highlighted in red.





**Appendix 2 Figure 2.** Results of comparison between the genomes of our ST7 with the genomes of CS100322 and SC070731, two novel ST7 types described by Dong et al. (7) as belonging to lineage III, with SC84 and SC19, two ST7 lineage I strains, and with P1-7 as reference strain. The figure was generated using Phandango, an interactive viewer for bacterial population genomics (<https://jameshadfield.github.io/phandango/#/>) using as input the file generated by Roary (gene\_presence\_absence.csv and accessory\_binary\_genes.fa.newick). Genes are shown as light blue bricks along the top and are sorted left to right by the proportion of isolates they are observed in. Presence (blue) and absence (white) of genes are plotted considering the phylogenetic placement of each isolates.

