

About the Author

Dr. Koh is a senior scientist at the Centre for Animal & Veterinary Sciences within the Animal & Veterinary Service of the National Parks Board, Singapore. Her research interests are focused on veterinary microbiology and animal disease biosurveillance.

References

1. World Organisation for Animal Health (WOAH). African swine fever (infection with African swine fever virus) [cited 2023 Jun 1]. https://www.woah.org/fileadmin/Home/eng/Health_standards/tahm/3.09.01_ASF.pdf
2. International Research Consortium on Animal Health. 2022 African Swine Fever Virus Research Review [cited 2023 Sep 13]. https://www.ars.usda.gov/gara/reports/ASFV-Report_draft_final_31-march-2022.pdf
3. Chapman DAG, Tcherepanov V, Upton C, Dixon LK. Comparison of the genome sequences of non-pathogenic and pathogenic African swine fever virus isolates. *J Gen Virol*. 2008;89:397–408. <https://doi.org/10.1099/vir.0.83343-0>
4. Gavrier-Widén D, Ståhl K, Dixon L. No hasty solutions for African swine fever. *Science*. 2020;367:622–4. <https://doi.org/10.1126/science.aaz8590>
5. Khoo CK, Norlina D, Roshaslinda D, Siti Suraya Hani MS, Zunaida B, Mohd Hasrul AH, et al. African swine fever in backyard pigs of Sabah state, East Malaysia, 2021. *Trop Biomed*. 2021;38:499–504. <https://doi.org/10.47665/tb.38.4.095>
6. Le VP, Jeong DG, Yoon SW, Kwon HM, Trinh TBN, Nguyen TL, et al. Outbreak of African swine fever, Vietnam, 2019. *Emerg Infect Dis*. 2019;25:1433–5. <https://doi.org/10.3201/eid2507.190303>
7. Kwak ML, Chavatte J-M, Chew KL, Lee BPY. Emergence of the zoonotic tick *Dermacentor (Indocentor) auratus* Supino, 1897 (Acari: Ixodidae) in Singapore. *Ticks Tick Borne Dis*. 2021;12:101574. <https://doi.org/10.1016/j.ttbdis.2020.101574>
8. Danecek P, Bonfield, JK, Liddle, J, Marshall, J, Ohan, V, Pollard, MO., et al. Twelve years of SAMtools and BCFtools. *Gigascience*. 2021;10:giab008. PubMed <https://doi.org/10.1093/gigascience/giab008>
9. Bastos ADS, Penrith M-L, Crucière C, Edrich JL, Hutchings G, Roger F, et al. Genotyping field strains of African swine fever virus by partial p72 gene characterisation. *Arch Virol*. 2003;148:693–706. <https://doi.org/10.1007/s00705-002-0946-8>
10. Gallardo C, Fernández-Pinero J, Pelayo V, Gazaev I, Markowska-Daniel I, Pridotkas G, et al. Genetic variation among African swine fever genotype II viruses, eastern and central Europe. *Emerg Infect Dis*. 2014;20:1544–7. <https://doi.org/10.3201/eid2009.140554>

Address for correspondence: Eileen Y. Koh, Animal and Veterinary Science, National Parks Board (NParks), 1 Cluny Rd, Singapore Botanic Gardens, Singapore 259569; email: eileen_koh@nparks.gov.sg

Substantial Diversity in Cocirculating Omicron Lineages in Hospital Setting, Porto Alegre, Brazil

Tiago F. Andreis, Vlademir V. Cantarelli, Marcelo B. da Silva, Mateus S. Helfer, Flávia R. Brust, GAIHN-HAI Team,¹ Alexandre P. Zavascki

Author affiliations: Hospital Moinhos de Vento, Porto Alegre, Brazil (T.F. Andreis, M.S. Helfer, A.P. Zavascki); Universidade Federal de Ciências da Saúde de Porto Alegre, Porto Alegre (V.V. Cantarelli); Unimed Vale dos Sinos, Novo Hamburgo, Brazil (M.B. da Silva); Universidade do Vale do Rio dos Sinos, São Leopoldo, Brazil (M.B. da Silva); Health Security Partners, Washington, DC, USA (F.R. Brust); Universidade Federal do Rio Grande do Sul, Porto Alegre (A.P. Zavascki)

DOI: <http://doi.org/10.3201/eid2912.230880>

We describe substantial variant diversity among 23 detected SARS-CoV-2 Omicron lineage viruses cocirculating among healthcare workers and inpatients (272 sequenced samples) from Porto Alegre, Brazil, during November 2022–January 2023. BQ.1 and related lineages (61.4%) were most common, followed by BE.9 (19.1%), first described in November 2022 in the Amazon region.

When SARS-CoV-2 variants of concern were first described, the epidemiologic situation was characterized by sequential waves of Alpha, Beta, Gamma, and Delta variants, with relatively few other variants cocirculating with the dominant variant of concern of each wave (1). The epidemiologic situation shifted with the emergence of the Omicron variant (B.1.1.529) in November 2021 (2). Distinct Omicron lineages rapidly emerged, causing successive, relatively narrow waves of infection associated with novel lineages that had pronounced immune escape and increased transmissibility (3).

The Global Action in Healthcare Network–Healthcare-associated Infection (GAIHN-HAI) module is a multinational network of healthcare facilities and laboratories developed by the Division of Healthcare Quality Promotion, National Center for Emerging and Zoonotic Infectious Diseases, US Centers for Disease Control and Prevention (Atlanta, GA, USA), to address emerging infectious disease threats in healthcare settings. The network began genomic surveillance of SARS-CoV-2 lineages affecting healthcare

¹Team members are listed at the end of this article.

workers (HCWs) and inpatients in Brazil in November 2022. We present initial findings from a tertiary-care COVID-19 reference hospital in Porto Alegre, the capital of the southernmost state of Brazil.

We conducted this surveillance study based on data from Hospital Moinhos de Vento, the first facility to join the Brazil GAIHN-HAI network. We obtained demographics and exposure risk factors by participant interview and invited into the study all HCWs and inpatients ≥ 18 years of age with COVID-19 diagnosed by real-time reverse transcription PCR that had a cycle threshold < 30 in any probe. We performed whole-genome sequencing (Appendix, <https://wwwnc.cdc.gov/EID/article/29/12/23-0880-App1.pdf>) and submitted all viral genome sequences to GISAID (<http://www.gisaid.org>) (Appendix Table 1) (4).

During November 2022–January 2023, we collected 552 deduplicated SARS-CoV-2 real-time reverse transcription PCR–positive specimens (360 [65.2%] HCWs, 192 [34.8%] inpatients). Of the 552 specimens, we excluded 124 (22.5%) because cycle threshold was > 30 and 156 (28.3%) because we were unable to obtain consent for, resulting in 272 (49.3%) samples sequenced and available for analysis. Analyzed samples consisted of 182 (66.9%) samples from HCWs and 90 (33.1%) from inpatients.

We identified 23 distinct lineages, all belonging to the Omicron variant (Table; Appendix Figure). BQ.1 and related lineages were most prevalent (61.4%), followed by BE.9 (19.1%) and others (19.5%). We detected BE.9 first, in epidemiologic

week 45 of 2022, and that lineage remained cocirculating with BQ.1 in a subdominant proportion of cases throughout the study period (Appendix Figure). We also noted genetic relatedness of other Omicron lineages (Figure).

We noted no difference in the distribution of lineages between HCWs and inpatients. Compared with other lineages, BE.9 was more common in female patients ($p = 0.027$), younger patients ($p = 0.017$), and patients reporting previous contact with a person infected with SARS-CoV-2 ($p \leq 0.001$). We noted vaccination status and report of a previous infection were similar among participants, regardless of lineage (Appendix Table 2).

Our findings from a select population of HCWs and inpatients from a hospital in southern Brazil revealed the cocirculation of 23 Omicron lineages over a relatively short (11-week) period. Although we observed BQ.1-related lineages most frequently, consistent with this lineage's recent predominance both globally (5) and in Brazil (3), the number of subvariants we observed represents a departure from the observed serial dominance common with earlier variants of concern (1,6). Generalizability of our findings is limited, but our observations are consistent with other global findings suggesting that Omicron has diversified to include multiple lineages, with adequate fitness allowing them to cocirculate among humans (2,3,7,8). One proposed explanation is that population-level variation in vaccination and previous infection has led to heterogeneous

Table. Identification of Omicron SARS-CoV-2 lineages among healthcare workers and inpatients in Porto Alegre, southern Brazil, November 2022–January 2023

Group	Lineage	No. cases	% Cases	Cumulative %
BQ.1, n = 167 (61.4%)	BQ.1.1	140	51.5	51.5
	BQ.1.1.18	14	5.1	56.6
	BQ.1.3	3	1.1	57.7
	BQ.1	3	1.1	58.8
	BQ.1.1.23	2	0.7	59.6
	BQ.1.1.15	1	0.4	59.9
	BQ.1.1.17	1	0.4	60.3
	BQ.1.1.22	1	0.4	60.7
	BQ.1.1.24	1	0.4	61.0
	BQ.1.1.4	1	0.4	61.4
BE.9, n = 52 (19.1%)	BE.9	52	19.1	80.5
BA.5, n = 15 (5.5%)	BA.5.3.1	6	2.2	82.7
	BA.5	4	1.5	84.2
	BA.5.2.1	3	1.1	85.3
	BA.5.1.27	2	0.7	86.0
Other lineages, n = 38 (14%)	BE.10	14	5.1	91.2
	CK.1	11	4.0	95.2
	DL.1	3	1.1	96.3
	XBB.1	3	1.1	97.4
	XBB.1.5	1	0.4	97.8
	BA.4.6	3	1.1	98.9
	BN.1.3.1	2	0.7	99.6
	BN.1.5	1	0.4	100.0
Total		272	100.0	100.0



Figure. Phylogenetic radial tree showing placement of 272 SARS-CoV-2 sequenced samples from healthcare workers and inpatients in Porto Alegre, southern Brazil, November 2022–January 2023. Data as of February 25, 2023.

immunologic background, leading to cocirculation of distinct lineages with varying proficiency for natural and induced immunogenic escape (3,7). However, recent research suggests that immune imprinting induces Omicron receptor-binding protein mutation convergence (9).

One unexpected finding of this study was the relatively high proportion of BE.9 lineages we ob-

served. BE.9 was first described in November 2022 in the northern Brazil state of Amazonas (7) and is characterized by a large, 244-nt deletion in the open reading frame (ORF) 7a gene at the position 27508–27751 and by mutations at spike:K444T, spike:N460K, spike:Y144del, ORF1a:V84del, and ORF1a:M85del (<https://github.com/cov-lineages/pango-designation/issues/1302>). Since its initial

description in Brazil, BE.9 has been reported in other countries, but only a small portion of those identified sequences have been submitted to GISAID (4). As for BE.9, most BE.10 cases ($\approx 75\%$) reported in GISAID are from Brazil (4). In contrast to other countries, where the recombinant XBB.1.5 has been circulating since August 2022 (10), we only detected XBB.1.5 in the study population starting January 19, 2023.

In conclusion, SARS-CoV-2 genomic surveillance at a hospital in southern Brazil found substantial diversity of Omicron lineages among HCWs and inpatients. Findings are specific to this facility and not generalizable to other hospitals or the population of Brazil. As countries globally adapt their national SARS-CoV-2 testing strategies to current COVID-19 epidemiology, they should consider focusing SARS-CoV-2 genomic surveillance strategies, along with infection trend monitoring on smaller, targeted populations such as HCWs and inpatients, to identify unusual epidemiologic events, characterize unusual viral transmission chains, and guide facility-level response measures.

The US Centers for Disease Control and Prevention GAIHN-HAI team comprises Matthew Westercamp, Valery Tashayev, Morgane Donadel, Reed Magleby, Emily Petersen, and Garrett Mahon.

Acknowledgments

We are grateful to the staff of the Laboratório de Genética e Biologia Molecular do Hospital Moinhos de Vento for the technical help.

This study was approved by the US CDC's human subjects' review committee, the ethics review committee at Hospital Moinhos de Vento, and the Brazilian Research Ethics Commission (Comissão Nacional de Ética em Pesquisa).

This work was funded by the US Centers for Disease Control and Prevention through a cooperative agreement with Health Security Partners. A.P.Z. is a research fellow of the National Council for Scientific and Technological Development, Ministry of Science and Technology, Brazil (308049/2022-5).

About the Author

Dr. Andreis is a molecular biologist at Laboratório de Genética e Biologia Molecular do Hospital Moinhos de Vento. His primary research interests are precision medicine, genomics, and SARS-CoV-2 surveillance.

References

- Li J, Lai S, Gao GF, Shi W. The emergence, genomic diversity and global spread of SARS-CoV-2. *Nature*. 2021;600:408–18. <https://doi.org/10.1038/s41586-021-04188-6>
- Viana R, Moyo S, Amoako DG, Tegally H, Scheepers C, Althaus CL, et al. Rapid epidemic expansion of the SARS-CoV-2 Omicron variant in southern Africa. *Nature*. 2022;603:679–86. <https://doi.org/10.1038/s41586-022-04411-y>
- Gangavarapu K, Latif AA, Mullen JL, Alkuzweny M, Hufbauer E, Tsueng G, et al.; GISAID Core and Curation Team. Outbreak.info genomic reports: scalable and dynamic surveillance of SARS-CoV-2 variants and mutations. *Nat Methods*. 2023;20:512–22. <https://doi.org/10.1038/s41592-023-01769-3>
- Elbe S, Buckland-Merrett G. Data, disease and diplomacy: GISAID's innovative contribution to global health. *Glob Chall*. 2017;1:33–46. <https://doi.org/10.1002/gch2.1018>
- World Health Organization. TAG-VE Statement on Omicron sublineages BQ.1 and XBB. 2022 [cited 2023 Apr 28]. <https://www.who.int/news/item/27-10-2022-tag-ve-statement-on-omicron-sublineages-bq.1-and-xbb>
- Giovanetti M, Slavov SN, Fonseca V, Wilkinson E, Tegally H, Patané JSL, et al. Genomic epidemiology of the SARS-CoV-2 epidemic in Brazil. *Nat Microbiol*. 2022;7:1490–500. <https://doi.org/10.1038/s41564-022-01191-z>
- Jung C, Kmiec D, Koepke L, Zech F, Jacob T, Sparrer KMJ, et al. Omicron: what makes the latest SARS-CoV-2 variant of concern so concerning? *J Virol*. 2022;96:e0207721. <https://doi.org/10.1128/jvi.02077-21>
- Luna N, Muñoz M, Ramírez AL, Patiño LH, Castañeda SA, Ballesteros N, et al. Genomic diversity of SARS-CoV-2 Omicron variant in South American countries. *Viruses*. 2022;14:1234. <https://doi.org/10.3390/v14061234>
- Cao Y, Jian F, Wang J, Yu Y, Song W, Yisimayi A, et al. Imprinted SARS-CoV-2 humoral immunity induces convergent Omicron RBD evolution. *Nature*. 2022;614:521–9. <https://doi.org/10.1038/s41586-022-05644-7>
- Wang Q, Iketani S, Li Z, Liu L, Guo Y, Huang Y, et al. Alarming antibody evasion properties of rising SARS-CoV-2 BQ and XBB subvariants. *Cell*. 2023;186:279–286.e8. <https://doi.org/10.1016/j.cell.2022.12.018>

Address for correspondence: Morgane Donadel, Centers for Disease Control and Prevention, 1600 Clifton Rd NE, Mailstop HI16-2, Atlanta, GA 30329-4027, USA; email: mxi4@cdc.gov

EID cannot ensure accessibility for supplementary materials supplied by authors. Readers who have difficulty accessing supplementary content should contact the authors for assistance.

Substantial Diversity in Cocirculating Omicron Lineages in Hospital Setting, Porto Alegre, Brazil

Appendix

Methods used for molecular testing and whole-genome sequencing and ethical considerations

Real-time RT-PCR

Real-time RT-PCRs were performed by using the commercial kits TaqMan 2019-nCoV Assay Kit v1 (Thermo Fisher Scientific, <https://www.thermofisher.com/us>), TaqCheck SARS-CoV-2 Fast PCR Assay (Thermo Fisher Scientific) or Xpert Xpress SARS-CoV-2 (Cepheid, <https://www.cepheid.com/en-US>).

Whole-Genome Sequencing

The viral genomic library was prepared by using the CleanPlex SARS-CoV-2 FLEX Kit and the SARS-CoV-2 Emerging Variants Panel Add-on v2 (Paragon Genomics, Inc, www.paragongenomics.com) following the manufacturer's instructions. The resulting libraries were sequenced on the Illumina MiSeq platform by using a V2 chemistry (Illumina, <https://www.illumina.com>).

Ethical approvals

The study was approved by the Brazilian National Health Council - Ministry of Health (CAAE: 59038722.0.1001.5330) and informed consent was obtained from all participants.

Appendix Table 1. GISAID accession number of all viral genome sequences from a study of healthcare workers and inpatients at Hospital Moinhos de Vento in Porto Alegre, southern Brazil, November 2022–January 2023

Lineage	GISAID accession number
BQ.1.1	EPI_ISL_15803844, EPI_ISL_15803846, EPI_ISL_16002026, EPI_ISL_15803851, EPI_ISL_15803853, EPI_ISL_16002028, EPI_ISL_16002021, EPI_ISL_16002008, EPI_ISL_16002025, EPI_ISL_16002022, EPI_ISL_16002017, EPI_ISL_16002027, EPI_ISL_16002019, EPI_ISL_16002014, EPI_ISL_16002012, EPI_ISL_16360410, EPI_ISL_16360411, EPI_ISL_16002015, EPI_ISL_16002024, EPI_ISL_16360412, EPI_ISL_16360413, EPI_ISL_16360415, EPI_ISL_16360416, EPI_ISL_16360418, EPI_ISL_16360419, EPI_ISL_16360420, EPI_ISL_16360421, EPI_ISL_16360422, EPI_ISL_16360423, EPI_ISL_16360425, EPI_ISL_16360429, EPI_ISL_16360383, EPI_ISL_16360430, EPI_ISL_16360432, EPI_ISL_16360434, EPI_ISL_16360435, EPI_ISL_16360436, EPI_ISL_16360395, EPI_ISL_16360428, EPI_ISL_16360399, EPI_ISL_17651884, EPI_ISL_16360437, EPI_ISL_16360438, EPI_ISL_16360439, EPI_ISL_16360440, EPI_ISL_17848718, EPI_ISL_16360403, EPI_ISL_16360385, EPI_ISL_16423445, EPI_ISL_16360404, EPI_ISL_16360386, EPI_ISL_16360387, EPI_ISL_16360389, EPI_ISL_16360405, EPI_ISL_16360408, EPI_ISL_16360393, EPI_ISL_16360394, EPI_ISL_16360409, EPI_ISL_17848770, EPI_ISL_17848775, EPI_ISL_17848758, EPI_ISL_16360398, EPI_ISL_17651886, EPI_ISL_17651885, EPI_ISL_17848790, EPI_ISL_17651894, EPI_ISL_16360362, EPI_ISL_16360366, EPI_ISL_17651887, EPI_ISL_16360363, EPI_ISL_17848761, EPI_ISL_17651889, EPI_ISL_17848762, EPI_ISL_16360365, EPI_ISL_17848722, EPI_ISL_17848724, EPI_ISL_17848765, EPI_ISL_17848774, EPI_ISL_17848764, EPI_ISL_17651891, EPI_ISL_17848768, EPI_ISL_16360373, EPI_ISL_16360369, EPI_ISL_16360370, EPI_ISL_16360372, EPI_ISL_17848793, EPI_ISL_17848769, EPI_ISL_17651892, EPI_ISL_16360374, EPI_ISL_17848776, EPI_ISL_16360371, EPI_ISL_17848794, EPI_ISL_16360375, EPI_ISL_17848771, EPI_ISL_17848772, EPI_ISL_17848777, EPI_ISL_17848773, EPI_ISL_16360377, EPI_ISL_16360381, EPI_ISL_16360376, EPI_ISL_17848778, EPI_ISL_16360380, EPI_ISL_17848779, EPI_ISL_17651893, EPI_ISL_16360382, EPI_ISL_17848780, EPI_ISL_17848791, EPI_ISL_17848785, EPI_ISL_17848786, EPI_ISL_17848792, EPI_ISL_17848726, EPI_ISL_17848730, EPI_ISL_18103841, EPI_ISL_17848729, EPI_ISL_17848727, EPI_ISL_17848789, EPI_ISL_17848731, EPI_ISL_17848736, EPI_ISL_17848734, EPI_ISL_18103818, EPI_ISL_17848746, EPI_ISL_17848737, EPI_ISL_17848742, EPI_ISL_17848741, EPI_ISL_17848744, EPI_ISL_17848738, EPI_ISL_17848757, EPI_ISL_17848739, EPI_ISL_17848740, EPI_ISL_17848748, EPI_ISL_17848751, EPI_ISL_17848752, EPI_ISL_17848754, EPI_ISL_17848800, EPI_ISL_17848753, EPI_ISL_17848803, EPI_ISL_17848759, EPI_ISL_17848797, EPI_ISL_17848805, EPI_ISL_17848798
BQ.1.1.18	EPI_ISL_15803847, EPI_ISL_16360414, EPI_ISL_16360396, EPI_ISL_16360401, EPI_ISL_16360388, EPI_ISL_16360407, EPI_ISL_16360391, EPI_ISL_17651890, EPI_ISL_17848743, EPI_ISL_17848766, EPI_ISL_17848781, EPI_ISL_16360379, EPI_ISL_17848787, EPI_ISL_17848750
BQ.1.3	EPI_ISL_16360431, EPI_ISL_16360402, EPI_ISL_16360397
BQ.1	EPI_ISL_17848799, EPI_ISL_17848795, EPI_ISL_17848733
BQ.1.1.23	EPI_ISL_17848721, EPI_ISL_16360426
BQ.1.1.15	EPI_ISL_17848728
BQ.1.1.17	EPI_ISL_16360384
BQ.1.1.22	EPI_ISL_17848782
BQ.1.1.24	EPI_ISL_17848725
BQ.1.1.4	EPI_ISL_17848755
BE.9	EPI_ISL_16014612, EPI_ISL_16423443, EPI_ISL_16423450, EPI_ISL_16014613, EPI_ISL_16423451, EPI_ISL_16423452, EPI_ISL_16423453, EPI_ISL_16423454, EPI_ISL_16423455, EPI_ISL_16423456, EPI_ISL_16423457, EPI_ISL_16423458, EPI_ISL_16423444, EPI_ISL_18050262, EPI_ISL_16423446, EPI_ISL_16423448, EPI_ISL_16423449, EPI_ISL_18103824, EPI_ISL_18103822, EPI_ISL_18103821, EPI_ISL_18103823, EPI_ISL_16423435, EPI_ISL_18103825, EPI_ISL_18103826, EPI_ISL_16423436, EPI_ISL_16423438, EPI_ISL_16423439, EPI_ISL_18103827, EPI_ISL_16423437, EPI_ISL_18103828, EPI_ISL_18103833, EPI_ISL_16423440, EPI_ISL_16423441, EPI_ISL_18050264, EPI_ISL_18103829, EPI_ISL_18103831, EPI_ISL_18103835, EPI_ISL_18103830, EPI_ISL_18103832, EPI_ISL_18103836, EPI_ISL_18103837, EPI_ISL_18103834, EPI_ISL_18103816, EPI_ISL_18103815, EPI_ISL_18103817, EPI_ISL_18103820, EPI_ISL_18103819, EPI_ISL_18103842, EPI_ISL_18103843, EPI_ISL_18103838, EPI_ISL_18103839, EPI_ISL_18103840,
BA.5.3.1	EPI_ISL_16360433, EPI_ISL_17848723, EPI_ISL_16360406, EPI_ISL_16423434, EPI_ISL_16360367, EPI_ISL_16423442
BA.5	EPI_ISL_16014611, EPI_ISL_16360424, EPI_ISL_17651888, EPI_ISL_17848760,
BA.5.2.1	EPI_ISL_15803849, EPI_ISL_16002023, EPI_ISL_16002009
BA.5.1.27	EPI_ISL_16423447, EPI_ISL_17848719
BE.10	EPI_ISL_15803842, EPI_ISL_15803843, EPI_ISL_15803845, EPI_ISL_15803848, EPI_ISL_15803852, EPI_ISL_16002010, EPI_ISL_16002011, EPI_ISL_16360390,

Lineage	GISAID accession number
	EPI_ISL_16360392, EPI_ISL_17848720, EPI_ISL_17848767, EPI_ISL_17848783, EPI_ISL_17848802, EPI_ISL_17848732
CK.1	EPI_ISL_15803850, EPI_ISL_16360400, EPI_ISL_16360364, EPI_ISL_18050263, EPI_ISL_16360378, EPI_ISL_17848784, EPI_ISL_17848749, EPI_ISL_17848745, EPI_ISL_17848735, EPI_ISL_17848801, EPI_ISL_17848796,
DL.1	EPI_ISL_16002018, EPI_ISL_16002016, EPI_ISL_16360368
XBB.1	EPI_ISL_16360427, EPI_ISL_17848763, EPI_ISL_17848788
XBB.1.5	EPI_ISL_16706637
BA.4.6	EPI_ISL_16002013, EPI_ISL_16002020, EPI_ISL_17848756
BN.1.3.1	EPI_ISL_16360417, EPI_ISL_17848759
BN.1.5	EPI_ISL_17848747

Appendix Table 2. Characteristics of healthcare workers and inpatients, Hospital Moinhos de Vento, Brazil, November 2022–January 2023


Characteristic	Total N (%)	BQ.1	BE.9	BA.5	Other lineages	p-value
Total	272 (100)	167 (61.4)	52 (19.1)	15 (5.5)	38 (14.0)	na
Inpatients	90 (33.1)	63 (37.7)	11 (21.2)	4 (26.7)	12 (31.6)	0.152
Healthcare workers	182 (66.9)	104 (62.3)	41 (78.8)	11 (73.3)	26 (68.4)	na
Female	195 (71.7)	113 (67.7)	46 (88.5)	11 (73.3)	25 (65.8)	0.027
Median age, y (IQR)	40.4 (31.0–64.4)	42.6 (32.0–72.2)	34.4 (27.5–43.4)	42.9 (34.0–54.9)	40.8 (28.8–62.4)	0.017*
Previous infection (Yes)	140 (51.5)	85 (50.9)	25 (48.1)	9 (60.0)	21 (55.3)	0.821
Symptoms Description†						0.193
Asymptomatic	17 (6.3)	15 (9.0)	1 (1.9)	0	1 (2.6)	na
Mild illness	222 (81.6)	126 (75.4)	48 (92.3)	14 (93.3)	34 (89.5)	na
Moderate to Severe Illness	32 (11.8)	25 (15.0)	3 (5.8)	1 (6.7)	3 (7.9)	na
Unknown	1 (0.4)	1 (0.6)	0	0	0	na
Vaccination						0.097
2 Doses	18 (6.6)	12 (7.2)	4 (7.7)	0	2 (5.3)	na
3 Doses	76 (27.9)	51 (30.5)	12 (23.1)	2 (13.3)	11 (28.9)	na
4 Doses	176 (64.7)	104 (62.3)	36 (69.2)	12 (80.0)	24 (63.2)	na
Not vaccinated	2 (0.7)	0	0	1 (6.7)	1 (2.6)	na
COVID-19 hospitalization	36 (13.2)	25 (15.0)	3 (5.8)	2 (13.3)	6 (15.8)	0.365
Link to known COVID-19 cases	78 (28.7)	39 (23.4)	28 (53.8)	5 (33.3)	6 (15.8)	<0.001

BQ.1.1, BQ.1.1.4, BQ.1.1.15, BQ.1.1.17, BQ.1.1.18, BQ.1.1.23, BQ.1.1.24 and BQ.1.3 are aggregated with BQ.1. BA.5.1.27, BA.5.2.1, BA.5.3.1 are aggregated with BA.5. Other lineages include BE.10, CK.1, DL.1, XBB.1, XBB.1.5, BA.4.6, BN.1.3.1, BN.1.5.

*Kruskal-Wallis test was used for continuous variables, the results are presented as median with interquartile ranges [IQR]. p ≤ 0.050 was considered statistically significant.

†Classification According to Clinical Spectrum of SARS-CoV-2 Infection - National Institutes of Health. Chi-square test with adjusted residual analysis was used for categorical variables.

na, not applicable.



Appendix Figure. Distribution of Omicron SARS-CoV-2 lineages among healthcare workers and inpatients at Hospital Moinhos de Vento in Porto Alegre, southern Brazil, November 2022–January 2023 (n = 272 specimens).