

Table 15. Genotyped Tuberculosis Cases with *Mycobacterium bovis*¹ by Origin of Birth: United States, 2004–2016

Year	Total genotyped cases	<i>Mycobacterium bovis</i> cases					
		Total		U.S.-born		Non-U.S.-born	
	No.	No. ²	(%)	No.	(%) ³	No.	(%) ³
2004	5,954	73	(1.2)	23	(31.5)	50	(68.5)
2005	7,500	80	(1.1)	22	(27.5)	58	(72.5)
2006	7,529	116	(1.5)	25	(21.6)	90	(77.6)
2007	8,428	113	(1.3)	17	(15.0)	95	(84.1)
2008	8,178	129	(1.6)	29	(22.5)	100	(77.5)
2009	7,718	113	(1.5)	27	(23.9)	86	(76.1)
2010	7,748	108	(1.4)	20	(18.5)	88	(81.5)
2011	7,620	117	(1.5)	30	(25.6)	87	(74.4)
2012	7,236	109	(1.5)	18	(16.5)	91	(83.5)
2013	7,063	96	(1.4)	22	(22.9)	74	(77.1)
2014	6,991	109	(1.6)	24	(22.0)	85	(78.0)
2015	7,194	125	(1.7)	30	(24.0)	95	(76.0)
2016	6,903	106	(1.5)	17	(16.0)	89	(84.0)

¹*M. bovis* cases were defined predominantly by spoligotyping results with missing spacers 3, 9, 16, and 39–43. Data exclude 118 cases of Bacillus Calmette-Guérin (BCG) *M. bovis*, which have x, y or z in the second MIRU position.

²This column reports all genotyped *M. bovis* cases, including those where origin of birth is unknown.

³Denominator is all *M. bovis* cases.