

Table 38. Extrapulmonary Tuberculosis Cases and Percentages by Site of Disease: Reporting Areas, 2016

Reporting area	Total extrapulm. cases ¹	Total extrapulm. sites ²	Site of disease															
			Pleural		Lymphatic		Bone or joint		Genitourinary		Meningeal		Peritoneal		Laryngeal		Other	
			No.	(%)	No.	(%)	No.	(%)	No.	(%)	No.	(%)	No.	(%)	No.	(%)	No.	(%)
United States	1,882	1,983	320	(16.1)	710	(35.8)	192	(9.7)	91	(4.6)	84	(4.2)	121	(6.1)	2	(0.1)	463	(23.3)
Alabama	11	11	2	(18.2)	2	(18.2)	1	(9.1)	0	(0)	0	(0)	1	(9.1)	1	(9.1)	4	(36.4)
Alaska	9	9	4	(44.4)	0	(0)	0	(0)	1	(11.1)	0	(0)	2	(22.2)	0	(0)	2	(22.2)
Arizona	35	35	9	(25.7)	11	(31.4)	1	(2.9)	5	(14.3)	0	(0)	2	(5.7)	0	(0)	7	(20.0)
Arkansas	17	17	4	(23.5)	7	(41.2)	2	(11.8)	0	(0)	0	(0)	0	(0)	1	(5.9)	3	(17.6)
California	359	376	64	(17.0)	127	(33.8)	32	(8.5)	17	(4.5)	13	(3.5)	26	(6.9)	0	(0)	97	(25.8)
Colorado	22	25	3	(12.0)	9	(36.0)	1	(4.0)	2	(8.0)	0	(0)	1	(4.0)	0	(0)	9	(36.0)
Connecticut	16	18	3	(16.7)	6	(33.3)	1	(5.6)	1	(5.6)	1	(5.6)	2	(11.1)	0	(0)	4	(22.2)
Delaware	3	3	0	(0)	1	(33.3)	0	(0)	1	(33.3)	1	(33.3)	0	(0)	0	(0)	0	(0)
District of Columbia	9	9	2	(22.2)	4	(44.4)	1	(11.1)	0	(0)	0	(0)	0	(0)	0	(0)	2	(22.2)
Florida	90	93	15	(16.1)	23	(24.7)	11	(11.8)	3	(3.2)	6	(6.5)	4	(4.3)	0	(0)	31	(33.3)
Georgia	60	62	13	(21.0)	19	(30.6)	9	(14.5)	2	(3.2)	0	(0)	5	(8.1)	0	(0)	14	(22.6)
Hawaii	14	15	4	(26.7)	5	(33.3)	0	(0)	2	(13.3)	2	(13.3)	1	(6.7)	0	(0)	1	(6.7)
Idaho	1	1	1	(100.0)	0	(0)	0	(0)	0	(0)	0	(0)	0	(0)	0	(0)	0	(0)
Illinois	84	89	19	(21.3)	36	(40.4)	8	(9.0)	2	(2.2)	5	(5.6)	3	(3.4)	0	(0)	16	(18.0)
Indiana	15	16	2	(12.5)	7	(43.8)	2	(12.5)	0	(0)	1	(6.3)	1	(6.3)	0	(0)	3	(18.8)
Iowa	13	16	2	(12.5)	5	(31.3)	2	(12.5)	0	(0)	1	(6.3)	1	(6.3)	0	(0)	5	(31.3)
Kansas	6	6	1	(16.7)	3	(50.0)	1	(16.7)	0	(0)	0	(0)	1	(16.7)	0	(0)	0	(0)
Kentucky	15	15	0	(0)	5	(33.3)	1	(6.7)	2	(13.3)	0	(0)	1	(6.7)	0	(0)	6	(40.0)
Louisiana	18	20	4	(20.0)	6	(30.0)	2	(10.0)	1	(5.0)	1	(5.0)	0	(0)	0	(0)	6	(30.0)
Maine	4	4	0	(0)	3	(75.0)	0	(0)	0	(0)	0	(0)	0	(0)	0	(0)	1	(25.0)
Maryland	77	82	15	(18.3)	34	(41.5)	8	(9.8)	3	(3.7)	1	(1.2)	6	(7.3)	0	(0)	15	(18.3)
Massachusetts	39	41	4	(9.8)	21	(51.2)	2	(4.9)	2	(4.9)	1	(2.4)	2	(4.9)	0	(0)	9	(22.0)
Michigan	53	56	6	(10.7)	16	(28.6)	3	(5.4)	1	(1.8)	1	(1.8)	1	(1.8)	0	(0)	28	(50.0)
Minnesota	55	60	4	(6.7)	35	(58.3)	8	(13.3)	2	(3.3)	1	(1.7)	2	(3.3)	0	(0)	8	(13.3)
Mississippi	13	14	4	(28.6)	4	(28.6)	1	(7.1)	0	(0)	2	(14.3)	1	(7.1)	0	(0)	2	(14.3)
Missouri	16	16	3	(18.8)	6	(37.5)	0	(0)	0	(0)	1	(6.3)	0	(0)	0	(0)	6	(37.5)
Nebraska	8	8	1	(12.5)	2	(25.0)	2	(25.0)	0	(0)	0	(0)	0	(0)	0	(0)	3	(37.5)
Nevada	16	18	3	(16.7)	4	(22.2)	2	(11.1)	2	(11.1)	2	(11.1)	1	(5.6)	0	(0)	4	(22.2)
New Hampshire	2	3	1	(33.3)	0	(0)	0	(0)	0	(0)	1	(33.3)	0	(0)	0	(0)	1	(33.3)
New Jersey	66	71	18	(25.4)	28	(39.4)	4	(5.6)	5	(7.0)	2	(2.8)	5	(7.0)	0	(0)	9	(12.7)
New Mexico	10	11	2	(18.2)	2	(18.2)	2	(18.2)	2	(18.2)	0	(0)	0	(0)	0	(0)	3	(27.3)

Table 38. (Con't) Extrapulmonary Tuberculosis Cases and Percentages by Site of Disease: Reporting Areas, 2016

Reporting area	Total extrapulm. cases ¹	Total extrapulm. sites ²	Site of disease															
			Pleural		Lymphatic		Bone or Joint		Genitourinary		Meningeal		Peritoneal		Laryngeal		Other	
			No.	(%)	No.	(%)	No.	(%)	No.	(%)	No.	(%)	No.	(%)	No.	(%)	No.	(%)
New York State ³	62	65	9	(13.8)	24	(36.9)	6	(9.2)	4	(6.2)	4	(6.2)	7	(10.8)	0	(0)	11	(16.9)
New York City	118	132	14	(10.6)	46	(34.8)	25	(18.9)	7	(5.3)	6	(4.5)	10	(7.6)	0	(0)	24	(18.2)
North Carolina	45	48	8	(16.7)	17	(35.4)	6	(12.5)	1	(2.1)	3	(6.3)	2	(4.2)	0	(0)	11	(22.9)
North Dakota	7	8	1	(12.5)	5	(62.5)	1	(12.5)	0	(0)	0	(0)	0	(0)	0	(0)	1	(12.5)
Ohio	39	39	4	(10.3)	15	(38.5)	7	(17.9)	2	(5.1)	2	(5.1)	3	(7.7)	0	(0)	6	(15.4)
Oklahoma	18	18	8	(44.4)	4	(22.2)	0	(0)	0	(0)	0	(0)	1	(5.6)	0	(0)	5	(27.8)
Oregon	22	22	3	(13.6)	10	(45.5)	1	(4.5)	0	(0)	0	(0)	2	(9.1)	0	(0)	6	(27.3)
Pennsylvania	45	46	4	(8.7)	14	(30.4)	5	(10.9)	4	(8.7)	2	(4.3)	7	(15.2)	0	(0)	10	(21.7)
Rhode Island	6	6	2	(33.3)	1	(16.7)	2	(33.3)	0	(0)	0	(0)	0	(0)	0	(0)	1	(16.7)
South Carolina	27	28	2	(7.1)	13	(46.4)	3	(10.7)	0	(0)	1	(3.6)	2	(7.1)	0	(0)	7	(25.0)
South Dakota	1	1	0	(0)	0	(0)	0	(0)	0	(0)	1	(100.0)	0	(0)	0	(0)	0	(0)
Tennessee	18	21	3	(14.3)	9	(42.9)	0	(0)	2	(9.5)	2	(9.5)	1	(4.8)	0	(0)	4	(19.0)
Texas	191	196	29	(14.8)	65	(33.2)	19	(9.7)	7	(3.6)	17	(8.7)	9	(4.6)	0	(0)	50	(25.5)
Utah	5	6	2	(33.3)	2	(33.3)	0	(0)	0	(0)	0	(0)	0	(0)	0	(0)	2	(33.3)
Vermont	1	1	0	(0)	1	(100.0)	0	(0)	0	(0)	0	(0)	0	(0)	0	(0)	0	(0)
Virginia	50	51	5	(9.8)	23	(45.1)	4	(7.8)	2	(3.9)	1	(2.0)	5	(9.8)	0	(0)	11	(21.6)
Washington	54	58	9	(15.5)	25	(43.1)	4	(6.9)	4	(6.9)	2	(3.4)	3	(5.2)	0	(0)	11	(19.0)
West Virginia	6	6	4	(66.7)	1	(16.7)	1	(16.7)	0	(0)	0	(0)	0	(0)	0	(0)	0	(0)
Wisconsin	11	11	0	(0)	4	(36.4)	1	(9.1)	2	(18.2)	0	(0)	0	(0)	0	(0)	4	(36.4)
Wyoming	0	0	0	...	0	...	0	...	0	...	0	...	0	...	0	...	0	...
American Samoa ⁴	0	0	0	...	0	...	0	...	0	...	0	...	0	...	0	...	0	...
Fed. States of Micronesia ⁴	7	7	2	(28.6)	3	(42.9)	1	(14.3)	0	(0)	0	(0)	1	(14.3)	0	(0)	0	(0)
Guam ⁴	3	3	0	(0)	1	(33.3)	0	(0)	0	(0)	0	(0)	0	(0)	0	(0)	2	(66.7)
Marshall Islands ⁴	40	40	23	(57.5)	7	(17.5)	1	(2.5)	1	(2.5)	0	(0)	7	(17.5)	0	(0)	1	(2.5)
N. Mariana Islands ⁴	0	0	0	...	0	...	0	...	0	...	0	...	0	...	0	...	0	...
Puerto Rico ⁴	0	0	0	...	0	...	0	...	0	...	0	...	0	...	0	...	0	...
Republic of Palau ⁴	2	2	1	(50.0)	1	(50.0)	0	(0)	0	(0)	0	(0)	0	(0)	0	(0)	0	(0)
U.S. Virgin Islands ⁴	0	0	0	...	0	...	0	...	0	...	0	...	0	...	0	...	0	...

¹Excludes cases with pulmonary site of disease and cases with site not stated.

²Patient might have more than one extrapulmonary site of disease.

³Excludes New York City.

⁴Not included in U.S. totals.

Note: Ellipses (...) indicate that the percentage cannot be calculated, and the denominator is 0.

See Technical Notes.