

Table 48. Tuberculosis Cases and Percentages, by Initial Drug Regimen: Reporting Areas, 2016

Reporting area	Total Cases	Cases in persons alive at diagnosis	Cases with information on initial drug regimen ¹		Percentage of cases in persons with initial drug regimen ^{2,3}		
			No.	(%)	IR	IRZ	IRZE ³
United States	9,272	8,968	8,872	(98.9)	(0.4)	(1.7)	(85.5)
Alabama	112	109	109	(100.0)	(0)	(2.8)	(89.0)
Alaska	57	56	56	(100.0)	(0)	(1.8)	(85.7)
Arizona	188	185	185	(100.0)	(0.5)	(1.1)	(92.4)
Arkansas	91	84	82	(97.6)	(0)	(1.2)	(95.1)
California	2,062	1,944	1,917	(98.6)	(0.3)	(0.9)	(89.2)
Colorado	64	60	59	(98.3)	(0)	(1.7)	(69.5)
Connecticut	52	52	52	(100.0)	(0)	(1.9)	(19.2)
Delaware	16	16	16	(100.0)	(0)	(0)	(68.8)
District of Columbia	25	25	25	(100.0)	(0)	(0)	(80.0)
Florida	639	622	620	(99.7)	(0.3)	(1.9)	(91.1)
Georgia	302	292	288	(98.6)	(0)	(2.1)	(33.7)
Hawaii	119	115	114	(99.1)	(0)	(0)	(91.2)
Idaho	18	16	16	(100.0)	(0)	(18.8)	(62.5)
Illinois	342	330	324	(98.2)	(0.3)	(4.3)	(85.2)
Indiana	109	108	108	(100.0)	(0)	(0.9)	(91.7)
Iowa	48	44	43	(97.7)	(0)	(0)	(90.7)
Kansas	39	38	36	(94.7)	(0)	(0)	(100.0)
Kentucky	91	88	88	(100.0)	(0)	(0)	(85.2)
Louisiana	127	123	123	(100.0)	(0)	(3.3)	(95.9)
Maine	23	23	23	(100.0)	(0)	(8.7)	(78.3)
Maryland	220	220	220	(100.0)	(0)	(1.8)	(86.4)
Massachusetts	190	184	183	(99.5)	(5.5)	(1.1)	(72.7)
Michigan	133	133	131	(98.5)	(2.3)	(3.8)	(88.5)
Minnesota	168	168	166	(98.8)	(0)	(4.2)	(79.5)
Mississippi	61	59	59	(100.0)	(0)	(10.2)	(72.9)
Missouri	101	96	96	(100.0)	(0)	(1.0)	(80.2)
Montana	4	4	4	(100.0)	(0)	(0)	(75.0)
Nebraska	28	28	28	(100.0)	(0)	(3.6)	(89.3)
Nevada	56	54	54	(100.0)	(1.9)	(1.9)	(87.0)
New Hampshire	15	15	15	(100.0)	(0)	(0)	(66.7)
New Jersey	294	284	281	(98.9)	(0.7)	(3.2)	(87.5)
New Mexico	39	39	38	(97.4)	(0)	(2.6)	(94.7)
New York State ⁴	203	199	199	(100.0)	(0.5)	(3.0)	(85.9)
New York City	565	553	546	(98.7)	(0.2)	(1.3)	(86.1)
North Carolina	219	212	212	(100.0)	(0)	(0.9)	(76.9)
North Dakota	22	22	22	(100.0)	(0)	(4.5)	(86.4)
Ohio	140	136	132	(97.1)	(0)	(0)	(75.8)
Oklahoma	78	71	71	(100.0)	(0)	(4.2)	(88.7)
Oregon	70	69	69	(100.0)	(2.9)	(1.4)	(89.9)
Pennsylvania	173	167	165	(98.8)	(0)	(0)	(89.7)
Rhode Island	12	12	12	(100.0)	(0)	(0)	(91.7)
South Carolina	102	98	98	(100.0)	(0)	(3.1)	(88.8)
South Dakota	12	11	11	(100.0)	(0)	(0)	(36.4)
Tennessee	103	100	100	(100.0)	(0)	(2.0)	(78.0)
Texas	1,250	1,230	1,213	(98.6)	(0.4)	(1.2)	(91.6)
Utah	20	20	20	(100.0)	(0)	(0)	(90.0)
Vermont	5	5	5	(100.0)	(20.0)	(0)	(40.0)
Virginia	205	201	200	(99.5)	(0)	(0)	(96.0)
Washington	205	196	189	(96.4)	(0)	(1.1)	(85.7)
West Virginia	14	13	13	(100.0)	(0)	(0)	(84.6)
Wisconsin	40	38	35	(92.1)	(0)	(2.9)	(88.6)
Wyoming	1	1	1	(100.0)	(0)	(0)	(100.0)
American Samoa ⁵	1	1	1	(100.0)	(0)	(0)	(100.0)
Fed. States of Micronesia ⁵	126	126	126	(100.0)	(0)	(0)	(98.4)
Guam ⁵	75	73	72	(98.6)	(0)	(2.8)	(91.7)
Marshall Islands ⁵	179	175	174	(99.4)	(0)	(4.6)	(92.0)
N. Mariana Islands ⁵	27	27	27	(100.0)	(0)	(22.2)	(7.4)
Puerto Rico ⁵	69	64	64	(100.0)	(0)	(15.6)	(76.6)
Republic of Palau ⁵	21	21	21	(100.0)	(0)	(0)	(100.0)
U.S. Virgin Islands ⁵	0	0	0

¹Includes persons who were alive at diagnosis and started on one or more drug.

²Overall U.S. percentage based on 52 reporting areas (50 states, New York City, and the District of Columbia).

³I, isoniazid; R, rifampin; Z, pyrazinamide; E, ethambutol. Cases with other drugs prescribed in addition to these regimens are excluded.

⁴Excludes New York City.

⁵Not included in U.S. totals.

Note: Excluding cases with no information on drug regimen, 75 (0.8%) persons were not started on any drugs, 10 (0.1%) were started on one drug, and 1,093 (12.2%) had an initial multiple drug regimen other than IR, IRZ, or IRZE.

Ellipses (...) indicate that the percentage cannot be calculated, and the denominator is 0.