

Table 1. Health insurance coverage status and coverage type for persons aged 19–25 by year, quarter, and region: United States, January 2010–June 2011.

| Year, quarter and region | Uninsured ¹ | Private health insurance coverage ² | Public health plan coverage ³ |
|-----------------------------|--------------------------|--|--|
| | Percent (standard error) | | |
| Total 2010 | 33.9 (0.73) | 51.0 (0.84) | 15.7 (0.55) |
| Quarter 1 | 33.8 (1.47) | 50.0 (1.69) | 16.8 (1.15) |
| Quarter 2 | 34.8 (1.40) | 51.5 (1.57) | 13.8 (0.94) |
| Quarter 3 | 35.6 (1.39) | 49.3 (1.50) | 15.8 (1.11) |
| Quarter 4 | 31.2 (1.33) | 53.3 (1.57) | 16.3 (1.08) |
| Total 2011 ⁴ | 28.8 (0.98) | 55.8 (1.16) | 16.1 (0.73) |
| Quarter 1 | 30.4 (1.33) | 53.2 (1.69) | 16.9 (1.12) |
| Quarter 2 | 27.3 (1.36) | 58.4 (1.62) | 15.2 (0.91) |
| Northeast 2010 | 25.6 (1.74) | 54.6 (2.12) | 20.4 (1.41) |
| Quarter 1 | 23.7 (2.48) | 55.4 (3.97) | 21.8 (3.46) |
| Quarter 2 | 28.1 (2.67) | 53.5 (4.06) | 18.5 (3.18) |
| Quarter 3 | 24.2 (3.65) | 54.3 (3.39) | 22.6 (3.20) |
| Quarter 4 | 26.3 (3.32) | 55.4 (2.73) | 18.9 (2.87) |
| Northeast 2011 ⁴ | 22.9 (2.58) | 58.8 (3.12) | 18.6 (1.58) |
| Quarter 1 | 28.6 (3.54) | 50.4 (4.02) | 21.0 (3.11) |
| Quarter 2 | 18.9 (3.10) | 64.9 (4.26) | 16.9 (2.19) |
| Midwest 2010 | 28.6 (1.36) | 56.3 (1.78) | 16.0 (1.22) |
| Quarter 1 | 30.3 (2.77) | 53.7 (3.37) | 16.7 (2.46) |
| Quarter 2 | 30.2 (2.88) | 55.1 (3.19) | 15.1 (1.98) |
| Quarter 3 | 29.3 (2.80) | 55.9 (3.11) | 15.6 (2.47) |
| Quarter 4 | 24.5 (2.85) | 60.5 (3.75) | 16.9 (2.48) |
| Midwest 2011 ⁴ | 21.1 (1.61) | 64.8 (2.25) | 14.8 (1.57) |
| Quarter 1 | 21.8 (2.42) | 62.0 (3.25) | 17.0 (2.40) |
| Quarter 2 | 20.4 (2.27) | 67.7 (2.72) | 12.4 (1.89) |
| South 2010 | 39.3 (1.41) | 47.8 (1.47) | 13.3 (0.77) |
| Quarter 1 | 37.8 (2.57) | 48.0 (2.95) | 14.7 (1.57) |
| Quarter 2 | 41.7 (2.68) | 46.3 (2.73) | 12.0 (1.45) |
| Quarter 3 | 44.4 (2.43) | 45.2 (2.36) | 10.9 (1.32) |
| Quarter 4 | 33.8 (2.29) | 51.6 (2.61) | 15.3 (1.49) |
| South 2011 ⁴ | 35.3 (1.87) | 49.8 (1.88) | 15.7 (1.19) |
| Quarter 1 | 34.4 (2.32) | 50.1 (2.69) | 16.1 (1.65) |
| Quarter 2 | 36.3 (2.34) | 49.4 (2.24) | 15.3 (1.60) |
| West 2010 | 36.2 (1.26) | 48.5 (1.53) | 15.6 (1.25) |
| Quarter 1 | 38.2 (3.15) | 45.5 (3.11) | 16.7 (2.46) |
| Quarter 2 | 34.0 (2.61) | 53.9 (2.87) | 12.1 (1.35) |
| Quarter 3 | 36.3 (2.47) | 46.0 (3.31) | 18.3 (2.46) |
| Quarter 4 | 36.9 (2.35) | 48.0 (3.33) | 15.6 (2.27) |
| West 2011 ⁴ | 30.6 (1.52) | 54.5 (2.33) | 15.9 (1.61) |
| Quarter 1 | 33.4 (2.61) | 51.5 (3.97) | 15.8 (2.36) |
| Quarter 2 | 28.0 (2.36) | 57.2 (3.31) | 15.9 (1.71) |

¹A person was defined as uninsured if he or she did not have any private health insurance, Medicare, Medicaid, Children's Health Insurance Program (CHIP), state-sponsored or other government-sponsored health plan, or military plan. A person was also defined as uninsured if he or she had only Indian Health Service coverage or had only a private plan that paid for one type of service, such as accidents or dental care.

²Excludes plans that paid for only one type of service, such as accidents or dental care.

³Includes Medicaid, CHIP, state-sponsored or other government-sponsored health plan, Medicare (disability), and military plans.

⁴Estimates are based on data collected from January through June.

NOTES: Data are based on household interviews of a sample of the civilian noninstitutionalized population. Coverage status is at the time of interview. The "Private health insurance" and "Public health plan" coverage categories are not mutually

exclusive, and a small number of persons had both types of coverage and are included in both categories. The 2011 health insurance estimates are being released prior to final data editing and weighting, to provide access to the most recent information from the National Health Interview Survey (NHIS). These estimates are provided along with comparable estimates from the 2010 NHIS. The resulting estimates for persons without health insurance are generally 0.1–0.3 percentage points lower than those based on the editing procedures used for the final data files. Occasionally, due to decisions made for the final data editing and weighting, estimates based on preliminary editing procedures may differ by more than 0.3 percentage points from estimates based on final files.

DATA SOURCE: CDC/NCHS, National Health Interview Survey, 2010–2011, Family Core component.

ACKNOWLEDGMENTS: This publication is a product of the NHIS Early Release Program. (<http://www.cdc.gov/nchs/nhis/releases.htm>). This table was produced by Michael E. Martinez and Robin A. Cohen of the Centers for Disease Control and Prevention's National Center for Health Statistics, Division of Health Interview Statistics.