

ADVISORY COMMITTEE ON IMMUNIZATION PRACTICES

VACCINES FOR CHILDREN PROGRAM

VACCINES TO PREVENT ROTAVIRUS GASTROENTERITIS

The purpose of this resolution is to add a newly licensed rotavirus vaccine to the Vaccines for Children Program.

VFC resolution 2/06-2 is repealed and replaced by the following:

Eligible Groups

Infants aged 6 weeks to 8 months

Recommended Schedule for Rotavirus Vaccines

<u>Dose</u>	<u>Rotateq</u> ® <u>Age</u>	<u>Rotarix</u> ® <u>Age</u>
Primary 1	2 months	2 months
Primary 2	4 months	4 months
Primary 3	6 months	-----

Dosage Intervals and Ages for Rotavirus Vaccines

	RV5 (Rotateq®; Merck)	RV1 (Rotarix®; GSK)*
Number of doses in series	3	2
Recommended ages for doses	2, 4, and 6 months	2 and 4 months
Minimum age for first dose	6 weeks	
Maximum age for first dose	14 weeks 6 days	
Interval between doses	4 weeks or more	
Maximum age for last dose	8 months 0 days	

Note: The use of brand names is not meant to preclude the use of other comparable licensed vaccines.

*Updated 11/29/2022 to include a new, oral dosing applicator only presentation of Rotarix licensed on 11/4/2022.

Recommended Dosages

Refer to product package inserts.

Contraindications and Precautions¹

The following conditions are contraindications to administration of rotavirus vaccine:

a. Serious Allergic Reaction to Vaccine Components

Severe allergic reaction (e.g., anaphylaxis) after a previous dose of rotavirus vaccine or to a vaccine component. Latex rubber is contained in the Rotarix® oral applicator, so infants with a severe (anaphylactic) allergy to latex should not receive Rotarix®. The Rotateq® dosing tube is latex-free.

b. Severe combined immunodeficiency (SCID)

c. History of intussusception

The following are precautions to administration of rotavirus vaccine:

a. Altered Immunocompetence

Practitioners should consider the potential risks and benefits of administering rotavirus vaccine to infants with known or suspected altered immunocompetence; consultation with an immunologist or infectious diseases specialist is advised. Children and adults who are immunocompromised because of congenital immunodeficiency, hematopoietic transplantation, or solid organ transplantation sometimes experience severe, prolonged and even fatal rotavirus gastroenteritis. However, no safety or efficacy data are available for the administration of rotavirus vaccine to infants who are potentially immunocompromised, including

- Infants with primary and acquired immunodeficiency states, including cellular immunodeficiencies; and hypogammaglobulinemic and dysgammaglobulinemic states.
- Infants with blood dyscrasias, leukemia, lymphomas of any type, or other malignant neoplasms affecting the bone marrow or lymphatic system
- Infants on immunosuppressive therapy (including high dose systemic corticosteroids);
- Infants who are HIV-exposed or infected. However, the following considerations support vaccination of HIV-exposed or infected infants:
 - the HIV diagnosis may not be established in infants born to HIV-infected mothers before the age of the first rotavirus vaccine dose; only 1.5%-3% of HIV-exposed infants in the US will be determined to be HIV-infected, and
 - Vaccine strains of rotavirus are considerably attenuated

b. Acute Gastroenteritis

In usual circumstances, rotavirus vaccine should not be administered to infants with acute, moderate to severe gastroenteritis until the condition improves. However, infants with mild acute gastroenteritis can be vaccinated, particularly if the delay in vaccination might be substantial and might make the child ineligible to receive vaccine (e.g., aged ≥ 15 weeks 0 days before the vaccine series is started). Rotavirus vaccine has not been studied among infants with concurrent acute gastroenteritis. In these infants, the immunogenicity and efficacy of rotavirus vaccine can theoretically be compromised. For example, infants who receive oral poliovirus vaccine (OPV) during an episode of acute gastroenteritis in some circumstances have diminished poliovirus antibody responses.

c. Moderate to Severe Illness

As with all other vaccines, the presence of a moderate or severe acute illness with or without fever is a precaution to administration of rotavirus vaccine. Infants with a moderate to severe acute illness should be vaccinated as soon as they have recovered from the acute phase of the illness. This precaution avoids superimposing any potential adverse effects of the vaccine on the underlying illness or mistakenly attributing a manifestation of the underlying illness to the vaccine. Vaccination should not be delayed because of the presence of mild respiratory tract illness or other mild acute illness with or without fever.

d. Preexisting Chronic Gastrointestinal Disease

Infants with preexisting gastrointestinal conditions (e.g., congenital malabsorption syndromes, Hirschsprung’s disease, short-gut syndrome) who are not undergoing immunosuppressive therapy should benefit from rotavirus vaccine vaccination, and ACIP considers the benefits to outweigh the theoretical risks. However, no data are available on the safety and efficacy of rotavirus vaccine for infants with preexisting chronic gastrointestinal disease.

¹Contraindications and Precautions updated on 2/10/2023 to align with the CDC Child and Adolescent Immunization Schedule and current ACIP recommendations.

Adopted and Effective: June 25, 2008

This document can be found on the CDC website at:
<https://www.cdc.gov/vaccines/programs/vfc/providers/resolutions.html>