


Norovirus in Healthcare Facilities Fact Sheet



General Information

Virology

Noroviruses (genus *Norovirus*, family *Caliciviridae*) are a group of related, single-stranded RNA, non-enveloped viruses that cause acute gastroenteritis in humans. Norovirus is the official genus name for the group of viruses provisionally described as “Norwalk-like viruses”. Currently, human noroviruses belong to one of three norovirus genogroups (GI, GII, or GIV), which are further divided into >25 genetic clusters. Over 75% of confirmed human norovirus infections are associated with genotype GII.

Clinical manifestations

The average incubation period for norovirus-associated gastroenteritis is 12 to 48 hours, with a median period of approximately 33 hours. Illness is characterized by nausea, acute-onset vomiting, and watery, non-bloody diarrhea with abdominal cramps. In addition, myalgia, malaise, and headache are commonly reported. Low-grade fever is present in about half of cases. Dehydration is the most common complication and may require intravenous replacement fluids. Symptoms usually last 24 to 60 hours. Up to 30% of infections may be asymptomatic.

Epidemiology of transmission

Noroviruses are highly contagious, with as few as 18 virus particles thought to be sufficient to cause infection. This pathogen is estimated to be the causative agent in over 21 million gastroenteritis cases every year in the United States, representing approximately 60% of all acute gastroenteritis cases from known pathogens. Noroviruses are transmitted primarily through the fecal-oral route, either by direct person-to-person spread or fecally contaminated food or water. Noroviruses can also spread via a droplet route from vomitus. These viruses are relatively stable in the environment and can survive freezing and heating to 60°C (140°F). In healthcare facilities, transmission can also occur through

hand transfer of the virus to the oral mucosa via contact with materials, fomites, and environmental surfaces that have been contaminated with either feces or vomitus.

Norovirus infections are seen in all age groups, although severe outcomes and longer durations of illness are most likely to be reported among the elderly. Among hospitalized persons who are immunocompromised or have significant medical comorbidities, norovirus infection can directly result in prolonged hospital stays, additional medical complications, and, rarely, death. There is currently no vaccine available for norovirus and, generally, no specific medical treatment is offered for norovirus infection apart from oral or intravenous repletion of volume.


The ease of its transmission, a very low infectious dose, a short incubation period, environmental persistence, and lack of durable immunity following infection enables norovirus to spread rapidly through confined populations. Healthcare facilities and other institutional settings (e.g., daycare centers, schools, etc.) are particularly at-risk for outbreaks because of increased person-to-person contact. Healthcare facilities managing outbreaks of norovirus gastroenteritis may experience significant costs relating to isolation precautions and personal protective equipment, ward closures, supplemental environmental cleaning, staff cohorting or replacement, and sick time.

Diagnosis of norovirus infection

Diagnosis of norovirus infection relies on the detection of viral RNA in the stools of affected persons, by use of reverse transcription-polymerase chain reaction (RT-PCR) assays. This technology is available at CDC and most state public health laboratories and should be considered in the event of outbreaks of gastroenteritis in healthcare facilities. Enzyme immune-assays may also be used for identification of norovirus outbreak but are not recommended for diagnosis of individuals. Identification of the virus can be best made from stool specimens taken within 48 to 72 hours after onset of symptoms, although positive results can be obtained by using RT-PCR on samples taken as long as 7 days after symptom onset. Because of the limited availability of timely and routine laboratory diagnostic methods, a clinical diagnosis of norovirus infection is often used, especially when other agents of gastroenteritis have been ruled out.



U.S. Department of Health and Human Services
Centers for Disease Control and Prevention



Measures to Limit Transmission*

Patient Cohorting and Isolation Precautions

Avoid exposure to vomitus or diarrhea. Place patients on Contact Precautions in a single occupancy room if they present with symptoms consistent with norovirus gastroenteritis.

Hand Hygiene

During outbreaks, use soap and water for hand hygiene after providing care or having contact with patients suspected or confirmed with norovirus gastroenteritis.

Patient Transfer and Ward Closure

Consider limiting transfers to those for which the receiving facility is able to maintain Contact Precautions; otherwise, it may be prudent to postpone transfers until patients no longer require Contact Precautions. During outbreaks, medically suitable individuals recovering from norovirus gastroenteritis can be discharged to their place of residence.

Diagnostics

In the absence of clinical laboratory diagnostics or in the case of delay in obtaining laboratory results, use Kaplan's clinical and epidemiologic criteria to identify a norovirus gastroenteritis outbreak.

Kaplan's Criteria

1. Vomiting in more than half of symptomatic cases and,
2. Mean (or median) incubation period of 24 to 48 hours and,
3. Mean (or median) duration of illness of 12 to 60 hours and,
4. No bacterial pathogen isolated in stool culture

Environmental Cleaning

Increase the frequency of cleaning and disinfection of patient care areas and frequently touched surfaces during outbreaks of norovirus gastroenteritis (e.g., increase ward/unit level cleaning to twice daily to maintain cleanliness, with frequently touched surfaces cleaned and disinfected three times daily using the US Environmental Protection Agency's list of approved products for healthcare settings (<http://www.epa.gov/oppad001/chemregindex.htm>).

Staff Leave and Policy

Develop and adhere to sick leave policies for healthcare personnel who have symptoms consistent with norovirus infection.

Exclude ill personnel from work for a minimum of 48 hours after the resolution of symptoms. Once personnel return to work, the importance of performing frequent hand hygiene should be reinforced, especially before and after each patient contact.

Establish protocols for staff cohorting in the event of an outbreak of norovirus gastroenteritis. Ensure staff care for one patient cohort on their ward and do not move between patient cohorts (e.g., patient cohorts may include symptomatic, asymptomatic exposed, or asymptomatic unexposed patient groups).

Communication and Notification

As with all outbreaks, notify appropriate local and state health departments, as required by state and local public health regulations, if an outbreak of norovirus gastroenteritis is suspected.

*Prevention and control recommendations taken from priority recommendations in the CDC HICPAC Guideline for the Prevention and Control of Norovirus Gastroenteritis Outbreaks in Healthcare Settings (<http://www.cdc.gov/hicpac/pdf/norovirus/Norovirus-Guideline-2011.pdf>)

Date last modified: September 6, 2011

Content source: Division of Healthcare Quality Promotion (DHQP), National Center for Preparedness, Detection, and Control of Infectious Diseases (NCEID)

Contact Us: Centers for Disease Control and Prevention
1600 Clifton Road, Atlanta, GA 30333, USA

1-800-CDC-INFO (1-800-232-4636)

TTY:888-232-6348,

24 hours/everyday at cdcinfo@cdc.gov (TTY)



U.S. Department of Health and Human Services
Centers for Disease Control and Prevention