

Number, rate, and standard error of deaths for discharges from short-stay hospitals, by age and selected first-listed diagnosis: United States, 2009

[Discharges of inpatients from nonfederal hospitals. Diagnostic groups and code numbers are based on the *International Classification of Diseases, Ninth Revision, Clinical Modification (ICD-9-CM)*]

First-listed diagnosis and ICD-9-CM code	All ages	Under	65 years	All ages	Under	65 years	
		65 years	and over		65 years	and over	
	Number of deaths in thousands				Rate per 100 discharges		
All conditions \1	786	237	548		2.2	1.1	3.9
Septicemia	038	134	38	96	18.6	13.9	21.5
Heart disease	391-392.0, 393-398, 402, 404, 410-416, 420-429	111	22	89	2.8	1.5	3.5
Acute myocardial infarction	410	34	*5	30	5.4	*1.9	7.6
Cardiac dysrhythmias	427	*11	*	*8	*1.4	*	*1.5
Congestive heart failure	428.0, 428.2-428.4	37	*	32	3.4	*	4.2
Malignant neoplasms	140-209.30, 230-234	73	26	47	5.5	3.8	7.2
Malignant neoplasm on trachea, bronchus, and lung	162, 176.4, 197.0, 197.3, 209.21	17	*	*12	8.9	*	*10.6
Respiratory failure	518.81, 518.83, 518.84	63	21	43	16.9	12.8	20.0
Cerebrovascular disease	430-438	51	*	38	5.2	*	5.8
Pneumonia	480-486	44	*9	35	3.9	*1.7	5.8
Nephritis, nephrotic syndrome and nephrosis	580-589	28	*	24	5.1	*	6.9
Fractures, all sites \2	800-829	19	*	14	1.8	*	2.4
Fracture of neck of femur	820	*5	*	*5	*1.9	*	*2.0
Pneumonitis due to solids and liquids	507	19	*	17	10.7	*	12.6
Standard error							
All conditions \1	87.6	30.7	61.2		0.2	0.1	0.4
Septicemia	038	16.6	7.0	12.4	2.3	2.5	2.8
Heart disease	391-392.0, 393-398, 402, 404, 410-416, 420-429	15.4	4.6	12.8	0.4	0.3	0.5
Acute myocardial infarction	410	5.7	*1.4	5.2	0.9	*0.6	1.3
Cardiac dysrhythmias	427	*2.7	*	*2.3	*0.3	*	*0.4
Congestive heart failure	428.0, 428.2-428.4	6.8	*	6.4	0.6	*	0.8
Malignant neoplasms	140-209.30, 230-234	10.7	5.3	7.6	0.8	0.8	1.2
Malignant neoplasm on trachea, bronchus, and lung	162, 176.4, 197.0, 197.3, 209.21	3.9	*	*3.3	2.0	*	*2.9
Respiratory failure	518.81, 518.83, 518.84	11.6	4.3	10.0	3.1	2.7	4.7
Cerebrovascular disease	430-438	8.8	*	6.5	0.9	*	1.0
Pneumonia	480-486	6.6	*2.3	5.6	0.6	*0.4	0.9
Nephritis, nephrotic syndrome and nephrosis	580-589	7.4	*	7.1	1.3	*	2.0
Fractures, all sites \2	800-829	4.3	*	3.9	0.4	*	0.7
Fracture of neck of femur	820	*1.4	*	*1.4	*0.5	*	*0.6
Pneumonitis due to solids and liquids	507	4.2	*	4.1	2.4	*	3.0

* Figure does not meet standard of reliability or precision.

\1 Includes data for deaths not shown in table.

\2 Excludes fractures coded as 733.1, pathologic fracture.