In 2017, Lesbian, Gay, and Bisexual Students Were at Greater Risk of Experiencing Several Risk Factors

| U.S. high school students who: | Lesbian, Gay, Bisexual (LGB) | Not Sure | Heterosexual | Increased Risk for LGB vs. Heterosexual Students |
|--|---------------------------------|-------------|--------------|---|
| Had 4 or more partners during their life | 14.7% | 9.9% | 9.1% | 62% more likely |
| Used a condom and effective hormonal birth control | 4.4% | 3.7% | 9.6% | 54% less likely |
| Used a condom at last sex | 39.9% | 44.1% | 56.1% | 29% less likely |
| Ever had sex | 48.4% | 28.4% | 39.1% | 24% more likely |
| Used effective hormonal birth control | 24.4% | 18.8% | 30.3% | 19% less likely |
| Were currently sexually active | 33.7% | 19.8% | 28.5% | 18% more likely |
| Ever injected illegal drugs | 3.4% | 6.1% | 1.0% | > 3x more likely |
| Ever misused prescription opioids | 24.3% | 17.7% | 12.9% | 88% more likely |
| Ever used illicit drugs | 23.1% | 26.8% | 12.3% | 88% more likely |
| Were forced to have sex | 21.9% | 13.1% | 5.4% | > 4x more likely |
| Experienced sexual dating violence | 15.8% | 14.1% | 5.5% | ~ 3x more likely |
| Experienced physical dating violence | 17.2% | 14.1% | 6.4% | ~ 3x more likely |
| Were electronically bullied | 27.1% | 22.0% | 13.3% | > 2x more likely |
| Were bullied at school | 33.0% | 24.3% | 17.1% | ~ 2x more likely |
| Were threatened or injured with a weapon at school | 9.4% | 11.1% | 5.4% | 74% more likely |
| Did not go to school because of safety concerns | 10.0% | 10.7% | 6.1% | 64% more likely |
| Were injured in a suicide attempt | 7.5% | 5.6% | 1.7% | > 4x more likely |
| Attempted suicide | 23.0% | 14.3% | 5.4% | > 4x more likely |
| Made a suicide plan | 38.0% | 25.6% | 10.4% | > 3x more likely |
| Seriously considered suicide | 47.7% | 31.8% | 13.3% | > 3x more likely |
| Felt sad or hopeless | 63.0% | 46.4% | 27.5% | > 2x more likely |