

south were clear for adults and for children (Figure, panel A); the level of ILI was 3–5× for children. The influenza subtypes causing the 3 peaks in the north were preceded by a peak of the same subtypes in the south. During winter 2006–07, the influenza subtype was seasonal H1N1 and to a lesser extent H3N2. In winter 2007–08, the virus was B/Yamagata; and in 2008–09, it was again seasonal influenza A (H1N1), which was almost absent in the south during April–December 2007. Antigenic characteristics of the influenza virus from the north were similar to those from the south in the same epidemic episode (2). Furthermore, influenza A (H3N2) was in southern China throughout the year, whereas in northern China, this subtype only showed a clear peak in the first 2 winters of the study period. Subtype B/Victoria and B (unsubtyped) were both in northern and southern China in irregular and low numbers. Data from the 3 northern provincial areas with year-round surveillance confirmed that influenza cases during April–September were negligible (data not shown).

The influenza subtypes of seasonal influenza A (H1N1) and B/Yamagata that have caused the past 3 summer peaks in southern China were followed by an epidemic of the same subtypes in northern China during the subsequent winter. This finding may indicate that these peaks are regular epidemic phenomena for seasonal influenza in China. Another possible explanation is that other subtypes were cocirculating with the predominant subtype at the time of epidemics.

The dual pattern of seasonal peaks for influenza is well-known for the Northern and Southern Hemispheres, but apparently it is also possible on 1 side of the equator. China is a large country with climatic differences between north and south. Although most of southern China is above the Tropic of Cancer, it is warmer and more humid than northern China (Figure, pan-

el B), which may explain the different seasonal patterns within mainland China (3). Knowledge of the dual patterns of influenza in China is relevant for determining effective control measures, and knowledge of the underlying mechanisms of such patterns is relevant to understanding the epidemiology of influenza in general.

This study was supported by the National Science Fund for Distinguished Young Scholars (grant no. 30725032), the Special Program for Prevention and Control of Infectious Diseases (grant no. 2008ZX10004-012), and the Natural Science Foundation of China (grants nos. 30810103903 and 30972521).

**Yue-Long Shu,¹ Li-Qun Fang,¹
Sake J. de Vlas, Yan Gao,
Jan Hendrik Richardus,
and Wu-Chun Cao¹**

Author affiliations: Chinese Center for Disease Control and Prevention, Beijing, People's Republic of China (Y.L. Shu, Y. Gao); Beijing Institute of Microbiology and Epidemiology, Beijing (L.-Q. Fang); University Medical Center Rotterdam, Rotterdam, the Netherlands (S.J. de Vlas, J.H. Richardus); and State Key Laboratory of Pathogen and Biosecurity, Beijing (W.-C. Cao)

DOI: 10.3201/eid1604.091578

References

1. World Health Organization. WHO recommended surveillance standards, Second edition [cited 2009 Aug 19]. http://www.who.int/csr/resources/publications/surveillance/WHO_CDS_CSR_ISR_99_2_EN/en/
2. Zhang Y, et al. Antigenic and genetic study of influenza virus circulated in China in 2006 [in Chinese]. *Zhonghua Shi Yan He Lin Chuang Bing Du Xue Za Zhi*. 2007;21:304–6.
3. Chan PK, Mok HY, Lee TC, Chu IM, Lam WY, Sung JJ. Seasonal influenza activity in Hong Kong and its association with meteorological variations. *J Med Virol*. 2009; 81:1797–806.

¹These authors contributed equally to this article.

Address for correspondence: Wu-Chun Cao, Beijing Institute of Microbiology and Epidemiology, State Key Laboratory of Pathogen and Biosecurity, 100071, Beijing, People's Republic of China; email: caowc@nic.bmi.ac.cn

Avian Influenza Prevalence in Pigs, Egypt

To the Editor: Since 1996, avian influenza virus (H5N1) has spread to ≥65 countries (1). The disease represents a serious threat for the poultry industry and public health. Egypt has the highest human infection and fatality rates outside Asia (2). Some isolates of influenza virus (H5N1) in Egypt are resistant to oseltamivir (3), and in others, virulent mutations have developed, leading to case-fatality rates of 100% (4).

Pigs have the largest epidemiologic role in the evolution of new influenza viruses (5). Recombination between the newly emerged influenza virus subtypes H1N1 and H5N1 in pigs would have catastrophic results. We therefore investigated the seroprevalence of influenza virus (H5N1) in pigs in Egypt.

In May 2008, we collected 1 serum sample and 1 nasal swab from each of 240 pigs (11 herds) in Cairo slums. May was selected because it directly follows the season of bird migration and the seasonal storms usually accompanied by airborne diseases. Cairo slums were selected because 1) pigs there feed on organic remains, including dead birds, and thus have a higher chance of becoming infected; 2) Cairo is at the base of the Nile Delta, where most subtype H5N1 foci occurred; and 3) Cairo is near Fayum, the main stop-over site for migrating birds.

To detect anti-avian influenza antibodies in the serum, we used hemagglutination inhibition (HI) assays with 2 inactivated antigens: subtype H5N2 from the Veterinary Laboratories Agency, UK; and a local subtype H5N1 prepared according to the protocol used in the central national laboratories. To detect viral RNA in the nasal swabs, we used real-time PCR, as was recommended for detection of influenza (H5N1) infection during outbreaks in Southeast Asia (6).

Although all nasal samples reacted negatively to influenza A/H5 by real-time PCR, only 4 serum samples showed positive results by HI when using subtype H5N2 antigen; titers were 32 for 3 samples and 64 for 1. Seven additional positive serum samples were detected when antigen prepared from local subtype H5N1 virus was used; titers ranged from 16 (6 samples) to 512 (1 sample). Also during this 2-week sampling period, titers of 32 for 3 samples and 128 for 1 were obtained. Seroprevalence rate of avian influenza for the 240 pigs was 1.67% and 4.6% when the nonlocal or local viral antigens, respectively, were used. Of the 11 positive pigs, 8 were from 1 herd and 3 were from 3 other herds.

Failure to detect viral RNA in the upper respiratory tract indicates the absence of acute infections in the investigated pigs. Inability of the virus to persist in the pigs was reported (7). Contrary to the HI results, results of routine examination of the 240 pigs found no abnormalities. Absence of clinical signs in infected pigs was reported (8) and was attributed to their low susceptibility to influenza (H5N1) (7). The results indicate that infection rate for pigs in Egypt is clearly higher than that for pigs in China and Vietnam (8,9). This increase may be attributed to different spatial and temporal factors leading to increased infection risk among sampled pigs, higher antigenicity of native isolates, or most probably to the disease situation in Egypt. The detection of 8 positive reactors from

1 herd indicates a subtype H5N1 focus there as was reported in Indonesia (8). The difference in the number of reactors when using different antigens indicates the difference in antigenicity. These data are supported by field observations regarding low protection level ($\approx 35\%$ in some reports) of imported vaccines (A. El-Sayed, unpub. data). The relatively low seroprevalence of avian influenza in pigs may be misleading because of the poor immunogenicity of some avian influenza lines and lack of sensitivity for detecting low titers of induced antibodies (10). It may be also explained by the use of a virus antigen other than that existing in the population, as was done in the present study.

Human risk for influenza (H5N1) infection in Egypt seems to be associated mainly with infected birds. It has not yet been associated with infected pigs.

Acknowledgments

We thank Elham Elebiary and Lamia Omar for their excellent help.

This study was financed by Cairo University.

**Amr El-Sayed,
Walid Awad, Adel Fayed,
Hans-Peter Hamann,
and Michael Zschöck**

Author affiliations: Cairo University, Giza, Egypt (A. El-Sayed, W. Awad, A. Fayed); and Hessian State Laboratory, Giessen, Germany (H.P. Hamann, M. Zschöck)

DOI: 10.3201/eid1604.091316

References

1. Center for Infectious Disease Research and Policy. Avian influenza (bird flu): implications for human disease [cited 2010 Jan 29]. http://www.cidrap.umn.edu/cidrap/content/influenza/avianflu/biofacts/avflu_human.html
2. World Health Organization. Confirmed human cases of avian influenza A/(H5N1); 2009 [cited 2009 Apr 4]. http://www.who.int/csr/disease/avian_influenza/country/en/

3. Kage B. Bird flu virus mutates into Tamiflu-resistant strain in Egypt; WHO does not expect pandemic [cited 2007 Jan 24]. http://www.flunews.org/1058_bird_flu_Tamiflu.html
4. Niman L. Third H5N1 fatality in Gharbiya cluster in Nile Delta. *Recombinomics commentary*, December 27, 2006 [cited 2010 Feb 11]. http://www.recombinomics.com/News/12270601/H5N1_Gharbiya_Fatal_3.html
5. Ma W, Kahn RE, Richt JA. The pig as a mixing vessel for influenza viruses: human and veterinary implications. *J Mol Genet Med*. 2008;3:158–66.
6. Tran TH, Nguyen L, Nguyen D, Luong S, Pham M, Nguyen C, et al. Avian influenza A (H5N1) in 10 patients in Vietnam. *N Engl J Med*. 2004;350:1179–88. DOI: 10.1056/NEJMoa040419
7. Lipatov AS, Kwon YK, Sarmento LV, Lager KM, Spackman E, Suarez DL, et al. Domestic pigs have low susceptibility to H5N1 highly pathogenic avian influenza viruses. *PLoS Pathog*. 2008;4:e1000102. DOI: 10.1371/journal.ppat.1000102
8. Cyranoski D. Bird flu spreads among Java's pigs. *Nature*. 2005;435:390–1. DOI: 10.1038/435390a
9. Choi YK, Nguyen TD, Ozaki H, Webby RJ, Puthavathana P, Buranathal C, et al. Studies of H5N1 influenza virus infection of pigs by using viruses isolated in Vietnam and Thailand in 2004. *J Virol*. 2005;79:10821–5. DOI: 10.1128/JVI.79.16.10821-10825.2005
10. de Jong MD, Hien TT. Avian influenza A (H5N1). *J Clin Virol*. 2006;35:2–13. DOI: 10.1016/j.jcv.2005.09.002

Address for correspondence: Amr El-Sayed, Faculty of Veterinary Medicine, Cairo University, Giza, Egypt; email: aaelsayed2000@yahoo.de

