

# *Rickettsia japonica* Infections in Humans, Zhejiang Province, China, 2015

Qunying Lu, Jianping Yu, Liquan Yu,  
Yanjuan Zhang, Yitao Chen,  
Meiai Lin, Xiaofei Fang

We investigated 16 Japanese spotted fever cases that occurred in southeastern China during September–October 2015. Patients had fever, rash, eschar, and lymphadenopathy. We confirmed 9 diagnoses and obtained 2 isolates with high identity to *Rickettsia japonica* strain YH. *R. japonica* infection should be considered for febrile patients in China.

*Rickettsia japonica* is a member of the spotted fever group rickettsiae (SFGR) that causes tickborne Japanese spotted fever (JSF). First reported in Japan's Tokushima Prefecture in 1984 (1,2), JSF has been recognized in multiple countries of Asia, including Japan, South Korea, the Philippines, and Thailand (3–5). In China, 4 species of SFGR have been reported to cause human infection: *R. heilongjiangensis*, *R. sibirica* subspecies *sibirica* BJ-90, *Candidatus R. tarasevichiae*, and *R. raoultii* (6). In this report, we investigated the causative agent of 16 JSF cases that occurred in southeastern China in late 2015.

## The Study

The ethics committee of the Zhejiang Province Center for Disease Control and Prevention, Hangzhou, China, approved this research. During September–October 2015, a total of 16 febrile patients were hospitalized at Linan First People's Hospital (Linan, China). All these patients lived in the Xitianmu Mountain area of Linan in Zhejiang Province. Besides fever (38.8°C–40.3°C), clinical signs of disease in these patients included rashes on the trunk and limbs (15/16) and an eschar (11/16) (Table). In those with eschar, lymphadenopathy was found at the site of the draining lymph node (6/11). Five patients had rash and no eschar. Laboratory results revealed that all patients' leukocyte levels were within reference ranges, but a high percentage of neutrophils (12/16 patients) and minor hepatic transaminase elevation (11/16 patients) were also observed. All 16 patients were treated with doxycycline

or azithromycin and were cured, and no patient experienced severe illness.

With patient consent, we collected acute-phase (n = 16) and convalescent-phase (n = 14) whole blood specimens and sent them to Zhejiang Province Center for Disease Control and Prevention for laboratory confirmation of *Rickettsia* infection. We extracted DNA from acute-phase blood specimens by using the DNeasy Blood & Tissue Kit (QIAGEN, Hilden, Germany). We used nested PCR to amplify the partial *groEL* genes of SFGR, typhus group rickettsiae, and *Orientia tsutsugamushi* bacteria (online Technical Appendix, <https://wwwnc.cdc.gov/EID/article/24/11/17-0044-Techapp1.pdf>). The targeted fragments (217 bp) were present in the blood specimens from 9 patients (Table). We sequenced these fragments and analyzed them using BLAST (<http://www.ncbi.nlm.nih.gov/BLAST>), and all had 100% identity to *R. japonica* YH prototype strain (GenBank accession no. NC016050) (7,8). All specimens were negative for typhus group rickettsiae and *O. tsutsugamushi* rickettsia DNA by PCR.

We also inoculated 200 µL of acute-phase blood specimens onto HL60 and DH82 cells in 25-mL flasks and cultured at 37°C. Cytopathic effect was not observed with inoculated HL60 cells, but inoculated DH82 cells exfoliated completely by 4 weeks of culture. We also performed indirect immunofluorescence assays (IFAs) every 2 days to assess SFGR growth (online Technical Appendix). Two of the inoculated cultures exhibited bright fluorescent apple-green, rod-shaped particles (Table) after 3 weeks of culture, confirming SFGR infection for 2 patients. We then extracted DNA from the 2 SFGR-positive cultures (LA4/2015 and LA16/2015) and amplified and sequenced a 2,493-bp fragment containing the full-length sequences of SFGR *groES* and *groEL* (GenBank accession nos. KY073364–5) and a 609-bp fragment containing the partial rickettsial *ompA* gene sequence (GenBank accession nos. KY347792–3; online Technical Appendix Table). These sequences were found to be 100% identical to the corresponding sequences of *R. japonica* YH.

We used IFAs with bacterial substrate antigens *R. japonica* (HL-60 cells infected with LA4/2015) and *R. rickettsii* (FOCUS Diagnostics Inc., Cyprus, CA, USA) to test patients for specific antibodies, and in all 16 patient serum samples, we detected SFGR IgG. All paired serum samples (n = 14) showed a >4-fold increase in titer against SFGR (Table). The 2 patients we did not receive

Author affiliations: Zhejiang Province Center for Disease Control and Prevention, Hangzhou, China (Q. Lu, Y. Zhang, X. Fang); Linan First People's Hospital, Linan, China (J. Yu, L. Yu); Zhejiang Chinese Medical University, Hangzhou (Y. Chen, M. Lin)

DOI: <https://doi.org/10.3201/eid2411.170044>

**Table.** Clinical data of febrile patients with Japanese spotted fever diagnoses, Linan, China, 2015\*

Patient no.	Age, y/sex	Tick bite	Rash	Eschar	Lymphadenopathy	Days after onset		PCR,†	IFA titer‡		Isolation
						AP	CP		AP	CP	
1	53/M	Not known	Yes	No	No	6	42	–	<64	4,096	–
2	62/M	Not known	Yes	Yes	No	5	58	+	<64	2,048	–
3	52/M	Yes	Yes	Yes	Yes	5	74	+	64	4,096	–
4	79/F	Not known	Yes	No	No	4	60	+	128	9,192	+
5	67/F	No	Yes	No	No	6	75	+	512	4,096	–
6	62/F	Yes	Yes	Yes	No	7	70	–	<64	1,024	–
7	53/F	Yes	No	Yes	No	7	57	–	64	1,024	–
8	63/F	Yes	Yes	Yes	Yes	10	67	–	<64	4,096	–
9	57/M	Yes	Yes	Yes	No	2	45	+	64	9,192	–
10	51/F	Yes	Yes	Yes	Yes	6	§	+	64	§	–
11	53/M	Yes	Yes	Yes	No	3	§	+	<64	§	–
12	45/F	Yes	Yes	Yes	Yes	5	70	+	<64	4,096	–
13	33/F	Yes	Yes	Yes	Yes	6	42	–	64	2,048	–
14	30/F	Not known	Yes	No	No	4	46	–	128	2,048	–
15	36/F	Not known	Yes	No	No	5	54	–	<64	2,048	–
16	67/F	Yes	Yes	Yes	Yes	0	60	+	<64	9,192	+

\*AP, acute phase; CP, convalescent phase; IFA, indirect immunofluorescence assay; +, positive test result; –, negative test result.

†PCR amplifying 217-bp sequence of *Rickettsia japonica groEL* gene.

‡Detection of IgG against spotted fever group rickettsiae.

§Convalescent-phase serum samples not obtained.

convalescent-phase serum specimens from were positive for *R. japonica* by PCR.

All serum specimens were negative for *O. tsutsugamushi* IgG. Some convalescent-phase serum specimens had low-titer reactions to *R. typhi* bacterial antigen.

## Conclusions

The 4 SFGR species *R. japonica*, *R. heilongjiangensis*, *R. rhipicephali*, and *R. massiliae* have been identified in *Haemaphysalis longicornis* and *Rhipicephalus haemaphysaloides* ticks in Zhejiang Province (9–11), indicating a potential for these species to infect humans in China. In our research, we determined the etiologic agent of 16 JSF cases and isolated 2 *R. japonica* rickettsiae. The prototype strain *R. japonica* YH was isolated in Japan in 1985 (1). After ≈30 years, only a few *R. japonica* isolates have been isolated from patients in China: 2 from our research and 1 from Li et al. (12). Our findings indicate that the full-length *groES* and *groEL* genes and the partial *ompA* gene sequences were 100% identical to *R. japonica* YH, suggesting that the *R. japonica* genome has been relatively conserved. Nine patients had clinically confirmed JSF, displaying fever, rash, eschar, and lymphadenopathy; these signs and symptoms were similar to those seen in JSF patients in Japan (13).

Of the vectorborne rickettsial diseases in China, human scrub typhus and murine typhus frequently occur in Zhejiang Province, and spotted fever group rickettsiosis probably occurs but has gone relatively unnoticed. Because the clinical symptoms of spotted fever and scrub typhus are similar, some SFGR infections have likely been diagnosed as scrub typhus. We found that the blood specimens of 7 febrile patients were negative for the targeted PCR fragments but showed a >4-fold increase in antibody titer to SFGR.

Although these results suggest SFGR infection, we cannot conclude these 7 patients were infected with *R. japonica*.

In summary, *R. japonica* infections occur in Zhejiang Province, China. These infections are likely more broadly distributed throughout the mainland areas than had been previously realized. Improvements in JSF clinical diagnosis and human epidemiologic surveillance are urgently needed in China.

## Acknowledgments

We thank our colleagues in the Linan First People's Hospital, Zhejiang Province, Linan, China, for their assistance in specimen collection.

## About the Author


Dr. Lu is a principal investigator at the Zhejiang Province Center for Disease Control and Prevention, Hangzhou, China. Her research interests include microbiology, epidemiology, and the ecology of tickborne diseases.

## References

1. Uchida T, Uchiyama T, Kumano K, Walker DH. *Rickettsia japonica* sp. nov., the etiological agent of spotted fever group rickettsiosis in Japan. *Int J Syst Bacteriol*. 1992;42:303–5. <http://dx.doi.org/10.1099/00207713-42-2-303>
2. Mahara F. Three Weil-Felix reaction OX2 positive cases with skin eruptions and high fever [in Japanese]. *J Anan Med Assoc*. 1984;68:4–7.
3. Camer GA, Alejandria M, Amor M, Satoh H, Muramatsu Y, Ueno H, et al. Detection of antibodies against spotted fever group *Rickettsia* (SFGR), typhus group *Rickettsia* (TGR), and *Coxiella burnetii* in human febrile patients in the Philippines. *Jpn J Infect Dis*. 2003;56:26–8. PubMed
4. Chung MH, Lee SH, Kim MJ, Lee JH, Kim ES, Lee JS, et al. Japanese spotted fever, South Korea. *Emerg Infect Dis*. 2006;12:1122–4. <http://dx.doi.org/10.3201/eid1207.051372>

5. Gaywee J, Sunyakumthorn P, Rodkvamtook W, Ruang-areerate T, Mason CJ, Sirisopana N. Human infection with *Rickettsia* sp. related to *R. japonica*, Thailand. *Emerg Infect Dis.* 2007;13:657–9. <http://dx.doi.org/10.3201/eid1304.060585>
6. Fang LQ, Liu K, Li XL, Liang S, Yang Y, Yao HW, et al. Emerging tick-borne infections in mainland China: an increasing public health threat. *Lancet Infect Dis.* 2015;15:1467–79. [http://dx.doi.org/10.1016/S1473-3099\(15\)00177-2](http://dx.doi.org/10.1016/S1473-3099(15)00177-2)
7. Dong X, El Karkouri K, Robert C, Raoult D, Fournier PE. Genomic analysis of *Rickettsia japonica* strain YHT. *J Bacteriol.* 2012;194:6992. <http://dx.doi.org/10.1128/JB.01928-12>
8. Matsutani M, Ogawa M, Takaoka N, Hanaoka N, Toh H, Yamashita A, et al. Complete genomic DNA sequence of the East Asian spotted fever disease agent *Rickettsia japonica*. *PLoS One.* 2013;8:e71861. <http://dx.doi.org/10.1371/journal.pone.0071861>
9. Sun J, Lin J, Gong Z, Chang Y, Ye X, Gu S, et al. Detection of spotted fever group rickettsiae in ticks from Zhejiang Province, China. *Exp Appl Acarol.* 2015;65:403–11. <http://dx.doi.org/10.1007/s10493-015-9880-9>
10. Meng Z, Jiang LP, Lu QY, Cheng SY, Ye JL, Zhan L. Detection of co-infection with Lyme spirochetes and spotted fever group rickettsiae in a group of *Haemaphysalis longicornis* [in Chinese]. *Zhonghua Liu Xing Bing Xue Za Zhi.* 2008;29:1217–20.
11. Cheng SY, Zheng SQ, Meng Z, Ye XD, Jiang LP, Wang ZQ, et al. Analysis of DNA sequences of spotted fever group *Rickettsia* detected from ticks in Zhejiang Province [in Chinese]. *Dis Surveil.* 2010;25:466–8.
12. Li J, Hu W, Wu T, Li HB, Hu W, Sun Y, et al. Japanese spotted fever endemic in East China, 2013. *Emerg Infect Dis.* 2018 Nov [cited 2018 Aug 27]. <https://doi.org/10.3201/eid2411.170264>
13. Mahara F. Japanese spotted fever: report of 31 cases and review of the literature. *Emerg Infect Dis.* 1997;3:105–11. <http://dx.doi.org/10.3201/eid0302.970203>

Address for correspondence: Qunying Lu, Zhejiang Province Center for Disease Control and Prevention, No. 3399 Binsheng Rd, Hangzhou, 310052 China; email: lqycdc@163.com



**TICKBORNE DISEASES OF THE UNITED STATES**  
*A Reference Manual for Health Care Providers*  
 Second Edition, 2014

U.S. Department of Health and Human Services  
 Centers for Disease Control and Prevention

**Recognize the signs of tickborne disease**

**Understand diagnostic testing and lab findings**

**Quickly find treatment recommendations**

**Order or download at**  
**[www.cdc.gov/pubs](http://www.cdc.gov/pubs)**

# *Rickettsia japonica* Infections in Humans, Zhejiang Province, China, 2015

## Technical Appendix

### Materials

#### Molecular Detection of Rickettsial Infection

For laboratory confirmation of *Rickettsia* infection, patient whole blood specimens were collected during the acute phase and sent to Zhejiang Province Center for Disease Control and Prevention, Hangzhou, China. DNA extraction and PCR were performed in a standard PCR laboratory and each step included negative controls. DNA was extracted from acute phase blood specimens and cell culture by using the DNeasy Blood & Tissue Kit (QIAGEN, Hilden, Germany) according to the manufacturer's instructions.

First, nested PCR of the *groEL* gene (217 bp) of spotted fever group rickettsiae (SFGR), typhus group rickettsiae, and *Orientia tsutsugamushi* bacteria were performed as described previously (1,2). PCR targeting the SFGR *ompA* gene was performed to identify the bacteria grown in cell culture as described previously (3). Another PCR was performed with primers specific to the nucleotide sequences of the full-length SFGR *groES* and *groEL* genes (Technical Appendix Table). The sequences were aligned and trimmed with MEGA 5.0 (<https://www.megasoftware.net/>) (4).

#### Indirect Immunofluorescence Assay (IFA) Detection for SFGR in Inoculated Cells

In total, 200  $\mu$ L of blood samples (collected with K2 EDTA tubes) from every patient were inoculated onto HL60 and DH82 cells and cultured at 37°C. Slides of inoculated and

noninoculated (negative control) cells were prepared and fixed in cold acetone for 7 minutes; IFA was used to determine whether the inoculated cells were infected (5). SFGR positivity of the 2 convalescent patient serum samples was confirmed by *Rickettsia* IFA IgG (FOCUS Diagnostics Inc., Cypress, CA, USA), and then, these positive serum samples were used as the primary antibody for SFGR-specific IFAs, with goat anti-human IgG (Sigma-Aldrich, St. Louis, MO, USA) serving as the secondary antibody.

## References

1. Park HS, Lee JH, Jeong EJ, Kim JE, Hong SJ, Park TK, et al. Rapid and simple identification of *Orientia tsutsugamushi* from other group rickettsiae by duplex PCR assay using *groEL* gene. *Microbiol Immunol.* 2005;49:545–9. [PubMed http://dx.doi.org/10.1111/j.1348-0421.2005.tb03760.x](http://dx.doi.org/10.1111/j.1348-0421.2005.tb03760.x)
2. Fu XP, Zhang JS, Shen XJ, Luan MC, Li ML, Zhang LJ. Surveillance on *Rickettsia* in epidemic areas of scrub typhus in Xinyang areas of Henan [in Chinese]. *Zhonghua Liu Xing Bing Xue Za Zhi.* 2007;28:547–9. [PubMed](#)
3. Zhang L, Jin J, Fu X, Raoult D, Fournier P-E. Genetic differentiation of Chinese isolates of *Rickettsia sibirica* by partial *ompA* gene sequencing and multispacer typing. *J Clin Microbiol.* 2006;44:2465–7. [PubMed http://dx.doi.org/10.1128/JCM.02272-05](http://dx.doi.org/10.1128/JCM.02272-05)
4. Tamura K, Peterson D, Peterson N, Stecher G, Nei M, Kumar S. MEGA5: molecular evolutionary genetics analysis using maximum likelihood, evolutionary distance, and maximum parsimony methods. *Mol Biol Evol.* 2011;28:2731–9. [PubMed http://dx.doi.org/10.1093/molbev/msr121](http://dx.doi.org/10.1093/molbev/msr121)
5. Uchida T, Uchiyama T, Kumano K, Walker DH. *Rickettsia japonica* sp. nov., the etiological agent of spotted fever group rickettsiosis in Japan. *Int J Syst Bacteriol.* 1992;42:303–5. [PubMed http://dx.doi.org/10.1099/00207713-42-2-303](http://dx.doi.org/10.1099/00207713-42-2-303)

**Technical Appendix Table.** Information on primers used for PCR and sequencing in study of febrile patients with Japanese spotted fever, Linan, China, 2015\*

Organism	Target gene	Primer	Sequence	Reference		
<i>Rickettsiaceae</i>	<i>groEL</i>	GRO1	5'-AAGAAGGMGTGATAAC-3'	(1)		
		GRO2	5'-ACTTCMGTAGCACC-3'			
<i>Rickettsia</i>	<i>groEL</i>	SF1	5'-GATAGAAGAAAAGCAATGATG-3'	(2)		
		SR2	5'-CAGCTATTTGAGATTTAATTTG-3'			
<i>Orientia tsutsugamushi</i>	<i>groEL</i>	TF1	5'-ATATATCACAGTACTTTGCAAC-3'			
		TR2	5'-GTTCTAACTTAGATGTATCAT-3'			
Spotted fever group rickettsiae	<i>groES</i> and <i>groEL</i>	F1	5'-CTCCGAATAGTTTAGGTAATTGGC-3'	This research		
		R1	5'-TTTCGGTRCCTGCCCATTTAC-3'			
		F2	5'-GGTGAAGAGAAAATAAGGTGGA-3'			
		R2	5'-CTGCTAAATCCATACCACGCTT-3'			
		F3	5'-TGCAGAGGTAGCHGGTGAYGG-3'			
		R3	5'-TAACATTTTCAAGCTTCATACCTA-3'			
		F4	5'-CGCTTGTAGTCAATAGATTACGTGG-3'			
		R4	5'-TACTAGATCTAYTRCATAACCTATCTT-3'			
		<i>ompA</i>	Rr190.70p		5'-ATGGCGAATATTTCTCCAAAA-3'	(3)
			Rr190-701R		5'-GTTCCGTTAATGGCAGCATCT-3'	

\*Primers GRO1 and GRO2 were used in the first round for nested PCR. SF1 and SR2, specific to *Rickettsia*, were used in the second round of PCR. TF1 and TR2 are specific to *Orientia tsutsugamushi*. Primers F1, R1, F2, R2, F3, R3, F4, and R4 refer to the sequence of *R. japonica* YH (GenBank accession no. NC\_016050).