

Large Nationwide Outbreak of Invasive Listeriosis Associated with Blood Sausage, Germany, 2018–2019

Sven Halbedel,¹ Hendrik Wilking,¹ Alexandra Holzer, Sylvia Kleta, Martin A. Fischer, Stefanie Lüth, Ariane Pietzka, Steliana Huhulescu, Raskit Lachmann, Amrei Krings, Werner Ruppitsch, Alexandre Leclercq, Rolf Kamphausen, Maylin Meincke, Christiane Wagner-Wiening, Matthias Contzen, Iris Barbara Kraemer, Sascha Al Dahouk, Franz Allerberger, Klaus Stark,² Antje Flieger²

Invasive listeriosis is a severe foodborne infection in humans and is difficult to control. Listeriosis incidence is increasing worldwide, but some countries have implemented molecular surveillance programs to improve recognition and management of listeriosis outbreaks. In Germany, routine whole-genome sequencing, core genome multilocus sequence typing, and single nucleotide polymorphism calling are used for subtyping of *Listeria monocytogenes* isolates from listeriosis cases and suspected foods. During 2018–2019, an unusually large cluster of *L. monocytogenes* isolates was identified, including 134 highly clonal, benzalkonium-resistant sequence type 6 isolates collected from 112 notified listeriosis cases. The outbreak was one of the largest reported in Europe during the past 25 years. Epidemiologic investigations identified blood sausage contaminated with *L. monocytogenes* highly related to clinical isolates; withdrawal of the product from the market ended the outbreak. We describe how epidemiologic investigations and complementary molecular typing of food isolates helped identify the outbreak vehicle.

Listeriosis is a severe, mainly foodborne, human infection associated with higher case-fatality and hospitalization rates than other bacterial

gastrointestinal pathogens (1). The causative agent, *Listeria monocytogenes*, occurs ubiquitously in the environment and disseminates into the food production chain. Patients develop either self-limiting noninvasive gastroenteritis or invasive listeriosis (2,3). Listeriosis adversely affects older and immunocompromised persons, as well as pregnant women, causing a severe invasive form of the disease that leads to sepsis, meningitis, and encephalitis, as well as neonatal infections and miscarriage (4). Case-fatality rates of invasive listeriosis are ≈30% for neurolisteriosis and even higher in septic patients (5). In Europe and North America, invasive listeriosis affects 0.3–0.6 persons/100,000 population/year (6,7).

L. monocytogenes forms hard-to-remove biofilms in food-processing plants, can acquire tolerance to sanitizers, and multiplies even at temperatures used for refrigeration (8). These properties complicate efficient prevention of *L. monocytogenes* contaminations in different types of ready-to-eat products, including dairy, meat, and fish, and in fruits and vegetables, all of which have been vehicles for listeriosis outbreaks in the past (9–12).

Author affiliations: Robert Koch Institute, Wernigerode, Germany (S. Halbedel, M.A. Fischer, A. Flieger); Robert Koch Institute, Berlin, Germany (H. Wilking, A. Holzer, R. Lachmann, A. Krings, M. Meincke, K. Stark); German Federal Institute for Risk Assessment, Berlin (S. Kleta, S. Lüth, S.A. Dahouk); Freie Universität Berlin, Berlin (S. Lüth); Austrian Agency for Health and Food Safety, Vienna, Austria (A. Pietzka, S. Huhulescu, W. Ruppitsch, F. Allerberger); European Centre for Disease Prevention and Control, Stockholm, Sweden (A. Krings, M. Meincke); Institut Pasteur, Paris, France (A. Leclercq); Ministry for Environment, Agriculture, Conservation and Consumer Protection

of the State of North Rhine-Westphalia, Düsseldorf, Germany (R. Kamphausen); State Health Office Baden-Wuerttemberg, Stuttgart, Germany (M. Meincke, C. Wagner-Wiening); Chemical and Veterinary Investigations Office, Fellbach, Germany (M. Contzen); Bavarian Health and Food Safety Authority, Oberschleißheim, Germany (I.B. Kraemer); Rheinisch-Westfälische Technische Hochschule, Aachen, Germany (S.A. Dahouk)

DOI: <https://doi.org/10.3201/eid2607.200225>

¹These authors contributed equally to this article.

²These authors contributed equally to this article.

Outbreaks of listeriosis are difficult to control for several reasons. First, case numbers are low, impairing the generation of valid hypotheses about possible food sources through patient interviews. Second, incubation time can be long, 1–67 days (13), and patients often are seriously ill, further complicating patient interviews. Third, the large variety of possible food sources makes pinpointing through patient interviews and follow-up tracing of food difficult. Moreover, listeriosis outbreaks can be geographically widespread due to long-distance food trade connections, e-commerce, and travel, thus hampering outbreak recognition by local authorities (10,14,15). In addition, listeriosis outbreaks can be protracted and last for several years (16), making it difficult to correctly identify affected patient groups and the common source of infection.

Nationwide systematic collection of *L. monocytogenes* isolates from human listeriosis cases and subtyping by using high-resolution whole-genome sequencing (WGS)-based typing techniques can aid in rapid and reliable detection of outbreak clusters (3,17–22), some of which were not detectable in the past. At the same time, systematic and on-demand typing of food-associated *L. monocytogenes* isolates assist in detecting outbreak sources. In a recent molecular surveillance study in France, one third of all isolates were grouped into WGS clusters, and most clusters contained <5 isolates (20). Larger outbreaks of invasive listeriosis occur, although infrequently, and 2 of the world's largest outbreaks in the recent past included 147 cases in a multistate outbreak associated with cantaloupes in the United States in 2011 (10) and 1,060 cases in an outbreak associated with French polony sausage in South Africa during 2017–2018 (11). Since August 2019, Spain has been experiencing another large listeriosis outbreak (23), but the scientific evaluation of this outbreak is ongoing.

We describe an exceptionally large nationwide outbreak that included 134 laboratory-confirmed *L. monocytogenes* isolates from 112 patients with epidemiologic investigations and complementary WGS-based typing of food isolates identifying the outbreak vehicle. This outbreak represents one of the largest outbreaks of invasive listeriosis in Europe documented in the scientific literature during the past 25 years.

Methods and Materials

Isolation, Growth, and Serotyping of *L. monocytogenes*

We isolated *L. monocytogenes* from 184 specimens from human cases and food sources (Appendix Table 1,

<http://wwwnc.cdc.gov/EID/article/26/7/20-0225-App1.pdf>). We performed routine *L. monocytogenes* cultures in brain heart infusion (BHI) broth, or on BHI or sheep blood agar plates at 37°C. We detected and enumerated *L. monocytogenes* from food samples according to International Organization for Standardization (ISO) methods EN ISO 11290–1:2017 and EN ISO 11290–2:2017 (24,25). We confirmed the species by using EN ISO 11290–1:2017 or matrix-assisted laser desorption/ionization time-of-flight mass spectrometry, and a previously described multiplex PCR (26). We used the GenElute Bacterial Genomic DNA Kit (SigmaAldrich, <https://www.sigmaaldrich.com>) or the QIAamp DNA Mini Kit (QIAGEN, <https://www.qiagen.com>) to isolate chromosomal DNA and determined molecular serogroups by using multiplex PCR (27).

Genome Sequencing, Multilocus Sequence Typing, and Core Genome MLST

We quantified DNA by using the Qubit dsDNA BR or HS Assay Kit and Qubit fluorometers (Invitrogen, <https://www.thermofisher.com>). We prepared libraries by using the Nextera XT DNA Library Prep Kit (Illumina, <https://www.illumina.com>) and sequenced isolates on MiSeq, HiSeq, or NextSeq sequencers (Illumina). We trimmed and assembled raw reads in SeqSphere (Ridom, <https://www.ridom.de>) by using the Velvet assembler. We extracted in silico serogroups, multilocus sequence types (STs), and 1,701 locus core genome multilocus sequence typing (cgMLST) complex types (CTs) by using SeqSphere and automated allele submission to the *L. monocytogenes* cgMLST server (<http://www.cgmlst.org/ncs/schema/690488>) (28). We deposited genome sequences in the European Nucleotide Archive (<https://www.ebi.ac.uk/ena>; accession numbers in Appendix Table 1). Coverage ranged between 22- and 116-fold (median 54-fold). We defined cgMLST clusters as groups of isolates with ≤ 10 different alleles between neighboring isolates. We used SeqSphere in the pairwise ignore missing values mode and an unweighted pair group method with arithmetic mean to generate phylogenetic trees.

Single-Nucleotide Polymorphism–Based Alignments

We used pipelines developed in-house to map sequencing reads, generate consensus sequences, calculate alignment, and filter single-nucleotide polymorphisms (SNPs) using an exclusion distance of 300 bps (17). We used the 10-092876-0769 LM12 genome (GenBank accession no. CP019625) a member of serogroup IVb, ST6, and CT6304, as the reference. We

generated maximum likelihood trees by using the Geneious 9.1.3 Tree Builder (<https://www.geneious.com>) and the randomized accelerated maximum likelihood plugin.

Virulome and Resistome Analyses and Susceptibility Testing

We included virulence and resistance genes of *L. monocytogenes* as target loci in SeqSphere task templates, as previously described (17,29). We extracted targets from assembled contigs by using SeqSphere and considered alleles present when identity was >90% and the query sequence aligned $\geq 99\%$ with the reference sequence.

We performed antimicrobial drug susceptibility testing by using a microdilution assay in a 96-well plate format adapted from a study by Noll et al. (30). All susceptibility testing was performed in accordance with European Committee on Antimicrobial Susceptibility Testing guidelines in Mueller Hinton fastidious (MH-F) broth (31).

We spread *L. monocytogenes* isolates on BHI agar plates and placed 6 mm cellulose discs loaded with 10 μ L of a 10 mg/mL aqueous benzalkonium chloride solution on top of the agar plate. We incubated plates at 37°C overnight, then determined growth inhibition zone diameters. We used the Student *t*-test to assess statistical significance.

Case-Control Study

We defined outbreak cases as patients reported to public health authorities with disease onset during August 2018–June 2019 and isolation of *L. monocytogenes* from normally sterile body fluids and confirmation by cgMLST and SNP analysis. *L. monocytogenes* isolates were sent to the Robert Koch Institute (Wernigerode, Germany) and notification and typing data were merged for investigation. After the outbreak was identified, patients were interviewed by using a standardized questionnaire on food consumption during the 2 weeks before illness onset, general eating habits, and food purchasing behaviors. These data identified 40 food items for inclusion in the case-control study.

We collaborated with a survey institute to contact and interview case-controls. We frequency matched case-controls to case-patients for age, gender, and federal state of residence. We considered food items with $p \leq 0.05$ consumed by $\geq 50\%$ of participants for multivariable analysis. We used a stepwise-backward approach for model formation to consecutively exclude food items that were no longer significantly associated from the multivariable model until only significantly associated foods and their confounders

remained. We determined risk measures, including odds, univariable, and multivariable ratios, in the statistical analysis.

Results

Molecular Surveillance

The binational German–Austrian Consultant Laboratory for *L. monocytogenes* collects and sequences genomes of isolates from approximately two thirds of all mandatorily notified listeriosis cases in Germany; 699 cases were notified in 2018 and 593 in 2019. A phylogenetically diverse cluster, designated Epsilon1, was identified by using cgMLST. Epsilon1 included 46 PCR serogroup IVb isolates belonging to ST6 and CT90, CT2981, CT3803, CT3805, CT3806, CT3921, CT4083, CT4465, CT6236, CT6331, CT7353, and CT7451, all of which had specific allelic profiles within a CT threshold of ≤ 10 different alleles (17,28). The Epsilon1 cluster included isolates collected from all over Germany during 2011–2019 with no apparent geographic concentration. Allelic distances between isolates varied from 0–25 (median 11). In autumn 2018, a sudden increase of CT4465 and CT7353 isolates belonging to the Epsilon1 cluster was detected. Furthermore, the number of listeriosis cases reported in calendar weeks 34–43, 46, 48, and 50 exceeded the median of the 5 previous years (Appendix Figure 1). To identify the outbreak clone among all incoming serogroup IVb isolates, we developed a clone-specific PCR (Appendix). Altogether, 134 clinical CT4465 and CT7353 isolates were collected during August 2018–April 2019. These isolates formed a remarkably homogenous cluster with 0–5 (median 0) different cgMLST alleles (Figure 1). In contrast, 2 CT4465 isolates collected earlier, in July 2017 and June 2018, differed in 9–15 alleles (Figure 1).

We mapped raw sequence reads of all Epsilon1 strains against the 10-092876-0769 LM12 genome, the most closely related complete genome available. SNP calling separated the Epsilon1 cluster into several sub-clusters, but all CT4465 and CT7353 isolates collected from August 2018 onwards formed a single cluster (Appendix Figure 2). This subcluster was named Epsilon1a, and SNP distances in this cluster ranged from 0–3 SNPs (median 0). The 2 earlier CT4465 isolates were separated from the Epsilon1a cluster by 6–10 SNPs difference (median 8). Thus, SNP calling supported detection of a cluster of closely related CT4465 and CT7353 strains. Of note, only 21–29 cgMLST alleles (median 26) and 8–12 SNPs (median 8) differed between the Epsilon1a clone and the outbreak strain from South Africa (11), CT5886 (Figure 1; Appendix Figure 2).

Case Cohort

We collected 134 isolates from 112 patients who met the case definition. Initial cases were reported in August 2018, and the outbreak peaked in September 2018 (Figure 2, panel A); the last notified case was in April 2019. Cases occurred in 11/16 federal states in Germany; most cases occurred in western and southern Germany (Figure 2, panel B). This outbreak and the assembled genome of 1 representative isolate (isolate no. 18-04540) were shared via the Epidemic Intelligence Information System platform of the European Centre for Disease Prevention and Control on October 23, 2018 (UI-516, <https://www.ecdc.europa.eu>). France, the only other country involved, reported an Epsilon1a listeriosis case in a patient who had traveled to and purchased food in Germany. Sequence data of isolate 18-04540 was submitted to the European Nucleotide Archive (<https://www.ebi.ac.uk/ena>; accession no. SAMEA5041142). However, the closest related isolate available at the National Center for Biotechnology Information (<https://www.ncbi.nlm.nih.gov>) pathogen detection pipeline was a 2016 clinical isolate from the Netherlands with an SNP distance of 12, which is clearly above the SNP distances observed in the Epsilon1a cluster.

One (0.9%) case-patient was pregnant, but the gestational age and health outcome of her newborn were not reported. The remaining 111 case-patients were 53–98 (median 79) years of age; 66 (59%) were men, 45 (41%) were women. Seven (6.3%) case-patients died, 2 of whom had listeriosis as the primary cause of death. The age distribution was not noticeably different from other notified listeriosis outbreaks. Of the 134 Epsilon1a isolates, 99 were from blood samples, 13 from cerebrospinal fluid, and 1 each from lymph nodes, ascites, sputum, pleura, joints, abscesses, or a superficial wound (Appendix Table 1). The isolation source was not reported for the remaining 15 isolates.

Properties of the Outbreak Clone

Virulome analysis revealed the presence of *Listeria* pathogenicity island 1 (LIPI-1) in all Epsilon1a outbreak isolates and detected the complete listeriolysin S-encoding LIPI-3 in 64% (Appendix Figure 3). However, we did not detect LIPI-4, which encodes a putative phosphotransferase system associated with neuroinfection (32). Epsilon1a clones carried the same complement of internalin genes as other serogroup IVb strains (Appendix Figure 3).

Susceptibility testing revealed sensitivity toward most clinically relevant antimicrobial drugs, but all tested isolates were fully resistant to ceftriaxone and daptomycin (Appendix Table 2), which

is consistent with the intrinsic resistance of *L. monocytogenes* and the absence of additional resistance determinants, as suggested by the resistome




Figure 1. Phylogenetic tree constructed by using unweighted pair group method with arithmetic mean and core genome multilocus sequence typing data of *Listeria monocytogenes* isolates from a large listeriosis outbreak, Germany. Green indicates clinical isolates of Epsilon1a subcluster; blue indicates food isolates of Epsilon1a subcluster; pink indicates isolates from the Epsilon1 cluster; violet indicates 2 complex type 4465 isolates not belonging to Epsilon1a from earlier listeriosis cases in July 2017 and June 2018; yellow indicates isolates from a listeriosis outbreak in South Africa (11); black indicates reference strain 10-092876-0769 LM12 used for SNP calling (Appendix Figure 2, <https://wwwnc.cdc.gov/EID/article/26/7/20-0225-App1.pdf>). Scale bar indicates allelic substitutions per site. SNP, single-nucleotide polymorphism.





Figure 2. Spatial and temporal distribution of cases during a large listeriosis outbreak, Germany. A) Number of *Listeria monocytogenes* isolates from subcluster Epsilon1a received by the consulting laboratory per week during the outbreak. B) Geographic distribution of laboratory-confirmed Epsilon1a cases in Germany during the outbreak. CW, calendar week.



approach (Appendix Figure 4). Further resistome analysis demonstrated the prevalence of the *emrC* gene, which is associated with benzalkonium chloride tolerance (Appendix Figure 4). In full agreement with this observation, Epsilon1a and Epsilon1 isolates demonstrated increased tolerance to benzalkonium chloride compared with ST6 or serogroup IVb isolates from other outbreak clusters (Figure 3).


Identification the Outbreak Vehicle

Our case-control study included 41 case-patients and 155 controls. A total of 40/41 (98%) case-patients reported that they purchased food in a single specific supermarket chain, compared with 99/154 controls (64.3%; odds ratio [OR] 22.5, 95% CI 2.9–174.9;

$p = 0.003$). No other supermarket chains were associated with outbreak cases, and we only included case-patients and controls that had purchased food from the specific supermarket chain in further analyses. In the fourth calendar week of 2019, we detected a strong association between cases and consumption of minced meat (OR 42.4, 95% CI 4.3–415.4; $p = 0.001$) and blood sausage (OR 23.1, 95% CI 4.3–123.5; $p < 0.001$; Table). Among case-patients, 90% reported consuming minced meat and 80% reported consuming blood sausage compared with 23% of controls who consumed minced meat and 45% who consumed blood sausage. None of the case-patients were vegetarians.

To perform risk-oriented screening, food samples were collected in supermarkets and the house-

Figure 3. Tolerance of isolates of *Listeria monocytogenes* from subcluster Epsilon1a in Germany to benzalkonium chloride. Three representative isolates from human listeriosis clusters Epsilon1a, Epsilon1, and distinct listeriosis clusters Lambda2 (ST2, CT2402), (17) Pi3 (ST217, CT5744), or Theta3 (ST249, CT4449) were tested for resistance to benzalkonium chloride by disc diffusion, along with 3 representative ST6 isolates, not belonging to Epsilon1. Epsilon1a and Epsilon1 isolates showed increased resistance to benzalkonium chloride. Circles, squares, and diamonds represent results of 3 independent replicates for 3 isolates per group. Asterisk indicates statistically significant differences to Epsilon1a ($p \leq 0.01$) calculated by using the Student *t*-test. CT, complex type; Eps1, Epsilon1; Eps1a, Epsilon1a; Lam2, Lambda2; nonEps1, nonEpsilon1; ST, sequence type.



holds of some patients, according to the results of the epidemiologic investigations. In 1 case, *L. monocytogenes* was detected in 3 open samples from a patient's refrigerator. Among these, sliced blood sausage purchased at the implicated supermarket chain showed the highest contamination ($>3 \times 10^6$ CFU/g). This finding led to another round of intensified screening of prepackaged blood sausage. In calendar week 7 of 2019, *L. monocytogenes* was found in an original sealed package of sliced blood sausage (<10 CFU/g) and in a second sample of blood sausage from the same manufacturer. In total, 5 isolates from patients' household food items and from blood sausage samples grouped with clinical Epsilon1a isolates after cgMLST (0–3 different alleles, median = 0) and SNP calling (0–2 SNPs, median = 0) (Figure 1; Appendix Figure 2).

The blood sausage was produced by a large meat and sausage manufacturer in Germany and sold in many parts of the country. The product was withdrawn from the market on February 12, 2019. The last clinical Epsilon1a isolate was collected on April 18, 2019. In contrast, Epsilon1 isolates not belonging to the Epsilon1a cluster caused disease even after the end of the Epsilon1a outbreak. The plant was cleaned and disinfected. Thereafter, *L. monocytogenes* was not detected from products or the production site among several hundred samples collected by food safety inspectors and $\approx 2,500$ control samples collected by the manufacturer.

Discussion

The Epsilon1a outbreak is the largest identified outbreak of listeriosis in Germany and one of the largest documented outbreaks of invasive listeriosis in Europe in >25 years. The last reported outbreak of invasive listeriosis of this size in Europe was during 1992–1993 when 247 patients were infected in France with a serotype IVb clone from contaminated pork tongue in aspic (33). Cantaloupe was the vehicle in the large outbreak in the United States in 2011 that was caused by 5 different clones (10). In contrast, the single clone that caused a large outbreak in South Africa showed strong clonality and the genomes of 326 isolates differed in ≤ 4 cgMLST alleles (11). Likewise, the Epsilon1a outbreak was caused by a single clone, and we observed high clonality among isolates. The mutation rate in the natural *L. monocytogenes* population is 2.6×10^{-7} substitutions per site per year (29). On average, 1 SNP/year can be expected for *L. monocytogenes* strains under natural conditions. The high clonality of Epsilon1a could imply that the outbreak clone only persisted in the production facility and did not undergo rapid multiplication.

Purchases in a particular supermarket chain and consumption of blood sausage were strongly associated with listeriosis in the case-control study, and the outbreak clone was identified in blood sausage samples from a patient's household and from the implicated supermarket chain. Blood sausage is heat-treated during production, so contamination likely

Table. Results of multivariable analysis of risk for infection by food consumption from a case-control study during listeriosis outbreak, Germany 2018–2019*

Food item	Odds ratio (95% CI)†	p value
Minced meat	42.4 (4.3–415.1)	0.001
Blood sausage	23.1 (4.3–123.5)	<0.001
Cold cuts, including roast pork and Kassler	15.4 (2.9–82.1)	0.001
Edamer cheese	7.3 (1.6–32.8)	0.009
Smoked ham‡	0.06 (0.0–0.4)	0.003
Hard cheese‡	0.2 (0.0–0.9)	0.038

*Includes 40/41 cases and 99/155 controls that confirmed shopping at the implicated supermarket chain.

†Adjusted for age, gender, and residence in northern or southern region.

‡Smoked ham and hard cheese are confounders for cold cuts and Edamer cheese.

occurred after production, possibly during slicing or packaging. The shelf-life of sliced blood sausage is several days to a few weeks (34) and the amount of *L. monocytogenes* found in unopened blood sausage samples was below the limit of 100 CFU/g. Storage beyond the anticipated shelf-life or insufficient refrigeration might have allowed *L. monocytogenes* to multiply inside the vehicle, which would only be prevented by a zero-tolerance policy.

Typically, pregnancy-related listeriosis accounts for ≈7% of all listeriosis cases (35). However, in this outbreak only 1/112 (0.8%) cases was in a pregnant woman. Official recommendations for pregnant women to apply special hygiene practices to sliced sausage products likely had an effect (36).

Analysis of the Epsilon1a genome has yielded some insights into the infectivity of this ST6 clone. *L. monocytogenes* ST6 clones were first isolated in 1990 (37), have caused various outbreaks in the past, including the large outbreak in South Africa (11), and are associated with increased rates of meningitis (38). The Epsilon1a clone and the outbreak strain from South Africa are closely related. Thus, 2 descendants of the same historic *L. monocytogenes* ancestor have spread globally and contaminated food production facilities on 2 different continents. The Epsilon1a clone carried the *emrC* gene, which presumably caused its increased tolerance to benzalkonium chloride (39). Europe banned use of benzalkonium chloride as a disinfectant in 2016 (40), but its past use might have selected tolerant strains.

The identification of this outbreak and its vehicle resulted from an efficient collaboration between public health and food safety authorities in Germany. Several requirements had to be met for successful outbreak clarification: development of a mandatory notification system for systematic patient interviews and an efficient questionnaire to generate hypotheses on possible food sources; implementation of a WGS-based molecular surveillance

program for reliable identification of outbreak clusters by public health authorities; systematic collection of food isolates from internal controls and on-demand investigations and use of a harmonized WGS-based subtyping methodology by food safety authorities; and a continuous exchange of information on outbreak clusters between the institutions involved. These prerequisites have identified the causative food vehicles for 5 of 6 large listeriosis clusters that occurred in Germany during 2014–2019, which likely would not have been possible before use of WGS in outbreak investigations. However, introduction of routine interviews of listeriosis patients, regardless of outbreaks, probably could further accelerate identification of outbreak vehicles. In our opinion, the Epsilon1a outbreak demonstrates how WGS-based pathogen surveillance combined with efficient interventions of the involved stakeholders can improve management and prevention of food-borne infectious diseases.

Acknowledgments

We thank Andrea Thürmer and her team for sequencing support, Birgitt Hahn and Simone Dumschat for excellent technical assistance, and Yvonne Pfeiffer for help with some experiments. We also thank Karan Gautam Kaval for the critical review of the manuscript.

This project was supported by the intensified molecular surveillance initiative of the Robert Koch Institute to A.F. and K.S. (2016–2018, grant no. 832133) and grants of the German Ministry of Health to S.H. (grant no. ZMVII-2518NIK703) and to A.F. and S.K. (grant no. MolTypList). The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

About the Author

Dr. Halbedel is a molecular microbiologist in the Division of Enteropathogenic Bacteria and *Legionella* at the Robert Koch Institute in Wernigerode, Germany, and the deputy head of the consultant laboratory for *Listeria*. His research focuses on physiology, virulence, and epidemiology of *Listeria monocytogenes*.

References

1. Werber D, Hille K, Frank C, Dehnert M, Altmann D, Müller-Nordhorn J, et al. Years of potential life lost for six major enteric pathogens, Germany, 2004–2008. *Epidemiol Infect.* 2013;141:961–8. <https://doi.org/10.1017/S0950268812001550>
2. Ooi ST, Lorber B. Gastroenteritis due to *Listeria monocytogenes*. *Clin Infect Dis.* 2005;40:1327–32. <https://doi.org/10.1086/429324>







3. Halbedel S, Prager R, Banerji S, Kleta S, Trost E, Nishanth G, et al. A *Listeria monocytogenes* ST2 clone lacking chitinase ChiB from an outbreak of non-invasive gastroenteritis. *Emerg Microbes Infect.* 2019;8:17–28. <https://doi.org/10.1080/22221751.2018.1558960>
4. Allerberger F, Wagner M. Listeriosis: a resurgent foodborne infection. *Clin Microbiol Infect.* 2010;16:16–23. <https://doi.org/10.1111/j.1469-0691.2009.03109.x>
5. Charlier C, Perrodeau É, Leclercq A, Cazenave B, Pilmis B, Henry B, et al.; MONALISA study group. Clinical features and prognostic factors of listeriosis: the MONALISA national prospective cohort study. *Lancet Infect Dis.* 2017;17:510–9. [https://doi.org/10.1016/S1473-3099\(16\)30521-7](https://doi.org/10.1016/S1473-3099(16)30521-7)
6. de Noordhout CM, Devleeschauwer B, Angulo FJ, Verbeke G, Haagsma J, Kirk M, et al. The global burden of listeriosis: a systematic review and meta-analysis. *Lancet Infect Dis.* 2014;14:1073–82. [https://doi.org/10.1016/S1473-3099\(14\)70870-9](https://doi.org/10.1016/S1473-3099(14)70870-9)
7. European Centre for Disease Prevention and Control. Surveillance atlas of infectious diseases. 2018 [cited 2020 Mar 16]. <https://atlas.ecdc.europa.eu/public/index.aspx>
8. Ferreira V, Wiedmann M, Teixeira P, Stasiewicz MJ. *Listeria monocytogenes* persistence in food-associated environments: epidemiology, strain characteristics, and implications for public health. *J Food Prot.* 2014;77:150–70. <https://doi.org/10.4315/0362-028X.JFP-13-150>
9. Swaminathan B, Gerner-Smith P. The epidemiology of human listeriosis. *Microbes Infect.* 2007;9:1236–43. <https://doi.org/10.1016/j.micinf.2007.05.011>
10. McCollum JT, Cronquist AB, Silk BJ, Jackson KA, O'Connor KA, Cosgrove S, et al. Multistate outbreak of listeriosis associated with cantaloupe. *N Engl J Med.* 2013;369:944–53. <https://doi.org/10.1056/NEJMoa1215837>
11. Smith AM, Tau NP, Smouse SL, Allam M, Ismail A, Ramalwa NR, et al. Outbreak of *Listeria monocytogenes* in South Africa, 2017–2018: laboratory activities and experiences associated with whole-genome sequencing analysis of isolates. *Foodborne Pathog Dis.* 2019;16:524–30. <https://doi.org/10.1089/fpd.2018.2586>
12. Stephan R, Althaus D, Kiefer S, Lehner A, Hatz C, Schmutz C, et al. Foodborne transmission of *Listeria monocytogenes* via ready-to-eat salad: A nationwide outbreak in Switzerland, 2013–2014. *Food Control.* 2015;57:14–7. <https://doi.org/10.1016/j.foodcont.2015.03.034>
13. Goulet V, King LA, Vaillant V, de Valk H. What is the incubation period for listeriosis? *BMC Infect Dis.* 2013;13:11. <https://doi.org/10.1186/1471-2334-13-11>
14. Heiman KE, Garalde VB, Gronostaj M, Jackson KA, Beam S, Joseph L, et al. Multistate outbreak of listeriosis caused by imported cheese and evidence of cross-contamination of other cheeses, USA, 2012. *Epidemiol Infect.* 2016;144:2698–708. <https://doi.org/10.1017/S095026881500117X>
15. European Centre for Disease Prevention and Control and European Food Safety Authority. Multi-country outbreak of *Listeria monocytogenes* sequence type 8 infections linked to consumption of salmon products – 25 October 2018. Stockholm and Parma: ECDC/EFSA; 2018 [cited 2019 Dec 17]. <https://www.ecdc.europa.eu/sites/default/files/documents/listeria-multi-country-outbreak-october-2018.pdf>
16. Ruppitsch W, Prager R, Halbedel S, Hyden P, Pietzka A, Huhulescu S, et al. Ongoing outbreak of invasive listeriosis, Germany, 2012 to 2015. *Euro Surveill.* 2015;20:30094. <https://doi.org/10.2807/1560-7917.ES.2015.20.50.30094>
17. Halbedel S, Prager R, Fuchs S, Trost E, Werner G, Flieger A. Whole-genome sequencing of recent *Listeria monocytogenes* isolates from Germany reveals population structure and disease clusters. *J Clin Microbiol.* 2018;56:e00119–18. <https://doi.org/10.1128/JCM.00119-18>
18. Kleta S, Hammerl JA, Dieckmann R, Malorny B, Borowiak M, Halbedel S, et al. Molecular tracing to find source of protracted invasive listeriosis outbreak, southern Germany, 2012–2016. *Emerg Infect Dis.* 2017;23:1680–3. <https://doi.org/10.3201/eid2310.161623>
19. Jackson BR, Tarr C, Strain E, Jackson KA, Conrad A, Carleton H, et al. Implementation of nationwide real-time whole-genome sequencing to enhance listeriosis outbreak detection and investigation. *Clin Infect Dis.* 2016;63:380–6. <https://doi.org/10.1093/cid/ciw242>
20. Moura A, Tourdjman M, Leclercq A, Hamelin E, Laurent E, Fredriksen N, et al. Real-time whole-genome sequencing for surveillance of *Listeria monocytogenes*, France. *Emerg Infect Dis.* 2017;23:1462–70. <https://doi.org/10.3201/eid2309.170336>
21. Kwong JC, Mercoulia K, Tomita T, Easton M, Li HY, Bulach DM, et al. Prospective whole-genome sequencing enhances national surveillance of *Listeria monocytogenes*. *J Clin Microbiol.* 2016;54:333–42. <https://doi.org/10.1128/JCM.02344-15>
22. Chen Y, Gonzalez-Escalona N, Hammack TS, Allard MW, Strain EA, Brown EW. Core genome multilocus sequence typing for identification of globally distributed clonal groups and differentiation of outbreak strains of *Listeria monocytogenes*. *Appl Environ Microbiol.* 2016;82:6258–72. <https://doi.org/10.1128/AEM.01532-16>
23. World Health Organization. Listeriosis–Spain, disease outbreak news. Geneva: the Organization; updated 2019 Sep 16 [cited 2019 Oct 1]. <https://www.who.int/csr/don/16-september-2019-listeriosis-spain>
24. International Organization for Standardization. ISO 11290-1:2017. Microbiology of the food chain – Horizontal method for the detection and enumeration of *Listeria monocytogenes* and of *Listeria* spp. – part 1: detection method. Geneva: The Organization; 2017 [cited 2019 Dec 17]. <https://www.iso.org/standard/60313.html>
25. International Organization for Standardization. ISO 11290-2:2017. Microbiology of the food chain – horizontal method for the detection and enumeration of *Listeria monocytogenes* and of *Listeria* spp. – part 2: enumeration method. Geneva: The Organization; 2017 [cited 2019 Dec 17]. <https://www.iso.org/standard/60314.html>
26. Bubert A, Hein I, Rauch M, Lehner A, Yoon B, Goebel W, et al. Detection and differentiation of *Listeria* spp. by a single reaction based on multiplex PCR. *Appl Environ Microbiol.* 1999;65:4688–92. <https://doi.org/10.1128/AEM.65.10.4688-4692.1999>
27. Kérouanton A, Marault M, Petit L, Grout J, Dao TT, Brisabois A. Evaluation of a multiplex PCR assay as an alternative method for *Listeria monocytogenes* serotyping. *J Microbiol Methods.* 2010;80:134–7. <https://doi.org/10.1016/j.mimet.2009.11.008>
28. Ruppitsch W, Pietzka A, Prior K, Bletz S, Fernandez HL, Allerberger F, et al. Defining and evaluating a core genome multilocus sequence typing scheme for whole-genome sequence-based typing of *Listeria monocytogenes*. *J Clin Microbiol.* 2015;53:2869–76. <https://doi.org/10.1128/JCM.01193-15>
29. Moura A, Criscuolo A, Pouseele H, Maury MM, Leclercq A, Tarr C, et al. Whole genome-based population biology and epidemiological surveillance of *Listeria monocytogenes*. *Nat Microbiol.* 2016;2:16185. <https://doi.org/10.1038/nmicrobiol.2016.185>

30. Noll M, Kleta S, Al Dahouk S. Antibiotic susceptibility of 259 *Listeria monocytogenes* strains isolated from food, food-processing plants and human samples in Germany. *J Infect Public Health*. 2018;11:572-7. <https://doi.org/10.1016/j.jiph.2017.12.007>
31. European Committee on Antimicrobial Susceptibility Testing. Breakpoint tables for interpretation of MICs and zone diameters. Version 9.0, 2019. Basal (Switzerland): EUCAST; 2019 Jan 1 [cited 2019 Dec 5]. http://www.eucast.org/fileadmin/src/media/PDFs/EUCAST_files/Breakpoint_tables/v_9.0_Breakpoint_Tables.pdf
32. Maury MM, Tsai YH, Charlier C, Touchon M, Chenal-Francisque V, Leclercq A, et al. Uncovering *Listeria monocytogenes* hypervirulence by harnessing its biodiversity. *Nat Genet*. 2016;48:308-13. <https://doi.org/10.1038/ng.3501>
33. Jacquet C, Catimel B, Brosch R, Buchrieser C, Dehaumont P, Goulet V, et al. Investigations related to the epidemic strain involved in the French listeriosis outbreak in 1992. *Appl Environ Microbiol*. 1995;61:2242-6. <https://doi.org/10.1128/AEM.61.6.2242-2246.1995>
34. Pereira JA, Silva P, Matos TJ, Patarata L. Shelf life determination of sliced Portuguese traditional blood sausage – Morcela de Arroz de Monchique through microbiological challenge and consumer test. *J Food Sci*. 2015;80:M642-8. <https://doi.org/10.1111/1750-3841.12782>
35. Allerberger F, Huhulescu S. Pregnancy related listeriosis: treatment and control. *Expert Rev Anti Infect Ther*. 2015;13:395-403. <https://doi.org/10.1586/14787210.2015.1003809>
36. Groeneveld M, Wichmann-Schauer H, Oehlschläger J, Werber D, Maschkowski G; Federal Institute of Agriculture and Food. Listeriosis and toxoplasmosis-safe eating during pregnancy, 2nd edition; B. Klein editor [in German]. Bonn, Germany: Bundesanstalt für Landwirtschaft und Ernährung; 2017 [cited 2019 Dec 17]. https://www.ble-medien-service.de/frontend/esddownload/index/id/513/on/0346_DL/act/dl
37. Cantinelli T, Chenal-Francisque V, Diancourt L, Frezal L, Leclercq A, Wirth T, et al. "Epidemic clones" of *Listeria monocytogenes* are widespread and ancient clonal groups. *J Clin Microbiol*. 2013;51:3770-9. <https://doi.org/10.1128/JCM.01874-13>
38. Koopmans MM, Brouwer MC, Bijlsma MW, Bovenkerk S, Keijzers W, van der Ende A, et al. *Listeria monocytogenes* sequence type 6 and increased rate of unfavorable outcome in meningitis: epidemiologic cohort study. *Clin Infect Dis*. 2013;57:247-53. <https://doi.org/10.1093/cid/cit250>
39. Kremer PH, Lees JA, Koopmans MM, Ferwerda B, Arends AW, Feller MM, et al. Benzalkonium tolerance genes and outcome in *Listeria monocytogenes* meningitis. *Clin Microbiol Infect*. 2017;23:265.e1-e7. PubMed <https://doi.org/10.1016/j.cmi.2016.12.008>
40. European Union. Commission implementing decision (EU) 2016/1950 of 4 November 2016 on the non-approval of certain biocidal active substances pursuant to Regulation (EU) No. 528/2012 of the European Parliament and of the Council. *Official Journal of the European Union* 2016. p. 16-20 [cited 2019 Dec 17]. https://www.helpdesk-biocides.fr/files/PDF/reglementation_europe/en_non_inscription_sa/20161104_decision_2016_1950_eu_non_approval_substances_bpr.pdf

Addresses for correspondence: Sven Halbedel or Antje Flieger, FG11 Division of Enteropathogenic Bacteria and Legionella, Robert Koch Institute, Burgstrasse 37, 38855 Wernigerode, Germany; email: halbedels@rki.de or fliegera@rki.de

EID Podcast: Emerging Infectious Diseases Cover Art

Byron Breedlove, managing editor of the journal, elaborates on aesthetic considerations and historical factors, as well as the complexities of obtaining artwork for Emerging Infectious Diseases.



Visit our website to listen:
**EMERGING
INFECTIOUS DISEASES**
<https://www2c.cdc.gov/podcasts/player.asp?f=8646224>

Large Nationwide Outbreak of Invasive Listeriosis Associated with Blood Sausage, Germany, 2018–2019

Appendix

Additional Methods

Diagnostic PCR for Identification of the Outbreak Clone

During this investigation, the median time from arrival of a sample in the consultant laboratory until genome sequencing results was 34.5 days. Therefore, we searched for DNA regions specific for Epsilon1a isolates to identify Epsilon1a clones by PCR and prioritize samples for genome sequencing and initiate patient interviews as soon as possible. For this purpose, the contigs of 3 Epsilon1a isolates (target) and of 50 nonEpsilon1a isolates of serogroup IVb (nontarget) were analyzed by the RUCS 1.0 algorithm, designed to identify primer pairs for unique core sequences in a target genome dataset and absent in a set of nontarget genomes (*1*). This approach identified a 262 bp fragment specific for the chosen target genomes that could be amplified by using the primers Eps-1-fw (AGTCGTCTTTAGTGCGCTGAA) and Eps-1-rev (TAGGTCTGTTGATGGCACCAC). The applicability of this primer set was tested by using 16 genome sequences of Epsilon1a and 12 of nonEpsilon1a serogroup IVb clones. Experimentally determined sensitivity of the PCR system was 94% and specificity was 75%. The 262 bp fragment is part of an open reading frame encoding a phage tail length tape measure protein, which was detected in 119/134 (89%) Epsilon1a strains according to genomic data. In contrast, among 662 analyzed serogroup IVb genomes, this phage tail open reading frame was found in only 20 (3%) ST6 genomes that did not belong to the Epsilon1a group. Thus, the Eps1a PCR was used to identify possible Epsilon1a isolates among the incoming serogroup IVb isolates to prioritize them for WGS. Of the 67 serogroup IVb isolates that had been tested by the Eps1a PCR during the outbreak, only 11 (16%) were false-positive.

References

1. Thomsen MCF, Hasman H, Westh H, Kaya H, Lund O. RUCS: rapid identification of PCR primers for unique core sequences. *Bioinformatics*. 2017;33:3917–21. [PubMed](#)
<https://doi.org/10.1093/bioinformatics/btx526>
2. Kropac AC, Eshwar AK, Stephan R, Tasara T. New insights on the role of the pLMST6 plasmid in *Listeria monocytogenes* biocide tolerance and virulence. *Front Microbiol*. 2019;10:1538. [PubMed](#)
<https://doi.org/10.3389/fmicb.2019.01538>

Appendix Table 1. *Listeria monocytogenes* isolates included from a large listeriosis outbreak, Germany, 2018–2019*

Isolate ID	Sample accession no.	Secondary accession no.	Study number	Type	Source	Cluster	Complex type
11–04869	SAMEA104485064	ERS2103006	PRJEB24496	Human	Blood	Epsilon1	90
16–00332	SAMEA104485223	ERS2103165	PRJEB24496	Human	Blood	Epsilon1	90
16–00478	SAMEA104485231	ERS2103173	PRJEB24496	Human	Blood	Epsilon1	90
16–00634	SAMEA104485236	ERS2103178	PRJEB24496	Human	CSF	Epsilon1	90
16–00830	SAMEA5770458	ERS3574002	PRJEB33238	Human	Blood	Epsilon1	90
16–00831	SAMEA104485244	ERS2103186	PRJEB24496	Human	Uterus	Epsilon1	90
16–00955	SAMEA104485248	ERS2103190	PRJEB24496	Human	Blood	Epsilon1	90
16–01401	SAMEA104485262	ERS2103204	PRJEB24496	Human	Blood	Epsilon1	90
16–01909	SAMEA104485285	ERS2103227	PRJEB24496	Human	Blood	Epsilon1	90
16–01911	SAMEA104485286	ERS2103228	PRJEB24496	Human	Blood	Epsilon1	90
16–02052	SAMEA104485291	ERS2103233	PRJEB24496	Human	Unknown	Epsilon1	3803
16–02281	SAMEA104485298	ERS2103240	PRJEB24496	Human	Blood	Epsilon1	3805
16–02328	SAMEA104485301	ERS2103243	PRJEB24496	Human	Unknown	Epsilon1	3806
16–02495	SAMEA104485307	ERS2103249	PRJEB24496	Human	Unknown	Epsilon1	90
16–02497	SAMEA104485309	ERS2103251	PRJEB24496	Human	Blood	Epsilon1	2981
16–02650	SAMEA104485313	ERS2103255	PRJEB24496	Human	Blood	Epsilon1	3921
16–03183	SAMEA104485347	ERS2103289	PRJEB24496	Human	CSF	Epsilon1	4083
16–04063	SAMEA5770459	ERS3574003	PRJEB33238	Human	Blood	Epsilon1	90
16–04236	SAMEA104485396	ERS2103338	PRJEB24496	Human	Blood	Epsilon1	90
16–04386	SAMEA5770460	ERS3574004	PRJEB33238	Human	Blood	Epsilon1	90
16–04399	SAMEA104485408	ERS2103350	PRJEB24496	Human	Blood	Epsilon1	90
16–04799	SAMEA104485422	ERS2103364	PRJEB24496	Human	Ascites	Epsilon1	90
16–04800	SAMEA104485423	ERS2103365	PRJEB24496	Human	Blood	Epsilon1	90
16–05014	SAMEA104485430	ERS2103372	PRJEB24496	Human	Ascites	Epsilon1	90
17–00454	SAMEA104485458	ERS2103400	PRJEB24496	Human	Blood	Epsilon1	4083
17–00659	SAMEA5769034	ERS3572580	PRJEB33238	Human	Blood	Epsilon1	2981
17–01077	SAMEA5769035	ERS3572581	PRJEB33238	Human	Blood	Epsilon1	90
17–03140	SAMEA5769036	ERS3572582	PRJEB33238	Human	Blood	Epsilon1	4465
17–05508	SAMEA5769037	ERS3572583	PRJEB33238	Human	Blood	Epsilon1	2981
17–06068	SAMEA5769038	ERS3572584	PRJEB33238	Human	Blood	Epsilon1	90
17–06319	SAMEA5769039	ERS3572585	PRJEB33238	Human	CSF	Epsilon1	90
17–06904	SAMEA5769040	ERS3572586	PRJEB33238	Human	Blood	Epsilon1	90
18–00080	SAMEA5769102	ERS3572648	PRJEB33238	Human	CSF	Epsilon1	90
18–00304	SAMEA5769041	ERS3572587	PRJEB33238	Human	Blood	Epsilon1	6236
18–00445	SAMEA5769103	ERS3572649	PRJEB33238	Human	Unknown	Epsilon1	6331
18–01855	SAMEA5769042	ERS3572588	PRJEB33238	Human	Blood	Epsilon1	90
18–02683	SAMEA5769104	ERS3572650	PRJEB33238	Human	Blood	Epsilon1	4465
18–02987	SAMEA5769105	ERS3572651	PRJEB33238	Human	GS	Epsilon1	90
18–03576	SAMEA5769106	ERS3572652	PRJEB33238	Human	Unknown	Epsilon1	2981
18–03577	SAMEA5769107	ERS3572653	PRJEB33238	Human	GS	Epsilon1	2981
18–04116	SAMEA5769108	ERS3572654	PRJEB33238	Human	Blood	Epsilon1a	4465
18–04317	SAMEA5769109	ERS3572655	PRJEB33238	Human	CSF	Epsilon1a	7353
18–04364	SAMEA5769110	ERS3572656	PRJEB33238	Human	Blood	Epsilon1a	4465
18–04365	SAMEA5769043	ERS3572589	PRJEB33238	Human	Blood	Epsilon1a	4465
18–04414	SAMEA5769111	ERS3572657	PRJEB33238	Human	CSF	Epsilon1a	7353
18–04434	SAMEA5769112	ERS3572658	PRJEB33238	Human	Blood	Epsilon1a	4465
18–04472	SAMEA5769113	ERS3572659	PRJEB33238	Human	Blood	Epsilon1	6331
18–04499	SAMEA5769114	ERS3572660	PRJEB33238	Human	Blood	Epsilon1a	4465

Isolate ID	Sample accession no.	Secondary accession no.	Study number	Type	Source	Cluster	Complex type
18-04500	SAMEA5769115	ERS3572661	PRJEB33238	Human	Blood	Epsilon1a	4465
18-04539	SAMEA5769116	ERS3572662	PRJEB33238	Human	Blood	Epsilon1a	4465
18-04540	SAMEA5041142	ERS2852884	PRJEB29295	Human	Blood	Epsilon1a	4465
18-04543	SAMEA5769044	ERS3572590	PRJEB33238	Human	Blood	Epsilon1a	4465
18-04581	SAMEA5769045	ERS3572591	PRJEB33238	Human	Blood	Epsilon1a	4465
18-04652	SAMEA5769046	ERS3572592	PRJEB33238	Human	Blood	Epsilon1a	4465
18-04653	SAMEA5769047	ERS3572593	PRJEB33238	Human	Blood	Epsilon1a	7353
18-04654	SAMEA5769048	ERS3572594	PRJEB33238	Human	Blood	Epsilon1a	4465
18-04655	SAMEA5769049	ERS3572595	PRJEB33238	Human	Blood	Epsilon1a	4465
18-04657	SAMEA5769050	ERS3572596	PRJEB33238	Human	Blood	Epsilon1a	7353
18-04772	SAMEA5769051	ERS3572597	PRJEB33238	Human	Blood	Epsilon1a	4465
18-04825	SAMEA5769052	ERS3572598	PRJEB33238	Human	Blood	Epsilon1a	7353
18-04826	SAMEA5769053	ERS3572599	PRJEB33238	Human	Blood	Epsilon1a	4465
18-04827	SAMEA5769117	ERS3572663	PRJEB33238	Human	Blood	Epsilon1a	7353
18-04850	SAMEA5769118	ERS3572664	PRJEB33238	Human	Blood	Epsilon1a	4465
18-04852	SAMEA5769119	ERS3572665	PRJEB33238	Human	Blood	Epsilon1a	4465
18-04897	SAMEA5769120	ERS3572666	PRJEB33238	Human	Blood	Epsilon1a	7353
18-04898	SAMEA5769121	ERS3572667	PRJEB33238	Human	Blood	Epsilon1a	4465
18-04954	SAMEA5769122	ERS3572668	PRJEB33238	Human	Blood	Epsilon1a	4465
18-04955	SAMEA5769123	ERS3572669	PRJEB33238	Human	Blood	Epsilon1a	4465
18-05034	SAMEA5769124	ERS3572670	PRJEB33238	Human	Blood	Epsilon1	7451
18-05035	SAMEA5769125	ERS3572671	PRJEB33238	Human	Blood	Epsilon1a	7353
18-05038	SAMEA5769126	ERS3572672	PRJEB33238	Human	Blood	Epsilon1a	4465
18-05084	SAMEA5769127	ERS3572673	PRJEB33238	Human	Blood	Epsilon1a	4465
18-05142	SAMEA5769128	ERS3572674	PRJEB33238	Human	Unknown	Epsilon1a	4465
18-05143	SAMEA5769129	ERS3572675	PRJEB33238	Human	Unknown	Epsilon1a	7353
18-05144	SAMEA5769130	ERS3572676	PRJEB33238	Human	Blood	Epsilon1a	4465
18-05199	SAMEA5769131	ERS3572677	PRJEB33238	Human	Blood	Epsilon1a	4465
18-05201	SAMEA5769132	ERS3572678	PRJEB33238	Human	Blood	Epsilon1a	4465
18-05202	SAMEA5769133	ERS3572679	PRJEB33238	Human	Blood	Epsilon1a	4465
18-05203	SAMEA5769134	ERS3572680	PRJEB33238	Human	Blood	Epsilon1a	4465
18-05237	SAMEA5769135	ERS3572681	PRJEB33238	Human	Blood	Epsilon1a	4465
18-05327	SAMEA5769136	ERS3572682	PRJEB33238	Human	Blood	Epsilon1a	4465
18-05328	SAMEA5769137	ERS3572683	PRJEB33238	Human	Blood	Epsilon1a	4465
18-05329	SAMEA5769138	ERS3572684	PRJEB33238	Human	Blood	Epsilon1a	4465
18-05393	SAMEA5769139	ERS3572685	PRJEB33238	Human	CSF	Epsilon1a	4465
18-05394	SAMEA5769140	ERS3572686	PRJEB33238	Human	Blood	Epsilon1a	4465
18-05396	SAMEA5769141	ERS3572687	PRJEB33238	Human	Blood	Epsilon1a	7353
18-05398	SAMEA5769054	ERS3572600	PRJEB33238	Human	Blood	Epsilon1a	7353
18-05449	SAMEA5769142	ERS3572688	PRJEB33238	Human	Blood	Epsilon1a	4465
18-05450	SAMEA5769143	ERS3572689	PRJEB33238	Human	Blood	Epsilon1a	4465
18-05496	SAMEA5769144	ERS3572690	PRJEB33238	Human	Blood	Epsilon1a	4465
18-05542	SAMEA5769145	ERS3572691	PRJEB33238	Human	CSF	Epsilon1a	4465
18-05544	SAMEA5769146	ERS3572692	PRJEB33238	Human	Blood	Epsilon1a	4465
18-05558	SAMEA5769147	ERS3572693	PRJEB33238	Human	Blood	Epsilon1a	4465
18-05655	SAMEA5769148	ERS3572694	PRJEB33238	Human	Blood	Epsilon1a	4465
18-05657	SAMEA5769149	ERS3572695	PRJEB33238	Human	CSF	Epsilon1a	4465
18-05658	SAMEA5769150	ERS3572696	PRJEB33238	Human	Blood	Epsilon1a	4465
18-05660	SAMEA5769151	ERS3572697	PRJEB33238	Human	Blood	Epsilon1a	4465
18-05714	SAMEA5769152	ERS3572698	PRJEB33238	Human	Blood	Epsilon1a	4465
18-05726	SAMEA5769153	ERS3572699	PRJEB33238	Human	Blood	Epsilon1a	7353
18-05748	SAMEA5769154	ERS3572700	PRJEB33238	Human	Blood	Epsilon1a	4465
18-05767	SAMEA5769155	ERS3572701	PRJEB33238	Human	Blood	Epsilon1a	4465
18-05768	SAMEA5769156	ERS3572702	PRJEB33238	Human	CSF	Epsilon1a	4465
18-05836	SAMEA5769157	ERS3572703	PRJEB33238	Human	Unknown	Epsilon1a	7353
18-05837	SAMEA5769055	ERS3572601	PRJEB33238	Human	Blood	Epsilon1a	7353
18-05951	SAMEA5769158	ERS3572704	PRJEB33238	Human	Blood	Epsilon1a	4465
18-05970	SAMEA5769159	ERS3572705	PRJEB33238	Human	CSF	Epsilon1a	4465
18-06023	SAMEA5769056	ERS3572602	PRJEB33238	Human	Unknown	Epsilon1a	7353
18-06024	SAMEA5769057	ERS3572603	PRJEB33238	Human	Ascites	Epsilon1a	4465
18-06035	SAMEA5769058	ERS3572604	PRJEB33238	Human	Blood	Epsilon1a	4465
18-06036	SAMEA5769059	ERS3572605	PRJEB33238	Human	Blood	Epsilon1a	4465
18-06121	SAMEA5769060	ERS3572606	PRJEB33238	Human	Blood	Epsilon1a	4465
18-06126	SAMEA5769061	ERS3572607	PRJEB33238	Human	Blood	Epsilon1a	4465
18-06127	SAMEA5769062	ERS3572608	PRJEB33238	Human	CSF	Epsilon1a	4465
18-06128	SAMEA5769063	ERS3572609	PRJEB33238	Human	Blood	Epsilon1a	4465
18-06129	SAMEA5769064	ERS3572610	PRJEB33238	Human	Blood	Epsilon1a	4465
18-06130	SAMEA5769065	ERS3572611	PRJEB33238	Human	Blood	Epsilon1a	7353

Isolate ID	Sample accession no.	Secondary accession no.	Study number	Type	Source	Cluster	Complex type
18-06131	SAMEA5769066	ERS3572612	PRJEB33238	Human	CSF	Epsilon1a	4465
18-06138	SAMEA5769067	ERS3572613	PRJEB33238	Human	Blood	Epsilon1a	4465
18-06170	SAMEA5769160	ERS3572706	PRJEB33238	Human	Unknown	Epsilon1a	4465
18-06263	SAMEA5769161	ERS3572707	PRJEB33238	Human	Blood	Epsilon1a	4465
18-06331	SAMEA5769162	ERS3572708	PRJEB33238	Human	Blood	Epsilon1a	4465
18-06438	SAMEA5769163	ERS3572709	PRJEB33238	Human	Unknown	Epsilon1a	4465
18-06540	SAMEA5769164	ERS3572710	PRJEB33238	Human	Blood	Epsilon1a	4465
18-06541	SAMEA5769165	ERS3572711	PRJEB33238	Human	CSF	Epsilon1a	4465
18-06646	SAMEA5769166	ERS3572712	PRJEB33238	Human	CSF	Epsilon1a	4465
18-06680	SAMEA5769167	ERS3572713	PRJEB33238	Human	Blood	Epsilon1a	4465
18-06776	SAMEA5769168	ERS3572714	PRJEB33238	Human	Blood	Epsilon1a	4465
18-06820	SAMEA5769169	ERS3572715	PRJEB33238	Human	Unknown	Epsilon1a	7353
18-06916	SAMEA5769170	ERS3572716	PRJEB33238	Human	Blood	Epsilon1a	4465
18-06954	SAMEA5769068	ERS3572614	PRJEB33238	Human	Blood	Epsilon1a	7353
18-06955	SAMEA5769069	ERS3572615	PRJEB33238	Human	Blood	Epsilon1a	4465
18-07018	SAMEA5769070	ERS3572616	PRJEB33238	Human	Blood	Epsilon1a	4465
18-07037	SAMEA5769071	ERS3572617	PRJEB33238	Human	Blood	Epsilon1a	4465
18-07092	SAMEA5769072	ERS3572618	PRJEB33238	Human	Unknown	Epsilon1a	4465
18-07157	SAMEA5769073	ERS3572619	PRJEB33238	Human	Blood	Epsilon1a	4465
18-07158	SAMEA5769074	ERS3572620	PRJEB33238	Human	Blood	Epsilon1a	4465
18-07267	SAMEA5769171	ERS3572717	PRJEB33238	Human	Blood	Epsilon1a	4465
18-07300	SAMEA5769172	ERS3572718	PRJEB33238	Human	Unknown	Epsilon1a	7353
19-00076	SAMEA5769173	ERS3572719	PRJEB33238	Human	Blood	Epsilon1a	4465
19-00080	SAMEA5769174	ERS3572720	PRJEB33238	Human	Blood	Epsilon1	90
19-00082	SAMEA5769175	ERS3572721	PRJEB33238	Human	Unknown	Epsilon1a	4465
19-00149	SAMEA5769176	ERS3572722	PRJEB33238	Human	CSF	Epsilon1a	4465
19-00151	SAMEA5769177	ERS3572723	PRJEB33238	Human	Blood	Epsilon1a	4465
19-00179	SAMEA5769178	ERS3572724	PRJEB33238	Human	Unknown	Epsilon1a	4465
19-00191	SAMEA5769179	ERS3572725	PRJEB33238	Human	Blood	Epsilon1a	4465
19-00240	SAMEA5769180	ERS3572726	PRJEB33238	Human	Unknown	Epsilon1a	4465
19-00278	SAMEA5769181	ERS3572727	PRJEB33238	Human	Blood	Epsilon1a	4465
19-00281	SAMEA5769182	ERS3572728	PRJEB33238	Human	Blood	Epsilon1a	7353
19-00312	SAMEA5769075	ERS3572621	PRJEB33238	Human	Blood	Epsilon1a	7353
19-00347	SAMEA5769076	ERS3572622	PRJEB33238	Human	Blood	Epsilon1a	4465
19-00419	SAMEA5769077	ERS3572623	PRJEB33238	Human	Blood	Epsilon1a	4465
19-00444	SAMEA5769078	ERS3572624	PRJEB33238	Human	Blood	Epsilon1a	4465
19-00499	SAMEA5769079	ERS3572625	PRJEB33238	Human	Blood	Epsilon1a	4465
19-00500	SAMEA5769080	ERS3572626	PRJEB33238	Human	Blood	Epsilon1a	4465
19-00520	SAMEA5769081	ERS3572627	PRJEB33238	Human	Blood	Epsilon1a	4465
19-00549	SAMEA5769082	ERS3572628	PRJEB33238	Human	Blood	Epsilon1a	4465
19-00582	SAMEA5769083	ERS3572629	PRJEB33238	Human	Blood	Epsilon1a	7353
19-00609	SAMEA5769084	ERS3572630	PRJEB33238	Human	Unknown	Epsilon1a	7353
19-00973	SAMEA5769183	ERS3572729	PRJEB33238	Human	Unknown	Epsilon1a	4465
19-00974	SAMEA5769184	ERS3572730	PRJEB33238	Human	Blood	Epsilon1a	7353
19-00998	SAMEA5769185	ERS3572731	PRJEB33238	Human	Blood	Epsilon1a	4465
19-01023	SAMEA5769186	ERS3572732	PRJEB33238	Human	Blood	Epsilon1a	4465
19-01108	SAMEA5769187	ERS3572733	PRJEB33238	Human	Blood	Epsilon1a	4465
19-01166	SAMEA5769188	ERS3572734	PRJEB33238	Human	Placenta	Epsilon1	90
19-01173	SAMEA5769189	ERS3572735	PRJEB33238	Human	Blood	Epsilon1	90
19-01197	SAMEA5769190	ERS3572736	PRJEB33238	Human	Blood	Epsilon1a	4465
19-01319	SAMEA5769191	ERS3572737	PRJEB33238	Human	Synovia	Epsilon1a	4465
19-01387	SAMEA5769192	ERS3572738	PRJEB33238	Human	PF	Epsilon1a	4465
19-01604	SAMEA5769193	ERS3572739	PRJEB33238	Human	Wound	Epsilon1a	4465
19-01607	SAMEA5769194	ERS3572740	PRJEB33238	Human	BA	Epsilon1a	4465
19-01930	SAMEA5769195	ERS3572741	PRJEB33238	Human	Blood	Epsilon1a	4465
19-01961	SAMEA5769196	ERS3572742	PRJEB33238	Human	Blood	Epsilon1	90
19-02578	SAMEA5769085	ERS3572631	PRJEB33238	Human	Blood	Epsilon1a	7353
19-02579	SAMEA5769086	ERS3572632	PRJEB33238	Human	Blood	Epsilon1a	7353
19-02581	SAMEA5769087	ERS3572633	PRJEB33238	Human	Blood	Epsilon1a	4465
19-02587	SAMEA5769088	ERS3572634	PRJEB33238	Human	Blood	Epsilon1a	7353
19-02590	SAMEA5769089	ERS3572635	PRJEB33238	Human	CSF	Epsilon1a	4465
19-02598	SAMEA5769090	ERS3572636	PRJEB33238	Human	CSF	Epsilon1a	4465
19-02600	SAMEA5769091	ERS3572637	PRJEB33238	Human	Blood	Epsilon1a	4465
19-LI00135-0	SAMEA5769092	ERS3572638	PRJEB33238	Food	-	Epsilon1a	4465
19-LI00136-0	SAMEA5769093	ERS3572639	PRJEB33238	Food	-	Epsilon1a	4465
19-LI00137-0	SAMEA5769094	ERS3572640	PRJEB33238	Food	-	Epsilon1a	4465
19-LI00138-0	SAMEA5769095	ERS3572641	PRJEB33238	Food	-	Epsilon1a	4465
19-LI00175-0	SAMEA5769096	ERS3572642	PRJEB33238	Food	-	Epsilon1a	4465

Isolate ID	Sample accession no.	Secondary accession no.	Study number	Type	Source	Cluster	Complex type
------------	----------------------	-------------------------	--------------	------	--------	---------	--------------

*Isolates from nonsterile materials were not included in the outbreak description. BA, brain abscess; CSF, cerebrospinal fluid; GS, gynecological swab; PF, pleural fluid.

Appendix Table 2. Antimicrobial susceptibility in 79 isolates of *Listeria monocytogenes* from listeriosis outbreak cluster Epsilon1a, Germany*

Antimicrobial drug	MIC, mg/L (no. isolates)													
	0.003	0.06	0.125	0.25	0.5	1.0	2.0	4.0	8.0	16.0	32	64	128	>128
Ampicillin†	ND	–	–	S (29)	S (50)	I	–	–	–	ND	ND	ND	ND	ND
Benzylpenicillin†	ND	–	–	S (32)	S (43)	I	–	–	–	ND	ND	ND	ND	ND
Ceftriaxone‡	ND	ND	ND	ND	ND	I	I	R	R	R	R	R	R (6)	R (73)
Ciprofloxacin§	ND	ND	ND	–	–	I (67)	R (12)	–	–	ND	ND	ND	ND	ND
Daptomycin§	ND	ND	ND	ND	–	I	R (9)	R (68)	R (2)	–	ND	ND	ND	ND
Erythromycin†	ND	ND	ND	S (79)¶	–	I	–	–	–	ND	ND	ND	ND	ND
Gentamicin§	ND	ND	ND	ND	S (65)¶	I (14)	–	–	–	–	ND	ND	ND	ND
Linezolid#	ND	ND	ND	ND	–	–	I	I (79)	–	–	ND	ND	ND	ND
Meropenem**	ND	–	S (79)	I	–	–	–	–	–	ND	ND	ND	ND	ND
Rifampin††	ND	I (79)¶¶	I	I	–	–	–	ND	ND	ND	ND	ND	ND	ND
Tetracycline‡	ND	ND	ND	–	–	I (4)	I (75)	–	–	ND	ND	ND	ND	ND
Tigecycline‡‡	R	S (1)	S (77)	S (1)	I	–	–	–	ND	ND	ND	ND	ND	ND
Cotrimoxazole§§	S	I	–	–	–	–	–	–	ND	ND	ND	ND	ND	ND
Vancomycin‡	(79)¶¶	ND	ND	ND	ND	S (74)¶¶	I (5)	–	–	–	ND	ND	ND	ND

*All 79 strains were tested against 14 antimicrobial drugs. I, intermediate; R, resistant; S, susceptible; –, no isolates in this MIC category.

†EUCAST breakpoint 1.0 mg/mL for *Listeria monocytogenes*.

‡EUCAST breakpoint 2.0 mg/mL for *Streptococcus pneumoniae*.

§EUCAST breakpoint 1.0 mg/mL for *Staphylococcus aureus*.

¶No observable growth at the lowest tested concentration.


#EUCAST breakpoint 4.0 mg/mL for *Streptococcus pneumoniae*.

**EUCAST breakpoint 0.25 mg/mL for *Listeria monocytogenes*.


††EUCAST breakpoint 0.5 mg/mL for *Streptococcus pneumoniae*.

‡‡EUCAST breakpoint 0.5 mg/mL for *Staphylococcus aureus*.


§§Trimethoprim/sulfamethoxazole; EUCAST breakpoint 0.06 mg/mL for *Listeria monocytogenes*.




Appendix Figure 1. Number of notified listeriosis cases during the 2018–2019 outbreak in Germany compared with minimum, median, and maximum reported case numbers per week during the reference period, 2013–2017.



Appendix Figure 2. Maximum likelihood tree of *Listeria monocytogenes* Epsilon1a outbreak cluster, Germany. SNP calling demonstrated phylogenetic relatedness of isolates after read mapping to the genome of the *L. monocytogenes* serogroup IVb reference strain 10-092876-0769 LM12 (2). SNP, single nucleotide polymorphism. Scale bar indicates nucleotide substitutions per site.



Appendix Figure 3. Virulome analysis of *Listeria monocytogenes* Epsilon1 and Epsilon1a isolates. Assembled genome sequences were searched for genes belonging to LIPI-1, LIPI-3, or LIPI-4; encoding internalins; or involved in adhesion, invasion, intracellular survival, chitin hydrolysis, gene regulation, protein anchoring to the cell surface, peptidoglycan modification, immunomodulation, or bile resistance. Genomes of strains EGD-e (accession no. NC_003210.1), F2365 (accession no. NC_002973.6), and CLIP80459 (accession no. NC_012488.1) were included for comparison. The variability in the presence or absence of chromosomally encoded virulence genes within the highly clonal Epsilon1a cluster likely is not real, but instead results from acceptance bias due to cutoff values used during allele calling. adh, adhesion; anch, protein anchoring; chi, chitin hydrolysis; I, immunomodulation; LIPI, *Listeria* pathogenicity island; PM, peptidoglycan modification.



Appendix Figure 4. Resistome analysis of clinical *L. monocytogenes* Epsilon1 and Epsilon1a isolates. Genes are listed at the top. Assembled genome sequences were searched for genes known to confer resistance to sanitizers, heavy metals, and antimicrobial drugs by using a task template in SeqSphere (Ridom, <https://www.ridom.de>). Genomes of strains EGD-e (accession no. NC_003210.1), L2624 (accession no. NZ_CP007686), FORC_049 (accession no. NZ_CP016629), 6179 (accession no. NZ_HG813249), LM201 (accession no. SAMN03267130), 2012–0070 (accession no. SAMN05715911), NCTC 10887 (accession no. SAMN06434199), 10–092876–0731 LM5 (NZ_CP019618), and 12754_4#74 (accession no. ERR564017), and pLMST6 (accession no. GCA_900164035.1) were included for comparison. Of note, the *emrC* gene is located on plasmid pLMST6 that occurs in certain ST6 strains (2). Plasmid loss might explain why it was not detected throughout the entire Epsilon1a population. cap, chloramphenicol; cfx, ceftriaxone; erm, erythromycin; flu, fluoroquinolones; fos, fosfomycin; lin, lincomycin; QACs, quaternary ammonium compounds; sm, streptomycin; tet, tetracycline, tmx, trimethoprim.