

# COVID-19 Outbreak on a Passenger Ship and Assessment of Response Measures, Greece, 2020

Sophia Hatzianastasiou,<sup>1</sup> Varvara A. Mouchtouri,<sup>1</sup> Androula Pavli, Maria Tseroni, Spyros Sapounas, Charalampos Vasileiou, Katerina Dadouli, Maria Kyritsi, Michalis Koureas, Panagiotis Prezerakos, Matthaios Speletas, Georgios Panagiotakopoulos, Sotirios Tsiodras,<sup>2</sup> Christos Hadjichristodoulou<sup>2</sup>

We describe response measures to an outbreak involving 128 (33.4%) coronavirus disease cases (46.1% asymptomatic) among 383 persons onboard a passenger ship. Multivariate analysis indicated that dining in certain rooms and bar areas, nationality, working department (for crew members), and quarantining onboard the ship were significantly associated with infection.

On March 7, 2020, a passenger ship (2,500-passenger and 1,606-bed capacity) with 33 crew members sailed from Piraeus, Greece, to Cesme, Turkey, where an additional 350 crew members embarked on March 8, 2020 (1). For 21 days, the ship sailed without any disembarkations or embarkations until the first suspected coronavirus disease (COVID-19) case was reported to the health authority of the Piraeus port on March 28, 2020. We describe results of the outbreak investigation, including risk factors for transmission of severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2).

## The Study

We collected data by completing standardized forms through interviews and medical examination of all

Author affiliations: Hellenic National Public Health Organization, Athens, Greece (S. Hatzianastasiou, A. Pavli, M. Tseroni, S. Sapounas, G. Panagiotakopoulos); University of Thessaly Faculty of Medicine, Larisa, Greece (V.A. Mouchtouri, K. Dadouli, M. Kyritsi, M. Koureas, M. Speletas, C. Hadjichristodoulou); European Union Healthy Gateways Joint Action, Larisa (V.A. Mouchtouri, C. Hadjichristodoulou); Department of Hygiene Inspections of the Port of Piraeus, Region of Attica, Athens (C. Vasileiou); University of Peloponnese Department of Nursing, Tripoli, Greece (P. Prezerakos); University of Patras Faculty of Medicine, Patra, Greece (G. Panagiotakopoulos); National and Kapodistrian University Medical School, Athens (S. Tsiodras)

DOI: <https://doi.org/10.3201/eid2707.210398>

travelers onboard and by reviewing the ship records and logs. We used descriptive statistics to analyze the study variables and performed univariate and multivariate analyses.

In conducting clinical management of cases, we followed guidelines from the Hellenic National Public Health Organization (NPHO) for health measures on travelers and repatriation, which were based on the European Union Healthy Gateways Joint Action advice for management of COVID-19 cases onboard ships (2) (Appendix Table 1, <https://wwwnc.cdc.gov/EID/article/27/7/21-0398-App1.pdf>). NPHO and the Piraeus Port Health Authority provided passengers with information about using medical facemasks at all times when outside of their cabins, as well as handwashing, physical distancing, and cleaning and disinfecting cabins; ship officers supervised.

Food preparation and laundry and cleaning services were halted; travelers were instructed to clean their own cabins and store used linens in plastic bags. Cleaning and disinfection of the terminal was done by a private company under supervision of the Piraeus Port Health Authority, after all travelers disembarked the ship at the port of Piraeus. A catering company provided packaged meals; personal hygiene supplies were also provided (including facemasks and hand sanitizer). Methods and results of the environmental sampling have been published elsewhere (3).

We collected oropharyngeal specimens from all travelers onboard. Molecular tests for SARS-CoV-2 detection were performed by using the Cobas SARS-CoV-2 test qualitative assay and the Cobas 6800/8800 System (La Roche, <https://www.roche.com>). Serologic tests were performed on blood

<sup>1</sup>These first authors contributed equally to this article.

<sup>2</sup>These senior authors contributed equally to this article..

specimens collected from 116 cases. Serum samples were initially tested with the Xiamen Boson Biotech (<https://www.bosonbio.com>) Rapid 2019-nCoV IgG/IgM Combo Test Card, a rapid lateral flow (immunochromatographic) test, and subsequently with the MAGLUMI800 chemiluminescence immunoassay (Snibe Diagnostic, <https://www.snibe.com>).

Our study was a public health response as part of activities of the Hellenic NPHO and local authorities (i.e., Piraeus Port Health Authority and Port Administration). Participants provided verbal informed consent for recording and processing of information during interviews, and written consent was obtained from participants for blood specimen analysis. All required ethics considerations were applied according to rules of the Hellenic NPHO and the Ministry of Health.

The first 3 symptomatic cases occurred on March 20 among travelers (passengers and crew) of different nationalities and working departments (hotel, dining room service, and housekeeping [cabin steward]). The peak of the outbreak occurred during March 30–April 1 (Figure). We conducted laboratory tests for SARS-CoV-2 and for antibodies during 3 follow-up examinations (Appendix Table 2).

Travelers who tested positive were isolated onboard (except the first case-patients, who were hospitalized, and 2 travelers who were isolated in hotels designated by the government of Greece for that purpose). All travelers onboard who tested negative were considered contacts and quarantined individually in quarantine facilities ashore (hotels designated by the government of Greece), except 36 crew members who tested negative but quarantined in separate decks and facilities onboard to ensure safe ship operation. We compiled characteristics of travelers,


hospitalizations, and quarantine measures (Appendix Table 2), symptom frequency (Appendix Table 3), and results of univariate analysis for testing risk factors (Appendix Table 4). No deaths occurred; 7 patients were hospitalized, including the first patient, who was intubated.

We conducted multivariate analysis, in the form of binary logistic regression, using SPSS 25.0 (IBM, <https://www.ibm.com>). For all analyses, we used a 5% significance level. Multivariate analysis results indicated that test-negative travelers quarantined at hotels had lower odds of SARS-CoV-2 infection than those who quarantined onboard the ship (odds ratio 0.07, 95% CI 0.01–0.58). Travelers of nationality A who worked in the entertainment department had higher odds of infection (odds ratio 3.54, 95% CI 1.03–12.16). Multivariate analysis indicated that dining in certain rooms and bar areas, nationality, working department (for crew members), and quarantine onboard the ship significantly associated with infection (Table).

## Conclusions

Our findings can be used in COVID-19 prevention and control preparedness plans for ports and ships. Ongoing transmission can occur onboard ships, affecting a large number of travelers, without any sign of symptoms for an extended period, especially in cases where most of the travelers are of young age. Assuming a serial interval number of 5 days and a reproduction number of 2.6, we could conclude that 1 infectious traveler embarked on the first or second day of the voyage, and 21 days later 120 travelers onboard had been infected (4,5).

Active or passive surveillance for COVID-19 based on symptoms onboard ships alone cannot



**Figure.** Epidemic curve for symptomatic coronavirus disease case-patients, by date of symptom onset, during an outbreak on a passenger ship, Greece, March 20–April 18, 2020. A total of 65 case-patients had a known date of symptom onset.

**Table.** Multivariate analysis of risk factors for SARS-CoV-2 infection during an outbreak on a passenger ship, Greece, March 20–April 18, 2020\*

Factor	Area of work (for crew members)			Consuming meals and drinks in galley/dining area/bar A‡
	Food and beverage	Housekeeping and hotel	Other†	
Nationality				
A			Referent	
B	0.00 (0.00–0.00)	NA	NA	NA
C	1.20 (0.51–2.81)	<b>0.11 (0.02–0.75)</b>	<b>0.03 (0.002–0.42)</b>	<b>0.49 (0.24–0.99)</b>
D	0.81 (0.36–1.83)	0.23 (0.05–1.14)	NA	0.54 (0.27–1.08)
E	0.90 (0.25–3.21)	0.96 (0.10–9.27)	NA	0.94 (0.32–2.78)
Other†	1.51 (0.41–5.62)	1.02 (0.14–7.20)	0.22 (0.02–2.98)	1.52 (0.59–3.91)
Consuming meals and drinks in galley/dining area/bar A‡	Referent	<b>2.71 (1.34–5.46)</b>	<b>4.44 (1.81–10.95)</b>	NA

\*A total of 120 passengers and crew tested positive during the first specimen collection and testing. Values in bold type are statistically significant. SARS-CoV-2, severe acute respiratory syndrome coronavirus 2; NA, not applicable.

†Entertainment, security, shop, hospital, other.

‡Meals and beverages were prepared and served in galley and dining room and café-bar A (for travelers who embarked at Cesme port) and in galley and dining room and café-bar B (for travelers who embarked at Piraeus port).

be effective for early COVID-19 outbreak detection because only a small proportion of cases can be identified when the disease has already spread widely. Screening of incoming travelers and pre-boarding and regular routine laboratory testing can be implemented in addition to surveillance (6). We advise laboratory diagnostic testing of passengers and crew for SARS-CoV-2 within 72 hours before embarkation and a second test the day of embarkation. In addition, for crew members, we advise a 10-day quarantine period before starting work and regular testing of all crew members on board every 7 days. The preparedness plans of ships and ports should ensure laboratory capacity for diagnosis of all persons onboard once an outbreak has been identified and frequent and regular testing until negative results are obtained in accordance with port policies (7).

Because of the large proportion of travelers who tested positive in this outbreak, the relevant authorities decided that COVID-19 case-patients would be isolated onboard the ship, whereas travelers who tested negative would disembark and quarantine individually in hotels. Individual quarantine and isolation in separate facilities was effective in preventing further spread (2,6,8–11). These measures contributed to preventing intra-cabin transmission, which was documented in other COVID-19 outbreaks onboard passenger ships (12,13). Halting food preparation and service, housekeeping activities in cabins, and laundry service stopped transmission onboard. This outcome is contrary to the outbreak management approach taken onboard another cruise ship, where crew continued working duties while passengers and crew were isolated in their cabins, resulting in further COVID-19 spread (11). In our study, logistic regression statistical analysis showed that quarantining travelers in hotels was a protective factor against

SARS-CoV-2 infection compared with quarantining onboard the ship.

Crew members who worked in the entertainment department, were of a certain nationality, and consumed meals and drinks at specific dining and bar areas had a higher risk for infection. Dining areas and bars can be settings for COVID-19 transmission because of congregation of persons and because facemasks are not used while eating or drinking (13). Avoiding self-service, encouraging service of meals in cabins, staggering meal times, and reconfiguring dining room seating to ensure physical distancing are recommended to avoid possible transmission within ships’ dining rooms (2,11,14).

A coordinated approach from a country’s central government, in cooperation with local port authorities, is needed to define the maximum response capacities of each port and the maximum number of ships that can be allowed to call. This approach will help to avoid confining travelers on ships for quarantine and isolation and reduce the risk for spread onboard when outbreaks occur, in line with the World Health Organization’s International Health Regulations, which are designed to prevent unnecessary interference with international traffic and trade (7).

**Acknowledgments**

We acknowledge the contribution of the Ministry of Health’s COVID-19 Taskforce, the Ministry of Shipping and Island Policy, the Hellenic National Public Health Organization, the General Secretariat for Civil Protection; the National Center for Blood Donation in Athens, the Laboratory of Hygiene and Epidemiology of the University of Thessaly, the Department of Hygiene Inspections of the Port of Piraeus, Region of Attica, Athens, the custom authorities, and the Piraeus Port Authority. Moreover, we thank the ship medical doctor and all ship officers and crew members for their contribution.

## About the Author

Dr. Hatzianastasiou is an internal medicine and infectious disease physician who works for the Department of Travel Medicine of the Hellenic National Public Health Organization. Her primary research interests include travel medicine. Dr. Mouchtouri is associate professor in the Department of Hygiene and Epidemiology of the University of Thessaly's Faculty of Medicine. She works for the European Union's Healthy Gateways Joint Action. Her primary research interests include the prevention and control of cross-border health threats.

## References

1. MarineTraffic. 2021. Eleftherios Venizelos travel information [in Greek] [cited 2021 Mar 1]. [https://www.marinetraffic.com/el/ais/details/ships/shipid:208272/mmsi:237628000/imo:7907673/vessel:el\\_venizelos](https://www.marinetraffic.com/el/ais/details/ships/shipid:208272/mmsi:237628000/imo:7907673/vessel:el_venizelos)
2. European Union Health Gateways. Advice for ship operators for preparedness and response to the outbreak of COVID-19. 2020 Feb 20 [cited 2021 Mar 1]. [https://www.healthygateways.eu/Portals/0/plcdocs/EU\\_HEALTHY\\_GATEWAYS\\_COVID-19\\_MARITIME\\_20\\_2\\_2020\\_FINAL.pdf](https://www.healthygateways.eu/Portals/0/plcdocs/EU_HEALTHY_GATEWAYS_COVID-19_MARITIME_20_2_2020_FINAL.pdf)
3. Mouchtouri VA, Koureas M, Kyritsi M, Vontas A, Kourentis L, Sapounas S, et al. Environmental contamination of SARS-CoV-2 on surfaces, air-conditioner and ventilation systems. *Int J Hyg Environ Health*. 2020;230:113599. <https://doi.org/10.1016/j.ijheh.2020.113599>
4. Du Z, Xu X, Wu Y, Wang L, Cowling BJ, Meyers LA. Serial interval of COVID-19 among publicly reported confirmed cases. *Emerg Infect Dis*. 2020;26:1341-3. <https://doi.org/10.3201/eid2606.200357>
5. Zhang S, Diao M, Yu W, Pei L, Lin Z, Chen D. Estimation of the reproductive number of novel coronavirus (COVID-19) and the probable outbreak size on the Diamond Princess cruise ship: a data-driven analysis. *Int J Infect Dis*. 2020;93:201-4. <https://doi.org/10.1016/j.ijid.2020.02.033>
6. European Union Health Gateways. Interim advice for restarting cruise ship operations after lifting restrictive measures in response to the COVID-19 pandemic. 2020 Jun 30 [cited 2021 Mar 1]. [https://www.healthygateways.eu/Portals/0/plcdocs/EU\\_HEALTHY\\_GATEWAYS\\_COVID-19\\_RESTARTING\\_CRUISES.pdf](https://www.healthygateways.eu/Portals/0/plcdocs/EU_HEALTHY_GATEWAYS_COVID-19_RESTARTING_CRUISES.pdf)
7. World Health Organization. International Health Regulations [cited 2021 Mar 1]. <https://www.who.int/health-topics/international-health-regulations>
8. Rocklöv J, Sjödin H, Wilder-Smith A. COVID-19 outbreak on the Diamond Princess cruise ship: estimating the epidemic potential and effectiveness of public health countermeasures. *J Travel Med*. 2020;27:27. <https://doi.org/10.1093/jtm/taaa030>
9. Gupta A, Kunte R, Goyal N, Ray S, Singh K. A comparative analysis of control measures on-board ship against COVID-19 and similar novel viral respiratory disease outbreak: quarantine ship or disembark suspects? *Med J Armed Forces India*. 2020. <https://doi.org/10.1016/j.mjafi.2020.06.003>
10. Mouchtouri VA, Dirksen-Fischer M, Hadjichristodoulou C. Health measures to travellers and cruise ships in response to COVID-19. *J Travel Med*. 2020;27:27. <https://doi.org/10.1093/jtm/taaa043>
11. World Health Organization. Interim guidance for operational considerations for managing COVID-19 cases/outbreak on board ships 2020 [cited 2021 Mar 1]. <https://www.who.int/publications-detail/operational-considerations-for-managing-covid-19-cases-outbreak-on-board-ships>.
12. Plucinski MM, Wallace M, Uehara A, Kurbatova EV, Tobolowsky FA, Schneider ZD, et al. COVID-19 in Americans aboard the Diamond Princess cruise ship. *Clin Infect Dis*. 2020;ciaa1180. <https://doi.org/10.1093/cid/ciaa1180>
13. Kakimoto K, Kamiya H, Yamagishi T, Matsui T, Suzuki M, Wakita T. Initial investigation of transmission of COVID-19 among crew members during quarantine of a cruise ship – Yokohama, Japan, February 2020. *MMWR Morb Mortal Wkly Rep*. 2020;69:312-3. <https://doi.org/10.15585/mmwr.mm6911e2>
14. Centers for Disease Control and Prevention. Interim guidance for ships on managing suspected or confirmed cases of coronavirus disease 2019 (COVID-19) 2021 [cited 2021 Mar 1]. <https://www.cdc.gov/quarantine/maritime/recommendations-for-ships.html>

---

Address for correspondence: Christos Hadjichristodoulou, Laboratory of Hygiene and Epidemiology, Faculty of Medicine, University of Thessaly, 22 Papakyriazi St, 41222, Larissa, Greece; email: xhatzi@uth.gr

# COVID-19 Outbreak on a Passenger Ship and Assessment of Response Measures, Greece, 2020

## Appendix

**Appendix Table 1.** Risk categorization of COVID-19 patients according to travelers' characteristics\*

Risk category†	Characteristics (demographics, signs and symptoms)
Low risk	Asymptomatic or Mild symptoms (fever <38.50 C and/or cough and/or sore throat) and No underlying conditions and Age <65 y and Respiratory rate ≤16/min or SaO2 >94%
Moderate risk	Symptoms: fever <38.50°C and/or cough and/or sore throat and Underlying condition or Age >65 y or Abnormal chest x-ray and Respiratory rate ≤16/min or SaO2 >94%
High risk	Symptoms: fever ≥38.5°C and/or cough and/or tiredness and/or dyspnea and Underlying condition or Age over >65 y or Abnormal chest x-ray

\*The ship medical doctor performed daily health monitoring of all persons on board, while four visits for medical examinations and specimen collection from all persons on board were conducted by the public health authority's team. A second medical and nursing team performed health monitoring of travelers quarantined in hotels.

†125 /128 patients were categorized into the low risk group, while two and one were categorized into the moderate and high risk group, respectively.

**Appendix Table 2.** Characteristics of symptomatic and asymptomatic travelers on board the passenger ship and laboratory results

Characteristics	Frequency (%)
Age, Male/Female, Total	
20–30	156 (52.9), 44 (50.6), 200 (52.4)
31–40	67 (22.7), 25 (28.7), 92 (24.1)
41–50	40 (13.6), 14 (16.1), 54 (14.1)
51–60	27 (9.2), 3 (3.4), 30 (7.9)
61–70	4 (1.4), 1 (1.1), 5 (1.3)
71–80	1 (0.3), 0 (0.0), 1 (0.3)
Total	295 (100.0), 87 (100.0), 382 (100.0)
Mean age (min-max, SD)	33.2 (19–73, 11.5), 31.8 (20–64, 9.8), 32.9 (19–73, 11.1)
Sex	
Female	87 (22.7)
Male	296 (77.3)
Total	383 (100)
Nationality	
A	152 (39.7)
B	36 (9.4)
C	73 (19.1)
D	82 (21.4)
E	17 (4.4)
Other	23 (6.0)
Total	383 (100.0)
Embarkation port	
Piraeus	33 (8.6)
Cesme	350 (91.4)
Total	383 (100.0)
Working department	
Food and beverage	211 (55.1)
Housekeeping and Hotel	103 (26.9)
Deck and Engine	31 (8.1)
Other*	38 (9.9)
Total	383 (100.0)
SARS-CoV-2 RT-PCR test results	
Positive	128 (33.4)
Negative	255 (66.6)
Total	383 (100.0)
Positive SARS-CoV-2 RT-PCR test results	
1 <sup>st</sup> examination 30–31/03/2020	120 (93.7)

Characteristics	Frequency (%)
2 <sup>nd</sup> examination 14/04/2020	45† (35.1)
3 <sup>rd</sup> examination 21–22/04/2020	16 (12.5)
4 <sup>th</sup> examination 04/05/2020	1 (0.8)
Total	128 (100.0)
IgG antibodies test results according to the number of days passed since the onset of symptoms	
Asymptomatic	19/55
Symptomatic <5 d	1/1
Symptomatic 6–10 d	4/21
Symptomatic 11–15 d	17/30
Symptomatic >15 d	5/9
Total	46/116 (39.6)
IgM antibodies test results according to the number of days passed since the onset of symptoms‡	
Asymptomatic	11/55
Symptomatic <5 d	1/1
Symptomatic 6–10 d	1/21
Symptomatic 11–15 d	5/30
Symptomatic >15 d	4/9
Total	22/116 (19.0)
Symptom status	
Symptomatic	69 (53.9)
Asymptomatic	59 (46.1)
Total	128 (100.0)
Hospitalizations	
	7 (5.5)
Deaths	
	0 (0.0)
Place of quarantine for contacts	
Ship	36 (13.7)
Hotel	227 (86.3)
Total	263 (100.0)
Place of isolation for cases	
Ship	117 (98.3)
Hotel	2 (1.7)
Total	119 (100.0)

\*Entertainment (N = 23), security (N = 8), shop (N = 4), supernumerary (N = 2), hospital (N = 1)

†Including 8 new cases

‡Clinical specimens were collected on April 10 2020.

**Appendix Table 3.** Symptoms frequency (N = 69)

Symptoms	Frequency	% Proportion (95%CI)
Low-grade fever (<38.5°C)	22	31.9 (21.2–44.2)
Fever (≥38.5°C)	15	21.7 (12.7–33.3)
Cough	23	33.3 (22.4–45.7)
Sore throat	12	17.4 (9.3–28.4)
Headache	12	17.4 (9.3–28.4)
Muscle aches and pains	10	14.5 (7.2–25.0)
Dyspnea with strenuous exercise	0	0 (0.0–5.2)
Dyspnea with no exercise	1	1.5 (0.0–7.8)
Diarrhea	5	7.3 (2.4–16.1)
Abdominal pain	3	4.4 (0.9–12.2)
Vomiting	0	0 (0.0–5.2)
Anosmia	14	20.3 (11.6–31.7)
Ageusia	1	1.5 (0.0–7.8)
Other	9	13.0 (6.1–23.3)

**Appendix Table 4.** Testing for risk factors of becoming infected using univariate analysis\* (N = 120 who tested positive during the first specimen collection)

Factors	Relative Risk (95%CI)	p value
Gender (Male/ Female)	1.18 (0.81–1.72)	0.392
Work position		
Bar service	Reference	
Dining room service	<b>2.33 (1.16–4.69)</b>	<b>0.007</b>
Cook	1.45 (0.70–3.01)	0.299
Cabin steward	0.54 (0.19–1.52)	0.234
Utility	0.50 (0.14–1.72)	0.310
Department of work		
Housekeeping and Hotel	Reference	
Food and beverage	<b>2.41 (1.49–3.91)</b>	<b>&lt;0.001</b>
Deck and Engine	1.04 (0.41–2.61)	0.999
Other†	<b>3.39 (1.97–5.83)</b>	<b>&lt;0.001</b>
Nationality		
B	Reference	
A	<b>3.08 (1.34–7.09)</b>	<b>0.001</b>
C	1.58 (0.63–3.97)	0.317
D	1.49 (0.60–3.73)	0.380
E	2.54 (0.90–7.17)	0.073
Other	<b>3.44 (1.37–8.63)</b>	<b>0.004</b>
Nationality and work department		
Food and beverage & Nationality A	Reference	
Entertainment & Nationality A	<b>1.78 (1.16–2.73)</b>	<b>0.042</b>
Food and beverage & Nationality D	0.81 (0.47–1.39)	0.552
Housekeeping & Nationality D	<b>0.29 (0.12–0.67)</b>	<b>0.001</b>
Food and beverage & Nationality C	1.00 (0.61–1.64)	0.999
Food and beverage & Nationality E	0.86 (0.37–1.97)	0.767
Port of embarkation (Cesme/ Piraeus)	2.17 (0.95–4.93)	0.018
Galley and dining place A/ Galley and dining place B	<b>2.39 (1.04–5.46)</b>	<b>0.018</b>

\*Means and standard deviations (SD) were reported for continuous variables, while frequencies and proportions with 95% Confidence Intervals (CI) were reported for categorical variables. Proportions' 95% CI were calculated based on binomial distribution. Data were also checked for deviation from normal distribution using the Shapiro-Wilk normality test. Categorical data were analyzed with the use of Chi-square tests and Fisher exact tests as appropriate; Student's t-tests and Mann-Whitney U tests were used for continuous data as appropriate. Univariate analysis was conducted by calculating the relative risk (RR) of becoming infected with SARS-CoV-2 with the following factors: gender, age, work position, department of work, nationality, port of embarkation, dining area, place of quarantine, and deck number.

†Entertainment, security, shop, supernumerary, hospital.