

About the Author

Miss Sun is a postgraduate student at The University of Hong Kong, Hong Kong, China. Her primary research interest is bioinformatics.

References

1. Webby R, Richt J. Influenza in swine. In: Webster RG, Monto AS, Braciale TJ, Lamb RA, editors. Textbook of influenza, 2nd edition. Chichester: Wiley Blackwell; 2013. p. 190–202.
2. Smith GJ, Vijaykrishna D, Bahl J, Lycett SJ, Worobey M, Pybus OG, et al. Origins and evolutionary genomics of the 2009 swine-origin H1N1 influenza A epidemic. *Nature*. 2009;459:1122–5. <https://doi.org/10.1038/nature08182>
3. Chauhan RP, Gordon ML. A systematic review analyzing the prevalence and circulation of influenza viruses in swine population worldwide. *Pathogens*. 2020;9:E355. <https://doi.org/10.3390/pathogens9050355>
4. Shi W, Li W, Li X, Haywood J, Ma J, Gao GF, et al. Phylogenetics of varied subtypes of avian influenza viruses in China: potential threat to humans. *Protein Cell*. 2014;5:253–7. <https://doi.org/10.1007/s13238-014-0036-1>
5. Gu H, Xie R, Adam DC, Tsui JL, Chu DK, Chang LDJ, et al. Genomic epidemiology of SARS-CoV-2 under an elimination strategy in Hong Kong. *Nat Commun*. 2022;13:736. <https://doi.org/10.1038/s41467-022-28420-7>
6. Al-Ghadeer H, Chu DKW, Rihan EMA, Abd-Allah EA, Gu H, Chin AWH, et al. Circulation of influenza A(H5N8) virus, Saudi Arabia. *Emerg Infect Dis*. 2018;24:1961–4. <https://doi.org/10.3201/eid2410.180846>
7. He P, Wang G, Mo Y, Yu Q, Xiao X, Yang W, et al. Novel triple-reassortant influenza viruses in pigs, Guangxi, China. *Emerg Microbes Infect*. 2018;7:85. <https://doi.org/10.1038/s41426-018-0088-z>
8. Cui H, Che G, de Jong MCM, Li X, Liu Q, Yang J, et al. The PB1 gene from H9N2 avian influenza virus showed high compatibility and increased mutation rate after reassorting with a human H1N1 influenza virus. *Virology*. 2022;19:20. <https://doi.org/10.1186/s12985-022-01745-x>
9. Sun H, Xiao Y, Liu J, Wang D, Li F, Wang C, et al. Prevalent Eurasian avian-like H1N1 swine influenza virus with 2009 pandemic viral genes facilitating human infection. *Proc Natl Acad Sci U S A*. 2020;117:17204–10. <https://doi.org/10.1073/pnas.1921186117>
10. Cheung JTL, Tsang TK, Yen HL, Perera RAPM, Mok CKP, Lin YP, et al. Determining existing human population immunity as part of assessing influenza pandemic risk. *Emerg Infect Dis*. 2022;28:977–85. <https://doi.org/10.3201/eid2805.211965>

Address for correspondence: Malik Peiris or Leo Poon, School of Public Health, The University of Hong Kong, 7 Sassoon Road, Hong Kong, China; email: malik@hku.hk or llmpoon@hku.hk

Persistent SARS-CoV-2 Alpha Variant Infection in Immunosuppressed Patient, France, February 2022

Slim Fourati, Guillaume Gautier, Myriam Chovelon, Alexandre Soulier, Melissa N'Debi, Vanessa Demontant, Céline Kennel, Christophe Rodriguez, Jean-Michel Pawlotsky

Author affiliations: Hôpital Henri Mondor (AP-HP), Université Paris-Est, Créteil, France (S. Fourati, A. Soulier, M. N'Debi, V. Demontant, C. Rodriguez, J.-M. Pawlotsky); William Morey General Hospital, Chalon-sur-Saône, France (G. Gautier, M. Chovelon, C. Kennel)

DOI: <https://doi.org/10.3201/eid2807.220467>

We describe persistent circulation of SARS-CoV-2 Alpha variant in an immunosuppressed patient in France during February 2022. The virus had a new pattern of mutation accumulation. The ongoing circulation of previous variants of concern could lead to reemergence of variants with the potential to propagate future waves of infection.

Immunosuppressed patients can have prolonged SARS-CoV-2 infection (1). Studies have reported the occurrence and selection of multiple mutations in the spike glycoprotein sequence in immunosuppressed patients with persistent SARS-CoV-2 infections (2–6). To date, intrahost mutations have been described essentially in the ancestral wild-type SARS-CoV-2 virus (3,5–8), especially during prolonged infection with variants of concern (VOCs) (9). Additional SARS-CoV-2 mutations in immunocompromised persons could enable increased virus transmissibility and immune evasion, shaping the emergence of new VOCs. We describe a new mutation accumulation pattern in SARS-CoV-2 Alpha virus in an immunosuppressed patient.

An 84-year-old woman with evolutive mantle cell lymphoma who was receiving maintenance rituximab and lenalidomide treatment was admitted to the hospital on May 17, 2021. She had asthenia, fever, and hypoxia (93% oxygen saturation). At admission (day 0), she tested positive for SARS-CoV-2 RNA (Figure). She had received 2 vaccine doses 84 and 66 days before admission. She did not have respiratory symptoms, but a chest computed tomography scan showed ground-glass opacities in her lungs. The patient was hospitalized and treated with corticosteroids for 10 days. She tested SARS-CoV-2-positive again on August 26, day 101 after her initial

positive test. On September 27 (day 133), she tested SARS-CoV-2-negative and was considered virologically cured. She received a vaccine booster (third)

dose at day 164 and a fourth dose on day 201, but we did not detect spike receptor binding domain antibodies at days 133, 201, or 210.




Figure. Timeline of SARS-CoV-2 diagnostic tests, hospitalizations, booster vaccination, and treatments for an immunosuppressed patient with persistent SARS-CoV-2 Alpha variant infection, France, 2022. RCHOP, combination therapy of rituximab, cyclophosphamide, doxorubicine, vincristine, and prednisone; RDHAC, combination therapy of rituximab, cytarabine, dexamethasone, and carboplatine.

Beginning in October 2021, the patient experienced intermittent fever and diarrhea because her lymphoma progressed, as shown by an abdominal computed tomography scan; she was started on a second-line treatment with ibrutinib (Figure). She was hospitalized again in early January 2022 (day 208) after clinicians discovered an evolutive lymph node mass. At admission, she tested SARS-CoV-2–positive despite the absence of COVID-19 symptoms. SARS-CoV-2 RNA levels from nasopharyngeal swab samples were high; cycle threshold values were 19 on day 208, 18 on day 228, and 22 on day 234.

At days 0, 101, 208, and 234, multiplex mutation-specific reverse transcription PCR revealed the absence of spike amino acid mutations E484Q, E484K, and L452R, and K417N was not detected on days 208 and 234, suggesting that the patient was not infected with Delta or Omicron variants, the 2 dominant variants in France at the time. These results suggested that the patient was infected with the Alpha variant, which was the dominant variant circulating when she first tested positive.

To determine whether the patient was infected with a new strain or reinfected with the same persistently replicating variant, we performed whole-genome sequencing on samples collected on days 208 (January 11, 2022) and 228 (January 31, 2022) and identified lineage B.1.1.7 (Alpha variant). Older samples were not available for sequencing. Our analysis revealed the presence of amino acid substitutions and mutations in addition to characteristics of the Alpha variant, including mutations in open reading frame (ORF) 1a (nonstructural protein 3, $n = 4$), spike protein ($n = 6$), matrix protein ($n = 1$), envelope protein ($n = 2$), ORF3a ($n = 1$), and ORF7 ($n = 2$) (Appendix Figure, <https://wwwnc.cdc.gov/EID/article/28/7/22-0467-App1.pdf>).

The SARS-CoV-2 mutational pattern in this immunosuppressed patient adds several new mutations to the Alpha variant characteristic. Although earlier samples were not available for sequencing, mutations in later samples align with an ongoing selection process. The mutations we observed share similarities with those observed in other VOCs and variants of interest, pointing to evolutionary convergence, such as spike del241–247, which also is found in part in the Beta variant. Several mutations that likely play a role in immune evasion were selected in the spike nucleocapsid terminal domain (e.g., K77E, S248F, and del14–18) and receptor-binding domain (L452M). These mutations have rarely been reported in isolates submitted to GISAID (<https://www.gisaid.org>),

suggesting that, when considered individually, they could be maladaptive.

In January 2022, the Alpha variant was no longer circulating in France, according to strains submitted to GISAID. Our case highlights the potential for persistence of supposedly extinct SARS-CoV-2 variants that might cause prolonged infection in immunocompromised patients and acquire adaptive mutations that confer increased transmissibility, antigenic divergence, and reduced pathogenicity, with obvious public health implications (1,3). Similar cases likely exist in other parts of the world because SARS-CoV-2 genome sequencing and reporting to GISAID are far from exhaustive.

Omicron-infected patients not immunized against older variants appear to mount a weak or no neutralizing response against variants that preceded Omicron, including VOCs (R.K. Suryawanshi et al., unpub. data, <https://doi.org/10.1101/2022.01.13.22269243>). Because the next dominant variant could emerge from a variant other than Omicron, ongoing circulation of older VOCs could feed reemergence of variants that eliminate Omicron, particularly in unvaccinated populations, emphasizing the crucial role of vaccination to prevent new SARS-CoV-2 waves.

In conclusion, this report highlights the need to reinforce precautions to avert nosocomial and community transmission involving immunocompromised patients, who might shed older SARS-CoV-2 variants longer. Prospective genomic surveillance for SARS-CoV-2 variants is needed in persons with prolonged infection, particularly in countries with many immunocompromised persons, such as countries with a high HIV prevalence and low vaccination rates.

Funding for this study was provided by the ministry of health of France as part of the surveillance program for SARS-CoV-2 genomic evolution.

S.F. has served as a speaker for GlaxoSmithKline, Abbvie, and Abbott Diagnostics. C.R. has served as an advisor and speaker for Illumina and Vela Diagnostics. J.-M.P. has served as an advisor and speaker for Abbvie, Gilead, Merck, Assembly Biosciences, and Arbutus. The remaining authors have no conflicts of interest to disclose.

About the Author

Prof. Fourati is a researcher at the University of East Paris, Créteil, France. He is also a clinical virologist in the Department of Microbiology, Henri Mondor Hospital, Créteil, France. His primary research interests include viral evolution and adaptation, viral hepatitis, and respiratory viruses, including SARS-CoV-2.

References

1. Corey L, Beyrer C, Cohen MS, Michael NL, Bedford T, Rolland M. SARS-CoV-2 variants in patients with immunosuppression. *N Engl J Med*. 2021;385:562–6. <https://doi.org/10.1056/NEJMs2104756>
2. Wang Y, Wang D, Zhang L, Sun W, Zhang Z, Chen W, et al. Intra-host variation and evolutionary dynamics of SARS-CoV-2 populations in COVID-19 patients. *Genome Med*. 2021;13:30. <https://doi.org/10.1186/s13073-021-00847-5>
3. Khateeb D, Gabrieli T, Sofer B, Hattar A, Cordela S, Chaouat A, et al. SARS-CoV-2 variants with reduced infectivity and varied sensitivity to the BNT162b2 vaccine are developed during the course of infection. *PLoS Pathog*. 2022;18:e1010242. <https://doi.org/10.1371/journal.ppat.1010242>
4. Weigang S, Fuchs J, Zimmer G, Schnepf D, Kern L, Beer J, et al. Within-host evolution of SARS-CoV-2 in an immunosuppressed COVID-19 patient as a source of immune escape variants. *Nat Commun*. 2021;12:6405. <https://doi.org/10.1038/s41467-021-26602-3>
5. Cele S, Karim F, Lustig G, San JE, Hermanus T, Tegally H, et al.; COMMIT-KZN Team. SARS-CoV-2 prolonged infection during advanced HIV disease evolves extensive immune escape. *Cell Host Microbe*. 2022;30:154–162.e5. <https://doi.org/10.1016/j.chom.2022.01.005>
6. Choi B, Choudhary MC, Regan J, Sparks JA, Padera RF, Qiu X, et al. Persistence and evolution of SARS-CoV-2 in an immunocompromised host. *N Engl J Med*. 2020;383:2291–3. <https://doi.org/10.1056/NEJMc2031364>
7. Pérez-Lago L, Aldámiz-Echevarría T, García-Martínez R, Pérez-Latorre L, Herranz M, Sola-Campoy PJ, et al.; on behalf of Gregorio Marañón Microbiology-Id Covid Study Group. Different within-host viral evolution dynamics in severely immunosuppressed cases with persistent SARS-CoV-2. *Biomedicines*. 2021;9:808. <https://doi.org/10.3390/biomedicines9070808>
8. Avanzato VA, Matson MJ, Seifert SN, Pryce R, Williamson BN, Anzick SL, et al. Case study: prolonged infectious SARS-CoV-2 shedding from an asymptomatic immunocompromised individual with cancer. *Cell*. 2020;183:1901–12.e9. <https://doi.org/10.1016/j.cell.2020.10.049>
9. Truffot A, Andréani J, Le Maréchal M, Caporossi A, Epaulard O, Germe R, et al. SARS-CoV-2 variants in immunocompromised patient given antibody monotherapy. *Emerg Infect Dis*. 2021;27:2725–8. <https://doi.org/10.3201/eid2710.211509>

Address for correspondence: Slim Fourati, Henri Mondor Hospital, 51 av du Marechal de Lattre de Tassigny, Créteil 94010, France; email: slim.fourati@aphp.fr

Increased Stability of SARS-CoV-2 Omicron Variant over Ancestral Strain

Alex Wing Hong Chin, Alison Man Yuk Lai, Malik Peiris, Leo Lit Man Poon

Author affiliations: The University of Hong Kong, Hong Kong, China (A.W.H. Chin, A.M.Y. Lai, M. Peiris, L.L.M. Poon); Centre for Immunology & Infection, Hong Kong (A.W.H. Chin, M. Peiris, L.L.M. Poon)

DOI: <https://doi.org/10.3201/eid2807.220428>

As of April 2022, the Omicron BA.1 variant of concern of SARS-CoV-2 was spreading quickly around the world and outcompeting other circulating strains. We examined its stability on various surfaces and found that this Omicron variant is more stable than its ancestral strain on smooth and porous surfaces.

The Omicron SARS-CoV-2 variant of concern (VOC) is highly transmissible in humans. As of April 2022, it has outcompeted other known variants and dominated in different regions (1). Its spike protein has >30 mutations compared with the ancestral strain (2). A 2022 structural study indicates the Omicron spike protein is more stable than that of the ancestral strain (3); this finding prompted us to hypothesize that Omicron VOC is also more stable on different surfaces. We previously showed that the ancestral SARS-CoV-2 strain can still be infectious at room temperature for several days on smooth surfaces and several hours on porous surfaces (4).

We used previously described ancestral SARS-CoV-2 (PANGO lineage A) and Omicron VOC (PANGO lineage BA.1) in this study (5,6). We tested their stability on different surfaces using our previously described protocol (4,7). In brief, we applied a 5- μ L droplet of each virus (10^7 50% tissue culture infectious dose [TCID₅₀]/mL) on different surfaces in triplicate. We incubated the treated surfaces at room temperature (21°C–22°C) for different time points as indicated and then immersed them in viral transport medium for 30 min to recover the residual infectious virus. We titrated the recovered virus by TCID₅₀ assays using Vero E6 cells, as described (4,7).

Compared with the ancestral SARS-CoV-2, the Omicron BA.1 variant was more stable on all surfaces we studied (Table). On day 4 postinoculation, we recovered no infectious ancestral SARS-CoV-2 from stainless steel, polypropylene sheet, or 2 of 3 glass

Persistent SARS-CoV-2 Alpha Variant Infection in Immunosuppressed Patient, France, February 2022


Appendix

Methods

We used GSD NovaType IV SARS-CoV-2 (Gold Standard Diagnostics, <https://www.goldstandarddiagnostics.com>), a multiplex mutation-specific reverse transcription PCR (RT-PCR) kit based on melting curve analyses, to search for the presence of Spike amino acid mutations E484Q, E484K, L452R, and beginning in January 2022, for K417N.

Full-length SARS-CoV-2 genome sequence analysis was performed from nasopharyngeal swab samples by using the COVIDSeq Test (Illumina, <https://www.illumina.com>). The sequences were demultiplexed and assembled as full-length genomes with the DRAGEN COVIDSeq Test Pipeline on a DRAGEN server (Illumina). Lineages were interpreted with Pangolin (<https://github.com/cov-lineages/pangolin>) and clades with Nextclade (Nextstrain, <https://clades.nextstrain.org>). All sequences were submitted to GISAID (<https://www.gisaid.org>) database and to GenBank (<https://www.ncbi.nlm.nih.gov/genbank>) under accession no. SUB11349926.

Our study protocol followed the ethical guidelines of the Declaration of Helsinki and was approved by our institutional review board. The patient provided informed consent.



Appendix Figure. SARS-CoV-2 genome mutational patterns relative to an Alpha variant consensus in an immunosuppressed patient with persistent SARS-CoV-2 Alpha variant infection, France, 2022. Sequence data are from nasopharyngeal samples collected at days 208 and 228. Black boxes indicate Alpha variant-defining mutations according to GISAID (<https://www.covariants.org/variants/20I.Alpha.V1>). Grey boxes indicate mutations other than those in the Alpha variant sequence. Red boxes indicate mutations in the patient's sequences at days 208 and 228. AF, allele frequency; E, extinction gene; M, matrix gene; N, nucleocapsid gene; ORF, open reading frame; S, spike gene.