

Pseudomonas guariconensis Necrotizing Fasciitis, United Kingdom

Edward J. Moseley, Jian Cheng Zhang,
O. Martin Williams

Author affiliations: UK Health Security Agency Specialist Microbiology and Laboratories, South West Region and Severn Infection Sciences, Bristol, UK (E.J. Moseley, J.C. Zhang, O.M. Williams); University Hospitals Bristol and Weston NHS Foundation Trust, Bristol (E.J. Moseley, O.M. Williams); Southmead Hospital, Bristol (J.C. Zhang)

DOI: <https://doi.org/10.3201/eid3001.231192>

We describe a case of necrotizing fasciitis in the United Kingdom in which *Pseudomonas guariconensis* was isolated from multiple blood culture and tissue samples. The organism carried a Verona integron-encoded metallo- β -lactamase gene and evidence of decreased susceptibility to β -lactam antimicrobial agents. Clinicians should use caution when treating infection caused by this rare pathogen.

A 67-year-old man in the United Kingdom was seen in the emergency department for right lower leg pain and swelling with associated fevers lasting 24 hours. He reported a right heel blister had formed 1 week earlier, after he purchased new footwear. His medical history included obesity, hypertension, atrial fibrillation, and left ventricular systolic dysfunction. He was a former smoker and had a 40 pack-year history.

On examination, the patient appeared alert and comfortable. He was febrile (38.0°C), tachycardic (124 beats/min) in atrial fibrillation, and had a stable blood pressure (108/72 mm Hg). His respiratory rate was 20 breaths/min, and oxygen saturation was 92% on room air. He had right leg swelling and erythema,

extending from the blister on his heel to his mid-calf. Blood test results showed leukocyte count was 15.03×10^9 cells/L (reference range 4.0–11.0 cells/L), neutrophils 13.44 (reference 1.5–8.0) $\times 10^9$ cells/L, and lymphocytes 0.38 (reference 1.0–4.0) $\times 10^9$ cells/L. Bilirubin was 41 (reference <21) $\mu\text{mol/L}$, albumin 34 (reference 35–50) g/L, and C-reactive protein 20 (reference <6.0) mg/L.

The patient was started on intravenous flucloxacillin (1g 4 \times /d) for lower limb cellulitis. Aerobic blood culture samples at admission were positive at 11.5 hours' incubation and cultures collected 8 hours after admission positive at 10.5 hours' incubation by BD Bactec FX system (Becton Dickinson, <https://www.bd.com>). Gram stain from the samples showed gram-negative bacilli, resulting in an immediate change of therapy to intravenous amoxicillin/clavulanic acid (1.2 g 3 \times /d) and gentamicin (400 mg; 5 mg/kg based on patient's ideal bodyweight). Direct extract from the first sample was tested by matrix-assisted laser desorption/ionization time-of-flight (MALDI-TOF) mass spectroscopy (Bruker Corporation, <https://www.bruker.com>), which identified *Pseudomonas guariconensis* with a score of 1.94 within 3 hours of the initial culture report. Growth on plates from the second blood culture was subsequently confirmed to be the same organism. Samples were sent to the UK Health Security Agency Antimicrobial Resistance and Healthcare Infection reference laboratory, which also confirmed *P. guariconensis* by MALDI-TOF mass spectroscopy with a score of 2.62.

The patient's treatment was changed to intravenous piperacillin/tazobactam (4.5 g 4 \times /d); gentamicin was continued. On review, no local features of necrotizing fasciitis were observed, and his leg appeared improved. The patient reported that he had been applying several over-the-counter creams of uncertain age to his blister since it had developed.

Table. Antimicrobial susceptibility data for blood culture isolates in a case of *Pseudomonas guariconensis* necrotizing fasciitis, United Kingdom*

Antimicrobial agent	Admission blood culture by ViTek 2† (MIC, mg/L)	8-h blood culture		Tissue culture by disc diffusion‡
		Disc diffusion‡	Gradient strip (MIC, mg/L)	
Ceftazidime	I (4)	I	ND	I
Ciprofloxacin	I (≤ 0.25)	I	ND	I
Gentamicin	S (≤ 1)	S	ND	S
Meropenem	I (4)	I	I (4)	I
Tobramycin	S (≤ 1)	S	ND	S
Piperacillin/tazobactam	R (64)	I	I (16)	I
Amikacin	S (≤ 2)	ND	ND	ND
Cefepime	I (2)	ND	ND	ND
Imipenem	I (1)	ND	ND	ND
Levofloxacin	I (0.5)	ND	ND	ND
Ticarcillin/clavulanic acid	R (≥ 128)	ND	ND	ND

*All breakpoints are European Committee on Antimicrobial Susceptibility Testing (EUCAST) clinical breakpoints version 11.0, except gentamicin disk diffusion testing, for which EUCAST version 9.0 was used. I, susceptible, increased exposure; ND, not done; R, resistant; S, susceptible.

†bioMérieux, <https://www.biomerieux.com>.

‡Using methods from EUCAST (1).

Attempts were made to recover the creams for culture but were unsuccessful.

Direct disk susceptibility testing was performed by using European Committee on Antimicrobial Susceptibility Testing (EUCAST) rapid antimicrobial susceptibility testing (AST) methodology, reading plates at 16–18 hours, which showed piperacillin/tazobactam susceptibility. Although unvalidated for this organism, AST suggested piperacillin/tazobactam susceptibility at increased exposure compared with EUCAST rapid AST breakpoints for *P. aeruginosa* and standard AST clinical breakpoints for *Pseudomonas* spp. (EUCAST criteria version 11.0), which was confirmed by standard EUCAST disk diffusion testing (1) from the second isolate. The patient's gentamicin was stopped. However, he remained tachycardic and hypotensive.

Confirmatory AST performed by using the VITEK 2 system and software version 9.02 (bioMérieux, <https://www.biomerieux.com>) produced a piperacillin/tazobactam MIC of 64 mg/L and meropenem MIC of 4 mg/L (Table). Piperacillin/tazobactam MIC by gradient strip testing performed on the second isolate was increased at 16 mg/L, particularly close to the EUCAST breakpoint, and meropenem MIC was increased at 4 mg/L. The patient's therapy was changed to 2 g intravenous ceftazidime (2 g 3×/d).

In-house multiplex PCR was performed using agarose gel electrophoresis for beta-lactamase genes (Appendix, <https://wwwnc.cdc.gov/EID/article/30/1/23-1192-App1.pdf>), based on previously published methodology (2–5). PCR detected a Verona integron-encoded metallo-β-lactamase enzyme, consistent with previously reported strains of this species (6,7).

The patient subsequently deteriorated and required inotropic and vasopressor support. He underwent above-knee amputation and debridement after fasciotomies, and exploration confirming necrotizing fasciitis. Tissue samples isolated pure growth *P. guariconensis*, and sensitivity testing by standard EUCAST disk methodology was consistent with previous samples (Table).

The patient remained in the critical care unit for 3 days and had high vasopressor requirements despite adequate antimicrobial drug therapy. He was deemed not stable for further surgery; life-sustaining treatment was withdrawn, and he died.

P. guariconensis is a gram-negative, strictly aerobic, non-spore-forming, rod-shaped bacterium that is motile by means of 2 polar flagella, is oxidase and catalase positive, and is indole and aesculin negative. *P. guariconensis* was described in 2013, isolated from rhizospheric soil of *Vigna unguiculata* (L.) Walp. (the cowpea) in Guárico, Venezuela (8). Isolates of the same species pro-

ducing novel carbapenemases have been reported from environmental samples taken in the Amazon Basin (9).

Reports of *P. guariconensis* human disease are rare; 1 case of infective endocarditis was reported in a patient with underlying lupus erythematosus (10). The rarity of reports likely reflects the recent description of the species and delays in updates to identification methodologies, such as MALDI-TOF databases. This case shows the pathogenic potential of *P. guariconensis* in an immunocompetent host and the degree of clinical suspicion required to exclude deep infection when isolating an unusual organism from a sterile site.

About the Author

Dr. Mosley is a specialty trainee in infectious diseases and medical microbiology in Bristol, United Kingdom. His research interests include antimicrobial stewardship, point of care testing, and medical education.

References

- Matuschek E, Brown DFJ, Kahlmeter G. Development of the EUCAST disk diffusion antimicrobial susceptibility testing method and its implementation in routine microbiology laboratories. *Clin Microbiol Infect*. 2014;20:O255–66. <https://doi.org/10.1111/1469-0691.12373>
- Hidalgo L, Hopkins KL, Gutierrez B, Ovejero CM, Shukla S, Douthwaite S, et al. Association of the novel aminoglycoside resistance determinant RmtF with NDM carbapenemase in Enterobacteriaceae isolated in India and the UK. *J Antimicrob Chemother*. 2013;68:1543–50. <https://doi.org/10.1093/jac/dkt078>
- Ellington MJ, Kistler J, Livermore DM, Woodford N. Multiplex PCR for rapid detection of genes encoding acquired metallo-beta-lactamases. *J Antimicrob Chemother*. 2006;59:321–2. <https://doi.org/10.1093/jac/dkl481>
- Poirel L, Héritier C, Tolün V, Nordmann P. Emergence of oxacillinase-mediated resistance to imipenem in *Klebsiella pneumoniae*. *Antimicrob Agents Chemother*. 2004;48:15–22. <https://doi.org/10.1128/AAC.48.1.15-22.2004>
- Yigit H, Queenan AM, Anderson GJ, Domenech-Sanchez A, Biddle JW, Steward CD, et al. Novel carbapenem-hydrolyzing β-lactamase, KPC-1, from a carbapenem-resistant strain of *Klebsiella pneumoniae*. *Antimicrob Agents Chemother*. 2001;45:1151–61. <https://doi.org/10.1128/AAC.45.4.1151-1161.2001>
- Public Health England. UK standards for microbiology investigations 60: detection of bacteria with carbapenem-hydrolyzing β-lactamases (carbapenemases). London: Public Health England; 2020.
- Adelowo OO, Vollmers J, Mäusezahl I, Kaster AK, Müller JA. Detection of the carbapenemase gene bla_{VM-5} in members of the *Pseudomonas putida* group isolated from polluted Nigerian wetlands. *Sci Rep*. 2018;8:15116. <https://doi.org/10.1038/s41598-018-33535-3>
- Toro M, Ramírez-Bahena MH, Cuesta MJ, Velázquez E, Peix A. *Pseudomonas guariconensis* sp. nov., isolated from rhizospheric soil. *Int J Syst Evol Microbiol*. 2013;63:4413–20. <https://doi.org/10.1099/ijs.0.051193-0>
- Souza CO, Cayó R, Lima KVB, Brasiliense DM, Streling AP, Siqueira AV, et al. Genetic and biochemical characterization

of BIM-1, a novel acquired subgroup B1 MBL found in a *Pseudomonas* sp. strain from the Brazilian Amazon region. *J Antimicrob Chemother.* 2023;78:1359–66. <https://doi.org/10.1093/jac/dkad077>

10. Okano H, Okado R, Ito H, Asakawa H, Nose K, Tsuruga S, et al. Ischemic hepatitis with infectious endocarditis: A case report. *Biomed Rep.* 2021;15:97. <https://doi.org/10.3892/br.2021.1473>

Address for correspondence: Edward J. Moseley, Health Security Agency Specialist Microbiology and Laboratories, South West Region and Severn Infection Sciences, University Hospitals Bristol and Weston NHS Foundation Trust, Bristol Royal Infirmary, Zone A Queens Bldg Level 8, Upper Maudlin Street, Bristol, BS2 8HW, England; email: edward.moseley@uhbw.nhs.uk

Rare *Spiroplasma* Bloodstream Infection in Patient after Surgery, China, 2022

Ningning Xiu,¹ Chao Yang,¹ Xiaowei Chen, Jianping Long, Pinghua Qu

Author affiliations: Dongguan Kanghua Hospital, Dongguan, China (N. Xiu, J. Long); Second Clinical Medical College of Guangzhou University of Chinese Medicine, Guangzhou, China (C. Yang, X. Chen, P. Qu); The Second Affiliated Hospital of Guangzhou University of Chinese Medicine, Guangdong Provincial Hospital of Traditional Chinese Medicine, Guangzhou (P. Qu)

DOI: <http://doi.org/10.3201/eid3001.230858>

We report a case of *Spiroplasma* bloodstream infection in a patient in China who developed pulmonary infection, acute respiratory distress syndrome, sepsis, and septic shock after emergency surgery for type A aortic dissection. One organism closely related to *Spiroplasma eriocheiris* was isolated from blood culture and identified by whole-genome sequencing.

Spiroplasma, a genus of bacteria in the phylum *Mycoplasmata*, is characterized by cell structures with no cell walls (1). *Spiroplasma* isolates have been primarily reported from plants, guts of insects, tick

tritulates, and crustaceans (2), although a few cases of *Spiroplasma* infection in humans have also been reported, causing cataracts and uveitis in infants and systemic infections in immunocompromised patients (3–6). We describe a rare *Spiroplasma* bloodstream infection in a patient after surgery for type A aortic dissection in China.

The case-patient, a 68-year-old man, underwent surgery to repair his aorta on June 3, 2022, and he developed a severe respiratory infection afterward while still hospitalized. Fibrobronchoscopy revealed extensive and severe airway erosion, with yellow and thick sputum adhering to the airway walls. A biopsy of a bronchial embolism was taken and sent for examination (Appendix Figure, panel A, <https://wwwnc.cdc.gov/EID/article/30/1/23-0858-App1.pdf>), and microscopic observation revealed a layered arrangement of thrombi mixed with neutrophils (Appendix Figure, panel B). On June 9, 2022, the patient's health began to deteriorate (Appendix Table). The patient was diagnosed with pulmonary infection, acute respiratory distress syndrome, sepsis, and septic shock.

Medical staff performed multiple tests on the patient to identify an infectious etiology to explain the patient's acute illness (Table). *Candida tropicalis* was cultured in bronchoalveolar lavage fluid (BALF) samples. Seven of 12 blood cultures tested positive (Bactec FX; Becton Dickinson, <https://www.bd.com/en-us>) for a microorganism that was isolated as rare colonies under conditions of 35°C and a 5% CO₂ atmosphere. Subcultures on Columbia blood agar showed pinpoint-size zones of hemolysis with no macroscopic colony growth at 4 days of incubation; however, Gram stain and Giemsa-Wright stain of the blood could not detect the presence of bacteria. Finally, metagenomic next-generation sequencing was performed on both the blood and BALF samples. Unique reads of *Spiroplasma eriocheiris* (n = 1,577 in BALF, n = 2,344 in blood), human alphaherpesvirus 1 (n = 66,185 in BALF, n = 1,942 in blood), and *Aspergillus fumigatus* (n = 7 in BALF, n = 12 in blood) were detected (Table). We have uploaded raw data to the National Center for Biotechnology Information Sequence Read Archive (BioProject no. PRJNA1021328).

We characterized the cultivated microorganism, designated DGKH1, by 16S rRNA gene sequencing and whole-genome sequencing analysis. Results of 16S rRNA gene phylogeny show DGKH1 is closely related to *S. eriocheiris* CCTCC M 207170^T (Figure). However, the average nucleotide identity value between the genomes of the 2 isolates was 94%, and the average digital DNA–DNA hybridization value between them was 56%, both of which were lower than

¹These authors contributed equally to this article.