

Appendix Table 1. Antigenic characterization of influenza (H3N2) viruses isolated from pigs*

Virus	Hyperimmune rabbit serum, titer				
	A/JXdh/312/2006	Sw/GD/7/2006	Sw/GD/211/2006	Sw/GD/968/2006	Dk/BJ/40/2004
A/JXdh/312/06 (H3N2)	640	160	≤10	≤10	≤10
Sw/GD/7/06 (H3N2)	160	1,280	≤10	≤10	≤10
Sw/SD/133/07 (H3N2)	160	640	≤10	≤10	≤10
Sw/SD/106/07 (H3N2)	160	640	≤10	≤10	≤10
Sw/FJ/43/07 (H3N2)	160	640	≤10	≤10	≤10
Sw/GD/211/06 (H3N2)	≤10	≤10	640	640	40
Sw/GD/968/06 (H3N2)	≤10	≤10	640	640	20
Sw/GD/223/06 (H3N2)	≤10	≤10	320	640	40
Sw/GD/423/06 (H3N2)	≤10	≤10	320	320	≤10
Sw/GD/811/06 (H3N2)	≤10	≤10	320	320	≤10
Dk/BJ/40/04 (H3N8)	≤10	≤10	≤10	≤10	1,280

*JXdh, Jiangxidonghu; sw, swine; GD, Guangdong; dk, duck; BJ, Beijing; SD, Shandong; FJ, Fujian. **Boldface** indicates hemagglutination-inhibition titers with homologous strains. Titers expressed as the reciprocal of the dilution end point given complete inhibition of hemagglutination.