


Estimated Number of AIDS Cases and Deaths among Adults and Adolescents with AIDS, 1985–2006—United States and Dependent Areas



Note. Data have been adjusted for reporting delays.




Reported AIDS Cases and Deaths, Cumulative through 2006—United States and Dependent Areas

| | Cases | Deaths | |
|------------------------|---------|---------|----|
| | | No. | % |
| Adults and adolescents | 983,343 | 544,015 | 55 |
| Children (<13 years) | 9,522 | 5,579 | 59 |
| Total | 992,865 | 549,594 | 55 |




Reported AIDS Cases, by Age and Sex, Cumulative through 2006—United States and Dependent Areas



* Excludes 3 persons of unknown sex.




Estimated Number of Adults and Adolescents Living with AIDS by Sex, 1993–2006—United States and Dependent Areas



Note. Data have been adjusted for reporting delays.




Estimated Number of Persons Living with AIDS, by Race/Ethnicity 1993–2006—United States and Dependent Areas



Note. Data have been adjusted for reporting delays.




Estimated Number of Asians/Pacific Islanders and of American Indians/Alaska Natives Living with AIDS 1993–2006—United States and Dependent Areas



Note. Data have been adjusted for reporting delays.




Estimated Number of Adults and Adolescents Living with AIDS by Region, 1993–2006—50 States and DC



Note. Data have been adjusted for reporting delays.




Proportion of AIDS Cases among Adults and Adolescents by Race/Ethnicity and Year of Diagnosis 1985–2006—United States and Dependent Areas



Note. Data have been adjusted for reporting delays.



Proportion of AIDS Cases among Adults and Adolescents by Transmission Category and Year of Diagnosis 1985–2006—United States and Dependent Areas




Note. Data have been adjusted for reporting delays and cases without risk factor information were proportionally redistributed.

* Heterosexual contact with a person known to have, or to be at high risk for, HIV infection.



Proportion of AIDS Cases among Adults and Adolescents by Sex and Transmission Category, Diagnosed in 2006—50 States and DC




Note. Data have been adjusted for reporting delays and cases without risk factor information were proportionally redistributed.

*Heterosexual contact with a person known to have, or to be at high risk for, HIV infection.

† Includes hemophilia, blood transfusion, perinatal exposure, and risk factor not reported or not identified.




AIDS Rates, Reported in 2006—United States and Dependent Areas





Proportion of AIDS Cases and Population by Race/Ethnicity, Reported in 2006—50 States and DC



* Includes 357 persons of unknown race or multiple races.



Estimated Number of AIDS Cases and Rates for Male Adults and Adolescents, by Race/Ethnicity 2006—50 States and DC

| <u>Race/Ethnicity</u> | <u>Cases</u> | <u>Rate (cases per 100,000 population)</u> |
|---|--------------|--|
| White, not Hispanic | 9,267 | 11.2 |
| Black, not Hispanic | 11,540 | 82.9 |
| Hispanic | 5,388 | 31.3 |
| Asian/Pacific Islander | 423 | 7.5 |
| <u>American Indian/ Alaska Native</u> | <u>118</u> | <u>12.2</u> |
| Total* | 26,989 | 22.4 |

Note. Data have been adjusted for reporting delays.

* Includes 254 male adults and adolescents of unknown race or multiple races.



Estimated Number of AIDS Cases and Rates for Female Adults and Adolescents, by Race/Ethnicity 2006—50 States and DC

| <u>Race/Ethnicity</u> | <u>Cases</u> | <u>Rate (cases per 100,000 population)</u> |
|-----------------------------------|--------------|--|
| White, not Hispanic | 1,659 | 1.9 |
| Black, not Hispanic | 6,391 | 40.4 |
| Hispanic | 1,516 | 9.5 |
| Asian/Pacific Islander | 95 | 1.6 |
| American Indian/ Alaska Native | 37 | 3.6 |
| <u>Total*</u> | <u>9,801</u> | <u>7.8</u> |



Note. Data have been adjusted for reporting delays.

* Includes 103 female adults and adolescents of unknown race or multiple races.





Proportion of AIDS Cases among Adults and Adolescents by Region and Size of Place of Residence Diagnosed in 2006—50 States and DC

| Region* | Metropolitan population | | Non-metropolitan area |
|------------------------|-------------------------|----------------|-----------------------|
| | >500,000 | 50,000–500,000 | |
| | % | % | % |
| Northeast N = 9,481 | 88 | 7 | 4 |
| Midwest N = 4,160 | 79 | 12 | 9 |
| South N = 17,085 | 76 | 12 | 11 |
| West N = 6,065 | 89 | 7 | 4 |

Note. Data have been adjusted for reporting delays.

* Region totals include persons whose metropolitan area of residence is unknown or missing.





Estimated Number and Proportion of AIDS Cases among Adults and Adolescents, by Transmission Category Diagnosed in 2006—50 States and DC

| <u>Transmission category</u> | <u>No.</u> | <u>%</u> |
|-------------------------------------|------------|----------|
| Male-to-male sexual contact | 16,001 | 43 |
| Injection drug use (IDU) | 6,795 | 18 |
| Male-to-male sexual contact and IDU | 1,803 | 5 |
| High-risk heterosexual contact* | 11,754 | 32 |
| <u>Other/not identified†</u> | <u>437</u> | 1 |
| Total | 36,790 | |



Note. Data have been adjusted for reporting delays and cases without risk factor information were proportionally redistributed.

* Heterosexual contact with a person known to have, or to be at high risk for, HIV infection.

† Includes hemophilia, blood transfusion, perinatal exposure, and risk factor not reported or not identified





Estimated Number and Proportion of AIDS Cases Associated with Injection Drug Use, by Transmission Category Diagnosed in 2006—United States and Dependent Areas

| | <u>No.</u> | <u>%</u> |
|--|--------------|----------|
| Heterosexual male injection drug user (IDU) | 4,686 | 44 |
| Female IDU | 2,467 | 23 |
| Male-to-male sexual contact and IDU | 1,844 | 17 |
| Heterosexual sex partners of IDUs | 1,703 | 16 |
| Children whose mothers are IDUs or sex partners of IDUs | 6 | <1 |
| <hr/> Total | <hr/> 10,706 | |



Note. Data have been adjusted for reporting delays and cases without risk factor information were proportionally redistributed.



Estimated Number and Proportion of AIDS Cases Associated with Injection Drug Use*, by Race/Ethnicity Diagnosed in 2006—United States and Dependent Areas

| <u>Race/Ethnicity</u> | <u>Cases</u> | <u>%</u> |
|-------------------------------|---------------|----------|
| White, not Hispanic | 2,665 | 25 |
| Black, not Hispanic | 5,366 | 50 |
| Hispanic | 2,385 | 22 |
| Asian/Pacific Islander | 105 | 1 |
| American Indian/Alaska Native | 61 | 1 |
| <u>Total[†]</u> | <u>10,706</u> | |

Note. Data have been adjusted for reporting delays and cases without risk factor information were proportionally redistributed.

* Includes injection drug users (IDUs), men who have sex with men and inject drugs, heterosexual sex partners of IDUs and children whose mothers are IDUs or sex partners of IDUs.

† Total includes 124 persons of unknown race or multiple races.





Reported AIDS Cases, by Age Group at Diagnosis Cumulative through 2006—United States and Dependent Areas

| Age (years) | AIDS | |
|--------------|----------------|----|
| | No. | % |
| <13 | 9,534 | <1 |
| 13–14 | 1,087 | <1 |
| 15–24 | 42,180 | 4 |
| 25–34 | 320,502 | 32 |
| 35–44 | 387,902 | 39 |
| 45–54 | 166,814 | 17 |
| 55–64 | 49,491 | 5 |
| ≥ 65 | 15,355 | 2 |
| Total | 992,865 | |













