


HIV AMONG YOUNG GAY AND BISEXUAL MEN

Young gay and bisexual men are severely affected by HIV...

About **1 in 5** (19%) of all new infections is among young men who have sex with men (MSM)




Young black MSM are most affected, accounting for **55%** of new infections among young MSM overall



8,800 total new HIV infections among MSM aged 13-24

...and new infections are on the rise

New infections increased **22%** in recent years among MSM aged 13-24



Many young gay and bisexual men are unaware of their HIV infection



In a 20-city study, half of HIV-positive young MSM aged 18-24 were unaware of their infection

1 in 4 not tested

In a 5-city study, 27% of young MSM aged 18-22 had not been tested in the last year, as CDC recommends



Combination of factors places young gay and bisexual men at increased risk for HIV



We can all take a stand against HIV

GET TESTED

GET TALKING

PROTECT YOURSELF

References

- CDC. Estimated HIV incidence among adults and adolescents in the United States, 2007–2010. HIV Surveillance Supplemental Report 2012;17(No. 4). <http://www.cdc.gov/hiv/topics/surveillance/resources/reports/#supplemental>. Published December 2012.
- Wejnert C, et al. HIV prevalence and awareness of infection in 2008 and 2011 among MSM – 20 cities in the United States. Presented at the Conference on Retroviruses and Opportunistic Infections (CROI 2013). Atlanta, GA, March 5, 2013.
- Oster, A et al. Trends in HIV Prevalence and HIV Testing Among Young MSM: Five United States Cities, 1994–2011. *AIDS Behavior*. 2013; DOI 10.1007/s10461-013-0566-1.



U.S. Department of Health and Human Services
Centers for Disease Control and Prevention