



Monitoring the

Nation's Health

SAFER • HEALTHIER • PEOPLE™

Vital and Health Statistics

Series 13, Number 151

September 2001

1999 National Hospital Discharge Survey: Annual Summary with Detailed Diagnosis and Procedure Data

DEPARTMENT OF HEALTH AND HUMAN SERVICES

Centers for Disease Control and Prevention • National Center for Health Statistics

Copyright information

All material appearing in this report is in the public domain and may be reproduced or copied without permission; citation as to source, however, is appreciated.

Suggested citation

Popovic JR. 1999 National Hospital Discharge Survey: Annual Summary with detailed diagnosis and procedure data. National Center for Health Statistics. Vital Health Stat 13(151). 2001.

Library of Congress Catalog Card Number 90-657616

**For sale by the U.S. Government Printing Office
Superintendent of Documents
Mail Stop: SSOP
Washington, DC 20402-9328
Printed on acid-free paper.**

Vital and Health Statistics

Series 13, Number 151

1999 National Hospital Discharge Survey: Annual Summary with Detailed Diagnosis and Procedure Data

Data From the National
Hospital Discharge Survey

U.S. DEPARTMENT OF HEALTH AND HUMAN SERVICES
Centers for Disease Control and Prevention
National Center for Health Statistics

Hyattsville, Maryland
September 2001
DHHS Publication No. (PHS) 2001-1722

National Center for Health Statistics

Edward J. Sondik, Ph.D., *Director*

Jack R. Anderson, *Deputy Director*

Jack R. Anderson, *Acting Associate Director for International Statistics*

Lawrence H. Cox, Ph.D., *Associate Director for Research and Methodology*

Jennifer H. Madans, Ph.D., *Acting Associate Director for Analysis, Epidemiology, and Health Promotion*

P. Douglas Williams, *Acting Associate Director for Data Standards, Program Development, and Extramural Programs*

Edward L. Hunter, *Associate Director for Planning, Budget, and Legislation*

Douglas L. Zinn, *Acting Associate Director for Management*

Charles J. Rothwell, *Associate Director for Data Processing and Services*

Division of Health Care Statistics

Linda K. Demlo, Ph.D., *Director*

Thomas McLemore, *Deputy Director*

Irma E. Arispe, Ph.D., *Associate Director for Science*

Robert Pokras, *Chief, Hospital Care Statistics Branch*

Cooperation of the U.S. Bureau of the Census

Under the legislation establishing the National Health Survey, the Public Health Service is authorized to use, insofar as possible, the services or facilities of other Federal, State, or private agencies.

In accordance with specifications established by the National Center for Health Statistics, the U.S. Bureau of the Census, under a contractual arrangement, participated in planning the surveys and collecting the data for the National Hospital Discharge Survey.

Contents

Abstract	1
Highlights.....	1
Introduction	2
Methods	3
Data Source	3
Estimation	4
Use of Tables.....	4
References	5
Appendix	187
Technical Notes	187
Abstract Form	187
Population Estimates	187
Reliability of Estimates.....	187
Definitions of Selected Terms.....	190
Changes in ICD–9–CM Codes	191

Detailed Tables

Demographic Tables

1. Number and rate of discharges from short-stay hospitals and of days of care, with average length of stay, by age: United States, selected years, 1970–99	7
2. Number, percent distribution, and rate of discharges from short-stay hospitals and of days of care, with average length of stay, by sex and age: United States, 1999	8
3. Number and rate of discharges from short-stay hospitals and days of care, with average length of stay, by sex, age, and geographic region: United States, 1999.....	9
4. Number of discharges from short-stay hospitals and of days of care, with average length of stay, by principal expected source of payment, sex, age, and geographic region: United States, 1999	11
5. Number and percent distribution of discharges from short-stay hospitals and of days of care, with average length of stay, by bed size of hospital, sex, age and geographic region: United States, 1999.....	14
6. Number and percent distribution of discharges from short-stay hospitals and of days of care, with average length of stay, by hospital ownership, sex, age, and geographic region: United States, 1999.....	16
7. Number and rate of deaths for discharges from short-stay hospitals, by sex and age: United States, 1999	17

Diagnosis Tables

8. Number and rate of discharges from short-stay hospitals and of days of care, with average length of stay, by selected first-listed diagnostic categories: United States, 1999	18
9. Number and rate of discharges from short-stay hospitals with average length of stay, for patients 65 years of age and over by age and selected first-listed diagnostic categories: United States, 1999	19
10. Number of discharges from short-stay hospitals, by age and first-listed diagnosis: United States, 1999	21
11. Rate of discharges from short-stay hospitals, by age and first-listed diagnosis: United States, 1999	22
12. Average length of stay for discharges from short-stay hospitals, by age and first-listed diagnosis: United States, 1999..	23
13. Number of discharges from short-stay hospitals, by sex and first-listed diagnosis: United States, 1999	24
14. Rate of discharges from short-stay hospitals, by sex and first-listed diagnosis: United States, 1999	25

15.	Average length of stay for discharges from short-stay hospitals, by sex and first-listed diagnosis: United States, 1999..	26
16.	Number of discharges from short-stay hospitals, by geographic region and first-listed diagnosis: United States, 1999 ..	27
17.	Rate of discharges from short-stay hospitals, by geographic region and first-listed diagnosis: United States, 1999	28
18.	Average length of stay for discharges from short-stay hospitals, by geographic region and first-listed diagnosis: United States, 1999.....	29
19.	Number of all-listed diagnoses for discharges from short-stay hospitals, by age and diagnosis: United States, 1999	30
20.	Number of all-listed diagnoses for discharges from short-stay hospitals, by sex and diagnosis: United States, 1999	31
21.	Number of all-listed diagnoses for discharges from short-stay hospitals, by geographic region and diagnosis: United States, 1999.....	32
22.	Number and rate of discharges from short-stay hospitals with human immunodeficiency virus diagnoses, by selected characteristics: United States, 1990–99	33
23.	Number and rate of deaths for discharges from short-stay hospitals, by age and selected first-listed diagnoses: United States, 1999.....	34

Procedure Tables

24.	Number of discharges from short-stay hospitals with and without procedures and percent with procedures, by selected characteristics: United States, 1999	34
25.	Number and rate of all-listed surgical and nonsurgical procedures for discharges from short-stay hospitals by selected procedure categories: United States, 1999.....	35
26.	Number and rate of all-listed surgical and nonsurgical procedures for patients 65 years of age and over discharged from short-stay hospitals, by age and selected procedure categories: United States, 1999	36
27.	Number and rate of discharges from short-stay hospitals with a coronary artery procedure and number and rate of procedures by sex, age, and region: United States, 1999	37
28.	Number of all-listed procedures for discharges from short-stay hospitals, by age and procedure category: United States, 1999	38
29.	Rate of all-listed procedures for discharges from short-stay hospitals, by age and procedure category: United States, 1999.....	39
30.	Number of all-listed procedures for discharges from short-stay hospitals, by sex and procedure category: United States, 1999.....	40
31.	Rate of all-listed procedures for discharges from short-stay hospitals, by sex and procedure category: United States, 1999.....	41
32.	Number of all-listed procedures for discharges from short-stay hospitals, by geographic region and procedure category: United States, 1999.....	42
33.	Rate of all-listed procedures for discharges from short-stay hospitals, by geographic region and procedure category: United States, 1999.....	43

Tables on Deliveries and Newborn Infants

34.	Number, percent distribution, and rate of discharges and days of care, with average length of stay for females with deliveries discharged from short-stay hospitals, by type of delivery, age, and geographic region: United States, 1999 ..	44
35.	Number and percent distribution of females with deliveries discharged from short-stay hospitals by length of stay and type of delivery: United States, selected years 1980–99	45
36.	Number and rate of selected obstetrical procedures for females with deliveries discharged from short-stay hospitals: United States, selected years 1980–99.....	46
37.	Number and rate of females with deliveries discharged from short-stay hospitals by type of delivery: United States, selected years 1980–99.....	46
38.	Number and rate of females with deliveries discharged from short-stay hospitals by type of delivery, age, and geographic region: United States, 1999.....	47
39.	Number, percent distribution, and average length of stay for newborn infants discharged from short-stay hospitals by sex, geographic region, and health status: United States, 1999	48
40.	Number and percent distribution of newborn infants discharged from short-stay hospitals by length of stay, and average length of stay by health status: United States, selected years 1980–99.....	48
41.	Number of all-listed diagnoses for sick newborn infants discharged from short-stay hospitals, by sex and selected diagnostic categories: United States, 1999	49

Diagnosis and Procedure Tables by ICD–9–CM Code

42.	Number of first-listed diagnoses for discharges from short-stay hospitals, by ICD–9–CM code, sex, age, and geographic region: United States, 1999.....	50
-----	---	----

43.	Number of days of care for discharges from short-stay hospitals, by ICD-9-CM code of first-listed diagnosis, sex, age, and geographic region: United States, 1999	78
44.	Number of all-listed diagnoses for discharges from short-stay hospitals, by ICD-9-CM code, sex, age, and geographic region: United States, 1999	106
45.	Number of all-listed procedures for discharges from short-stay hospitals, by ICD-9-CM code, sex, age, and geographic region: United States, 1999.....	166

Appendix Tables

I.	Civilian population by selected characteristics: United States, 1999.....	187
II.	Estimated parameters for approximate relative standard error equations for National Hospital Discharge Survey statistics by selected characteristics, 1999.....	190
III.	Code numbers for procedures considered nonsurgical in the National Hospital Discharge Survey by ICD-9-CM category, 1999.....	192
IV.	Changes in diagnosis codes	193
V.	Changes in procedure codes.....	202

Appendix Figure

I.	Medical abstract for the National Hospital Discharge Survey, 1999.....	188
----	--	-----

Objectives

This report presents 1999 national estimates and selected trend data on the use of non-Federal short-stay hospitals in the United States. Estimates are provided by demographic characteristics of patients discharged, geographic region, ownership, and bed size of hospitals, principal expected source of payment, conditions diagnosed, and surgical and nonsurgical procedures performed. Measurements of hospital use include number and rate of discharges and days of care, and the average length of stay. Estimates of first-listed diagnoses, days of care, all-listed diagnoses, and all-listed procedures are presented according to their code number in the *International Classification of Diseases, 9th Revision, Clinical Modification (ICD-9-CM)*.

Methods

The estimates are based on data collected through the National Hospital Discharge Survey. The survey has been conducted annually by the National Center for Health Statistics since 1965. In 1999 data were collected for approximately 300,000 discharges. Of the 487 eligible non-Federal short-stay hospitals in the sample, 458 (94 percent) responded to the survey.

Results

An estimated 32.1 million inpatients were discharged from non-Federal short-stay hospitals in 1999. These patients used an estimated 160.1 million days of care. Patients 65 years of age and over accounted for 40 percent of discharges and used 48 percent of days of care. More than half of all patients under age 65 in 1999 had a principal expected source of payment from a private source. Patients discharged from nonprofit hospitals accounted for nearly 78 percent of all discharges. Heart disease accounted for 4.5 million discharges, and deliveries made up 3.8 million discharges. An estimated 3.7 million newborn infants were discharged from short-stay hospitals after average stays of 3.2 days.

Keywords: hospitalization • inpatient • diagnoses • procedures

1999 National Hospital Discharge Survey: Annual Summary with Detailed Diagnosis and Procedure Data

by Jennifer R. Popovic, M.A., Division of Health Care Statistics

Highlights

Patient characteristics

- During 1999 an estimated 32.1 million inpatients were discharged from non-Federal short-stay hospitals. These patients used an estimated 160.1 million days of care.
- Patients 65 years of age and over comprised 12 percent of the population in 1999, but they accounted for 40 percent of discharges and used 48 percent of days of care (figure 1).
- In 1970 average lengths of stay ranged from 4.7 days (under 15 years) to 12.6 days (65 years and over). In 1999 the range was from 3.7 days (15 to 44 years) to 6.1 (65 years and over) (figure 2).
- During 1999 more than half of all patients under age 65 had a principal expected source of payment from a private source (Blue Cross/Blue Shield, HMO/PPO, or another private source) (figure 3).
- Deaths accounted for 3 percent of hospital discharges, with patients age 65 and over accounting for 78 percent of these.

Hospital characteristics

- Of the four geographic regions, the South made up 37 percent of the 160.1 million total days of care used in the United States during 1999.
- In 1999 almost half of inpatients were discharged from hospitals with less than 200 beds, while about 10 percent were discharged from hospitals with 500 or more beds.
- Nonprofit hospitals accounted for nearly 78 percent of hospital discharges in 1999. The proportion ranged from 94 percent in the Northeast to 61 percent in the South.

Diagnoses

- Heart disease accounted for 4.5 million discharges, and deliveries made up 3.8 million discharges.
- Heart disease was the most common diagnosis among inpatients 65 years of age and over, accounting for 23 percent of their discharges.
- Four diagnostic categories comprised half of all inpatient deaths: heart disease (21 percent), malignant neoplasms (12 percent), pneumonia (9 percent), and cerebrovascular disease (8 percent) (figure 4).

NOTE: This report was prepared in the Division of Health Care Statistics. Lola Jean Kozak provided editorial comments. Rong Cai of the OAO Corporation did the computer programing for the report and produced the estimated parameters for approximate relative standard error equations. This report was edited by Demarius V. Miller, typeset by Annette F. Holman, and the graphics produced by Jarmila G. Ogburn of the Publications Branch, Division of Data Services.




Figure 1. Percent distribution of the civilian population, hospital discharges, and hospital days of care by age: United States, 1999




Figure 2. Average length of stay in days by age: United States, selected years 1970-1999

short-stay hospitals after average stays of 3.2 days.

- In 1999 about 38 percent of newborn infants had at least one illness or risk-related diagnosis.
- In 1980 lengths of hospital stays averaged 3.2 days among well newborns and 7.1 days among sick newborns (those with at least one illness or risk-related diagnosis). In 1999 well newborn stays averaged 2.0 days, compared with 5.1 days for sick newborns (figure 5).
- More than half the diagnoses for sick newborn infants were hemolytic disease of fetus or newborn, respiratory distress syndrome and other respiratory conditions, disorders relating to short gestation and unspecified low birthweight, or congenital anomalies.

Introduction

This report presents data from the 1999 National Hospital Discharge Survey (NHDS). The survey has been conducted continuously by the National Center for Health Statistics (NCHS) since 1965. National estimates of hospital use derived from the NHDS are published for each calendar year by NCHS. This report provides a summary of the 1999 data, including estimates of diagnoses and procedures by the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM) code number (1).

The NHDS is the principal source for national data on the characteristics of patients discharged from non-Federal short-stay hospitals. Data from the NHDS are used to examine important topics in public health (2-6) and for a variety of planning, administrative, and evaluation activities by governmental, scientific, academic, and commercial institutions.

Estimates of the number of diagnoses and procedures shown in this report are for hospital inpatients only. Data on ambulatory medical care services are collected in other NCHS surveys. The National Ambulatory Medical Care Survey obtains information on patients' visits to

Procedures

- One or more procedures were reported for 62 percent of discharges.

Four surgical procedures were performed 1 million times or more: cardiac catheterization, repair of current obstetric laceration, episiotomy, and removal of coronary artery obstruction

and insertion of stent(s).

Three nonsurgical procedures were performed 1 million times or more: arteriography and angiocardiology, respiratory therapy, and diagnostic ultrasound.

Newborn infants

- An estimated 3.7 million newborn infants were discharged from




Figure 3. Percent distribution of discharges under age 65 by principal expected source of payment: United States, 1999

physicians' offices (7). The National Hospital Ambulatory Medical Care Survey collects data on visits to hospital outpatient and emergency departments (8,9). Data on ambulatory surgery performed in hospitals and in freestanding ambulatory surgery centers were collected by the National Survey of Ambulatory Surgery from 1994–96 (10).

Methods

Data Source

NHDS data are collected from a sample of inpatient records acquired from a national probability sample of hospitals. Because persons with multiple discharges during the year can be sampled more than

once, NHDS produces estimates for discharges, not persons. Only hospitals with an average length of stay of fewer than 30 days for all patients, general hospitals, or children's general hospitals are included in the survey. Federal, military, and Department of Veterans Affairs hospitals, as well as hospital units of institutions (such as prison hospitals), and hospitals with fewer than six beds staffed for patient use, are excluded. Beginning in 1988, the sampling frame for hospitals has been the SMG Hospital Market Database (11). The 1988 sample was updated in 1991, 1994, and 1997, to allow the inclusion of hospitals that opened or changed their eligibility status since the previous sample update.

NHDS originally used a two-stage design, but it was redesigned in 1988 and implemented a stratified three-stage design (12). Both the original and the redesigned samples include the largest

hospitals with certainty and sample noncertainty hospitals. The first stage of the current three-stage design consists of the selection of 112 primary sampling units (PSU's) that comprise a probability sample of PSU's used in the 1985–94 National Health Interview Survey (13). The second stage consists of the selection of noncertainty hospitals from the sample PSU's. At the third stage, a systematic random sample of discharges within hospitals is selected.

For 1999 the sample consisted of 513 hospitals, 26 of which were out of scope (ineligible) because they had gone out of business or otherwise failed to meet the criteria for the NHDS universe. Of the 487 in-scope (eligible) hospitals, 458 (94 percent) responded to the survey. Data were collected for approximately 300,000 discharges from the responding hospitals.

Two data collection procedures are used for the survey. One is a manual system in which sample selection and transcription of information from medical records are performed by hospital staff or by staff of the U.S. Bureau of the Census on behalf of NCHS. Completed forms are sent to NCHS for medical coding, data entry, editing, and estimation.

The other data collection procedure is an automated system in which NCHS purchases electronic files containing medical record data from commercial organizations, State data systems, hospitals, or hospital associations. Discharges from these files are systematically sampled by NCHS. Approximately 40 percent of respondent hospitals provided data through the automated system for the 1999 survey.

The NHDS includes data elements relating to patient demographic characteristics and medical information, as well as administrative and hospital-related data elements. The medical abstract form shown as [figure I](#) in the appendix collects patient date of birth or age, ZIP Code, dates of admission and discharge, discharge status, sex, race, ethnicity, marital status, expected source of payment, final medical diagnoses, procedures performed, and dates of surgery. Medical data are coded according to the *International Classification of Diseases*,





Figure 4. Percent distribution of deaths for discharges from short-stay hospitals by selected first-listed diagnosis: United States, 1999



¹With at least one illness or risk-related diagnosis, ICD-9-CM codes 001-999.

²Without any illness or risk-related diagnoses.

Figure 5. Average length of stay for newborn infants by health status: United States, selected years, 1980-1999

9th Revision, Clinical Modification (ICD-9-CM) (1). Administrative and hospital data collected include patient medical record numbers, number of hospital beds, type of ownership, and geographic region of the hospital. The patient medical record number, date of birth, and ZIP Code are confidential information that are not available to the public.

For the manual data collection, an ongoing quality control program is undertaken for the medical coding and data entry from abstracts to electronic form. Approximately 10 percent of the abstracts are independently recoded, with discrepancies resolved by the chief coder. The overall error rate for records manually coded for the 1999 survey year was 0.6 percent for medical (ICD-9-CM) coding and data entry, and 0.3 percent for demographic coding and data entry.

Estimation

Because they are sample data, the survey data must be inflated or weighted to produce national estimates. The estimation procedure produces essentially unbiased national estimates and has three basic components:

1. inflation by reciprocals of the probabilities of sample selection,
2. adjustment for nonresponse, and
3. population weighting ratio adjustments.

These three components of the final weight are described in more detail in another report (12). Information about the standard errors of statistics for the 1999 NHDS is found in the appendix of this report.

Use of Tables

Estimates of hospital use by age and sex of discharged patients and geographic region of hospitals are shown in [tables 1-3](#). Estimates of hospital use by principal expected source of payment, hospital bed size, and hospital ownership are shown in [tables 4-6](#). The number and rate of hospital deaths are in [table 7](#). Diagnostic data are included in [tables 8-23](#) and information about procedures is found in

tables 24–33. Tables 34–41 present data on women with deliveries and newborn infants. Estimates of diagnoses, procedures, and days of care by diagnosis are shown by ICD–9–CM code in tables 42–45. Figures 1–4 illustrate selected highlights from the tables. Definitions of terms used in these tables are presented in the appendix.

The diagnostic data in tables 8–21 and procedure data in tables 28–33 are presented by chapter of the ICD–9–CM. Within these chapters, subcategories of diagnoses or procedures are shown. These specific categories are selected because of their large estimates or because they were of special interest. Although diagnoses assigned ICD–9–CM codes E800–E999 (supplementary classification of external causes of injury and poisoning) are included in the NHDS, these diagnoses are not well reported in the survey and therefore are excluded from this report. A detailed discussion of injury diagnoses and E-codes has been published (14).

Table 27 shows data for coronary artery bypass graft (CABG) surgery and removal of coronary artery obstruction and insertion of stent(s). These procedures are often assigned more than one code for a single operation. CABG surgery frequently is assigned multiple codes to indicate the number and type of bypasses. Beginning with the 1996 NHDS data, changes to the ICD–9–CM coding system allowed patients to have a separate code for insertion of stent(s) along with a code for removal of coronary artery obstruction. As shown in table 27, the number of these procedures greatly exceeds the number of inpatients who had one or more of the procedures.

Tables 7 and 23 show rates of death per 100 discharges. In contrast, mortality rates are ratios of deaths to the population. Mortality rates include deaths that occur outside of hospitals and are for the underlying cause of death, which might not be the same as the first-listed diagnosis for a patient discharged dead. Mortality rates are published regularly by NCHS in *National Vital Statistics Reports* (15) and in *Vital Statistics of the United States* (16).

Data for newborn infants are included only in tables 39–41. Because estimates of newborn infants are based on a sample and because they do not include out-of-hospital births, these estimates may not agree with data on births published in *National Vital Statistics Reports* (17) and *Vital Statistics of the United States* (18), which are based on a complete count of births.

Tables 42–44 provide information by three-digit, four-digit, and five-digit ICD–9–CM diagnostic codes. Table 45 shows two-, three-, and four-digit procedure codes. Only ICD–9–CM code numbers appear in tables 42–45. Specific titles and more detailed information about the coding system can be found in appropriate volumes of the ICD–9–CM (1). A specific diagnosis or procedure code appears in tables 42, 44, and 45 if the estimate of the total for the code is 5,000 or more. The codes that appear in table 43 for days of care are identical to those in table 42 for first-listed diagnoses.

Estimates in all tables are rounded to the nearest thousand. Totals may include data for categories that are not shown individually in the tables. For these reasons, figures within tables do not always add to the totals. Rates, percents, and average lengths of stay are calculated from unrounded figures and may not precisely agree with measures calculated from rounded data.

Because of low reliability, estimates in tables 1–41 with relative standard errors of more than 30 percent or that are based on fewer than 30 records are not presented; only an asterisk (*) appears in the tables. Estimates based on fewer than 60 records are preceded by an asterisk (*) to indicate that they also have low reliability.

Only an asterisk is shown in tables 42, 44, and 45 if the estimate of diagnosis or procedure is less than 5,000, and only an asterisk is shown in table 43 for estimates of days of care derived from fewer than 5,000 estimated discharges. These estimates generally have a relative standard error of more than 30 percent or a sample size of less than 30. Estimates of diagnoses and procedures of less than 9,000 are preceded by an asterisk because they are

generally based on fewer than 60 records. Days of care estimates in table 43 that are derived from fewer than 9,000 estimated discharges are also preceded by asterisks.

The population estimates used to compute rates are shown in table I in the appendix. These estimates are adjusted for the 1990 census underenumeration. Populations used to compute rates in reports of 1990–96 of NHDS data were not adjusted for the underenumeration.

References

1. International Classification of Diseases, 9th Revision, Clinical Modification, 6th edition. U.S. Department of Health and Human Services, National Center for Health Statistics, Health Care Financing Administration. 1998.
2. Kozak LJ, Hall MJ, Owings MF. Trends in avoidable hospitalizations, 1980–1998. *Health Affairs* 20(2): 225–32. 2001.
3. Simonsen L, Fukuda K, Schonberger LB, Cox NJ. The impact of influenza epidemics on hospitalizations. *J Infect Dis* 181(3): 831–37. 2000.
4. Curtin SC, Kozak LJ, Gregory KD. U.S. cesarean and VBAC rates stalled in the mid-1990s. *Birth* 27(1): 54–7. 2000.
5. Feinglass J, Brown JL, LoSasso A et al. Rates of lower-extremity amputation and arterial reconstruction in the United States, 1979 to 1996. *Am J Public Health* 89(8):1222–27. 1999.
6. Xia Z, Roberts RO, Schottenfeld D et al. Trends in prostatectomy for benign prostatic hyperplasia among black and white men in the United States: 1980 to 1994. *Urology* 53(6): 1154–59. 1999.
7. Woodwell DA. National Ambulatory Medical Care Survey: 1998 summary. Advance Data from vital and health statistics; no 315. Hyattsville, Maryland: National Center for Health Statistics. 2000. <http://www.cdc.gov/nchs/>
8. Slusarcick AL, McCaig LF. National Hospital Ambulatory Medical Care Survey: 1998 outpatient department summary. Advance data from vital and health statistics; no 317. Hyattsville, Maryland: National Center for Health Statistics. 2000. <http://www.cdc.gov/nchs/>
9. McCaig LF. National Hospital Ambulatory Medical Care Survey: 1998

- emergency department summary. Advance data from vital and health statistics; no 313. Hyattsville, Maryland: National Center for Health Statistics. 2000. <http://www.cdc.gov/nchs/>
10. Hall MJ, Lawrence L. Ambulatory surgery in the United States, 1996. Advance data from vital and health statistics; no. 300. Hyattsville, Maryland: National Center for Health Statistics. 1998. <http://www.cdc.gov/nchs/>
11. SMG Marketing Group, Inc. Hospital Market Database. Chicago: Healthcare Information Specialists. 1987, April 1991, April 1994, April 1997.
12. Dennison CF, Pokras R. Design and Operation of the National Hospital Discharge Survey: 1988 Redesign. National Center for Health Statistics. Vital and Health Stat 1(39). 2000. <http://www.cdc.gov/nchs/>
13. Massey JT, Moore TF, Parsons VL, Tadros W. Design and estimation for the National Health Interview Survey, 1985–94. National Center for Health Statistics. Vital Health Stat 2(110). 1989.
14. Hall MJ, Owings MF. Hospitalizations for Injury: United States, 1996. Advance data from vital and health statistics; no. 318. Hyattsville, Maryland: National Center for Health Statistics. 2000. <http://www.cdc.gov/nchs/>
15. Murphy SL. Deaths: final data for 1998. National vital statistics reports 48(11). Hyattsville, Maryland: National Center for Health Statistics. 2000. <http://www.cdc.gov/nchs/>
16. National Center for Health Statistics. Vital statistics of the United States, 1991, vol II, mortality, part a. Washington, DC: Public Health Service. 1996.
17. Ventura SJ, Martin JA, Curtin SC. Births: final data for 1999. National vital statistics reports 49(1). Hyattsville, Maryland: National Center for Health Statistics. 2001. <http://www.cdc.gov/nchs/>
18. National Center for Health Statistics. Vital statistics of the United States, 1997, part I, natality. May 4, 2000. <http://www.cdc.gov/nchs/>
19. Bieler GS, Williams RL. Analyzing survey data using SUDAAN release 7.5. Research Triangle Park, NC: Research Triangle Institute. 1997.
20. Hansen MH, Hurwitz WN, Madow WG. Sample survey methods and theory, vol 1. New York: John Wiley & Sons. Wiley Classics Library Edition. 1993.
21. Cochran WG. Sampling techniques, 3rd ed. New York: John Wiley & Sons. 1977.

Table 1. Number and rate of discharges from short-stay hospitals and of days of care, with average length of stay, by age: United States, selected years, 1970–99

[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants]

Age	1970	1975	1980	1985	1990	1995	1999
Number of discharges in thousands							
All ages	29,127	34,043	37,832	35,056	30,788	30,722	32,132
Under 15 years	3,873	3,826	3,672	2,972	2,412	2,405	2,458
15–44 years	12,664	14,171	15,635	13,966	11,799	10,593	10,092
45–64 years	6,693	8,391	8,660	7,610	6,244	6,168	6,899
65 years and over	5,897	7,654	9,864	10,508	10,333	11,556	12,683
65–74 years	3,163	4,057	4,943	5,011	4,689	4,832	4,883
75–84 years	2,099	2,728	3,611	3,969	3,949	4,590	5,227
85 years and over	635	869	1,310	1,528	1,694	2,134	2,572
Rate of discharges per 1,000 population ¹							
All ages	144.3	159.2	167.7	148.4	122.3	115.7	116.6
Under 15 years	66.8	70.4	71.6	57.7	43.1	40.4	40.8
15–44 years	154.6	152.4	150.1	125.0	99.3	87.8	82.7
45–64 years	159.6	191.8	194.8	170.8	135.5	118.5	116.9
65 years and over	293.3	337.3	383.7	369.8	334.1	347.7	370.4
65–74 years	253.2	291.5	315.8	297.3	261.6	260.0	270.6
75–84 years	339.5	392.1	463.9	446.4	395.7	415.6	433.9
85 years and over	443.7	477.1	576.4	572.9	560.6	592.5	620.0
Number of days of care in thousands							
All ages	226,445	262,389	274,508	226,217	197,422	164,627	160,128
Under 15 years	18,189	17,554	16,191	13,554	11,655	10,715	11,026
15–44 years	71,605	80,687	81,951	67,397	54,062	40,825	37,236
45–64 years	62,419	75,393	71,008	53,541	42,153	34,207	34,595
65 years and over	74,232	88,754	105,358	91,726	89,552	78,880	77,270
65–74 years	37,873	44,963	49,253	41,090	37,422	31,310	28,474
75–84 years	27,633	32,906	40,355	36,024	35,926	31,974	32,257
85 years and over	8,726	10,886	15,750	14,612	16,204	15,597	16,539
Rate of days of care per 1,000 population ¹							
All ages	1,121.6	1,227.3	1,216.7	957.7	784.0	620.2	581.1
Under 15 years	313.9	322.9	315.7	263.0	208.4	179.9	183.0
15–44 years	874.0	867.9	786.8	603.3	454.9	338.4	305.1
45–64 years	1,488.8	1,723.0	1,596.9	1,201.6	914.4	657.5	586.3
65 years and over	3,691.9	3,910.6	4,098.4	3,228.0	2,895.6	2,373.7	2,256.8
65–74 years	3,031.6	3,230.8	3,147.0	2,437.3	2,087.8	1,684.7	1,578.1
75–84 years	4,469.2	4,729.2	5,183.7	4,052.2	3,599.5	2,894.7	2,677.6
85 years and over	6,102.1	5,978.1	6,932.1	5,478.7	5,361.7	4,330.6	3,986.7
Average length of stay in days							
All ages	7.8	7.7	7.3	6.5	6.4	5.4	5.0
Under 15 years	4.7	4.6	4.4	4.6	4.8	4.5	4.5
15–44 years	5.7	5.7	5.2	4.8	4.6	3.9	3.7
45–64 years	9.3	9.0	8.2	7.0	6.8	5.5	5.0
65 years and over	12.6	11.6	10.7	8.7	8.7	6.8	6.1
65–74 years	12.0	11.1	10.0	8.2	8.0	6.5	5.8
75–84 years	13.2	12.1	11.2	9.1	9.1	7.0	6.2
85 years and over	13.7	12.5	12.0	9.6	9.6	7.3	6.4

¹Rates for 1990, 1995, and 1999 are based on U.S. Bureau of the Census estimates of the civilian population that have been adjusted for net underenumeration in the 1990 census. See appendix.

Table 2. Number, percent distribution, and rate of discharges from short-stay hospitals and of days of care, with average length of stay, by sex and age: United States, 1999

[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants]

Sex and age	Discharges			Days of care			Average length of stay in days
	Number in thousands	Percent distribution	Rate per 1,000 population	Number in thousands	Percent distribution	Rate per 1,000 population	
Both sexes							
All ages	32,132	100.0	116.6	160,128	100.0	581.1	5.0
Under 15 years	2,458	7.7	40.8	11,026	6.9	183.0	4.5
Under 1 year	827	2.6	208.5	4,391	2.7	1,106.9	5.3
1–4 years	729	2.3	46.6	2,484	1.6	158.7	3.4
5–14 years	902	2.8	22.2	4,152	2.6	102.2	4.6
15–44 years	10,092	31.4	82.7	37,236	23.3	305.1	3.7
15–19 years	1,209	3.8	60.0	4,326	2.7	214.6	3.6
20–24 years	1,644	5.1	89.8	5,136	3.2	280.7	3.1
25–34 years	3,829	11.9	99.3	12,738	8.0	330.5	3.3
35–44 years	3,411	10.6	75.7	15,036	9.4	333.9	4.4
45–64 years	6,899	21.5	116.9	34,595	21.6	586.3	5.0
45–54 years	3,386	10.5	94.5	16,150	10.1	451.0	4.8
55–64 years	3,513	10.9	151.4	18,445	11.5	795.1	5.3
65 years and over	12,683	39.5	370.4	77,270	48.3	2,256.8	6.1
65–74 years	4,883	15.2	270.6	28,474	17.8	1,578.1	5.8
75–84 years	5,227	16.3	433.9	32,257	20.1	2,677.6	6.2
85 years and over	2,572	8.0	620.0	16,539	10.3	3,986.7	6.4
Male							
All ages	12,748	100.0	94.7	68,719	100.0	510.4	5.4
Under 15 years	1,398	11.0	45.3	6,366	9.3	206.5	4.6
Under 1 year	480	3.8	236.9	2,475	3.6	1,221.1	5.2
1–4 years	407	3.2	50.9	1,338	1.9	167.3	3.3
5–14 years	510	4.0	24.5	2,553	3.7	122.7	5.0
15–44 years	2,715	21.3	44.5	13,177	19.2	216.1	4.9
15–19 years	322	2.5	31.2	1,481	2.2	143.4	4.6
20–24 years	298	2.3	32.4	1,516	2.2	164.5	5.1
25–34 years	777	6.1	40.8	3,575	5.2	187.4	4.6
35–44 years	1,318	10.3	59.0	6,605	9.6	295.6	5.0
45–64 years	3,390	26.6	118.5	17,266	25.1	603.4	5.1
45–54 years	1,613	12.7	92.0	7,896	11.5	450.5	4.9
55–64 years	1,777	13.9	160.3	9,371	13.6	845.2	5.3
65 years and over	5,245	41.1	368.4	31,910	46.4	2,241.1	6.1
65–74 years	2,310	18.1	283.5	13,361	19.4	1,639.7	5.8
75–84 years	2,172	17.0	447.6	13,544	19.7	2,791.0	6.2
85 years and over	762	6.0	616.0	5,005	7.3	4,044.8	6.6
Female							
All ages	19,384	100.0	137.6	91,409	100.0	648.8	4.7
Under 15 years	1,060	5.5	36.0	4,660	5.1	158.4	4.4
Under 1 year	347	1.8	178.9	1,916	2.1	987.6	5.5
1–4 years	322	1.7	42.1	1,145	1.3	149.7	3.6
5–14 years	392	2.0	19.7	1,599	1.7	80.6	4.1
15–44 years	7,377	38.1	120.8	24,059	26.3	394.0	3.3
15–19 years	887	4.6	90.2	2,845	3.1	289.5	3.2
20–24 years	1,345	6.9	148.2	3,621	4.0	398.7	2.7
25–34 years	3,052	15.7	156.7	9,163	10.0	470.6	3.0
35–44 years	2,093	10.8	92.3	8,431	9.2	371.6	4.0
45–64 years	3,508	18.1	115.4	17,329	19.0	570.1	4.9
45–54 years	1,773	9.1	97.0	8,255	9.0	451.4	4.7
55–64 years	1,736	9.0	143.3	9,074	9.9	749.2	5.2
65 years and over	7,438	38.4	371.9	45,360	49.6	2,267.9	6.1
65–74 years	2,573	13.3	260.0	15,113	16.5	1,527.3	5.9
75–84 years	3,055	15.8	424.6	18,714	20.5	2,601.1	6.1
85 years and over	1,810	9.3	621.7	11,534	12.6	3,962.0	6.4

Table 3. Number and rate of discharges from short-stay hospitals and days of care, with average length of stay, by sex, age, and geographic region: United States, 1999

[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants]

Sex, age, and region	Discharges		Days of care		Average length of stay in days
	Number in thousands	Rate per 1,000 population	Number in thousands	Rate per 1,000 population	
Both sexes					
All ages:					
United States	32,132	116.6	160,128	581.1	5.0
Northeast	6,952	133.2	39,814	762.5	5.7
Midwest	7,368	115.9	33,988	534.7	4.6
South	12,007	123.0	59,635	610.9	5.0
West	5,805	93.4	26,692	429.6	4.6
Under 15 years:					
United States	2,458	40.8	11,026	183.0	4.5
Northeast	532	49.6	2,760	257.4	5.2
Midwest	503	36.6	1,984	144.3	3.9
South	876	40.8	3,930	183.3	4.5
West	548	38.2	2,352	163.9	4.3
15–44 years:					
United States	10,092	82.7	37,236	305.1	3.7
Northeast	2,057	90.1	9,269	406.0	4.5
Midwest	2,178	77.8	7,513	268.3	3.5
South	3,901	90.7	14,068	327.2	3.6
West	1,957	69.4	6,386	226.4	3.3
45–64 years:					
United States	6,899	116.9	34,595	586.3	5.0
Northeast	1,511	131.9	8,516	743.1	5.6
Midwest	1,490	108.8	6,900	503.6	4.6
South	2,718	128.7	13,372	633.3	4.9
West	1,179	92.6	5,807	456.0	4.9
65 years and over:					
United States	12,683	370.4	77,270	2,256.8	6.1
Northeast	2,852	395.9	19,269	2,675.1	6.8
Midwest	3,198	394.1	17,590	2,168.0	5.5
South	4,513	373.7	28,265	2,340.6	6.3
West	2,121	309.7	12,147	1,774.0	5.7
Male					
All ages					
United States	12,748	94.7	68,719	510.4	5.4
Northeast	2,919	115.4	17,738	701.5	6.1
Midwest	2,977	96.0	14,669	472.9	4.9
South	4,583	96.9	24,883	525.9	5.4
West	2,269	73.1	11,429	368.4	5.0
Under 15 years:					
United States	1,398	45.3	6,366	206.5	4.6
Northeast	302	55.0	1,620	295.1	5.4
Midwest	287	40.8	1,132	160.9	3.9
South	499	45.5	2,288	208.8	4.6
West	310	42.2	1,326	180.4	4.3
15–44 years:					
United States	2,715	44.5	13,177	216.1	4.9
Northeast	654	57.7	3,798	334.8	5.8
Midwest	606	43.3	2,652	189.5	4.4
South	1,003	47.3	4,790	225.8	4.8
West	452	31.4	1,938	134.5	4.3
45–64 years:					
United States	3,390	118.5	17,266	603.4	5.1
Northeast	756	137.2	4,241	769.8	5.6
Midwest	754	113.0	3,557	533.5	4.7
South	1,303	128.4	6,580	648.2	5.0
West	578	91.9	2,889	459.5	5.0

Table 3. Number and rate of discharges from short-stay hospitals and days of care, with average length of stay, by sex, age, and geographic region: United States, 1999—Con.

[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants]

Sex, age, and region	Discharges		Days of care		Average length of stay in days
	Number in thousands	Rate per 1,000 population	Number in thousands	Rate per 1,000 population	
65 years and over:					
United States	5,245	368.4	31,910	2,241.1	6.1
Northeast	1,207	409.8	8,079	2,743.0	6.7
Midwest	1,330	400.4	7,328	2,205.5	5.5
South	1,778	356.2	11,226	2,248.9	6.3
West	929	312.0	5,277	1,771.6	5.7
Female					
All ages:					
United States	19,384	137.6	91,409	648.8	4.7
Northeast	4,033	149.8	22,076	819.8	5.5
Midwest	4,391	134.9	19,319	593.6	4.4
South	7,424	147.6	34,751	690.7	4.7
West	3,535	113.6	15,263	490.6	4.3
Under 15 years:					
United States	1,060	36.0	4,660	158.4	4.4
Northeast	230	44.0	1,140	217.8	5.0
Midwest	216	32.2	852	126.9	3.9
South	377	35.9	1,642	156.7	4.4
West	238	33.9	1,027	146.6	4.3
15–44 years:					
United States	7,377	120.8	24,059	394.0	3.3
Northeast	1,403	122.2	5,471	476.4	3.9
Midwest	1,571	112.2	4,861	347.0	3.1
South	2,898	133.0	9,278	426.0	3.2
West	1,505	109.1	4,448	322.5	3.0
45–64 years:					
United States	3,508	115.4	17,329	570.1	4.9
Northeast	756	127.0	4,275	718.5	5.7
Midwest	737	104.7	3,344	475.2	4.5
South	1,415	129.1	6,792	619.5	4.8
West	601	93.2	2,918	452.5	4.9
65 years and over:					
United States	7,438	371.9	45,360	2,267.9	6.1
Northeast	1,644	386.3	11,189	2,628.2	6.8
Midwest	1,867	389.8	10,262	2,142.0	5.5
South	2,735	386.0	17,039	2,405.2	6.2
West	1,191	308.0	6,870	1,775.9	5.8

Table 4. Number of discharges from short-stay hospitals and of days of care, with average length of stay, by principal expected source of payment, sex, age and geographic region: United States 1999

[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants]

	All principal expected sources of payment ¹	Medicare	Medicaid	Worker's compensation and other government payments	HMO/PPO	Blue Cross/ Blue Shield and other private insurance	Self-pay	Other payments and no charge
Number of discharges in thousands								
United States:								
All	32,132	12,529	4,319	632	4,497	7,492	1,545	746
Male	12,748	5,272	1,357	360	1,619	2,906	729	337
Female	19,384	7,257	2,962	273	2,879	4,586	816	409
Under 15 years	2,458	19	899	43	406	741	134	*
15-44 years	10,092	544	2,376	295	2,103	3,345	936	332
45-64 years	6,899	1,169	849	213	1,347	2,616	419	181
65 years and over	12,683	10,797	194	81	641	790	56	45
Northeast	6,952	2,658	919	94	1,108	1,640	309	*
Midwest	7,368	3,285	681	104	971	1,849	300	95
South	12,007	4,677	1,777	252	1,636	2,524	701	*
West	5,805	1,909	942	183	782	1,480	236	204
Northeast								
Male	2,919	1,125	345	65	435	675	160	*
Female	4,033	1,533	574	29	673	965	149	*
Under 15 years	532	*	140	*	*	186	33	*
15-44 years	2,057	119	508	43	425	688	182	*
45-64 years	1,511	248	215	30	313	578	76	*
65 years and over	2,852	2,289	56	*	260	188	18	*
Midwest								
Male	2,977	1,391	223	57	337	745	146	42
Female	4,391	1,894	458	47	635	1,104	153	53
Under 15 years	*	*	143	*	55	*	*	*
15-44 years	2,178	128	384	46	498	853	191	41
45-64 years	1,490	238	135	38	329	638	71	21
65 years and over	3,198	2,917	19	*	90	127	9	*
South								
Male	4,583	1,915	511	140	540	962	313	*
Female	7,424	2,761	1,266	113	1,097	1,562	388	161
Under 15 years	876	*11	379	13	118	198	52	*
15-44 years	3,901	231	972	120	847	1,122	420	123
45-64 years	2,718	525	354	93	474	952	213	68
65 years and over	4,513	3,909	*	27	197	252	16	17
West								
Male	2,269	840	277	99	307	524	110	78
Female	3,535	1,069	665	84	474	955	126	*
Under 15 years	*	*	*	14	*	126	*	*
15-44 years	1,957	66	512	87	333	682	143	110
45-64 years	1,179	158	146	52	232	449	60	*
65 years and over	2,121	1,683	47	*	95	223	*	*
Number of days of care in thousands								
United States:								
All	160,128	77,443	20,898	3,034	17,523	29,867	6,201	3,458
Male	68,719	32,999	8,414	1,666	7,035	12,855	3,168	1,763
Female	91,409	44,444	12,484	1,367	10,488	17,012	3,033	1,695
Under 15 years	11,026	69	4,683	194	1,552	2,926	481	*
15-44 years	37,236	3,568	9,689	1,139	6,589	10,967	3,373	1,345
45-64 years	34,595	7,486	5,208	946	5,931	11,635	1,960	855
65 years and over	77,270	66,319	1,318	*	3,451	4,339	388	240
Northeast	39,814	18,544	5,861	489	4,994	7,340	1,505	*
Midwest	33,988	18,273	3,012	379	3,607	6,876	1,055	445
South	59,635	29,468	7,965	1,045	6,060	10,060	2,816	*
West	26,692	11,157	4,060	*	2,863	5,590	825	737

Table 4. Number of discharges from short-stay hospitals and of days of care, with average length of stay, by principal expected source of payment, sex, age and geographic region: United States 1999—Con.

[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants]

	All principal expected sources of payment ¹	Medicare	Medicaid	Worker's compensation and other government payments	HMO/PPO	Blue Cross/Blue Shield and other private insurance	Self-pay	Other payments and no charge
Northeast								
Male	17,738	7,850	2,774	342	2,096	3,295	819	*
Female	22,076	10,694	3,087	147	2,897	4,045	686	*
Under 15 years	2,760	*	*	*	*	830	*	*
15–44 years	9,269	920	2,929	195	1,487	2,518	770	*
45–64 years	8,516	1,811	1,449	183	1,526	2,845	422	*
65 years and over	19,269	15,809	447	*	1,501	1,148	*	*
Midwest								
Male	14,669	7,868	1,241	219	1,384	3,040	574	205
Female	19,319	10,406	1,771	160	2,224	3,836	482	240
Under 15 years	*	*	*	*	156	*	*	*
15–44 years	7,513	838	1,481	156	1,532	2,624	573	179
45–64 years	6,900	1,359	792	141	1,428	2,679	325	102
65 years and over	17,590	16,055	97	*	491	667	*	*
South								
Male	24,883	12,298	2,932	591	2,305	4,297	1,341	*
Female	34,751	17,170	5,033	454	3,755	5,763	1,475	*
Under 15 years	3,930	*37	1,794	*	453	747	181	*
15–44 years	14,068	1,384	3,641	428	2,590	3,699	1,566	*
45–64 years	13,372	3,191	2,046	400	1,998	4,222	982	283
65 years and over	28,265	24,856	484	*	1,019	1,393	87	97
West								
Male	11,429	4,983	1,468	514	1,250	2,223	435	348
Female	15,263	6,174	2,592	*	1,613	3,367	390	*
Under 15 years	*	*	*	*	*	*	*	*
15–44 years	6,386	427	1,638	360	980	2,125	464	315
45–64 years	5,807	1,125	921	223	979	1,890	231	293
65 years and over	12,147	9,599	290	*	440	1,131	*	*
Average length of stay								
United States:								
All	5.0	6.2	4.8	4.8	3.9	4.0	4.0	4.6
Male	5.4	6.3	6.2	4.6	4.3	4.4	4.3	5.2
Female	4.7	6.1	4.2	5.0	3.6	3.7	3.7	4.1
Under 15 years	4.5	3.6	5.2	4.5	3.8	3.9	3.6	5.4
15–44 years	3.7	6.6	4.1	3.9	3.1	3.3	3.6	4.1
45–64 years	5.0	6.4	6.1	4.4	4.4	4.4	4.7	4.7
65 years and over	6.1	6.1	6.8	9.3	5.4	5.5	7.0	5.3
Northeast	5.7	7.0	6.4	5.2	4.5	4.5	4.9	5.1
Midwest	4.6	5.6	4.4	3.7	3.7	3.7	3.5	4.7
South	5.0	6.3	4.5	4.1	3.7	4.0	4.0	5.1
West	4.6	5.8	4.3	6.1	3.7	3.8	3.5	3.6
Northeast								
Male	6.1	7.0	8.0	5.3	4.8	4.9	5.1	5.2
Female	5.5	7.0	5.4	5.1	4.3	4.2	4.6	5.0
Under 15 years	5.2	*	*	*7.0	4.3	4.5	4.3	4.4
15–44 years	4.5	7.7	5.8	4.6	3.5	3.7	4.2	5.3
45–64 years	5.6	7.3	6.7	6.1	4.9	4.9	5.6	5.9
65 years and over	6.8	6.9	8.0	5.1	5.8	6.1	9.2	4.2
Midwest								
Male	4.9	5.7	5.6	3.9	4.1	4.1	3.9	4.8
Female	4.4	5.5	3.9	3.4	3.5	3.5	3.1	4.5
Under 15 years	3.9	*	4.5	3.6	2.9	3.9	2.9	4.5
15–44 years	3.5	6.6	3.9	3.4	3.1	3.1	3.0	4.4
45–64 years	4.6	5.7	5.9	3.7	4.3	4.2	4.6	4.8
65 years and over	5.5	5.5	5.1	5.1	5.5	5.2	*	6.9

Table 4. Number of discharges from short-stay hospitals and of days of care, with average length of stay, by principal expected source of payment, sex, age and geographic region: United States 1999—Con.

[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants]

	All principal expected sources of payment ¹	Medicare	Medicaid	Worker's compensation and other government payments	HMO/PPO	Blue Cross/ Blue Shield and other private insurance	Self-pay	Other payments and no charge
South								
Male	5.4	6.4	5.7	4.2	4.3	4.5	4.3	5.8
Female	4.7	6.2	4.0	4.0	3.4	3.7	3.8	4.5
Under 15 years	4.5	*3.3	4.7	4.3	3.8	3.8	3.5	6.6
15–44 years	3.6	6.0	3.7	3.6	3.1	3.3	3.7	4.4
45–64 years	4.9	6.1	5.8	4.3	4.2	4.4	4.6	4.1
65 years and over	6.3	6.4	6.7	6.0	5.2	5.5	5.4	5.7
West								
Male	5.0	5.9	5.3	5.2	4.1	4.2	3.9	4.5
Female	4.3	5.8	3.9	*	3.4	3.5	3.1	3.1
Under 15 years	4.3	*	5.1	4.9	3.8	3.5	3.5	3.1
15–44 years	3.3	6.5	3.2	4.1	2.9	3.1	3.2	2.9
45–64 years	4.9	7.1	6.3	4.3	4.2	4.2	3.8	4.8
65 years and over	5.7	5.7	6.2	15.5	4.6	5.1	*4.9	4.6

¹Includes discharges for which no expected source of payment was provided.

* Figure does not meet standard of reliability or precision.

NOTE: HMO/PPO is defined as Health Maintenance Organization and Preferred Provider Organization.

Table 5. Number and percent distribution of discharges from short-stay hospitals and of days of care, with average length of stay, by bed size of hospital, sex, age and geographic region: United States, 1999

[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants]

	All bed sizes	6-99	100-199	200-299	300-499	500 or more
Number of discharges in thousands						
All	32,132	6,878	8,443	5,935	7,747	3,129
Sex						
Male	12,748	2,616	3,277	2,382	3,147	1,325
Female	19,384	4,262	5,166	3,552	4,600	1,804
Age						
Under 15 years	2,458	441	744	517	520	236
15-44 years	10,092	2,042	2,633	1,733	2,566	1,119
45-64 years	6,899	1,264	1,703	1,299	1,828	805
65 years and over	12,683	3,132	3,363	2,385	2,833	969
Region						
Northeast	6,952	1,239	1,389	1,489	1,860	976
Midwest	7,368	2,280	1,713	1,353	1,231	792
South	12,007	2,640	3,164	2,077	3,016	1,111
West	5,805	720	2,177	1,016	1,640	251
Percent distribution of patients discharged						
All	100.0	21.4	26.3	18.5	24.1	9.7
Sex						
Male	100.0	20.5	25.7	18.7	24.7	10.4
Female	100.0	22.0	26.7	18.3	23.7	9.3
Age						
Under 15 years	100.0	18.0	30.3	21.1	21.2	9.6
15-44 years	100.0	20.2	26.1	17.2	25.4	11.1
45-64 years	100.0	18.3	24.7	18.8	26.5	11.7
65 years and over	100.0	24.7	26.5	18.8	22.3	7.6
Region						
Northeast	100.0	17.8	20.0	21.4	26.8	14.0
Midwest	100.0	30.9	23.2	18.4	16.7	10.8
South	100.0	22.0	26.4	17.3	25.1	9.3
West	100.0	12.4	37.5	17.5	28.3	4.3
Number of days of care in thousands						
Total	160,128	31,131	40,238	30,884	40,255	17,620
Sex						
Male	68,719	12,838	16,642	13,207	17,910	8,123
Female	91,409	18,293	23,597	17,677	22,345	9,497
Age						
Under 15 years	11,026	1,961	3,079	2,223	2,573	1,190
15-44 years	37,236	7,022	8,579	6,573	10,000	5,063
45-64 years	34,595	5,572	8,037	6,687	9,582	4,717
65 years and over	77,270	16,576	20,543	15,402	18,099	6,650
Region						
Northeast	39,814	6,876	7,982	8,380	10,765	5,810
Midwest	33,988	9,127	8,141	6,697	5,968	4,053
South	59,635	12,013	14,909	10,708	15,617	6,387
West	26,692	3,114	9,205	5,099	7,904	1,370
Percent distribution of days of care						
All	100.0	19.4	25.1	19.3	25.1	11.0
Sex						
Male	100.0	18.7	24.2	19.2	26.1	11.8
Female	100.0	20.0	25.8	19.3	24.4	10.4

Table 5. Number and percent distribution of discharges from short-stay hospitals and of days of care, with average length of stay, by bed size of hospital, sex, age and geographic region: United States, 1999—Con.

[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants]

	All bed sizes	6–99	100–199	200–299	300–499	500 or more
Age						
Under 15 years	100.0	17.8	27.9	20.2	23.3	10.8
15–44 years	100.0	18.9	23.0	17.7	26.9	13.6
45–64 years	100.0	16.1	23.2	19.3	27.7	13.6
65 years and over	100.0	21.5	26.6	19.9	23.4	8.6
Region						
Northeast	100.0	17.3	20.1	21.0	27.0	14.6
Midwest	100.0	26.9	24.0	19.7	17.6	11.9
South	100.0	20.1	25.0	18.0	26.2	10.7
West	100.0	11.7	34.5	19.1	29.6	5.1
Average length of stay in days						
Total	5.0	4.5	4.8	5.2	5.2	5.6
Sex						
Male	5.4	4.9	5.1	5.5	5.7	6.1
Female	4.7	4.3	4.6	5.0	4.9	5.3
Age						
Under 15 years	4.5	4.4	4.1	4.3	4.9	5.0
15–44 years	3.7	3.4	3.3	3.8	3.9	4.5
45–64 years	5.0	4.4	4.7	5.1	5.2	5.9
65 years and over	6.1	5.3	6.1	6.5	6.4	6.9
Region						
Northeast	5.7	5.6	5.7	5.6	5.8	6.0
Midwest	4.6	4.0	4.8	5.0	4.8	5.1
South	5.0	4.6	4.7	5.2	5.2	5.8
West	4.6	4.3	4.2	5.0	4.8	5.5

Table 6. Number and percent distribution of discharges from short-stay hospitals and of days of care, with average length of stay, by hospital ownership, sex, age and geographic region: United States, 1999

[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants]

	All	Proprietary	Government	Nonprofit
		Number of discharges in thousands		
All	32,132	3,349	3,896	24,887
Sex				
Male	12,748	1,243	1,542	9,962
Female	19,384	2,106	2,354	14,924
Age				
Under 15 years	2,458	173	307	1,978
15-44 years	10,092	1,147	1,431	7,515
45-64 years	6,899	645	906	5,348
65 years and over	12,683	1,384	1,252	10,046
Region				
Northeast	6,952	101	312	6,540
Midwest	7,368	160	551	6,657
South	12,007	2,414	2,292	7,300
West	5,805	673	742	4,390
		Percent distribution of patients discharged		
All	100.0	10.4	12.1	77.5
Sex				
Male	100.0	9.8	12.1	78.1
Female	100.0	10.9	12.1	77.0
Age				
Under 15 years	100.0	7.1	12.5	80.5
15-44 years	100.0	11.4	14.2	74.5
45-64 years	100.0	9.4	13.1	77.5
65 years and over	100.0	10.9	9.9	79.2
Region				
Northeast	100.0	1.5	4.5	94.1
Midwest	100.0	2.2	7.5	90.4
South	100.0	20.1	19.1	60.8
West	100.0	11.6	12.8	75.6
		Number of days of care in thousands		
Total	160,128	16,639	18,942	124,547
Sex				
Male	68,719	6,646	8,398	53,676
Female	91,409	9,994	10,544	70,871
Age				
Under 15 years	11,026	670	1,221	9,135
15-44 years	37,236	4,142	5,567	27,527
45-64 years	34,595	3,103	4,607	26,884
65 years and over	77,270	8,723	7,547	61,001
Region				
Northeast	39,814	490	2,058	37,266
Midwest	33,988	827	2,469	30,692
South	59,635	12,252	10,848	36,534
West	26,692	3,071	3,567	20,055
		Percent distribution of days of care		
All	100.0	10.4	11.8	77.8
Sex				
Male	100.0	9.7	12.2	78.1
Female	100.0	10.9	11.5	77.5

Table 6. Number and percent distribution of discharges from short-stay hospitals and of days of care, with average length of stay, by hospital ownership, sex, age and geographic region: United States, 1999—Con.

[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants]

	All	Proprietary	Government	Nonprofit
Age				
Under 15 years	100.0	6.1	11.1	82.8
15–44 years	100.0	11.1	15.0	73.9
45–64 years	100.0	9.0	13.3	77.7
65 years and over	100.0	11.3	9.8	78.9
Region				
Northeast	100.0	1.2	5.2	93.6
Midwest	100.0	2.4	7.3	90.3
South	100.0	20.5	18.2	61.3
West	100.0	11.5	13.4	75.1
Average length of stay in days				
Total	5.0	5.0	4.9	5.0
Sex				
Male	5.4	5.3	5.4	5.4
Female	4.7	4.7	4.5	4.7
Age				
Under 15 years	4.5	3.9	4.0	4.6
15–44 years	3.7	3.6	3.9	3.7
45–64 years	5.0	4.8	5.1	5.0
65 years and over	6.1	6.3	6.0	6.1
Region				
Northeast	5.7	4.9	6.6	5.7
Midwest	4.6	5.2	4.5	4.6
South	5.0	5.1	4.7	5.0
West	4.6	4.6	4.8	4.6

Table 7. Number and rate of deaths for discharges from short-stay hospitals, by sex and age: United States, 1999

[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants]

Age	Number of deaths in thousands			Rate per 100 discharges		
	Both sexes	Male	Female	Both sexes	Male	Female
All ages	851	416	435	2.6	3.3	2.2
Under 65 years	191	107	84	1.0	1.4	0.7
Under 15 years	13	7	7	0.5	0.5	0.6
15–44 years	46	25	21	0.5	0.9	0.3
45–64 years	132	75	57	1.9	2.2	1.6
65 years and over	660	309	351	5.2	5.9	4.7
65–74 years	189	96	94	3.9	4.1	3.6
75–84 years	273	141	132	5.2	6.5	4.3
85 years and over	197	72	125	7.7	9.4	6.9

Table 8. Number and rate of discharges from short-stay hospitals and of days of care, with average length of stay, by selected first-listed diagnostic categories: United States, 1999[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Diagnostic groupings and code numbers are based on the *International Classification of Diseases, 9th Revision, Clinical Modification (ICD-9-CM)*]

Diagnostic category and ICD-9-CM code	Discharges		Days of care		Average length of stay in days
	Number in thousands	Rate per 10,000 population	Number in thousands	Rate per 10,000 population	
All conditions ¹	32,132	1,166.2	160,128	5,811.5	5.0
Heart disease 391-392.0,393-398,402,404,410-416,420-429	4,465	162.0	20,939	759.9	4.7
Acute myocardial infarction 410	829	30.1	4,608	167.2	5.6
Coronary atherosclerosis 414.0	1,153	41.9	4,481	162.6	3.9
Other ischemic heart disease 411-413,414.1-414.9	280	10.2	831	30.2	3.0
Cardiac dysrhythmias 427	704	25.6	2,541	92.2	3.6
Congestive heart failure 428.0	962	34.9	5,296	192.2	5.5
Females with deliveries V27	3,810	138.3	9,421	341.9	2.5
Pneumonia 480-486	1,378	50.0	8,295	301.1	6.0
Psychoses 290-299	1,309	47.5	11,001	399.3	8.4
Malignant neoplasms 140-208,230-234	1,274	46.2	8,962	325.3	7.0
Malignant neoplasm of large intestine and rectum 153-154,197.5	166	6.0	1,529	55.5	9.2
Malignant neoplasm of trachea, bronchus, and lung 162,176.4,197.0,197.3	164	6.0	1,205	43.7	7.3
Malignant neoplasm of breast 174-175,198.81	106	3.9	292	10.6	2.7
Fractures, all sites ² 800-829	1,002	36.4	5,472	198.6	5.5
Fracture of neck of femur ² 820	323	11.7	2,115	76.8	6.5
Cerebrovascular disease 430-438	961	34.9	5,218	189.4	5.4
Diabetes mellitus 250	545	19.8	2,938	106.6	5.4
Chronic bronchitis 491	539	19.6	2,816	102.2	5.2
Asthma 493	478	17.4	1,540	55.9	3.2
Volume depletion 276.5	477	17.3	1,886	68.4	4.0
Osteoarthritis and allied disorders 715	434	15.7	1,962	71.2	4.5
Benign neoplasms 210-229	366	13.3	1,301	47.2	3.6
Cellulitis and abscess 681-682	358	13.0	1,864	67.7	5.2
Cholelithiasis 574	357	13.0	1,377	50.0	3.9
Septicemia 038	341	12.4	2,880	104.5	8.4
Intervertebral disc disorder 722	341	12.4	1,018	36.9	3.0

¹Includes data for diagnostic conditions not shown in table.²Excludes fractures coded as 733.1, pathologic fracture.

Table 9. Number and rate of discharges from short-stay hospitals with average length of stay, for patients 65 years of age and over by age and selected first-listed diagnostic categories: United States, 1999[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Diagnostic groupings and code numbers are based on the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM)]

Diagnostic category and ICD-9-CM code	65 years and over	65-74 years	75-84 years	85 years and over
Discharges in thousands				
All conditions ¹	12,683	4,883	5,227	2,572
Heart disease 391-392.0,393-398,402,404,410-416,420-429	2,889	1,154	1,192	543
Acute myocardial infarction 410	509	204	210	95
Coronary atherosclerosis 414.0	648	344	253	50
Other ischemic heart disease 411-413,414.1-414.9	151	65	56	30
Cardiac dysrhythmias 427	491	186	212	94
Congestive heart failure 428.0	757	221	327	209
Pneumonia 480-486	810	240	336	234
Malignant neoplasms 140-208,230-234	729	344	291	93
Malignant neoplasm of large intestine and rectum 153-154,197.5	115	44	49	23
Malignant neoplasm of trachea, bronchus, and lung 162,176.4,197.0,197.3	104	52	42	10
Cerebrovascular disease 430-438	704	245	310	149
Fractures, all sites ² 800-829	531	117	206	209
Fracture of neck of femur ² 820	292	44	113	135
Chronic bronchitis 491	369	175	154	39
Osteoarthritis and allied disorders 715	303	161	119	23
Volume depletion 276.5	247	61	115	71
Septicemia 038	241	67	106	68
Psychoses 290-299	212	92	79	42
Diabetes mellitus 250	199	100	80	19
Diverticula of intestine 562	145	51	53	41
Cholelithiasis 574	133	63	52	18
Cellulitis and abscess 681-682	126	45	46	35
Discharge rate per 10,000 population				
All conditions ¹	3,704.1	2,706.4	4,338.9	6,200.2
Heart disease 391-392.0,393-398,402,404,410-416,420-429	843.7	639.5	989.4	1,309.0
Acute myocardial infarction 410	148.6	113.2	174.2	228.4
Coronary atherosclerosis 414.0	189.2	190.9	210.1	120.8
Other ischemic heart disease 411-413,414.1-414.9	44.0	36.0	46.4	72.0
Cardiac dysrhythmias 427	143.5	103.1	175.6	225.9
Congestive heart failure 428.0	221.1	122.6	271.1	504.0
Pneumonia 480-486	236.5	132.8	279.2	563.2
Malignant neoplasms 140-208,230-234	212.8	190.7	241.9	223.8
Malignant neoplasm of large intestine and rectum 153-154,197.5	33.7	24.5	40.4	54.6
Malignant neoplasm of trachea, bronchus, and lung 162,176.4,197.0,197.3	30.5	29.0	34.7	24.6
Cerebrovascular disease 430-438	205.5	135.6	257.3	359.3
Fractures, all sites ² 800-829	155.2	64.9	170.7	503.1
Fracture of neck of femur ² 820	85.2	24.3	93.6	325.7
Chronic bronchitis 491	107.8	97.2	127.9	95.0
Osteoarthritis and allied disorders 715	88.6	89.5	99.0	54.7
Volume depletion 276.5	72.2	33.7	95.6	171.9
Septicemia 038	70.5	37.1	88.2	164.1
Psychoses 290-299	62.0	50.8	65.5	100.6
Diabetes mellitus 250	58.0	55.3	66.4	45.5
Diverticula of intestine 562	42.3	28.5	43.7	98.1
Cholelithiasis 574	38.8	35.1	43.1	42.7
Cellulitis and abscess 681-682	36.9	24.8	38.6	84.3
Average length of stay in days				
All conditions ¹	6.1	5.8	6.2	6.4
Heart disease 391-392.0,393-398,402,404,410-416,420-429	5.0	4.8	5.2	5.3
Acute myocardial infarction 410	6.2	6.1	6.1	6.8
Coronary atherosclerosis 414.0	4.2	4.3	4.1	4.7
Other ischemic heart disease 411-413,414.1-414.9	3.2	3.2	3.1	3.5
Cardiac dysrhythmias 427	3.9	3.6	3.9	4.5
Congestive heart failure 428.0	5.6	5.2	6.0	5.2
Pneumonia 480-486	6.6	6.0	6.8	7.0
Malignant neoplasms 140-208,230-234	7.3	6.7	7.5	8.9
Malignant neoplasm of large intestine and rectum 153-154,197.5	9.8	8.9	9.7	11.9
Malignant neoplasm of trachea, bronchus, and lung 162,176.4,197.0,197.3	7.6	7.8	7.1	7.9
Cerebrovascular disease 430-438	5.4	5.1	5.3	6.1

Table 9. Number and rate of discharges from short-stay hospitals with average length of stay, for patients 65 years of age and over by age and selected first-listed diagnostic categories: United States, 1999—Con.[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Diagnostic groupings and code numbers are based on the *International Classification of Diseases, 9th Revision, Clinical Modification (ICD-9-CM)*]

Diagnostic category and ICD-9-CM code	65 years and over	65-74 years	75-84 years	85 years and over
Fractures, all sites ² 800-829	6.3	5.9	6.3	6.6
Fracture of neck of femur ² 820	6.6	6.6	6.5	6.8
Chronic bronchitis 491	5.5	5.2	5.7	6.4
Osteoarthritis and allied disorders 715	4.7	4.8	4.6	4.5
Volume depletion 276.5	5.0	4.2	5.5	4.9
Septicemia 038	8.8	10.5	8.6	7.3
Psychoses 290-299	10.4	10.8	10.3	9.8
Diabetes mellitus 250	6.3	6.2	6.6	5.5
Diverticula of intestine 562	6.1	6.7	6.1	5.4
Cholelithiasis 574	5.1	4.4	5.5	6.5
Cellulitis and abscess 681-682	6.3	5.4	6.6	7.1

* Figure does not meet standard of reliability or precision.

¹Includes data for diagnostic conditions not shown in table.²Excludes fractures coded as 733.1, pathologic fracture.

Table 10. Number of discharges from short-stay hospitals, by age and first-listed diagnosis: United States, 1999[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Diagnostic groupings and code numbers are based on the *International Classification of Diseases, 9th Revision, Clinical Modification (ICD-9-CM)*]

Category of first-listed diagnosis and ICD-9-CM code	All ages	Under 15 years	15-44 years	45-64 years	65 years and over
	Number in thousands				
All conditions	32,132	2,458	10,092	6,899	12,683
Infectious and parasitic diseases 001-139	826	173	186	133	334
Septicemia 038	341	19	26	55	241
Neoplasms 140-239	1,687	36	283	565	802
Malignant neoplasms 140-208,230-234	1,274	23	119	405	729
Malignant neoplasm of large intestine and rectum 153-154,197.5	166	*	*7	42	115
Malignant neoplasm of trachea, bronchus, and lung 162,176.4,197.0,197.3	164	*	5	54	104
Malignant neoplasm of breast 174-175,198.81	106	*	13	42	52
Benign neoplasms 210-229	366	*	156	148	50
Endocrine, nutritional and metabolic diseases, and immunity disorders 240-279	1,393	159	264	351	618
Diabetes mellitus 250	545	21	138	187	199
Volume depletion 276.5	477	115	52	63	247
Diseases of the blood and blood-forming organs 280-289	368	57	96	74	141
Mental disorders 290-319	2,018	103	1,167	484	264
Psychoses 290-299	1,309	44	731	321	212
Alcohol dependence syndrome 303	164	*	92	60	11
Diseases of the nervous system and sense organs 320-389	503	89	115	112	188
Diseases of the circulatory system 390-459	6,344	25	417	1,797	4,105
Heart disease 391-392.0,393-398,402,404,410-416,420-429	4,465	15	263	1,299	2,889
Acute myocardial infarction 410	829	*	46	274	509
Coronary atherosclerosis 414.0	1,153	*	49	455	648
Other ischemic heart disease 411-413,414.1-414.9	280	*	25	104	151
Cardiac dysrhythmias 427	704	*	53	156	491
Congestive heart failure 428.0	962	*	30	173	757
Cerebrovascular disease 430-438	961	*2	40	214	704
Diseases of the respiratory system 460-519	3,689	741	430	686	1,832
Acute bronchitis and bronchiolitis 466	298	204	19	20	56
Pneumonia 480-486	1,378	211	130	228	810
Chronic bronchitis 491	539	*	16	153	369
Asthma 493	478	190	122	94	73
Diseases of the digestive system 520-579	3,122	221	747	830	1,324
Appendicitis 540-543	260	59	143	39	19
Noninfectious enteritis and colitis 555-558	272	63	68	49	91
Diverticula of intestine 562	239	*	23	70	145
Cholelithiasis 574	357	*	114	109	133
Diseases of the genitourinary system 580-629	1,719	81	568	428	641
Calculus of kidney and ureter 592	192	*	82	75	34
Complications of pregnancy, childbirth, and the puerperium ¹ 630-677	495	*	491	*	...
Diseases of the skin and subcutaneous tissue 680-709	518	*	133	134	182
Cellulitis and abscess 681-682	358	32	94	105	126
Diseases of the musculoskeletal system and connective tissue 710-739	1,543	39	326	481	697
Osteoarthritis and allied disorders 715	434	*	13	118	303
Intervertebral disc disorders 722	341	*	144	136	60
Congenital anomalies 740-759	182	130	34	11	8
Certain conditions originating in the perinatal period 760-779	170	170	*	*	*
Symptoms, signs, and ill-defined conditions 780-799	313	58	110	77	69
Injury and poisoning 800-999	2,565	228	787	526	1,024
Fractures, all sites ² 800-829	1,002	64	245	162	531
Fracture of neck of femur ² 820	323	*	9	21	292
Poisonings 960-989	182	17	105	37	23
Supplementary classifications V01-V82	4,676	77	3,939	207	453
Females with deliveries V27	3,810	7	3,797	*6	...

* Figure does not meet standard of reliability or precision.

... Category not applicable.

¹First-listed diagnosis for females with deliveries is coded V27, shown under "Supplementary classifications."²Excludes fractures coded as 733.1, pathologic fracture.

Table 11. Rate of discharges from short-stay hospitals, by age and first-listed diagnosis: United States, 1999[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Diagnostic groupings and code numbers are based on the *International Classification of Diseases, 9th Revision, Clinical Modification (ICD-9-CM)*]

Category of first-listed diagnosis and ICD-9-CM code	All ages	Rate per 10,000 population			
		Under 15 years	15-44 years	45-64 years	65 years and over
All conditions	1,166.2	408.0	827.1	1,169.0	3,704.1
Infectious and parasitic diseases 001-139	30.0	28.7	15.2	22.6	97.6
Septicemia 038	12.4	3.2	2.1	9.3	70.5
Neoplasms 140-239	61.2	6.0	23.2	95.8	234.4
Malignant neoplasms 140-208,230-234	46.2	3.7	9.7	68.5	212.8
Malignant neoplasm of large intestine and rectum 153-154,197.5	6.0	*	*0.6	7.2	33.7
Malignant neoplasm of trachea, bronchus, and lung 162,176.4,197.0,197.3	6.0	*	0.4	9.2	30.5
Malignant neoplasm of breast 174-175,198.81	3.9	*	1.0	7.1	15.2
Benign neoplasms 210-229	13.3	*	12.8	25.2	14.7
Endocrine, nutritional and metabolic diseases, and immunity disorders 240-279	50.5	26.5	21.6	59.5	180.5
Diabetes mellitus 250	19.8	3.5	11.3	31.7	58.0
Volume depletion 276.5	17.3	19.1	4.2	10.7	72.2
Diseases of the blood and blood-forming organs 280-289	13.4	9.5	7.9	12.5	41.1
Mental disorders 290-319	73.2	17.1	95.6	82.1	77.1
Psychoses 290-299	47.5	7.4	59.9	54.4	62.0
Alcohol dependence syndrome 303	6.0	*	7.5	10.2	3.2
Diseases of the nervous system and sense organs 320-389	18.3	14.7	9.4	18.9	54.9
Diseases of the circulatory system 390-459	230.2	4.1	34.2	304.5	1,199.0
Heart disease 391-392.0,393-398,402,404,410-416,420-429	162.0	2.4	21.5	220.1	843.7
Acute myocardial infarction 410	30.1	*	3.7	46.4	148.6
Coronary atherosclerosis 414.0	41.9	*	4.0	77.1	189.2
Other ischemic heart disease 411-413,414.1-414.9	10.2	*	2.0	17.7	44.0
Cardiac dysrhythmias 427	25.6	*	4.3	26.4	143.5
Congestive heart failure 428.0	34.9	*	2.5	29.4	221.1
Cerebrovascular disease 430-438	34.9	*0.4	3.3	36.3	205.5
Diseases of the respiratory system 460-519	133.9	122.9	35.3	116.3	535.0
Acute bronchitis and bronchiolitis 466	10.8	33.8	1.5	3.4	16.2
Pneumonia 480-486	50.0	35.1	10.6	38.6	236.5
Chronic bronchitis 491	19.6	*	1.3	25.9	107.8
Asthma 493	17.4	31.5	10.0	15.9	21.2
Diseases of the digestive system 520-579	113.3	36.6	61.2	140.6	386.8
Appendicitis 540-543	9.4	9.8	11.7	6.6	5.6
Noninfectious enteritis and colitis 555-558	9.9	10.5	5.6	8.3	26.6
Diverticula of intestine 562	8.7	*	1.9	11.9	42.3
Cholelithiasis 574	13.0	*	9.3	18.4	38.8
Diseases of the genitourinary system 580-629	62.4	13.5	46.6	72.6	187.2
Calculus of kidney and ureter 592	7.0	*	6.7	12.7	9.8
Complications of pregnancy, childbirth, and the puerperium ¹ 630-677	18.0	*	40.3	*	...
Diseases of the skin and subcutaneous tissue 680-709	18.8	*	10.9	22.6	53.1
Cellulitis and abscess 681-682	13.0	5.4	7.7	17.8	36.9
Diseases of the musculoskeletal system and connective tissue 710-739	56.0	6.5	26.7	81.6	203.6
Osteoarthritis and allied disorders 715	15.7	*	1.0	19.9	88.6
Intervertebral disc disorders 722	12.4	*	11.8	23.1	17.5
Congenital anomalies 740-759	6.6	21.6	2.8	1.8	2.3
Certain conditions originating in the perinatal period 760-779	6.2	28.2	*	*	*
Symptoms, signs, and ill-defined conditions 780-799	11.4	9.6	9.0	13.0	20.1
Injury and poisoning 800-999	93.1	37.9	64.5	89.1	299.1
Fractures, all sites ² 800-829	36.4	10.6	20.1	27.4	155.2
Fracture of neck of femur ² 820	11.7	*	0.8	3.5	85.2
Poisonings 960-989	6.6	2.8	8.6	6.2	6.7
Supplementary classifications V01-V82	169.7	12.8	322.8	35.1	132.4
Females with deliveries V27	138.3	1.2	311.2	*0.9	...

* Figure does not meet standard of reliability or precision.

... Category not applicable.

¹First-listed diagnosis for females with deliveries is coded V27, shown under "Supplementary classifications."²Excludes fractures coded as 733.1, pathologic fracture.

Table 12. Average length of stay for discharges from short-stay hospitals, by age and first-listed diagnosis: United States, 1999[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Diagnostic groupings and code numbers are based on the *International Classification of Diseases, 9th Revision, Clinical Modification (ICD-9-CM)*]

Category of first-listed diagnosis and ICD-9-CM code	All ages	Under 15 years	15-44 years	45-64 years	65 years and over
		Average length of stay in days			
All conditions	5.0	4.5	3.7	5.0	6.1
Infectious and parasitic diseases 001-139	6.4	3.0	5.8	7.6	8.0
Septicemia 038	8.4	4.2	7.8	8.8	8.8
Neoplasms 140-239	6.2	7.4	4.3	5.8	7.2
Malignant neoplasms 140-208,230-234	7.0	10.0	5.7	6.7	7.3
Malignant neoplasm of large intestine and rectum 153-154,197.5	9.2	*	*8.7	7.7	9.8
Malignant neoplasm of trachea, bronchus, and lung 162,176.4,197.0,197.3	7.3	*	6.1	7.0	7.6
Malignant neoplasm of breast 174-175,198.81	2.7	*	2.7	2.7	2.8
Benign neoplasms 210-229	3.6	2.4	3.2	3.3	5.8
Endocrine, nutritional and metabolic diseases, and immunity disorders 240-279	4.7	2.9	3.8	4.9	5.5
Diabetes mellitus 250	5.4	2.7	4.2	5.6	6.3
Volume depletion 276.5	4.0	2.3	2.8	3.7	5.0
Diseases of the blood and blood-forming organs 280-289	4.8	4.2	5.1	4.9	4.8
Mental disorders 290-319	7.5	11.7	6.7	7.2	9.9
Psychoses 290-299	8.4	10.4	7.5	8.8	10.4
Alcohol dependence syndrome 303	5.3	*	5.5	4.7	6.4
Diseases of the nervous system and sense organs 320-389	5.1	3.9	5.0	4.7	6.0
Diseases of the circulatory system 390-459	4.9	6.1	4.0	4.3	5.2
Heart disease 391-392.0,393-398,402,404,410-416,420-429	4.7	6.5	3.8	4.1	5.0
Acute myocardial infarction 410	5.6	*	3.8	4.7	6.2
Coronary atherosclerosis 414.0	3.9	*	3.2	3.4	4.2
Other ischemic heart disease 411-413,414.1-414.9	3.0	*	2.2	2.8	3.2
Cardiac dysrhythmias 427	3.6	*	2.4	3.1	3.9
Congestive heart failure 428.0	5.5	*	5.8	5.2	5.6
Cerebrovascular disease 430-438	5.4	*	5.4	5.5	5.4
Diseases of the respiratory system 460-519	5.4	3.1	4.6	5.3	6.6
Acute bronchitis and bronchiolitis 466	3.3	2.9	3.1	3.5	4.6
Pneumonia 480-486	6.0	3.9	6.0	5.7	6.6
Chronic bronchitis 491	5.2	*	4.3	4.6	5.5
Asthma 493	3.2	2.3	3.2	3.7	5.0
Diseases of the digestive system 520-579	4.8	3.6	3.7	4.7	5.7
Appendicitis 540-543	3.5	3.4	2.9	3.7	7.2
Noninfectious enteritis and colitis 555-558	4.1	2.5	4.1	5.0	4.7
Diverticula of intestine 562	5.6	*	4.7	5.0	6.1
Cholelithiasis 574	3.9	*4.7	2.7	3.5	5.1
Diseases of the genitourinary system 580-629	3.8	3.2	2.8	3.5	5.0
Calculus of kidney and ureter 592	2.3	*	2.1	2.2	3.0
Complications of pregnancy, childbirth, and the puerperium ¹ 630-677	2.6	*	2.6	*	...
Diseases of the skin and subcutaneous tissue 680-709	5.7	4.7	4.6	5.8	6.9
Cellulitis and abscess 681-682	5.2	3.4	4.1	5.5	6.3
Diseases of the musculoskeletal system and connective tissue 710-739	4.3	4.3	3.3	3.7	5.2
Osteoarthritis and allied disorders 715	4.5	*	3.5	4.2	4.7
Intervertebral disc disorders 722	3.0	*	2.8	2.7	4.2
Congenital anomalies 740-759	5.7	6.3	3.8	5.5	4.9
Certain conditions originating in the perinatal period 760-779	9.4	9.4	*	*	*
Symptoms, signs, and ill-defined conditions 780-799	3.2	3.2	2.3	2.7	5.1
Injury and poisoning 800-999	5.4	4.4	4.4	5.4	6.3
Fractures, all sites ² 800-829	5.5	2.9	4.4	5.2	6.3
Fracture of neck of femur ² 820	6.5	*	5.4	5.7	6.6
Poisonings 960-989	2.5	2.3	2.1	2.9	4.2
Supplementary classifications V01-V82	3.7	4.3	2.6	8.3	11.2
Females with deliveries V27	2.5	2.5	2.5	*4.2	...

* Figure does not meet standard of reliability or precision.

... Category not applicable.

¹First-listed diagnosis for females with deliveries is coded V27, shown under "Supplementary classifications."²Excludes fractures coded as 733.1, pathologic fracture.

Table 13. Number of discharges from short-stay hospitals, by sex and first-listed diagnosis: United States, 1999[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Diagnostic groupings and code numbers are based on the *International Classification of Diseases, 9th Revision, Clinical Modification (ICD-9-CM)*]

Category of first-listed diagnosis and ICD-9-CM code	Both sexes	Male	Female
All conditions	32,132	12,748	19,384
Infectious and parasitic diseases 001-139	826	385	441
Septicemia 038	341	146	195
Neoplasms 140-239	1,687	655	1,032
Malignant neoplasms 140-208,230-234	1,274	596	678
Malignant neoplasm of large intestine and rectum 153-154,197.5	166	80	86
Malignant neoplasm of trachea, bronchus, and lung 162,176.4,197.0,197.3	164	91	73
Malignant neoplasm of breast 174-175,198.81	106	*	106
Benign neoplasms 210-229	366	44	322
Endocrine, nutritional and metabolic diseases, and immunity disorders 240-279	1,393	577	816
Diabetes mellitus 250	545	262	283
Volume depletion 276.5	477	186	292
Diseases of the blood and blood-forming organs 280-289	368	162	206
Mental disorders 290-319	2,018	1,048	970
Psychoses 290-299	1,309	640	669
Alcohol dependence syndrome 303	164	119	45
Diseases of the nervous system and sense organs 320-389	503	226	278
Diseases of the circulatory system 390-459	6,344	3,161	3,183
Heart disease 391-392.0,393-398,402,404,410-416,420-429	4,465	2,302	2,163
Acute myocardial infarction 410	829	481	348
Coronary atherosclerosis 414.0	1,153	703	450
Other ischemic heart disease 411-413,414.1-414.9	280	133	147
Cardiac dysrhythmias 427	704	329	375
Congestive heart failure 428.0	962	430	532
Cerebrovascular disease 430-438	961	434	527
Diseases of the respiratory system 460-519	3,689	1,742	1,947
Acute bronchitis and bronchiolitis 466	298	153	145
Pneumonia 480-486	1,378	672	707
Chronic bronchitis 491	539	228	311
Asthma 493	478	190	288
Diseases of the digestive system 520-579	3,122	1,353	1,769
Appendicitis 540-543	260	148	112
Noninfectious enteritis and colitis 555-558	272	107	165
Diverticula of intestine 562	239	93	146
Cholelithiasis 574	357	110	247
Diseases of the genitourinary system 580-629	1,719	534	1,185
Calculus of kidney and ureter 592	192	118	73
Complications of pregnancy, childbirth, and the puerperium ¹ 630-677	495	...	495
Diseases of the skin and subcutaneous tissue 680-709	518	255	263
Cellulitis and abscess 681-682	358	183	175
Diseases of the musculoskeletal system and connective tissue 710-739	1,543	675	868
Osteoarthritis and allied disorders 715	434	169	265
Intervertebral disc disorders 722	341	179	161
Congenital anomalies 740-759	182	94	88
Certain conditions originating in the perinatal period 760-779	170	102	68
Symptoms, signs, and ill-defined conditions 780-799	313	146	168
Injury and poisoning 800-999	2,565	1,248	1,317
Fractures, all sites ² 800-829	1,002	419	583
Fracture of neck of femur ² 820	323	81	243
Poisonings 960-989	182	80	102
Supplementary classifications V01-V82	4,676	386	4,290
Females with deliveries V27	3,810	...	3,810

* Figure does not meet standard of reliability or precision.

... Category not applicable.

¹First-listed diagnosis for females with deliveries is coded V27, shown under "Supplementary classifications."²Excludes fractures coded as 733.1, pathologic fracture.

Table 14. Rate of discharges from short-stay hospitals, by sex and first-listed diagnosis: United States, 1999[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Diagnostic groupings and code numbers are based on the *International Classification of Diseases, 9th Revision, Clinical Modification (ICD-9-CM)*]

Category of first-listed diagnosis and ICD-9-CM code	Both sexes	Rate per 10,000 population	
		Male	Female
All conditions	1,166.2	946.8	1,375.8
Infectious and parasitic diseases 001-139	30.0	28.6	31.3
Septicemia 038	12.4	10.9	13.8
Neoplasms 140-239	61.2	48.6	73.2
Malignant neoplasms 140-208,230-234	46.2	44.3	48.2
Malignant neoplasm of large intestine and rectum 153-154,197.5	6.0	5.9	6.1
Malignant neoplasm of trachea, bronchus, and lung 162,176.4,197.0,197.3	6.0	6.8	5.2
Malignant neoplasm of breast 174-175,198.81	3.9	*	7.5
Benign neoplasms 210-229	13.3	3.3	22.8
Endocrine, nutritional and metabolic diseases, and immunity disorders 240-279	50.5	42.9	57.9
Diabetes mellitus 250	19.8	19.4	20.1
Volume depletion 276.5	17.3	13.8	20.7
Diseases of the blood and blood-forming organs 280-289	13.4	12.0	14.7
Mental disorders 290-319	73.2	77.8	68.9
Psychoses 290-299	47.5	47.5	47.5
Alcohol dependence syndrome 303	6.0	8.8	3.2
Diseases of the nervous system and sense organs 320-389	18.3	16.8	19.7
Diseases of the circulatory system 390-459	230.2	234.8	225.9
Heart disease 391-392.0,393-398,402,404,410-416,420-429	162.0	170.9	153.5
Acute myocardial infarction 410	30.1	35.7	24.7
Coronary atherosclerosis 414.0	41.9	52.2	32.0
Other ischemic heart disease 411-413,414.1-414.9	10.2	9.9	10.4
Cardiac dysrhythmias 427	25.6	24.4	26.6
Congestive heart failure 428.0	34.9	31.9	37.8
Cerebrovascular disease 430-438	34.9	32.2	37.4
Diseases of the respiratory system 460-519	133.9	129.4	138.2
Acute bronchitis and bronchiolitis 466	10.8	11.4	10.3
Pneumonia 480-486	50.0	49.9	50.2
Chronic bronchitis 491	19.6	16.9	22.1
Asthma 493	17.4	14.1	20.5
Diseases of the digestive system 520-579	113.3	100.5	125.5
Appendicitis 540-543	9.4	11.0	7.9
Noninfectious enteritis and colitis 555-558	9.9	7.9	11.7
Diverticula of intestine 562	8.7	6.9	10.3
Cholelithiasis 574	13.0	8.2	17.5
Diseases of the genitourinary system 580-629	62.4	39.7	84.1
Calculus of kidney and ureter 592	7.0	8.8	5.2
Complications of pregnancy, childbirth, and the puerperium ¹ 630-677	18.0	...	35.1
Diseases of the skin and subcutaneous tissue 680-709	18.8	18.9	18.7
Cellulitis and abscess 681-682	13.0	13.6	12.4
Diseases of the musculoskeletal system and connective tissue 710-739	56.0	50.2	61.6
Osteoarthritis and allied disorders 715	15.7	12.5	18.8
Intervertebral disc disorders 722	12.4	13.3	11.5
Congenital anomalies 740-759	6.6	7.0	6.3
Certain conditions originating in the perinatal period 760-779	6.2	7.6	4.8
Symptoms, signs, and ill-defined conditions 780-799	11.4	10.8	11.9
Injury and poisoning 800-999	93.1	92.7	93.5
Fractures, all sites ² 800-829	36.4	31.1	41.4
Fracture of neck of femur ² 820	11.7	6.0	17.2
Poisonings 960-989	6.6	5.9	7.2
Supplementary classifications V01-V82	169.7	28.6	304.5
Females with deliveries V27	138.3	...	270.4

* Figure does not meet standard of reliability or precision.

... Category not applicable.

¹First-listed diagnosis for females with deliveries is coded V27, shown under "Supplementary classifications."²Excludes fractures coded as 733.1, pathologic fracture.

Table 15. Average length of stay for discharges from short-stay hospitals, by sex and first-listed diagnosis: United States, 1999[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Diagnostic groupings and code numbers are based on the *International Classification of Diseases, 9th Revision, Clinical Modification (ICD-9-CM)*]

Category of first-listed diagnosis and ICD-9-CM code	Both sexes	Average length of stay in days	
		Male	Female
All conditions	5.0	5.4	4.7
Infectious and parasitic diseases 001-139	6.4	6.5	6.3
Septicemia 038	8.4	8.8	8.2
Neoplasms 140-239	6.2	7.0	5.7
Malignant neoplasms 140-208,230-234	7.0	7.2	6.9
Malignant neoplasm of large intestine and rectum 153-154,197.5	9.2	8.9	9.5
Malignant neoplasm of trachea, bronchus, and lung 162,176.4,197.0,197.3	7.3	7.0	7.8
Malignant neoplasm of breast 174-175,198.81	2.7	*	2.8
Benign neoplasms 210-229	3.6	4.7	3.4
Endocrine, nutritional and metabolic diseases, and immunity disorders 240-279	4.7	4.8	4.7
Diabetes mellitus 250	5.4	5.5	5.3
Volume depletion 276.5	4.0	3.5	4.3
Diseases of the blood and blood-forming organs 280-289	4.8	4.6	5.0
Mental disorders 290-319	7.5	7.5	7.5
Psychoses 290-299	8.4	8.4	8.4
Alcohol dependence syndrome 303	5.3	5.1	5.7
Diseases of the nervous system and sense organs 320-389	5.1	5.2	5.0
Diseases of the circulatory system 390-459	4.9	4.7	5.0
Heart disease 391-392.0,393-398,402,404,410-416,420-429	4.7	4.5	4.9
Acute myocardial infarction 410	5.6	5.3	6.0
Coronary atherosclerosis 414.0	3.9	3.7	4.1
Other ischemic heart disease 411-413,414.1-414.9	3.0	2.9	3.1
Cardiac dysrhythmias 427	3.6	3.4	3.8
Congestive heart failure 428.0	5.5	5.4	5.6
Cerebrovascular disease 430-438	5.4	5.4	5.5
Diseases of the respiratory system 460-519	5.4	5.3	5.6
Acute bronchitis and bronchiolitis 466	3.3	3.1	3.5
Pneumonia 480-486	6.0	5.9	6.1
Chronic bronchitis 491	5.2	4.9	5.5
Asthma 493	3.2	2.8	3.5
Diseases of the digestive system 520-579	4.8	4.7	4.9
Appendicitis 540-543	3.5	3.7	3.2
Noninfectious enteritis and colitis 555-558	4.1	3.9	4.2
Diverticula of intestine 562	5.6	5.3	5.9
Cholelithiasis 574	3.9	4.5	3.6
Diseases of the genitourinary system 580-629	3.8	4.4	3.6
Calculus of kidney and ureter 592	2.3	2.1	2.7
Complications of pregnancy, childbirth, and the puerperium ¹ 630-677	2.6	...	2.6
Diseases of the skin and subcutaneous tissue 680-709	5.7	5.8	5.6
Cellulitis and abscess 681-682	5.2	5.1	5.3
Diseases of the musculoskeletal system and connective tissue 710-739	4.3	4.2	4.4
Osteoarthritis and allied disorders 715	4.5	4.6	4.5
Intervertebral disc disorders 722	3.0	2.9	3.1
Congenital anomalies 740-759	5.7	5.3	6.2
Certain conditions originating in the perinatal period 760-779	9.4	9.4	9.5
Symptoms, signs, and ill-defined conditions 780-799	3.2	3.1	3.2
Injury and poisoning 800-999	5.4	5.3	5.5
Fractures, all sites ² 800-829	5.5	5.3	5.6
Fracture of neck of femur ² 820	6.5	7.0	6.4
Poisonings 960-989	2.5	2.7	2.4
Supplementary classifications V01-V82	3.7	8.9	3.2
Females with deliveries V27	2.5	...	2.5

* Figure does not meet standard of reliability or precision.

... Category not applicable.

¹First-listed diagnosis for females with deliveries is coded V27, shown under "Supplementary classifications."²Excludes fractures coded as 733.1, pathologic fracture.

Table 16. Number of discharges from short-stay hospitals, by geographic region and first-listed diagnosis: United States, 1999[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Diagnostic groupings and code numbers are based on the *International Classification of Diseases, 9th Revision, Clinical Modification (ICD-9-CM)*]

Category of first-listed diagnosis and ICD-9-CM code	United States	North-east	Midwest	South	West
	Number in thousands				
All conditions	32,132	6,952	7,368	12,007	5,805
Infectious and parasitic diseases 001-139	826	198	171	318	139
Septicemia 038	341	82	72	131	56
Neoplasms 140-239	1,687	396	373	599	318
Malignant neoplasms 140-208,230-234	1,274	312	290	436	236
Malignant neoplasm of large intestine and rectum 153-154,197.5	166	42	43	52	30
Malignant neoplasm of trachea, bronchus, and lung 162,176.4,197.0,197.3	164	36	39	58	32
Malignant neoplasm of breast 174-175,198.81	106	31	25	36	15
Benign neoplasms 210-229	366	75	73	142	77
Endocrine, nutritional and metabolic diseases, and immunity disorders 240-279	1,393	316	329	518	229
Diabetes mellitus 250	545	119	115	214	97
Volume depletion 276.5	477	106	130	171	71
Diseases of the blood and blood-forming organs 280-289	368	88	83	136	61
Mental disorders 290-319	2,018	615	475	718	210
Psychoses 290-299	1,309	405	318	458	127
Alcohol dependence syndrome 303	164	28	40	*	*
Diseases of the nervous system and sense organs 320-389	503	105	123	190	86
Diseases of the circulatory system 390-459	6,344	1,485	1,523	2,328	1,009
Heart disease 391-392.0,393-398,402,404,410-416,420-429	4,465	1,080	1,071	1,608	705
Acute myocardial infarction 410	829	207	208	266	148
Coronary atherosclerosis 414.0	1,153	274	268	415	197
Other ischemic heart disease 411-413,414.1-414.9	280	65	59	119	37
Cardiac dysrhythmias 427	704	186	166	246	107
Congestive heart failure 428.0	962	214	258	364	126
Cerebrovascular disease 430-438	961	193	246	360	162
Diseases of the respiratory system 460-519	3,689	771	901	1,378	640
Acute bronchitis and bronchiolitis 466	298	52	65	123	58
Pneumonia 480-486	1,378	252	362	523	242
Chronic bronchitis 491	539	113	149	207	69
Asthma 493	478	136	107	154	81
Diseases of the digestive system 520-579	3,122	649	717	1,157	598
Appendicitis 540-543	260	48	64	75	74
Noninfectious enteritis and colitis 555-558	272	65	61	105	41
Diverticula of intestine 562	239	57	55	88	39
Cholelithiasis 574	357	82	70	129	77
Diseases of the genitourinary system 580-629	1,719	330	382	693	314
Calculus of kidney and ureter 592	192	34	52	79	27
Complications of pregnancy, childbirth, and the puerperium ¹ 630-677	495	89	90	220	96
Diseases of the skin and subcutaneous tissue 680-709	518	113	119	209	77
Cellulitis and abscess 681-682	358	90	89	121	57
Diseases of the musculoskeletal system and connective tissue 710-739	1,543	310	369	556	308
Osteoarthritis and allied disorders 715	434	95	120	130	90
Intervertebral disc disorders 722	341	61	82	138	59
Congenital anomalies 740-759	182	53	*	*	*
Certain conditions originating in the perinatal period 760-779	170	27	*	50	*
Symptoms, signs, and ill-defined conditions 780-799	313	57	50	148	57
Injury and poisoning 800-999	2,565	541	587	938	500
Fractures, all sites ² 800-829	1,002	205	245	356	196
Fracture of neck of femur ² 820	323	67	75	124	58
Poisonings 960-989	182	35	39	72	36
Supplementary classifications V01-V82	4,676	809	1,016	1,786	1,065
Females with deliveries V27	3,810	636	809	1,454	912

* Figure does not meet standard of reliability or precision.

¹First-listed diagnosis for females with deliveries is coded V27, shown under "Supplementary classifications."²Excludes fractures coded as 733.1, pathologic fracture.

Table 17. Rate of discharges from short-stay hospitals, by geographic region and first-listed diagnosis: United States, 1999[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Diagnostic groupings and code numbers are based on the *International Classification of Diseases, 9th Revision, Clinical Modification (ICD-9-CM)*]

Category of first-listed diagnosis and ICD-9-CM code	United States	North-east	Midwest	South	West
All conditions	1,166.2	1,331.6	1,159.1	1,229.9	934.2
Infectious and parasitic diseases 001-139	30.0	37.9	26.9	32.6	22.3
Septicemia 038	12.4	15.7	11.4	13.4	9.0
Neoplasms 140-239	61.2	75.9	58.7	61.3	51.2
Malignant neoplasms 140-208,230-234	46.2	59.8	45.6	44.6	38.0
Malignant neoplasm of large intestine and rectum 153-154,197.5	6.0	8.0	6.7	5.3	4.8
Malignant neoplasm of trachea, bronchus, and lung 162,176.4,197.0,197.3	6.0	6.9	6.1	5.9	5.1
Malignant neoplasm of breast 174-175,198.81	3.9	5.9	3.9	3.7	2.4
Benign neoplasms 210-229	13.3	14.3	11.5	14.5	12.3
Endocrine, nutritional and metabolic diseases, and immunity disorders 240-279	50.5	60.5	51.8	53.1	36.9
Diabetes mellitus 250	19.8	22.8	18.2	21.9	15.5
Volume depletion 276.5	17.3	20.2	20.4	17.6	11.3
Diseases of the blood and blood-forming organs 280-289	13.4	16.9	13.1	13.9	9.8
Mental disorders 290-319	73.2	117.7	74.7	73.5	33.9
Psychoses 290-299	47.5	77.6	50.0	46.9	20.5
Alcohol dependence syndrome 303	6.0	5.4	6.3	*	*
Diseases of the nervous system and sense organs 320-389	18.3	20.0	19.4	19.5	13.8
Diseases of the circulatory system 390-459	230.2	284.4	239.5	238.4	162.4
Heart disease 391-392.0,393-398,402,404,410-416,420-429	162.0	206.9	168.5	164.7	113.5
Acute myocardial infarction 410	30.1	39.6	32.7	27.3	23.8
Coronary atherosclerosis 414.0	41.9	52.4	42.2	42.5	31.6
Other ischemic heart disease 411-413,414.1-414.9	10.2	12.4	9.3	12.2	5.9
Cardiac dysrhythmias 427	25.6	35.6	26.1	25.2	17.2
Congestive heart failure 428.0	34.9	41.1	40.5	37.3	20.3
Cerebrovascular disease 430-438	34.9	37.0	38.6	36.9	26.1
Diseases of the respiratory system 460-519	133.9	147.6	141.7	141.1	102.9
Acute bronchitis and bronchiolitis 466	10.8	9.9	10.2	12.6	9.4
Pneumonia 480-486	50.0	48.2	56.9	53.5	39.0
Chronic bronchitis 491	19.6	21.7	23.5	21.2	11.1
Asthma 493	17.4	26.0	16.8	15.8	13.0
Diseases of the digestive system 520-579	113.3	124.4	112.8	118.5	96.2
Appendicitis 540-543	9.4	9.1	10.0	7.6	12.0
Noninfectious enteritis and colitis 555-558	9.9	12.4	9.7	10.7	6.6
Diverticula of intestine 562	8.7	10.9	8.6	9.0	6.2
Cholelithiasis 574	13.0	15.7	11.0	13.2	12.3
Diseases of the genitourinary system 580-629	62.4	63.2	60.1	71.0	50.5
Calculus of kidney and ureter 592	7.0	6.5	8.1	8.1	4.4
Complications of pregnancy, childbirth, and the puerperium ¹ 630-677	18.0	17.1	14.2	22.5	15.4
Diseases of the skin and subcutaneous tissue 680-709	18.8	21.7	18.7	21.4	12.3
Cellulitis and abscess 681-682	13.0	17.2	14.1	12.4	9.2
Diseases of the musculoskeletal system and connective tissue 710-739	56.0	59.5	58.1	57.0	49.5
Osteoarthritis and allied disorders 715	15.7	18.1	18.8	13.3	14.4
Intervertebral disc disorders 722	12.4	11.6	13.0	14.1	9.5
Congenital anomalies 740-759	6.6	10.1	*	*	*
Certain conditions originating in the perinatal period 760-779	6.2	5.3	*	5.2	*
Symptoms, signs, and ill-defined conditions 780-799	11.4	11.0	7.9	15.2	9.2
Injury and poisoning 800-999	93.1	103.6	92.3	96.1	80.5
Fractures, all sites ² 800-829	36.4	39.2	38.6	36.5	31.5
Fracture of neck of femur ² 820	11.7	12.8	11.8	12.7	9.3
Poisonings 960-989	6.6	6.6	6.2	7.3	5.8
Supplementary classifications V01-V82	169.7	154.9	159.9	183.0	171.4
Females with deliveries V27	138.3	121.7	127.2	148.9	146.8

* Figure does not meet standard of reliability or precision.

¹First-listed diagnosis for females with deliveries is coded V27, shown under "Supplementary classifications."²Excludes fractures coded as 733.1, pathologic fracture.

Table 18. Average length of stay for discharges from short-stay hospitals, by geographic region and first-listed diagnosis: United States, 1999[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Diagnostic groupings and code numbers are based on the *International Classification of Diseases, 9th Revision, Clinical Modification (ICD-9-CM)*]

Category of first-listed diagnosis and ICD-9-CM code	United States	North-east	Midwest	South	West
Average length of stay in days					
All conditions	5.0	5.7	4.6	5.0	4.6
Infectious and parasitic diseases 001-139	6.4	8.0	5.2	6.3	5.9
Septicemia 038	8.4	10.0	7.1	8.7	7.4
Neoplasms 140-239	6.2	7.0	5.8	6.0	6.1
Malignant neoplasms 140-208,230-234	7.0	7.6	6.5	7.0	6.9
Malignant neoplasm of large intestine and rectum 153-154,197.5	9.2	9.6	9.6	9.1	8.5
Malignant neoplasm of trachea, bronchus, and lung 162,176.4,197.0,197.3	7.3	8.0	6.2	7.5	7.8
Malignant neoplasm of breast 174-175,198.81	2.7	2.7	2.4	2.8	3.2
Benign neoplasms 210-229	3.6	4.3	3.3	3.2	3.7
Endocrine, nutritional and metabolic diseases, and immunity disorders 240-279	4.7	5.6	4.3	4.4	5.0
Diabetes mellitus 250	5.4	6.4	4.6	4.9	6.0
Volume depletion 276.5	4.0	4.8	4.0	3.7	3.3
Diseases of the blood and blood-forming organs 280-289	4.8	4.9	4.3	5.0	4.9
Mental disorders 290-319	7.5	9.2	6.5	6.8	7.1
Psychoses 290-299	8.4	10.1	7.1	7.8	8.6
Alcohol dependence syndrome 303	5.3	6.1	6.1	4.3	6.0
Diseases of the nervous system and sense organs 320-389	5.1	5.2	4.9	5.4	4.7
Diseases of the circulatory system 390-459	4.9	5.4	4.6	4.9	4.4
Heart disease 391-392.0,393-398,402,404,410-416,420-429	4.7	5.2	4.4	4.7	4.3
Acute myocardial infarction 410	5.6	6.0	5.0	6.0	4.9
Coronary atherosclerosis 414.0	3.9	4.1	3.6	3.9	3.7
Other ischemic heart disease 411-413,414.1-414.9	3.0	3.4	2.5	3.0	2.7
Cardiac dysrhythmias 427	3.6	4.1	3.4	3.6	3.1
Congestive heart failure 428.0	5.5	6.4	5.3	5.4	5.0
Cerebrovascular disease 430-438	5.4	6.0	4.9	5.6	5.3
Diseases of the respiratory system 460-519	5.4	6.3	4.6	5.6	5.2
Acute bronchitis and bronchiolitis 466	3.3	3.8	2.9	3.6	2.8
Pneumonia 480-486	6.0	7.1	5.1	6.3	5.7
Chronic bronchitis 491	5.2	5.8	4.7	5.4	5.0
Asthma 493	3.2	3.6	2.7	3.4	2.8
Diseases of the digestive system 520-579	4.8	5.4	4.6	4.9	4.5
Appendicitis 540-543	3.5	3.7	3.2	3.9	3.2
Noninfectious enteritis and colitis 555-558	4.1	4.3	4.1	4.0	3.9
Diverticula of intestine 562	5.6	6.3	5.4	5.6	5.1
Cholelithiasis 574	3.9	4.2	4.1	3.5	3.9
Diseases of the genitourinary system 580-629	3.8	4.2	3.4	3.8	4.1
Calculus of kidney and ureter 592	2.3	2.5	1.9	2.5	2.3
Complications of pregnancy, childbirth, and the puerperium ¹ 630-677	2.6	2.6	3.0	2.6	2.3
Diseases of the skin and subcutaneous tissue 680-709	5.7	5.7	5.0	6.5	4.9
Cellulitis and abscess 681-682	5.2	5.3	4.6	5.8	4.7
Diseases of the musculoskeletal system and connective tissue 710-739	4.3	4.5	4.1	4.4	4.2
Osteoarthritis and allied disorders 715	4.5	5.0	4.3	4.6	4.3
Intervertebral disc disorders 722	3.0	2.9	3.2	2.9	2.9
Congenital anomalies 740-759	5.7	5.6	6.0	5.1	6.8
Certain conditions originating in the perinatal period 760-779	9.4	8.6	9.2	10.5	9.0
Symptoms, signs, and ill-defined conditions 780-799	3.2	2.5	2.6	3.4	3.8
Injury and poisoning 800-999	5.4	5.9	4.8	5.6	5.0
Fractures, all sites ² 800-829	5.5	6.3	4.9	5.5	5.2
Fracture of neck of femur ² 820	6.5	7.5	6.2	6.5	5.8
Poisonings 960-989	2.5	2.9	2.6	2.4	2.3
Supplementary classifications V01-V82	3.7	4.0	3.9	3.8	3.2
Females with deliveries V27	2.5	2.7	2.5	2.5	2.3

* Figure does not meet standard of reliability or precision.

¹First-listed diagnosis for females with deliveries is coded V27, shown under "Supplementary classifications."²Excludes fractures coded as 733.1, pathologic fracture.

Table 19. Number of all-listed diagnoses for discharges from short-stay hospitals, by age and diagnosis: United States, 1999[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Diagnostic groupings and code numbers are based on the *International Classification of Diseases, 9th Revision, Clinical Modification (ICD-9-CM)*]

All-listed diagnosis and ICD-9-CM code	All ages	Under 15 years	15-44 years	45-64 years	65 years and over
	Number in thousands				
All conditions	135,765	6,393	33,320	30,008	66,044
Infectious and parasitic diseases 001-139	3,554	465	884	701	1,504
Septicemia 038	692	50	78	134	429
Neoplasms 140-239	4,432	126	623	1,378	2,305
Malignant neoplasms 140-208,230-234	3,455	102	302	1,039	2,012
Malignant neoplasm of large intestine and rectum 153-154,197.5	274	*	15	69	187
Malignant neoplasm of trachea, bronchus, and lung 162,176.4,197.0,197.3	481	*	*	153	299
Malignant neoplasm of breast 174-175,198.81	172	*	20	66	86
Benign neoplasms 210-229	834	17	304	313	201
Endocrine, nutritional and metabolic diseases, and immunity disorders 240-279	14,053	536	1,702	3,916	7,899
Diabetes mellitus 250	4,571	30	461	1,494	2,586
Volume depletion 276.5	2,171	330	304	317	1,220
Diseases of the blood and blood-forming organs 280-289	4,170	176	849	861	2,284
Mental disorders 290-319	8,927	284	3,846	2,312	2,486
Psychoses 290-299	3,028	65	1,054	596	1,313
Alcohol dependence syndrome 303	724	*	362	276	85
Diseases of the nervous system and sense organs 320-389	3,519	403	574	717	1,825
Diseases of the circulatory system 390-459	29,558	119	1,527	7,134	20,778
Heart disease 391-392.0,393-398,402,404,410-416,420-429	17,999	74	754	4,045	13,126
Acute myocardial infarction 410	981	*	53	307	620
Coronary atherosclerosis 414.0	4,200	*	116	1,147	2,929
Other ischemic heart disease 411-413,414.1-414.9	2,454	*	101	773	1,575
Cardiac dysrhythmias 427	3,773	21	152	605	2,994
Congestive heart failure 428.0	3,105	*	75	504	2,515
Cerebrovascular disease 430-438	1,977	8	72	398	1,498
Diseases of the respiratory system 460-519	10,508	1,235	1,274	2,156	5,844
Acute bronchitis and bronchiolitis 466	460	249	48	46	117
Pneumonia 480-486	2,135	274	214	368	1,280
Chronic bronchitis 491	1,064	*	36	269	755
Asthma 493	1,347	315	377	310	345
Diseases of the digestive system 520-579	8,898	478	1,699	2,413	4,309
Appendicitis 540-543	296	64	159	48	25
Noninfectious enteritis and colitis 555-558	577	139	130	105	203
Diverticula of intestine 562	526	*	30	123	372
Cholelithiasis 574	624	*3	165	181	275
Diseases of the genitourinary system 580-629	7,198	178	1,917	1,703	3,399
Calculus of kidney and ureter 592	273	*	111	98	60
Complications of pregnancy, childbirth, and the puerperium ¹ 630-677	8,350	17	8,312	*	...
Diseases of the skin and subcutaneous tissue 680-709	1,833	147	354	440	892
Cellulitis and abscess 681-682	791	51	166	225	349
Diseases of the musculoskeletal system and connective tissue 710-739	5,054	72	755	1,279	2,949
Osteoarthritis and allied disorders 715	1,281	*	36	242	1,002
Intervertebral disc disorders 722	489	*	176	192	121
Congenital anomalies 740-759	623	368	134	64	58
Certain conditions originating in the perinatal period 760-779	487	485	*	*	*
Symptoms, signs, and ill-defined conditions 780-799	7,144	593	1,482	1,832	3,238
Injury and poisoning 800-999	6,038	445	1,936	1,294	2,363
Fractures, all sites ² 800-829	1,599	87	457	270	785
Fracture of neck of femur ² 820	380	*	12	28	338
Poisonings 960-989	337	29	196	69	43
Supplementary classifications V01-V82	11,418	267	5,452	1,789	3,911
Females with deliveries V27	3,810	7	3,797	*6	...

* Figure does not meet standard of reliability or precision.

... Category not applicable.

¹First-listed diagnosis for females with deliveries is coded V27, shown under "Supplementary classifications."²Excludes fractures coded as 733.1, pathologic fracture.

Table 20. Number of all-listed diagnoses for discharges from short-stay hospitals, by sex and diagnosis: United States, 1999[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Diagnostic groupings and code numbers are based on the *International Classification of Diseases, 9th Revision, Clinical Modification (ICD-9-CM)*]

All-listed diagnosis and ICD-9-CM code	Both sexes	Number in thousands	
		Male	Female
All conditions	135,765	54,901	80,864
Infectious and parasitic diseases 001-139	3,554	1,552	2,002
Septicemia 038	692	329	363
Neoplasms 140-239	4,432	1,888	2,544
Malignant neoplasms 140-208,230-234	3,455	1,691	1,764
Malignant neoplasm of large intestine and rectum 153-154,197.5	274	133	141
Malignant neoplasm of trachea, bronchus, and lung 162,176.4,197.0,197.3	481	265	216
Malignant neoplasm of breast 174-175,198.81	172	*	171
Benign neoplasms 210-229	834	139	695
Endocrine, nutritional and metabolic diseases, and immunity disorders 240-279	14,053	5,801	8,252
Diabetes mellitus 250	4,571	2,063	2,508
Volume depletion 276.5	2,171	877	1,294
Diseases of the blood and blood-forming organs 280-289	4,170	1,683	2,487
Mental disorders 290-319	8,927	4,367	4,560
Psychoses 290-299	3,028	1,347	1,681
Alcohol dependence syndrome 303	724	521	203
Diseases of the nervous system and sense organs 320-389	3,519	1,533	1,986
Diseases of the circulatory system 390-459	29,558	13,999	15,560
Heart disease 391-392.0,393-398,402,404,410-416,420-429	17,999	8,900	9,099
Acute myocardial infarction 410	981	556	425
Coronary atherosclerosis 414.0	4,200	2,311	1,889
Other ischemic heart disease 411-413,414.1-414.9	2,454	1,356	1,098
Cardiac dysrhythmias 427	3,773	1,795	1,978
Congestive heart failure 428.0	3,105	1,347	1,759
Cerebrovascular disease 430-438	1,977	892	1,085
Diseases of the respiratory system 460-519	10,508	5,030	5,478
Acute bronchitis and bronchiolitis 466	460	230	231
Pneumonia 480-486	2,135	1,052	1,083
Chronic bronchitis 491	1,064	492	572
Asthma 493	1,347	487	860
Diseases of the digestive system 520-579	8,898	3,754	5,144
Appendicitis 540-543	296	163	133
Noninfectious enteritis and colitis 555-558	577	224	353
Diverticula of intestine 562	526	197	328
Cholelithiasis 574	624	202	422
Diseases of the genitourinary system 580-629	7,198	2,429	4,769
Calculus of kidney and ureter 592	273	156	117
Complications of pregnancy, childbirth, and the puerperium ¹ 630-677	8,350	...	8,350
Diseases of the skin and subcutaneous tissue 680-709	1,833	885	948
Cellulitis and abscess 681-682	791	396	394
Diseases of the musculoskeletal system and connective tissue 710-739	5,054	1,810	3,244
Osteoarthritis and allied disorders 715	1,281	425	856
Intervertebral disc disorders 722	489	243	246
Congenital anomalies 740-759	623	300	324
Certain conditions originating in the perinatal period 760-779	487	304	183
Symptoms, signs, and ill-defined conditions 780-799	7,144	3,156	3,988
Injury and poisoning 800-999	6,038	3,056	2,982
Fractures, all sites ² 800-829	1,599	717	882
Fracture of neck of femur ² 820	380	103	276
Poisonings 960-989	337	142	194
Supplementary classifications V01-V82	11,418	3,356	8,062
Females with deliveries V27	3,810	...	3,810

* Figure does not meet standard of reliability or precision.

... Category not applicable.

¹First-listed diagnosis for females with deliveries is coded V27, shown under "Supplementary classifications."²Excludes fractures coded as 733.1, pathologic fracture.

Table 21. Number of all-listed diagnoses for discharges from short-stay hospitals, by geographic region and diagnosis: United States, 1999[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Diagnostic groupings and code numbers are based on the *International Classification of Diseases, 9th Revision, Clinical Modification (ICD-9-CM)*]

All-listed diagnosis and ICD-9-CM code	United States	North-east	Midwest	South	West
	Number in thousands				
All conditions	135,765	29,665	33,022	48,432	24,646
Infectious and parasitic diseases 001-139	3,554	788	778	1,340	647
Septicemia 038	692	165	141	268	118
Neoplasms 140-239	4,432	1,073	1,042	1,531	785
Malignant neoplasms 140-208,230-234	3,455	865	830	1,160	600
Malignant neoplasm of large intestine and rectum 153-154,197.5	274	65	69	94	46
Malignant neoplasm of trachea, bronchus, and lung 162,176.4,197.0,197.3	481	114	120	174	73
Malignant neoplasm of breast 174-175,198.81	172	47	43	59	23
Benign neoplasms 210-229	834	174	175	322	163
Endocrine, nutritional and metabolic diseases, and immunity disorders 240-279	14,053	3,112	3,559	4,961	2,421
Diabetes mellitus 250	4,571	1,063	1,051	1,722	735
Volume depletion 276.5	2,171	459	563	783	366
Diseases of the blood and blood-forming organs 280-289	4,170	908	981	1,483	798
Mental disorders 290-319	8,927	2,241	2,193	3,088	1,405
Psychoses 290-299	3,028	797	732	1,084	414
Alcohol dependence syndrome 303	724	195	180	227	121
Diseases of the nervous system and sense organs 320-389	3,519	765	871	1,222	661
Diseases of the circulatory system 390-459	29,558	6,933	7,432	10,439	4,755
Heart disease 391-392.0,393-398,402,404,410-416,420-429	17,999	4,347	4,556	6,239	2,857
Acute myocardial infarction 410	981	240	248	316	177
Coronary atherosclerosis 414.0	4,200	1,042	1,058	1,442	658
Other ischemic heart disease 411-413,414.1-414.9	2,454	623	576	854	401
Cardiac dysrhythmias 427	3,773	909	976	1,243	645
Congestive heart failure 428.0	3,105	694	859	1,145	408
Cerebrovascular disease 430-438	1,977	400	504	734	339
Diseases of the respiratory system 460-519	10,508	2,280	2,563	3,844	1,821
Acute bronchitis and bronchiolitis 466	460	85	104	182	90
Pneumonia 480-486	2,135	420	526	818	373
Chronic bronchitis 491	1,064	220	269	420	155
Asthma 493	1,347	369	321	410	248
Diseases of the digestive system 520-579	8,898	1,929	2,114	3,271	1,584
Appendicitis 540-543	296	54	73	87	82
Noninfectious enteritis and colitis 555-558	577	129	127	235	86
Diverticula of intestine 562	526	125	134	185	81
Cholelithiasis 574	624	145	135	219	126
Diseases of the genitourinary system 580-629	7,198	1,498	1,702	2,708	1,291
Calculus of kidney and ureter 592	273	57	75	102	39
Complications of pregnancy, childbirth, and the puerperium ¹ 630-677	8,350	1,422	1,822	3,082	2,024
Diseases of the skin and subcutaneous tissue 680-709	1,833	429	428	679	296
Cellulitis and abscess 681-682	791	192	195	270	133
Diseases of the musculoskeletal system and connective tissue 710-739	5,054	1,049	1,340	1,744	920
Osteoarthritis and allied disorders 715	1,281	266	394	424	197
Intervertebral disc disorders 722	489	83	129	189	88
Congenital anomalies 740-759	623	168	121	186	*
Certain conditions originating in the perinatal period 760-779	487	85	*	147	*
Symptoms, signs, and ill-defined conditions 780-799	7,144	1,522	1,848	2,530	1,244
Injury and poisoning 800-999	6,038	1,243	1,458	2,103	1,234
Fractures, all sites ² 800-829	1,599	311	407	569	311
Fracture of neck of femur ² 820	380	81	87	149	62
Poisonings 960-989	337	58	80	126	72
Supplementary classifications V01-V82	11,418	2,219	2,672	4,074	2,453
Females with deliveries V27	3,810	636	809	1,454	912

* Figure does not meet standard of reliability or precision.

¹First-listed diagnosis for females with deliveries is coded V27, shown under "Supplementary classifications."²Excludes fractures coded as 733.1, pathologic fracture.

Table 22. Number and rate of discharges from short-stay hospitals with human immunodeficiency virus diagnoses, by selected characteristics: United States, 1990–99

[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Data are for discharges with at least one of the following *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD–9–CM) codes: 042–044, 279.19, 795.8 (prior to 1995); 042, V08 (beginning in 1995)]

Characteristic	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999
Number in thousands										
All HIV discharges ¹	146	165	194	225	234	249	227	178	189	180
Sex										
Male	114	127	153	174	177	183	165	123	132	121
Female	32	37	41	51	57	66	62	55	57	59
Age										
Under 25 years	14	16	16	14	18	17	12	11	9	8
25–29 years	20	25	26	37	20	26	18	17	15	13
30–34 years	38	38	42	46	50	48	44	32	30	41
35–39 years	36	36	53	51	56	59	52	40	42	34
40–44 years	17	25	30	40	44	49	49	38	43	34
45 years and over	21	25	27	37	45	51	53	40	51	49
Region										
Northeast	66	61	75	94	91	95	86	69	69	70
Midwest	20	26	23	28	33	27	25	*	23	20
South	37	44	56	64	78	87	77	72	71	67
West	24	33	40	39	32	41	39	19	26	23
Rate per 100,000 population ²										
All HIV discharges ¹	58.1	64.6	75.2	86.4	88.9	93.9	84.7	65.8	69.1	65.3
Sex										
Male	93.4	102.5	121.7	136.8	137.7	141.6	125.7	93.1	98.8	89.5
Female	24.7	28.7	31.0	38.4	42.3	48.4	45.4	39.7	40.7	42.1
Age										
Under 25 years	14.8	16.5	17.4	15.0	19.2	17.6	12.6	11.3	9.1	8.1
25–29 years	95.1	118.3	129.1	185.5	103.9	132.2	91.3	86.7	77.0	70.5
30–34 years	172.8	170.7	186.3	204.7	223.3	216.2	203.6	153.6	145.2	208.3
35–39 years	179.0	176.3	249.7	235.0	254.0	262.7	227.8	176.5	185.4	149.1
40–44 years	94.8	130.9	156.3	207.2	222.9	241.4	233.1	175.2	194.4	152.0
45 years and over	27.4	32.1	33.2	44.8	53.9	60.2	60.6	45.5	55.6	52.9
Region										
Northeast	128.8	119.7	146.5	182.7	176.4	183.3	164.9	132.3	133.4	135.0
Midwest	33.6	43.4	37.1	44.8	52.8	43.5	40.5	*	36.2	30.7
South	42.1	49.5	62.3	70.5	85.2	93.1	82.0	75.1	73.1	68.4
West	44.1	60.8	71.6	68.9	54.8	69.5	65.2	30.9	42.0	37.0

* Figure does not meet standard of reliability or precision.

¹HIV is human immunodeficiency virus.

²Rates are based on the U.S. Bureau of the Census estimates of the civilian population that have been adjusted for net underenumeration in the 1990 census. See appendix.

Table 23. Number and rate of deaths for discharges from short-stay hospitals, by age and selected first-listed diagnoses: United States, 1999[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Diagnostic groupings and code numbers are based on the *International Classification of Diseases, 9th Revision, Clinical Modification (ICD-9-CM)*]

First-listed diagnosis and ICD-9-CM code	All ages	Under 65 years	65 years and over	Rate per 100 discharges		
				All ages	Under 65 years	65 years and over
	Number of deaths in thousands			Rate per 100 discharges		
All conditions ¹	851	191	660	2.6	1.0	5.2
Heart disease 391-392.0,393-398,402,404,410-416,420-429	182	28	154	4.1	1.8	5.3
Acute myocardial infarction 410	78	14	64	9.5	4.4	12.7
Coronary atherosclerosis 414.0	14	*	12	1.2	*	1.9
Cardiac dysrhythmias 427	18	*5	13	2.6	*2.3	2.7
Congestive heart failure 428.0	48	*3	46	5.0	*1.3	6.0
Malignant neoplasms 140-208,230-234	99	33	66	7.8	6.1	9.0
Malignant neoplasm of trachea, bronchus, and lung 162,176.4,197.0,197.3	24	8	16	14.9	13.5	15.7
Pneumonia 480-486	80	10	70	5.8	1.8	8.6
Cerebrovascular disease 430-438	67	14	54	7.0	5.4	7.6
Septicemia 038	61	13	48	18.0	13.2	19.9
Fractures, all sites ² 800-829	17	*	13	1.7	*	2.4
Chronic bronchitis 491	14	*	12	2.5	*	3.3
Volume depletion 276.5	13	*	11	2.8	*	4.4

*Figure does not meet standard of reliability or precision.

¹Includes data for deaths not shown in table.²Excludes fractures coded as 733.1, pathologic fracture.**Table 24. Number of discharges from short-stay hospitals with and without procedures and percent with procedures, by selected characteristics: United States, 1999**

[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants]

Characteristic	All discharges	Discharges without procedures	Discharges with procedures			
			Total	With surgical procedures	Total	With surgical procedures
	Number in thousands		Percent			
All patients	32,132	12,322	19,810	14,444	61.7	45.0
Age						
Under 15 years	2,458	1,338	1,120	620	45.6	25.2
15-44 years	10,092	2,856	7,236	5,697	71.7	56.4
45-64 years	6,899	2,502	4,397	3,289	63.7	47.7
65 years and over	12,683	5,625	7,057	4,839	55.6	38.2
65-74 years	4,883	1,902	2,981	2,174	61.0	44.5
75-84 years	5,227	2,351	2,876	1,943	55.0	37.2
85 years and over	2,572	1,372	1,200	722	46.6	28.1
Sex						
Male	12,748	5,285	7,463	5,175	58.5	40.6
Female	19,384	7,036	12,347	9,269	63.7	47.8
Region						
Northeast	6,952	2,420	4,532	3,013	65.2	43.3
Midwest	7,368	3,179	4,189	3,167	56.9	43.0
South	12,007	4,973	7,033	5,351	58.6	44.6
West	5,805	1,749	4,056	2,914	69.9	50.2

NOTE: See appendix for definition of surgical procedures.

Table 25. Number and rate of all-listed surgical and nonsurgical procedures for discharges from short-stay hospitals, by selected procedure categories: United States, 1999[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Procedure groupings and code numbers are based on the *International Classification of Diseases, 9th Revision, Clinical Modification (ICD-9-CM)*]

Procedure category and ICD-9-CM code	Number in thousands	Rate per 10,000 population
Surgical procedures ¹	23,833	865.0
Cardiac catheterization37.21-37.23	1,271	46.1
Repair of current obstetric laceration ²75.5-75.6	1,116	40.5
Removal of coronary artery obstruction and insertion of stent(s) ³36.0	1,069	38.8
Episiotomy with or without forceps or vacuum extraction ²72.1,72.21,72.31,72.71,73.6	1,048	38.0
Cesarean section ²74.0-74.2,74.4,74.99	841	30.5
Artificial rupture of membranes ²73.0	809	29.3
Reduction of fracture76.7,79.0-79.3	628	22.8
Open reduction of fracture with internal fixation79.3	431	15.6
Hysterectomy68.3-68.7,68.9	616	22.4
Coronary artery bypass graft ³36.1	571	20.7
Endoscopy of small intestine with biopsy45.14,45.16	531	19.3
Oophorectomy and salpingo-oophorectomy65.3-65.6	466	16.9
Cholecystectomy51.2	428	15.5
Insertion, replacement, removal, and revision of pacemaker leads or device37.7-37.8	336	12.2
Debridement of wound, infection, or burn86.22,86.28	332	12.0
Bilateral destruction or occlusion of fallopian tubes66.2-66.3	329	12.0
Excision or destruction of intervertebral disc80.5	313	11.4
Lysis of peritoneal adhesions54.5	299	10.9
Appendectomy, excluding incidental47.0	290	10.5
Total knee replacement81.54	267	9.7
Partial excision of large intestine45.7	251	9.1
Partial excision of bone76.2-76.3,77.6-77.8	232	8.4
Prostatectomy60.2-60.6	192	7.0
Nonsurgical procedures ⁴	17,482	634.5
Arteriography and angiocardiology using contrast material88.4-88.5	2,034	73.8
Respiratory therapy93.9,96.7	1,117	40.5
Diagnostic ultrasound88.7	1,022	37.1
Computerized axial tomography87.03,87.41,87.71,88.01,88.38	871	31.6
Hemodialysis39.95	458	16.6
Insertion of endotracheal tube96.04	445	16.2
Endoscopy of small intestine without biopsy45.11-45.13	391	14.2
Endoscopy of large intestine without biopsy45.21-45.24	355	12.9
Spinal tap03.31	310	11.3
Injection or infusion of cancer chemotherapeutic substance99.25	211	7.6

¹Includes data for surgical procedures not shown in table. All procedures except those listed under "nonsurgical procedures" in the appendix are surgical procedures.²See table 33 for rate per 100 deliveries.³See table 27 for comparison of number of procedures with number of discharges with procedures.⁴Includes data for nonsurgical procedures not shown in table. The ICD-9-CM codes for nonsurgical procedures are listed in the appendix.

Table 26. Number and rate of all-listed surgical and nonsurgical procedures for patients 65 years of age and over discharged from short-stay hospitals, by age and selected procedure categories: United States, 1999[Discharges of inpatients from non-Federal hospitals. Procedure groupings and code numbers are based on the *International Classification of Diseases, 9th Revision, Clinical Modification (ICD-9-CM)*]

Procedure category and ICD-9-CM code	65 years and over	65-74 years	75-84 years	85 years and over
Surgical procedures ¹	8,069	3,763	3,223	1,084
Cardiac catheterization 37.21-37.23	618	345	241	32
Removal of coronary artery obstruction and insertion of stent(s) ² 36.0	504	284	192	28
Coronary artery bypass graft ² 36.1	322	194	119	9
Endoscopy of small intestine with biopsy 45.14,45.16	313	117	133	63
Insertion, replacement, removal, and revision of pacemaker leads or device 37.7-37.8	285	96	121	68
Reduction of fracture 76.7,79.0-79.3	267	67	103	97
Open reduction of fracture with internal fixation 79.3	203	50	76	76
Total knee replacement 81.54	182	103	69	11
Cholecystectomy 51.2	158	76	60	21
Prostatectomy 60.2-60.6	140	74	52	14
Debridement of wound, infection, or burn 86.22,86.28	121	46	52	23
Nonsurgical procedures ³	7,398	3,008	3,072	1,318
Arteriography and angiocardiology using contrast material 88.4-88.5	1,006	544	399	62
Diagnostic ultrasound 88.7	551	221	223	108
Respiratory therapy 93.9,96.7	509	208	215	86
Computerized axial tomography 87.03,87.41,87.71,88.01,88.38	453	172	177	104
Endoscopy of large intestine without biopsy 45.21-45.24	245	79	105	61
Insertion of endotracheal tube 96.04	244	102	104	39
Endoscopy of small intestine without biopsy 45.11-45.13	239	81	104	54
Hemodialysis 39.95	226	112	100	14
		Rate per 10,000 population		
Surgical procedures ¹	2,356.6	2,085.3	2,674.9	2,612.4
Cardiac catheterization 37.21-37.23	180.4	191.4	199.8	76.1
Removal of coronary artery obstruction and insertion of stent(s) ² 36.0	147.3	157.4	159.6	67.4
Coronary artery bypass graft ² 36.1	94.1	107.5	98.7	22.6
Endoscopy of small intestine with biopsy 45.14,45.16	91.5	65.0	110.2	152.6
Insertion, replacement, removal, and revision of pacemaker leads or device 37.7-37.8	83.2	53.0	100.5	164.7
Reduction of fracture 76.7,79.0-79.3	77.9	37.0	85.4	234.3
Open reduction of fracture with internal fixation 79.3	59.2	28.0	63.2	183.4
Total knee replacement 81.54	53.3	57.1	57.0	26.1
Cholecystectomy 51.2	46.0	42.3	49.7	51.5
Prostatectomy 60.2-60.6	41.0	41.1	43.4	33.1
Debridement of wound, infection, or burn 86.22,86.28	35.5	25.4	43.5	55.6
Nonsurgical procedures ³	2,160.6	1,667.1	2,550.0	3,176.4
Arteriography and angiocardiology using contrast material 88.4-88.5	293.7	301.6	331.5	149.7
Diagnostic ultrasound 88.7	161.0	122.2	185.2	259.4
Respiratory therapy 93.9,96.7	148.6	115.0	178.7	207.2
Computerized axial tomography 87.03,87.41,87.71,88.01,88.38	132.2	95.5	146.9	249.6
Endoscopy of large intestine without biopsy 45.21-45.24	71.7	44.0	86.9	147.8
Insertion of endotracheal tube 96.04	71.4	56.6	85.9	93.2
Endoscopy of small intestine without biopsy 45.11-45.13	69.7	44.7	86.0	131.2
Hemodialysis 39.95	65.9	61.9	82.7	34.4

¹Includes data for surgical procedures not shown in table. All procedures except those listed under "nonsurgical procedures" in the appendix are surgical procedures.²See table 24 for a comparison of number of procedures with number of discharges with procedures.³Includes data for nonsurgical procedures not shown in table. The ICD-9-CM codes for nonsurgical procedures are listed in the appendix.

Table 27. Number and rate of discharges from short-stay hospitals with a coronary artery procedure and number and rate of procedures by sex, age, and region: United States, 1999[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Procedure groupings and code numbers are based on the *International Classification of Diseases, 9th Revision, Clinical Modification (ICD-9-CM)*]

Sex, age, and region	Removal of coronary artery obstruction and insertion of stents(s) ¹		Coronary artery bypass graft ²	
	Number in thousands	Rate per 10,000 population	Number in thousands	Rate per 10,000 population
Discharges with procedures				
Total	599	21.7	355	12.9
Sex				
Male	396	29.4	238	17.7
Female	203	14.4	117	8.3
Age				
Under 15 years	*	*	*	*
15-44 years	33	2.7	10	9
45-64 years	281	47.6	137	23.2
65 years and over	285	83.2	207	60.5
65-74 years	160	88.5	124	69.0
75-84 years	110	91.3	76	62.7
85 years and over	15	36.7	7	17.4
Region				
Northeast	122	23.4	84	16.2
Midwest	159	25.0	84	13.2
South	203	20.8	113	11.6
West	114	18.4	74	11.8
Procedures				
Total	1,069	38.8	571	20.7
Sex				
Male	708	52.6	389	28.9
Female	361	25.6	182	12.9
Age				
Under 15 years	*	*	*	*
15-44 years	58	4.8	16	1.3
45-64 years	505	85.7	232	39.4
65 years and over	504	147.3	322	94.1
65-74 years	284	157.4	194	107.5
75-84 years	192	159.6	119	98.7
85 years and over	28	67.4	9	22.6
Region				
Northeast	226	43.2	147	28.1
Midwest	292	45.9	133	20.9
South	361	36.9	178	18.3
West	191	30.7	113	18.2

* Figure does not meet standard of reliability or precision.

¹ICD-9-CM code 36.0.²ICD-9-CM code 36.1.

NOTE: The number of procedures exceeds the number of discharges because these procedures are often assigned more than one code for a single operation. See Methods section.

Table 28. Number of all-listed procedures for discharges from short-stay hospitals, by age and procedure category: United States, 1999[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Procedure groupings and code numbers are based on the *International Classification of Diseases, 9th Revision, Clinical Modification (ICD-9-CM)*]

Procedure category and ICD-9-CM code	All ages	Under 15 years	15-44 years	45-64 years	65 years and over	
						Number in thousands
All procedures	41,315	2,077	13,791	9,981	15,467	
Operations on the nervous system	01-05	1,048	212	337	228	270
Spinal tap	03.31	310	144	71	45	50
Operations on the endocrine system	06-07	87	*4	29	33	20
Operations on the eye	08-16	110	10	31	28	40
Operations on the ear	18-20	54	27	11	*8	*7
Operations on the nose, mouth, and pharynx	21-29	284	67	101	60	56
Operations on the respiratory system	30-34	1,027	61	178	281	507
Bronchoscopy with or without biopsy	33.21-33.24,33.27	268	16	34	80	138
Operations on the cardiovascular system	35-39	6,133	179	569	2,170	3,215
Removal of coronary artery obstruction and insertion of stent(s) ¹	36.0	1,069	*	58	505	504
Coronary artery bypass graft ¹	36.1	571	*	16	232	322
Cardiac catheterization	37.21-37.23	1,271	10	96	547	618
Insertion, replacement, removal, and revision of pacemaker leads or device	37.7-37.8	336	*	*7	41	285
Hemodialysis	39.95	458	*	70	161	226
Operations on the hemic and lymphatic system	40-41	348	25	53	113	157
Operations on the digestive system	42-54	5,189	217	1,111	1,385	2,476
Endoscopy of small intestine with or without biopsy	45.11-45.14,45.16	922	18	128	224	552
Endoscopy of large intestine with or without biopsy	45.21-45.25	542	*	59	119	357
Partial excision of large intestine	45.7	251	*	21	82	146
Appendectomy, excluding incidental	47.0	290	65	158	42	24
Cholecystectomy	51.2	428	*	130	138	158
Lysis of peritoneal adhesions	54.5	299	6	127	83	83
Operations on the urinary system	55-59	966	40	225	299	402
Cystoscopy with or without biopsy	57.31-57.33	189	*	34	50	100
Operations on the male genital organs	60-64	277	32	10	66	170
Prostatectomy	60.2-60.6	192	*	*	49	140
Operations on the female genital organs	65-71	2,084	*	1,206	622	247
Oophorectomy and salpingo-oophorectomy	65.3-65.6	466	*	193	216	55
Bilateral destruction or occlusion of fallopian tubes	66.2-66.3	329	*	328	*	*
Hysterectomy	68.3-68.7,68.9	616	*	317	241	58
Obstetrical procedures	72-75	6,174	11	6,155	*	...
Episiotomy with or without forceps or vacuum extraction	72.1,72.21,72.31,72.71,73.6	1,048	*	1,044	*	...
Artificial rupture of membranes	73.0	809	*	806	*	...
Cesarean section	74.0-74.2,74.4,74.99	841	*	838	*	...
Repair of current obstetric laceration	75.5-75.6	1,116	*	1,113	*	...
Operations on the musculoskeletal system	76-84	3,219	163	903	921	1,231
Partial excision of bone	76.2-76.3,77.6-77.8	232	8	77	85	62
Reduction of fracture	76.7,79.0-79.3	628	46	201	114	267
Open reduction of fracture with internal fixation	79.3	431	13	132	83	203
Excision or destruction of intervertebral disc	80.5	313	*	132	134	46
Total hip replacement	81.51	168	*	12	54	102
Total knee replacement	81.54	267	*	*5	80	182
Operations on the integumentary system	85-86	1,351	*	408	374	430
Debridement of wound, infection, or burn	86.22,86.28	332	18	98	95	121
Miscellaneous diagnostic and therapeutic procedures	87-99	12,964	878	2,463	3,384	6,239
Computerized axial tomography	87.03,87.41,87.71,88.01,88.38	871	36	175	208	453
Arteriography and angiocardiology using contrast material	88.4-88.5	2,034	15	187	825	1,006
Diagnostic ultrasound	88.7	1,022	*	180	250	551
Respiratory therapy	93.9,96.7	1,117	224	154	231	509
Insertion of endotracheal tube	96.04	445	40	69	92	244
Injection or infusion of cancer chemotherapeutic substance	99.25	211	42	39	62	68

* Figure does not meet standard of reliability or precision.

... Category not applicable.

¹See table 27 for comparison of number of procedures with number of discharges with procedures.

Table 29. Rate of all-listed procedures for discharges from short-stay hospitals, by age and procedure category: United States, 1999[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Procedure groupings and code numbers are based on the *International Classification of Diseases, 9th Revision, Clinical Modification (ICD-9-CM)*]

Procedure category and ICD-9-CM code	All ages	Under 15 years	15-44 years	45-64 years	65 years and over	
						Rate per 10,000 population
All procedures	1,499.4	344.7	1,130.1	1,691.3	4,517.2	
Operations on the nervous system	01-05	38.0	35.2	27.6	38.7	78.9
Spinal tap	03.31	11.3	23.9	5.8	7.6	14.7
Operations on the endocrine system	06-07	3.1	*0.7	2.3	5.7	6.0
Operations on the eye	08-16	4.0	1.7	2.6	4.8	11.7
Operations on the ear	18-20	2.0	4.5	0.9	*1.4	*2.1
Operations on the nose, mouth, and pharynx	21-29	10.3	11.1	8.3	10.2	16.3
Operations on the respiratory system	30-34	37.3	10.2	14.6	47.6	148.0
Bronchoscopy with or without biopsy	33.21-33.24,33.27	9.7	2.7	2.8	13.6	40.4
Operations on the cardiovascular system	35-39	222.6	29.8	46.6	367.7	938.9
Removal of coronary artery obstruction and insertion of stent(s) ¹	36.0	38.8	*	4.8	85.7	147.3
Coronary artery bypass graft ¹	36.1	20.7	*	1.3	39.4	94.1
Cardiac catheterization	37.21-37.23	46.1	1.7	7.9	92.8	180.4
Insertion, replacement, removal, and revision of pacemaker leads or device	37.7-37.8	12.2	*	*0.5	7.0	83.2
Hemodialysis	39.95	16.6	*	5.7	27.2	65.9
Operations on the hemic and lymphatic system	40-41	12.6	4.2	4.3	19.1	46.0
Operations on the digestive system	42-54	188.3	36.1	91.0	234.7	723.0
Endoscopy of small intestine with or without biopsy	45.11-45.14,45.16	33.5	3.0	10.5	37.9	161.2
Endoscopy of large intestine with or without biopsy	45.21-45.25	19.7	*	4.8	20.2	104.4
Partial excision of large intestine	45.7	9.1	*	1.8	13.8	42.6
Appendectomy, excluding incidental	47.0	10.5	10.9	13.0	7.1	7.1
Cholecystectomy	51.2	15.5	*	10.6	23.4	46.0
Lysis of peritoneal adhesions	54.5	10.9	0.9	10.4	14.1	24.3
Operations on the urinary system	55-59	35.0	6.6	18.4	50.7	117.3
Cystoscopy with or without biopsy	57.31-57.33	6.9	*	2.8	8.5	29.2
Operations on the male genital organs	60-64	10.1	5.3	0.8	11.2	49.6
Prostatectomy	60.2-60.6	7.0	*	*	8.4	41.0
Operations on the female genital organs	65-71	75.6	*	98.8	105.4	72.1
Oophorectomy and salpingo-oophorectomy	65.3-65.6	16.9	*	15.8	36.6	16.1
Bilateral destruction or occlusion of fallopian tubes	66.2-66.3	12.0	*	26.9	*	*
Hysterectomy	68.3-68.7,68.9	22.4	*	26.0	40.9	17.1
Obstetrical procedures	72-75	224.1	1.8	504.4	*	...
Episiotomy with or without forceps or vacuum extraction	72.1,72.21,72.31,72.71,73.6	38.0	*	85.6	*	...
Artificial rupture of membranes	73.0	29.3	*	66.1	*	...
Cesarean section	74.0-74.2,74.4,74.99	30.5	*	68.7	*	...
Repair of current obstetric laceration	75.5-75.6	40.5	*	91.2	*	...
Operations on the musculoskeletal system	76-84	116.8	27.1	74.0	156.0	359.5
Partial excision of bone	76.2-76.3,77.6-77.8	8.4	1.3	6.3	14.5	18.0
Reduction of fracture	76.7,79.0-79.3	22.8	7.6	16.4	19.4	77.9
Open reduction of fracture with internal fixation	79.3	15.6	2.2	10.8	14.0	59.2
Excision or destruction of intervertebral disc	80.5	11.4	*	10.8	22.7	13.5
Total hip replacement	81.51	6.1	*	1.0	9.1	29.7
Total knee replacement	81.54	9.7	*	*0.4	13.5	53.3
Operations on the integumentary system	85-86	49.0	*	33.4	63.4	125.4
Debridement of wound, infection, or burn	86.22,86.28	12.0	2.9	8.0	16.1	35.5
Miscellaneous diagnostic and therapeutic procedures	87-99	470.5	145.7	201.9	573.4	1,822.3
Computerized axial tomography	87.03,87.41,87.71,88.01,88.38	31.6	5.9	14.4	35.2	132.2
Arteriography and angiocardiology using contrast material	88.4-88.5	73.8	2.6	15.4	139.8	293.7
Diagnostic ultrasound	88.7	37.1	*	14.7	42.3	161.0
Respiratory therapy	93.9,96.7	40.5	37.1	12.6	39.1	148.6
Insertion of endotracheal tube	96.04	16.2	6.6	5.7	15.6	71.4
Injection or infusion of cancer chemotherapeutic substance	99.25	7.6	7.0	3.2	10.5	19.8

* Figure does not meet standard of reliability or precision.

... Category not applicable.

¹See table 27 for comparison of number of procedures with number of discharges with procedures.

Table 30. Number of all-listed procedures for discharges from short-stay hospitals, by sex and procedure category: United States, 1999[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Procedure groupings and code numbers are based on the *International Classification of Diseases, 9th Revision, Clinical Modification (ICD-9-CM)*]

Procedure category and ICD-9-CM code	Both sexes	Male	Female
All procedures	41,315	16,370	24,945
Operations on the nervous system 01-05	1,048	475	572
Spinal tap 03.31	310	158	153
Operations on the endocrine system 06-07	87	20	67
Operations on the eye 08-16	110	59	51
Operations on the ear 18-20	54	32	22
Operations on the nose, mouth, and pharynx 21-29	284	175	109
Operations on the respiratory system 30-34	1,027	573	455
Bronchoscopy with or without biopsy 33.21-33.24,33.27	268	156	113
Operations on the cardiovascular system 35-39	6,133	3,498	2,634
Removal of coronary artery obstruction and insertion of stent(s) ¹ 36.0	1,069	708	361
Coronary artery bypass graft ¹ 36.1	571	389	182
Cardiac catheterization 37.21-37.23	1,271	758	513
Insertion, replacement, removal, and revision of pacemaker leads or device 37.7-37.8	336	159	176
Hemodialysis 39.95	458	227	231
Operations on the hemic and lymphatic system 40-41	348	170	178
Operations on the digestive system 42-54	5,189	2,189	3,000
Endoscopy of small intestine with or without biopsy 45.11-45.14,45.16	922	402	520
Endoscopy of large intestine with or without biopsy 45.21-45.25	542	208	335
Partial excision of large intestine 45.7	251	115	136
Appendectomy, excluding incidental 47.0	290	158	132
Cholecystectomy 51.2	428	137	292
Lysis of peritoneal adhesions 54.5	299	56	243
Operations on the urinary system 55-59	966	461	505
Cystoscopy with or without biopsy 57.31-57.33	189	101	88
Operations on the male genital organs 60-64	277	277	...
Prostatectomy 60.2-60.6	192	192	...
Operations on the female genital organs 65-71	2,084	...	2,084
Oophorectomy and salpingo-oophorectomy 65.3-65.6	466	...	466
Bilateral destruction or occlusion of fallopian tubes 66.2-66.3	329	...	329
Hysterectomy 68.3-68.7,68.9	616	...	616
Obstetrical procedures 72-75	6,174	...	6,174
Episiotomy with or without forceps or vacuum extraction 72.1,72.21,72.31,72.71,73.6	1,048	...	1,048
Artificial rupture of membranes 73.0	809	...	809
Cesarean section 74.0-74.2,74.4,74.99	841	...	841
Repair of current obstetric laceration 75.5-75.6	1,116	...	1,116
Operations on the musculoskeletal system 76-84	3,219	1,569	1,650
Partial excision of bone 76.2-76.3,77.6-77.8	232	124	109
Reduction of fracture 76.7,79.0-79.3	628	285	343
Open reduction of fracture with internal fixation 79.3	431	179	252
Excision or destruction of intervertebral disc 80.5	313	167	146
Total hip replacement 81.51	168	73	95
Total knee replacement 81.54	267	105	162
Operations on the integumentary system 85-86	1,351	603	749
Debridement of wound, infection, or burn 86.22,86.28	332	188	144
Miscellaneous diagnostic and therapeutic procedures 87-99	12,964	6,269	6,695
Computerized axial tomography 87.03,87.41,87.71,88.01,88.38	871	408	463
Arteriography and angiocardiology using contrast material 88.4-88.5	2,034	1,145	889
Diagnostic ultrasound 88.7	1,022	451	571
Respiratory therapy 93.9,96.7	1,117	560	557
Insertion of endotracheal tube 96.04	445	227	219
Injection or infusion of cancer chemotherapeutic substance 99.25	211	116	95

... Category not applicable.

¹See table 27 for comparison of number of procedures with number of discharges with procedures.

Table 31. Rate of all-listed procedures for discharges from short-stay hospitals, by sex and procedure category: United States, 1999[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Procedure groupings and code numbers are based on the *International Classification of Diseases, 9th Revision, Clinical Modification (ICD-9-CM)*]

Procedure category and ICD-9-CM code	Both sexes	Rate per 10,000 population	
		Male	Female
All procedures	1,499.4	1,215.8	1,770.5
Operations on the nervous system 01-05	38.0	35.3	40.6
Spinal tap 03.31	11.3	11.7	10.8
Operations on the endocrine system 06-07	3.1	1.5	4.8
Operations on the eye 08-16	4.0	4.4	3.6
Operations on the ear 18-20	2.0	2.4	1.6
Operations on the nose, mouth, and pharynx 21-29	10.3	13.0	7.7
Operations on the respiratory system 30-34	37.3	42.5	32.3
Bronchoscopy with or without biopsy 33.21-33.24,33.27	9.7	11.6	8.0
Operations on the cardiovascular system 35-39	222.6	259.8	187.0
Removal of coronary artery obstruction and insertion of stent(s) ¹ 36.0	38.8	52.6	25.6
Coronary artery bypass graft ¹ 36.1	20.7	28.9	12.9
Cardiac catheterization 37.21-37.23	46.1	56.3	36.4
Insertion, replacement, removal, and revision of pacemaker leads or device 37.7-37.8	12.2	11.8	12.5
Hemodialysis 39.95	16.6	16.9	16.4
Operations on the hemic and lymphatic system 40-41	12.6	12.7	12.6
Operations on the digestive system 42-54	188.3	162.6	212.9
Endoscopy of small intestine with or without biopsy 45.11-45.14,45.16	33.5	29.9	36.9
Endoscopy of large intestine with or without biopsy 45.21-45.25	19.7	15.4	23.8
Partial excision of large intestine 45.7	9.1	8.6	9.7
Appendectomy, excluding incidental 47.0	10.5	11.8	9.4
Cholecystectomy 51.2	15.5	10.2	20.7
Lysis of peritoneal adhesions 54.5	10.9	4.1	17.3
Operations on the urinary system 55-59	35.0	34.2	35.8
Cystoscopy with or without biopsy 57.31-57.33	6.9	7.5	6.2
Operations on the male genital organs 60-64	10.1	20.6	...
Prostatectomy 60.2-60.6	7.0	14.3	...
Operations on the female genital organs 65-71	75.6	...	147.9
Oophorectomy and salpingo-oophorectomy 65.3-65.6	16.9	...	33.1
Bilateral destruction or occlusion of fallopian tubes 66.2-66.3	12.0	...	23.4
Hysterectomy 68.3-68.7,68.9	22.4	...	43.8
Obstetrical procedures 72-75	224.1	...	438.2
Episiotomy with or without forceps or vacuum extraction 72.1,72.21,72.31,72.71,73.6	38.0	...	74.4
Artificial rupture of membranes 73.0	29.3	...	57.4
Cesarean section 74.0-74.2,74.4,74.99	30.5	...	59.7
Repair of current obstetric laceration 75.5-75.6	40.5	...	79.2
Operations on the musculoskeletal system 76-84	116.8	116.5	117.1
Partial excision of bone 76.2-76.3,77.6-77.8	8.4	9.2	7.7
Reduction of fracture 76.7,79.0-79.3	22.8	21.2	24.3
Open reduction of fracture with internal fixation 79.3	15.6	13.3	17.9
Excision or destruction of intervertebral disc 80.5	11.4	12.4	10.4
Total hip replacement 81.51	6.1	5.4	6.8
Total knee replacement 81.54	9.7	7.8	11.5
Operations on the integumentary system 85-86	49.0	44.8	53.2
Debridement of wound, infection, or burn 86.22,86.28	12.0	14.0	10.2
Miscellaneous diagnostic and therapeutic procedures 87-99	470.5	465.6	475.2
Computerized axial tomography 87.03,87.41,87.71,88.01,88.38	31.6	30.3	32.9
Arteriography and angiocardiology using contrast material 88.4-88.5	73.8	85.0	63.1
Diagnostic ultrasound 88.7	37.1	33.5	40.5
Respiratory therapy 93.9,96.7	40.5	41.6	39.5
Insertion of endotracheal tube 96.04	16.2	16.8	15.5
Injection or infusion of cancer chemotherapeutic substance 99.25	7.6	8.6	6.7

... Category not applicable.

¹See table 27 for comparison of number of procedures with number of discharges with procedures.

Table 32. Number of all-listed procedures for discharges from short-stay hospitals, by geographic region and procedure category: United States, 1999[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Procedure groupings and code numbers are based on the *International Classification of Diseases, 9th Revision, Clinical Modification (ICD-9-CM)*]

Procedure category and ICD-9-CM code	United States	North-east	Midwest	South	West
	Number in thousands				
All procedures	41,315	9,797	9,025	14,072	8,421
Operations on the nervous system 01-05	1,048	222	284	323	219
Spinal tap 03.31	310	72	73	95	69
Operations on the endocrine system 06-07	87	22	16	29	20
Operations on the eye 08-16	110	16	17	59	18
Operations on the ear 18-20	54	*	7	20	*12
Operations on the nose, mouth, and pharynx 21-29	284	81	51	83	68
Operations on the respiratory system 30-34	1,027	262	221	360	184
Bronchoscopy with or without biopsy 33.21-33.24,33.27	268	77	59	89	43
Operations on the cardiovascular system 35-39	6,133	1,400	1,381	2,241	1,111
Removal of coronary artery obstruction and insertion of stent(s) ¹ 36.0	1,069	226	292	361	191
Coronary artery bypass graft ¹ 36.1	571	147	133	178	113
Cardiac catheterization 37.21-37.23	1,271	269	290	512	199
Insertion, replacement, removal, and revision of pacemaker leads or device 37.7-37.8	336	85	71	134	47
Hemodialysis 39.95	458	95	78	189	96
Operations on the hemic and lymphatic system 40-41	348	96	81	109	63
Operations on the digestive system 42-54	5,189	1,119	1,205	1,846	1,019
Endoscopy of small intestine with or without biopsy 45.11-45.14,45.16	922	194	231	331	166
Endoscopy of large intestine with or without biopsy 45.21-45.25	542	125	132	198	88
Partial excision of large intestine 45.7	251	50	71	86	43
Appendectomy, excluding incidental 47.0	290	51	70	88	81
Cholecystectomy 51.2	428	94	86	159	89
Lysis of peritoneal adhesions 54.5	299	61	60	119	59
Operations on the urinary system 55-59	966	242	207	351	166
Cystoscopy with or without biopsy 57.31-57.33	189	52	48	63	26
Operations on the male genital organs 60-64	277	60	68	94	56
Prostatectomy 60.2-60.6	192	39	49	65	39
Operations on the female genital organs 65-71	2,084	320	426	905	433
Oophorectomy and salpingo-oophorectomy 65.3-65.6	466	70	99	205	92
Bilateral destruction or occlusion of fallopian tubes 66.2-66.3	329	35	64	165	65
Hysterectomy 68.3-68.7,68.9	616	93	140	265	118
Obstetrical procedures 72-75	6,174	1,162	1,445	2,068	1,498
Episiotomy with or without forceps or vacuum extraction 72.1,72.21,72.31,72.71,73.6	1,048	181	234	430	203
Artificial rupture of membranes 73.0	809	126	232	220	230
Cesarean section 74.0-74.2,74.4,74.99	841	143	170	335	194
Repair of current obstetric laceration 75.5-75.6	1,116	208	259	360	289
Operations on the musculoskeletal system 76-84	3,219	700	750	1,137	631
Partial excision of bone 76.2-76.3,77.6-77.8	232	48	52	85	48
Reduction of fracture 76.7,79.0-79.3	628	146	149	219	113
Open reduction of fracture with internal fixation 79.3	431	97	103	151	80
Excision or destruction of intervertebral disc 80.5	313	54	69	135	56
Total hip replacement 81.51	168	41	47	44	36
Total knee replacement 81.54	267	56	78	79	54
Operations on the integumentary system 85-86	1,351	323	283	539	206
Debridement of wound, infection, or burn 86.22,86.28	332	84	78	121	49
Miscellaneous diagnostic and therapeutic procedures 87-99	12,964	3,758	2,584	3,907	2,715
Computerized axial tomography 87.03,87.41,87.71,88.01,88.38	871	360	*	189	200
Arteriography and angiocardiology using contrast material 88.4-88.5	2,034	429	485	825	294
Diagnostic ultrasound 88.7	1,022	375	*	237	256
Respiratory therapy 93.9,96.7	1,117	302	185	295	335
Insertion of endotracheal tube 96.04	445	109	95	151	91
Injection or infusion of cancer chemotherapeutic substance 99.25	211	67	41	63	39

* Figure does not meet standard of reliability or precision.

¹See table 27 for comparison of number of procedures with number of discharges with procedures.

Table 33. Rate of all-listed procedures for discharges from short-stay hospitals, by geographic region and procedure category: United States, 1999[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Procedure groupings and code numbers are based on the *International Classification of Diseases, 9th Revision, Clinical Modification (ICD-9-CM)*]

Procedure category and ICD-9-CM code	United States	North-east	Midwest	South	West
All procedures	1,499.4	1,876.4	1,419.7	1,441.5	1,355.3
Operations on the nervous system 01-05	38.0	42.4	44.6	33.1	35.3
Spinal tap 03.31	11.3	13.9	11.6	9.7	11.2
Operations on the endocrine system 06-07	3.1	4.1	2.4	3.0	3.2
Operations on the eye 08-16	4.0	3.1	2.6	6.0	3.0
Operations on the ear 18-20	2.0	*	1.1	2.0	*2.0
Operations on the nose, mouth, and pharynx 21-29	10.3	15.6	8.1	8.5	11.0
Operations on the respiratory system 30-34	37.3	50.2	34.7	36.9	29.7
Bronchoscopy with or without biopsy 33.21-33.24,33.27	9.7	14.8	9.3	9.1	6.9
Operations on the cardiovascular system 35-39	222.6	268.1	217.2	229.6	178.8
Removal of coronary artery obstruction and insertion of stent(s) ¹ 36.0	38.8	43.2	45.9	36.9	30.7
Coronary artery bypass graft ¹ 36.1	20.7	28.1	20.9	18.3	18.2
Cardiac catheterization 37.21-37.23	46.1	51.6	45.7	52.4	32.1
Insertion, replacement, removal, and revision of pacemaker leads or device 37.7-37.8	12.2	16.3	11.1	13.7	7.5
Hemodialysis 39.95	16.6	18.2	12.3	19.4	15.4
Operations on the hemic and lymphatic system 40-41	12.6	18.3	12.7	11.2	10.1
Operations on the digestive system 42-54	188.3	214.3	189.5	189.1	164.0
Endoscopy of small intestine with or without biopsy 45.11-45.14,45.16	33.5	37.2	36.4	33.9	26.7
Endoscopy of large intestine with or without biopsy 45.21-45.25	19.7	23.9	20.7	20.3	14.1
Partial excision of large intestine 45.7	9.1	9.7	11.2	8.8	7.0
Appendectomy, excluding incidental 47.0	10.5	9.8	10.9	9.0	13.1
Cholecystectomy 51.2	15.5	18.0	13.6	16.3	14.4
Lysis of peritoneal adhesions 54.5	10.9	11.6	9.5	12.1	9.6
Operations on the urinary system 55-59	35.0	46.4	32.5	36.0	26.7
Cystoscopy with or without biopsy 57.31-57.33	6.9	10.0	7.6	6.4	4.1
Operations on the male genital organs 60-64	10.1	11.5	10.7	9.6	9.0
Prostatectomy 60.2-60.6	7.0	7.5	7.6	6.6	6.3
Operations on the female genital organs 65-71	75.6	61.2	67.1	92.7	69.7
Oophorectomy and salpingo-oophorectomy 65.3-65.6	16.9	13.5	15.5	21.0	14.8
Bilateral destruction or occlusion of fallopian tubes 66.2-66.3	12.0	6.8	10.1	16.9	10.5
Hysterectomy 68.3-68.7,68.9	22.4	17.8	22.1	27.1	19.1
Obstetrical procedures 72-75	224.1	222.6	227.4	211.9	241.1
Episiotomy with or without forceps or vacuum extraction 72.1,72.21,72.31,72.71,73.6	38.0	34.7	36.8	44.0	32.7
Artificial rupture of membranes 73.0	29.3	24.2	36.5	22.5	37.1
Cesarean section 74.0-74.2,74.4,74.99	30.5	27.3	26.8	34.3	31.1
Repair of current obstetric laceration 75.5-75.6	40.5	39.8	40.8	36.9	46.5
Operations on the musculoskeletal system 76-84	116.8	134.1	118.0	116.5	101.6
Partial excision of bone 76.2-76.3,77.6-77.8	8.4	9.2	8.1	8.7	7.7
Reduction of fracture 76.7,79.0-79.3	22.8	27.9	23.5	22.5	18.3
Open reduction of fracture with internal fixation 79.3	15.6	18.6	16.2	15.5	12.8
Excision or destruction of intervertebral disc 80.5	11.4	10.3	10.9	13.8	8.9
Total hip replacement 81.51	6.1	7.9	7.5	4.5	5.7
Total knee replacement 81.54	9.7	10.7	12.2	8.1	8.8
Operations on the integumentary system 85-86	49.0	61.9	44.5	55.2	33.2
Debridement of wound, infection, or burn 86.22,86.28	12.0	16.0	12.3	12.4	7.9
Miscellaneous diagnostic and therapeutic procedures 87-99	470.5	719.7	406.5	400.3	437.0
Computerized axial tomography 87.03,87.41,87.71,88.01,88.38	31.6	68.9	*	19.3	32.2
Arteriography and angiocardiology using contrast material 88.4-88.5	73.8	82.2	76.4	84.5	47.3
Diagnostic ultrasound 88.7	37.1	71.8	*	24.3	41.1
Respiratory therapy 93.9,96.7	40.5	57.8	29.0	30.2	54.0
Insertion of endotracheal tube 96.04	16.2	20.9	14.9	15.4	14.6
Injection or infusion of cancer chemotherapeutic substance 99.25	7.6	12.9	6.4	6.5	6.3

* Figure does not meet standard of reliability or precision.

¹See table 27 for comparison of number of procedures with number of discharges with procedures.

Table 34. Number, percent distribution, and rate of discharges and days of care, with average length of stay for females with deliveries discharged from short-stay hospitals, by type of delivery, age, and geographic region: United States, 1999[Discharges of inpatients from non-Federal hospitals. Data are for discharges 10–54 years of age with *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD–9–CM) code V27, females with deliveries]

Age and region	Discharges			Days of care			Average length of stay in days
	Number in thousands	Percent distribution	Rate per 1,000 population	Number in thousands	Percent distribution	Rate per 1,000 population	
All deliveries							
10–54 years	3,810	100.0	42.7	9,421	100.0	105.7	2.5
10–19 years	473	12.4	24.1	1,145	12.2	58.3	2.4
20–24 years	920	24.2	101.3	2,147	22.8	236.5	2.3
25–29 years	1,058	27.8	112.3	2,639	28.0	280.3	2.5
30–34 years	858	22.5	85.4	2,136	22.7	212.5	2.5
35–39 years	418	11.0	36.7	1,112	11.8	97.5	2.7
40–54 years	83	2.2	2.8	242	2.6	8.2	2.9
All regions	3,810	100.0	42.7	9,421	100.0	105.7	2.5
Northeast	636	16.7	37.7	1,736	18.4	103.0	2.7
Midwest	809	21.2	39.4	1,995	21.2	97.1	2.5
South	1,454	38.2	45.8	3,581	38.0	112.8	2.5
West	912	23.9	45.6	2,109	22.4	105.4	2.3
Vaginal deliveries¹							
10–54 years	2,969	100.0	33.3	6,232	100.0	69.9	2.1
10–19 years	395	13.3	20.1	827	13.3	42.1	2.1
20–24 years	754	25.4	83.0	1,550	24.9	170.6	2.1
25–29 years	829	27.9	88.0	1,765	28.3	187.4	2.1
30–34 years	649	21.9	64.6	1,355	21.7	134.8	2.1
35–39 years	291	9.8	25.5	630	10.1	55.3	2.2
40–54 years	52	1.7	1.7	105	1.7	3.6	2.0
All regions	2,969	100.0	33.3	6,232	100.0	69.9	2.1
Northeast	493	16.6	29.2	1,136	18.2	67.4	2.3
Midwest	639	21.5	31.1	1,368	21.9	66.6	2.1
South	1,119	37.7	35.3	2,358	37.8	74.3	2.1
West	719	24.2	35.9	1,371	22.0	68.5	1.9
Cesarean deliveries²							
10–54 years	841	100.0	9.4	3,189	100.0	35.8	3.8
10–19 years	78	9.3	4.0	318	10.0	16.2	4.1
20–24 years	167	19.8	18.4	598	18.7	65.8	3.6
25–29 years	229	27.2	24.3	874	27.4	92.9	3.8
30–34 years	209	24.9	20.8	781	24.5	77.7	3.7
35–39 years	127	15.1	11.1	481	15.1	42.2	3.8
40–54 years	31	3.7	1.1	136	4.3	4.6	4.3
All regions	841	100.0	9.4	3,189	100.0	35.8	3.8
Northeast	143	17.0	8.5	600	18.8	35.6	4.2
Midwest	170	20.2	8.3	627	19.7	30.5	3.7
South	335	39.8	10.6	1,223	38.4	38.5	3.7
West	193	23.0	9.7	738	23.2	36.9	3.8

¹Discharges of females with deliveries except those with cesarean deliveries.²Discharges of females with deliveries with ICD–9–CM procedure codes 74.0–74.2, 74.4, or 74.99.

Table 35. Number and percent distribution of females with deliveries discharged from short-stay hospitals by length of stay and type of delivery: United States, selected years 1980–99[Discharges of inpatients from non-Federal hospitals. Data are for discharges with *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD–9–CM) code V27, females with deliveries]

Length of stay and type of delivery	1980	1985	1990	1995	1999
Length of stay					
Number in thousands					
All deliveries	3,762	3,854	4,025	3,766	3,810
Less than 1 day	23	34	37	36	14
1 day	216	373	623	1,368	603
2 days	747	1,037	1,649	1,418	1,883
3 days	1,205	1,140	900	606	859
4 days	691	586	468	193	279
5–7 days	726	598	282	100	126
8 days or more	154	86	67	46	45
Vaginal deliveries ¹	3,143	2,977	3,080	2,981	2,969
Less than 1 day	23	33	35	34	13
1 day	215	371	620	1,362	599
2 days	743	1,030	1,624	1,271	1,767
3 days	1,185	1,054	618	228	488
4 days	618	338	106	39	54
5–7 days	317	125	55	27	33
8 days or more	42	26	23	21	16
Cesarean deliveries ²	619	877	945	785	841
Less than 1 day	*	*	*	*	*
1 day	*	*	*	*6	*
2 days	*	*	25	147	117
3 days	20	86	283	379	372
4 days	73	249	361	154	225
5–7 days	409	473	227	73	93
8 days or more	112	60	44	25	29
Percent distribution					
All deliveries	100.0	100.0	100.0	100.0	100.0
Less than 1 day	0.6	0.9	0.9	1.0	0.4
1 day	5.7	9.7	15.5	36.3	15.8
2 days	19.9	26.9	41.0	37.7	49.4
3 days	32.0	29.6	22.4	16.1	22.6
4 days	18.4	15.2	11.6	5.1	7.3
5–7 days	19.3	15.5	7.0	2.7	3.3
8 days or more	4.1	2.2	1.7	1.2	1.2
Vaginal deliveries ¹	100.0	100.0	100.0	100.0	100.0
Less than 1 day	0.7	1.1	1.1	1.1	0.4
1 day	6.8	12.5	20.1	45.7	20.2
2 days	23.6	34.6	52.7	42.6	59.5
3 days	37.7	35.4	20.1	7.6	16.4
4 days	19.7	11.4	3.4	1.3	1.8
5–7 days	10.1	4.2	1.8	0.9	1.1
8 days or more	1.3	0.9	0.7	0.7	0.5
Cesarean deliveries ²	100.0	100.0	100.0	100.0	100.0
Less than 1 day	*	*	*	*	*
1 day	*	*	*	*0.8	*
2 days	*	*	2.6	18.7	13.9
3 days	3.2	9.8	29.9	48.3	44.2
4 days	11.8	28.4	38.2	19.6	26.7
5–7 days	66.1	53.9	24.0	9.3	11.0
8 days or more	18.1	6.8	4.7	3.2	3.5
Average length of stay in days					
All deliveries	3.8	3.3	2.8	2.1	2.5
Vaginal deliveries ¹	3.2	2.7	2.3	1.7	2.1
Cesarean deliveries ²	6.5	5.2	4.5	3.6	3.8

* Figure does not meet standard of reliability or precision.

¹Discharges of females with deliveries except those with cesarean deliveries.²Discharges of females with deliveries with ICD–9–CM procedure codes 74.0–74.2, 74.4, or 74.99.

Table 36. Number and rate of selected obstetrical procedures for females with deliveries discharged from short-stay hospitals: United States, selected years 1980–99[Discharges from non-Federal hospitals. Procedure groupings and code numbers are based on the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD–9–CM). Data are for selected procedures performed on discharges with code V27, females with deliveries]

Procedure category and ICD–9–CM code	1980	1985	1990	1995	1999
Number of procedures in thousands					
All obstetrical procedures 72–75	3,526	4,239	6,660	6,282	6,104
Episiotomy 72.1,72.21,72.31,72.71,73.6	2,012	1,820	1,714	1,410	1,048
Forceps delivery ¹ 72.0–72.4,72.51,72.53,72.6,73.3	555	376	269	176	126
Vacuum extraction ¹ 72.7	22	66	190	279	254
Artificial rupture of membranes 73.0	120	315	691	752	808
Medical induction of labor 73.4	42	66	225	366	469
Manually assisted delivery 73.5	70	193	750	866	802
Cesarean section 74.0–74.2,74.4,74.99	619	877	945	785	841
Repair of current obstetric laceration 75.5–75.6	350	543	789	961	1,113
Rate per 100 deliveries ²					
All obstetrical procedures 72–75	93.7	110.0	165.5	166.8	160.2
Artificial rupture of membranes 73.0	3.2	8.2	17.2	20.0	21.2
Medical induction of labor 73.4	1.1	1.7	5.6	9.7	12.3
Cesarean section 74.0–74.2,74.4,74.99	16.5	22.7	23.5	20.8	22.1
Rate per 100 vaginal deliveries ³					
Episiotomy 72.1,72.21,72.31,72.71,73.6	64.0	61.1	55.6	47.2	35.2
Forceps delivery ¹ 72.0–72.4,72.51,72.53,72.6,73.3	17.6	12.5	8.6	5.8	4.2
Vacuum extraction ¹ 72.7	0.7	2.2	6.1	9.2	8.4
Manually assisted delivery 73.5	2.2	6.5	24.3	29.0	27.0
Repair of current obstetric laceration 75.5–75.6	11.1	18.1	25.5	32.0	37.3

¹Forceps delivery and vacuum extraction with episiotomy are also included in the episiotomy category.²Procedures per 100 discharges of females with deliveries.³Procedures for females with vaginal deliveries per 100 discharges with vaginal deliveries. Vaginal deliveries are all deliveries except those with a cesarean section.**Table 37. Number and rate of females with deliveries discharged from short-stay hospitals by type of delivery: United States, selected years 1980–99**[Discharges of inpatients from non-Federal hospitals. Data are for discharges with *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD–9–CM) code V27, females with deliveries]

Type of delivery	1980	1985	1990	1995	1999
Number in thousands					
Females with deliveries	3,762	3,854	4,025	3,766	3,810
Previous cesarean delivery	192	324	426	413	438
No previous cesarean delivery	3,571	3,530	3,600	3,353	3,372
Vaginal deliveries	3,143	2,977	3,080	2,981	2,969
Vaginal birth after cesarean delivery (VBAC)	*6	21	87	147	119
Other vaginal deliveries	3,137	2,956	2,993	2,835	2,850
Cesarean deliveries	619	877	945	785	841
No previous cesarean delivery (primary)	434	574	606	519	522
Previous cesarean delivery (repeat)	185	303	339	266	319
Rate per 100 deliveries					
Cesarean deliveries per 100 deliveries	16.5	22.7	23.5	20.8	22.1
Primary cesarean deliveries per 100 deliveries without previous cesarean delivery	12.1	16.3	16.8	15.5	15.5
Repeat cesarean deliveries per 100 cesarean deliveries	29.9	34.6	35.9	33.9	37.9
Vaginal birth after cesarean deliveries (VBAC) per 100 deliveries with previous cesarean delivery	*3.4	6.6	20.4	35.5	27.2

* Figure does not meet standard of reliability or precision.

NOTE: Previous cesarean deliveries are discharges with ICD–9–CM diagnostic code 654.2. Cesarean deliveries are discharges with ICD–9–CM procedure codes 74.0–74.2, 74.4 or 74.99.

Table 38. Number and rate of females with deliveries discharged from short-stay hospitals by type of delivery, age, and geographic region: United States, 1999[Discharges of inpatients from non-Federal hospitals. Data are for discharges 10–54 years of age with *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD–9–CM) code V27, females with deliveries]

Age and region	Cesarean deliveries			
	Total ¹	Primary ²	Repeat ³	VBAC ^{4,5}
Number in thousands				
10–54 years	841	522	319	119
10–19 years	78	69	9	*6
20–24 years	167	120	47	18
25–29 years	229	134	95	36
30–34 years	209	118	92	33
35–39 years	127	63	64	21
40–54 years	31	20	12	*5
All regions	841	522	319	119
Northeast	143	85	57	20
Midwest	170	104	67	29
South	335	213	122	42
West	193	120	73	28
Rate per 100 deliveries				
10–54 years	22.1	15.5	37.9	27.2
10–19 years	16.5	15.0	11.6	*41.0
20–24 years	18.1	14.0	28.1	27.4
25–29 years	21.6	14.4	41.6	27.3
30–34 years	24.4	16.0	43.8	26.6
35–39 years	30.4	18.9	50.4	24.7
40–54 years	37.9	29.6	37.6	*29.2
All regions	22.1	15.5	37.9	27.2
Northeast	22.5	15.3	40.1	26.3
Midwest	21.0	14.5	39.1	30.1
South	23.0	16.5	36.5	25.4
West	21.2	14.9	37.8	27.9

* Figure does not meet standard of reliability or precision.

¹Discharges with ICD–9–CM procedure codes 74.0–74.2, 74.4, or 74.99. Rate per 100 total deliveries.²Cesarean deliveries without ICD–9–CM diagnostic code 654.2. Rate per 100 deliveries without previous cesarean delivery.³Cesarean deliveries with ICD–9–CM diagnostic code 654.2. Rate per 100 cesarean deliveries.⁴VBAC is vaginal birth after cesarean delivery.⁵Vaginal deliveries with ICD–9–CM diagnostic code 654.2. Rate per 100 deliveries with previous cesarean delivery.

Table 39. Number, percent distribution, and average length of stay for newborn infants discharged from short-stay hospitals by sex, geographic region, and health status: United States, 1999

[Discharges of inpatients from non-Federal hospitals]

Sex, region and health status	Discharges		Average length of stay in days
	Number in thousands	Percent distribution	
All newborn infants	3,726	100.0	3.2
Sex			
Male	1,893	50.8	3.2
Female	1,833	49.2	3.1
Region			
Northeast	697	18.7	3.5
Midwest	801	21.5	2.8
South	1,341	36.0	3.4
West	888	23.8	3.0
Health status			
Well ¹	2,294	61.6	2.0
Male	1,116	30.0	2.0
Female	1,177	31.6	2.0
Sick ²	1,433	38.4	5.1
Male	777	20.8	5.0
Female	656	17.6	5.2

¹Without any illness or risk-related diagnoses.

²With at least one illness or risk-related diagnosis, ICD-9-CM codes 001–999.

Table 40. Number and percent distribution of newborn infants discharged from short-stay hospitals by length of stay, and average length of stay by health status: United States, selected years 1980–99

Length of stay and health status	1980	1985	1990	1995	1999
Length of stay					
Number in thousands					
All newborn infants	3,824	3,794	3,869	3,631	3,726
Less than 1 day	69	84	79	87	45
1 day	270	428	693	1,523	771
2 days	842	1,057	1,624	1,145	1,900
3 days	1,175	1,034	636	486	539
4 days	618	528	447	134	196
5–7 days	633	497	232	106	109
8 days or more	218	167	157	149	167
Percent distribution					
All newborn infants	100.0	100.0	100.0	100.0	100.0
Less than 1 day	1.8	2.2	2.0	2.4	1.2
1 day	7.1	11.3	17.9	41.9	20.7
2 days	22.0	27.9	42.0	31.5	51.0
3 days	30.7	27.3	16.4	13.4	14.5
4 days	16.2	13.9	11.6	3.7	5.3
5–7 days	16.5	13.1	6.0	2.9	2.9
8 days or more	5.7	4.4	4.1	4.1	4.5
Average length of stay in days					
All newborn infants	4.3	3.8	3.3	2.8	3.2
Well ¹	3.2	2.8	2.3	1.7	2.0
Sick ²	7.1	5.7	4.8	4.4	5.1

¹Without any illness or risk-related diagnoses.

²With at least one illness or risk-related diagnosis, ICD-9-CM codes 001–999.

Table 41. Number of all-listed diagnoses for sick newborn infants discharged from short-stay hospitals, by sex and selected diagnostic categories: United States, 1999

[Discharges of inpatients from non-Federal hospitals. Diagnostic groupings and code numbers are based on the *International Classification of Diseases, 9th Revision, Clinical Modification (ICD-9-CM)*]

Diagnostic category and ICD-9-CM code	Both sexes	Male	Female
		Number in thousands	
Sick newborn diagnoses 001-999	2,747	1,542	1,205
Congenital anomalies 740-759	258	147	110
Disorders relating to short gestation and unspecified low birthweight (prematurity) 765	281	148	133
Disorders relating to long gestation and high birthweight 766	201	132	69
Birth trauma 767	123	66	57
Injuries to scalp 767.1	78	43	35
Respiratory distress syndrome and other respiratory conditions of fetus and newborn 769-770	390	222	168
Transitory tachypnea of newborn 770.6	114	68	46
Infections specific to the perinatal period 771	85	48	36
Hemolytic disease of fetus or newborn, due to isoimmunization, and other perinatal jaundice 773-774	525	284	241
Neonatal jaundice associated with preterm delivery 774.2	118	63	55
Unspecified fetal and neonatal jaundice 774.6	342	188	155
Endocrine and metabolic disturbances specific to the fetus and newborn 775	140	80	60

Table 42. Number of first-listed diagnoses for discharges from short-stay hospitals, by ICD-9-CM code, sex, age, and geographic region: United States, 1999

[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table: see Methods section]

ICD-9-CM code	Sex			Age				Region			
	Total	Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	North-east	Midwest	South	West
Number of first-listed diagnoses in thousands											
All codes	32,132	12,748	19,384	2,458	10,092	6,899	12,683	6,952	7,368	12,007	5,805
001-139	826	385	441	173	186	133	334	198	171	318	139
003	*6	*	*	*	*	*	*	*	*	*	*
008	121	54	68	53	22	15	32	24	27	48	22
008.4	28	*7	21	*	*	*7	16	*7	*8	*9	*
008.45	25	*6	19	*	*	*6	14	*6	*7	*8	*
008.6	29	17	12	28	*	*	*	*	9	10	*7
008.61	28	17	11	27	*	*	-	*	9	10	*6
008.8	60	27	33	21	17	*7	15	13	10	27	10
009	15	*7	*7	*	*	*	*5	*	*	*6	*
009.0	11	*5	*5	*	*	*	*	*	*	*	*
011	*7	*5	*	*	*	*	*	*	*	*	*
011.9	*5	*	*	*	*	*	*	*	*	*	*
034	13	*6	*7	*6	*	*	*	*	*5	*	*
034.0	11	*5	*5	*5	*	*	*	*	*	*	*
038	341	146	195	19	26	55	241	82	72	131	56
038.0	26	11	16	*	*	*	18	*5	*7	9	*
038.1	50	28	22	*	*5	11	33	12	12	19	*7
038.10	10	*6	*	*	*	*	*8	*	*	*	*
038.11	29	18	12	*	*	*7	17	*6	*5	12	*6
038.19	11	*	*6	*	*	*	*8	*	*	*	*
038.2	13	*6	*7	*	*	*	*7	*	*	*	*
038.4	87	35	52	*	*5	19	60	24	21	26	17
038.40	*8	*	*5	*	*	*	*	*	*	*	*
038.42	51	17	34	*	*	*9	37	13	14	16	*8
038.49	20	9	11	*	*	*6	13	*	*	*6	*6
038.8	10	*	*8	*	*	*	*8	*	*	*	*
038.9	152	65	88	11	11	18	112	33	29	69	22
041	11	*6	*6	*	*	*	*	*	*	*	*
041.1	*5	*	*	*	*	*	*	*	-	*	*
042	70	49	21	*	52	16	*	28	*9	24	9
047	30	15	15	12	14	*	*	*	*	12	*8
047.9	27	14	13	11	13	*	*	*	*	12	*7
052	*7	*	*	*5	*	-	*	*	*	*	*
053	13	*	10	*	*	*	*8	*	*	*	*
053.9	*8	*	*6	*	*	*	*	*	*	*	*
054	10	*	*8	*6	*	*	*	*	*	*6	*
070	17	*9	*8	*	*9	*6	*	*	*	*7	*
070.5	*7	*	*	-	*	*	*	*	*	*	*
075	*7	*	*	*	*5	*	-	*	*	*	*
079	69	30	39	40	*9	*8	12	14	14	26	14
079.9	65	28	37	37	*9	*7	12	14	14	25	12
079.99	65	28	37	37	*9	*7	12	14	14	25	12
112	15	*7	*9	*	*	*	*9	*5	*	*6	*
117	*7	*	*	*	*	*	*	*	*	*	*
135	*6	*	*	-	*	*	*	*	*	*	*
136	10	*5	*	*	*5	*	*	*	*	*	*
136.3	*5	*	*	*	*	*	*	*	*	*	*
140-239	1,687	655	1,032	36	283	565	802	396	373	599	318
150	15	11	*	-	*	*7	*8	*	*	*6	*
150.5	*6	*	*	-	*	*	*	*	*	*	*
151	29	19	10	-	*	*8	19	*5	*6	*9	*8
151.0	*8	*7	*	-	*	*	*6	*	*	*	*

See footnotes and notes at end of table.

Table 42. Number of first-listed diagnoses for discharges from short-stay hospitals, by ICD-9-CM code, sex, age, and geographic region: United States, 1999—Con.[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table: see Methods section]

ICD-9-CM code	Sex		Age				Region				
	Total	Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	North-east	Midwest	South	West
Number of first-listed diagnoses in thousands											
153	117	53	64	*	*6	28	83	27	32	38	20
153.1	11	*6	*	—	*	*	9	*	*	*	*
153.2	10	*	*7	—	*	*	*6	*	*	*	*
153.3	28	13	15	—	*	*8	18	*8	*7	*7	*5
153.4	22	*9	14	—	*	*	18	*6	*	*8	*
153.6	17	*8	10	—	*	*	14	*	*7	*	*
153.9	15	*7	*7	*	*	*	9	*	*	*8	*
154	45	25	20	*	*	13	29	13	10	13	*9
154.0	17	9	*8	—	*	*	12	*5	*	*5	*
154.1	24	15	10	—	*	*8	16	*6	*7	*7	*
155	14	9	*5	*	*	*6	*7	*	*	*5	*
155.0	10	*6	*	*	*	*	*	*	*	*	*
156	12	*	*8	—	*	*	9	*	*	*	*
156.1	*6	*	*	—	—	*	*6	*	*	*	*
157	31	12	18	—	*	*9	19	*7	*5	12	*6
157.0	15	*7	*8	—	*	*6	9	*	*	*7	*
157.9	10	*	*6	—	*	*	*8	*	*	*	*
161	*8	*6	*	—	*	*	*	*	*	*	*
162	146	83	63	*	*	48	96	31	35	53	27
162.2	*9	*	*	—	*	*	*6	*	*	*	*
162.3	46	25	21	—	*	12	33	13	*8	14	11
162.4	*6	*	*	—	—	*	*	*	*	*	*
162.5	20	12	*8	*	*	*7	13	*	*	*7	*
162.8	16	9	*7	—	*	*5	10	*7	*5	*	*
162.9	49	29	21	*	*	17	31	*	14	25	*7
170	*6	*	*	*	*	*	*	*	*	*	*
171	*6	*	*	*	*	*	*	*	*	*	*
173	*7	*	*	*	*	*	*	*	*	*	*
174	106	—	106	—	13	42	51	31	24	36	15
174.2	*6	—	*6	—	*	*	*	*	*	*	*
174.4	21	—	21	—	*	*9	10	*7	*	*6	*
174.8	29	—	29	—	*	13	11	12	*6	*6	*
174.9	36	—	36	—	*	12	20	*7	*9	18	*
180	17	—	17	*	*7	*7	*	*	*	*7	*
180.8	*6	—	*6	—	*	*	*	*	*	*	*
180.9	*7	—	*7	*	*	*	*	*	*	*	*
182	30	—	30	—	*	12	15	*7	*8	9	*5
182.0	29	—	29	—	*	12	14	*7	*8	9	*
183	20	—	20	*	*	*9	*9	*	*	*7	*
183.0	20	—	20	*	*	*9	*9	*	*	*7	*
184	*7	—	*7	*	*	*	*	*	*	*	*
185	96	96	—	—	*	33	61	21	19	36	20
188	41	30	11	—	*	*6	34	13	11	13	*
188.2	*8	*6	*	—	*	*	*7	*	*	*	*
188.8	10	*7	*	—	—	*	*9	*	*	*	*
188.9	13	*7	*5	—	*	*	10	*	*	*6	*
189	30	17	13	*	*	*9	19	*6	*8	*9	*7
189.0	23	14	9	*	*	*8	13	*5	*6	*7	*5
191	26	15	10	*	*6	*9	*8	*6	*7	*7	*5
191.1	*5	*	*	*	*	*	*	*	*	*	*
191.2	*6	*	*	*	*	*	*	*	*	*	*
193	17	*	14	*	*9	*5	*	*	*	*6	*
195	*6	*	*	*	*	*	*	*	*	*	*

See footnotes and notes at end of table.

Table 42. Number of first-listed diagnoses for discharges from short-stay hospitals, by ICD-9-CM code, sex, age, and geographic region: United States, 1999—Con.

[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table: see Methods section]

ICD-9-CM code	Sex			Age				Region			
	Total	Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	North-east	Midwest	South	West
Number of first-listed diagnoses in thousands											
196	13	10	*	—	*	*	*6	*	*	*	*
196.0	*6	*	*	—	*	*	*	*	*	*	*
197	104	38	66	*	*8	33	62	25	23	37	20
197.0	18	*8	10	*	*	*6	*8	*	*	*	*
197.2	27	*6	21	*	*	10	16	*7	*	10	*
197.4	*5	*	*	—	*	*	*	*	*	*	*
197.6	14	*6	*8	—	*	*	9	*	*	*	*
197.7	30	14	16	*	*	*9	19	*6	*8	12	*
198	98	47	51	*	*	36	57	31	26	25	15
198.3	38	19	19	*	*	17	19	13	*8	12	*
198.5	40	20	19	*	*	12	26	12	13	*8	*7
198.8	12	*	*8	—	*	*	*7	*	*	*	*
198.89	11	*	*7	—	*	*	*6	*	*	*	*
199	30	13	18	*	*	11	17	*6	*8	9	*7
199.0	17	*8	9	*	*	*7	*8	*	*	*5	*
199.1	13	*5	*8	—	*	*	9	*	*	*	*
200	9	*	*6	*	*	*	*5	*	*	*	*
200.0	*6	*	*	—	*	*	*	*	*	*	*
202	36	20	17	*	*	14	20	*8	*7	13	*8
202.8	32	17	15	*	*	12	18	*8	*6	12	*6
202.80	16	10	*6	*	*	*6	10	*	*	*6	*
202.83	*5	*	*	—	*	*	*	*	*	*	*
203	22	13	9	—	*	*	17	*	*7	*6	*
203.0	21	13	*8	—	*	*	16	*	*7	*	*
203.00	20	12	*8	—	*	*	15	*	*6	*	*
204	16	*8	*8	*6	*	*	*6	*5	*	*6	*
204.0	10	*	*5	*6	*	*	*	*	*	*	*
204.00	*8	*	*	*5	*	*	*	*	*	*	*
204.1	*6	*	*	—	*	*	*5	*	*	*	*
204.10	*6	*	*	—	*	*	*5	*	*	*	*
205	21	12	10	*	*	*	11	*5	*	*8	*
205.0	17	*9	*8	*	*	*	9	*	*	*7	*
205.00	16	*8	*8	*	*	*	9	*	*	*7	*
210	*6	*	*	*	*	*	*	*	*	*	*
210.2	*6	*	*	—	*	*	*	*	*	*	*
211	31	14	17	*	*	9	20	*6	*6	12	*7
211.3	24	12	13	*	*	*7	16	*	*	9	*6
218	215	—	215	—	116	95	*	39	45	89	42
218.0	29	—	29	—	14	15	*	*8	*5	11	*
218.1	65	—	65	—	36	28	*	14	12	26	13
218.2	24	—	24	—	14	*9	*	*	*	9	*6
218.9	97	—	97	—	52	44	*	13	23	42	19
220	38	—	38	*	17	14	*5	*8	*8	12	10
225	22	*8	15	*	*6	*9	*8	*6	*	*8	*
225.2	15	*	11	—	*	*6	*7	*	*	*6	*
226	*8	*	*	*	*	*	*	*	*	*	*
227	13	*5	*7	*	*	*7	*	*	*	*	*
227.1	*6	*	*	—	*	*	*	*	*	*	*
227.3	*5	*	*	*	*	*	*	*	*	*	*
228	*9	*	*5	*5	*	*	*	*	*	*6	*
228.0	*8	*	*5	*5	*	*	*	*	*	*5	*
233	30	*	30	—	14	9	*6	*5	*5	13	*6
233.0	11	—	11	—	*	*	*5	*	*	*5	*
233.1	18	—	18	—	14	*	*	*	*	*8	*
235	*8	*	*	*	*	*	*5	*	*	*	*
235.2	*6	*	*	—	*	*	*	*	*	*	*

See footnotes and notes at end of table.

Table 42. Number of first-listed diagnoses for discharges from short-stay hospitals, by ICD-9-CM code, sex, age, and geographic region: United States, 1999—Con.[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table: see Methods section]

ICD-9-CM code	Sex		Age				Region				
	Total	Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	North-east	Midwest	South	West
Number of first-listed diagnoses in thousands											
237	*6	*	*	*	*	*	*	*	*	*	*
238	13	*5	*8	*	*	*	10	*	*	*5	*
238.7	*9	*	*5	*	*	*	*7	*	*	*	*
239	16	*	12	*	*	*7	*6	*	*	11	*
239.6	*6	*	*	*	*	*	*	*	*	*	*
240-279	1,393	577	816	159	264	351	618	316	329	518	229
241	15	*	14	*	*	*8	*	*	*	*	*
241.1	*8	*	*7	*	*	*	*	*	*	*	*
242	11	*	*8	*	*5	*	*	*	*	*5	*
242.9	*6	*	*	*	*	*	*	*	*	*	*
242.90	*5	*	*	*	*	*	*	*	*	*	*
244	15	*	12	*	*	*	*8	*	*	*	*
244.9	14	*	12	*	*	*	*8	*	*	*	*
250	545	262	283	21	138	187	199	119	115	214	97
250.0	163	68	95	9	33	64	57	32	29	77	25
250.00	64	29	35	*	*8	30	25	11	11	34	*8
250.01	25	10	15	*	*7	*8	*6	*5	*	12	*
250.02	44	18	27	*	9	17	18	9	*7	19	9
250.03	29	11	18	*	*8	*9	*9	*7	*6	12	*
250.1	101	51	50	10	61	22	*8	22	27	35	17
250.10	14	*7	*8	*	*9	*	*	*	*	*6	*
250.11	33	16	17	*5	22	*5	*	*8	10	*8	*8
250.12	14	*6	*8	*	*5	*5	*	*	*	*6	*
250.13	40	22	18	*	25	*9	*	*8	10	16	*6
250.2	14	*7	*6	*	*	*5	*6	*	*	*	*
250.22	*7	*	*	*	*	*	*	*	*	*	*
250.3	*7	*	*	*	*	*	*	*	*	*	*
250.4	31	16	15	—	*	14	12	*6	*	11	*8
250.40	13	*7	*6	—	*	*7	*5	*	*	*5	*
250.41	10	*	*5	—	*	*	*	*	*	*	*
250.42	*6	*	*	—	*	*	*	*	*	*	*
250.5	*6	*	*	—	*	*	*	*	*	*	*
250.6	53	22	31	*	11	22	19	12	12	17	13
250.60	26	10	16	—	*5	10	11	*5	*	*8	*8
250.61	12	*6	*6	*	*	*6	*	*	*	*	*
250.62	*8	*	*	—	*	*	*	*	*	*	*
250.63	*7	*	*	—	*	*	*	*	*	*	*
250.7	63	36	27	—	*	21	38	16	12	25	9
250.70	31	18	13	—	*	*9	21	*8	*7	12	*
250.71	17	*9	*9	—	*	*7	9	*	*	*7	*
250.72	*9	*6	*	—	*	*	*5	*	*	*	*
250.73	*6	*	*	—	*	*	*	*	*	*	*
250.8	95	49	46	*	10	33	50	21	25	33	16
250.80	43	23	20	—	*	14	26	10	11	14	*8
250.81	23	13	10	*	*	9	9	*5	*6	*8	*
250.82	16	*6	9	—	*	*6	10	*	*	*7	*
250.83	13	*7	*6	*	*	*7	*	*	*	*	*
250.9	12	*8	*	*	*5	*	*	*	*	*7	*
251	11	*5	*6	*	*	*	*	*	*	*5	*
251.2	*8	*	*	*	*	*	*	*	*	*	*
252	*5	*	*	—	*	*	*	*	*	*	*
253	12	*6	*7	*	*	*	*8	*	*	*	*
253.6	9	*	*5	*	*	*	*8	*	*	*	*
255	*6	*	*	*	*	*	*	*	*	*	*
261	*5	*	*	*	*	*	*	*	*	*	*

See footnotes and notes at end of table.

Table 42. Number of first-listed diagnoses for discharges from short-stay hospitals, by ICD-9-CM code, sex, age, and geographic region: United States, 1999—Con.

[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table: see Methods section]

ICD-9-CM code	Sex			Age				Region			
	Total	Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	North-east	Midwest	South	West
Number of first-listed diagnoses in thousands											
263	18	*7	11	*	*	*	14	*	*	*7	*
263.9	17	*7	10	*	*	*	13	*	*	*7	*
272	29	13	16	*	*8	13	*8	10	*	10	*
272.0	14	*6	*8	—	*	*7	*	*7	*	*	*
272.4	10	*5	*	—	*	*	*	*	*	*	*
274	11	*5	*5	—	*	*	*7	*	*	*	*
274.0	*7	*	*	—	*	*	*	*	*	*	*
275	18	*8	10	*	*	*	11	*	*	*8	*
275.4	16	*7	*9	*	*	*	10	*	*	*7	*
275.42	11	*5	*6	*	*	*	*8	*	*	*6	*
276	621	235	386	120	69	98	334	136	163	224	98
276.0	*8	*	*	*	*	*	*6	*	*	*	*
276.1	65	20	45	*	*	13	46	15	15	20	16
276.2	*7	*	*	*	*	*	*	*	*	*	*
276.5	477	186	292	115	52	63	247	106	130	171	71
276.7	21	*8	13	*	*	*	12	*	*	10	*
276.8	32	10	23	*	*5	12	15	*6	*8	15	*
277	16	*7	9	*7	*6	*	*	*6	*	*	*
277.0	12	*5	*7	*6	*5	*	—	*	*	*	*
277.00	12	*5	*6	*6	*5	*	—	*	*	*	*
278	33	*8	25	*	16	15	*	*8	*8	10	*8
278.0	30	*8	22	*	14	14	*	*7	*7	*9	*7
278.00	*6	*	*	*	*	*	*	*	*	*	*
278.01	24	*6	18	—	12	11	*	*6	*6	*7	*5
280-289	368	162	206	57	96	74	141	88	83	136	61
280	47	15	32	*	*7	*8	30	13	12	15	*6
280.0	22	*6	15	—	*	*	15	*8	*	*6	*
280.9	25	*8	16	*	*	*5	14	*	*7	9	*
281	*6	*	*	*	*	*	*	*	*	*	*
282	68	31	37	17	47	*	*	21	12	30	*
282.6	60	28	32	15	42	*	*	18	11	28	*
282.62	54	25	29	11	40	*	*	16	9	25	*
283	*6	*	*	*	*	*	*	*	*	*	*
284	32	18	14	*6	*	*9	13	*8	*7	*8	*8
284.8	28	15	12	*	*	*8	11	*6	*7	*6	*8
285	73	27	47	*	*9	18	43	13	19	31	10
285.1	13	*5	*8	—	*	*	*9	*	*	*	*
285.9	56	21	36	*	*6	14	32	10	14	27	*6
286	18	*8	10	*	*	*	11	*	*	*8	*
286.9	*5	*	*	*	*	*	*	*	*	*	*
287	37	18	20	*6	*9	10	12	9	*6	11	11
287.3	14	*7	*7	*	*	*	*	*	*	*	*
287.4	*5	*	*	*	*	*	*	*	*	*	*
287.5	11	*5	*6	*	*	*	*	*	*	*	*
288	57	29	28	12	*8	15	23	14	15	20	9
288.0	49	24	25	10	*6	14	19	12	12	16	*8
288.8	*7	*	*	*	*	*	*	*	*	*	*
289	24	12	12	*7	*9	*	*	*	*	9	*5
289.2	*8	*	*	*	*	*	*	*	*	*	*
289.8	*5	*	*	*	*	*	*	*	*	*	*

See footnotes and notes at end of table.

Table 42. Number of first-listed diagnoses for discharges from short-stay hospitals, by ICD-9-CM code, sex, age, and geographic region: United States, 1999—Con.[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table: see Methods section]

ICD-9-CM code	Sex			Age				Region			
	Total	Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	North-east	Midwest	South	West
Number of first-listed diagnoses in thousands											
290-319	2,018	1,048	970	103	1,167	484	264	615	475	718	210
290	30	11	18	*	—	*	28	*7	10	10	*
290.0	*6	*	*	—	—	—	*6	*	*	*	—
290.2	*7	*	*5	*	—	—	*7	*	*	*	*
290.4	12	*6	*6	—	—	*	11	*	*	*6	*
290.40	*5	*	*	—	—	*	*	*	*	*	*
291	108	83	25	—	59	40	9	31	40	18	19
291.0	15	12	*	—	*8	*6	*	*6	*	*	*
291.2	*5	*	*	—	*	*	*	*	*	*	*
291.8	82	64	19	—	48	29	*5	25	34	11	13
291.81	78	59	18	—	45	27	*5	24	31	11	12
292	64	36	29	*	44	12	*8	30	10	16	*7
292.0	36	22	14	*	27	*7	*	23	*5	*5	*
292.8	22	11	11	—	12	*	*7	*6	*	*8	*
292.81	10	*	*6	—	*	*	*7	*	*	*	*
292.84	*9	*6	*	—	*8	*	—	*	*	*	*
293	37	20	17	*	11	*9	17	17	9	*8	*
293.0	19	*7	11	*	*	*	14	*	*7	*	*
293.8	17	11	*6	—	10	*	*	12	*	*	*
293.83	14	9	*	—	*9	*	*	12	*	*	*
294	30	12	18	*	*	*	24	9	*8	*8	*
294.8	24	*8	16	*	*	*	21	*8	*7	*6	*
295	260	147	112	*	160	75	24	91	49	90	29
295.3	86	50	36	*	50	26	10	29	13	32	11
295.30	23	13	10	—	15	*6	*	*	*7	*8	*
295.32	15	9	*6	—	*7	*	*	*7	*	*6	*
295.34	47	28	19	*	28	14	*5	18	*5	18	*6
295.6	25	17	*8	*	15	*8	*	11	*	*7	*
295.60	*5	*	*	*	*	*	*	*	*	*	*
295.62	*7	*5	*	—	*	*	*	*	*	*	—
295.64	12	*9	*	—	*7	*	*	*	*	*	*
295.7	116	57	59	*	76	31	9	42	25	40	10
295.70	76	36	40	*	49	21	*6	21	20	28	*6
295.74	37	20	17	—	25	9	*	19	*	11	*
295.9	26	19	*8	*	15	*9	*	*8	*6	*9	*
295.90	17	11	*5	*	*8	*7	*	*	*5	*7	*
295.94	*8	*6	*	—	*6	*	*	*6	*	*	*
296	697	287	410	39	415	162	81	187	177	280	53
296.2	200	79	120	19	116	36	29	38	55	95	12
296.20	95	37	58	*8	57	16	13	20	29	39	*7
296.22	25	10	14	*	15	*5	*	*	*	21	*
296.23	48	20	28	*5	26	*8	*8	*8	14	24	*
296.24	31	12	19	*	17	*6	*5	9	10	11	*
296.3	245	87	158	*8	141	60	36	72	65	90	19
296.30	72	24	49	*	38	20	12	24	20	24	*
296.32	17	*7	10	*	10	*	*	*	*6	*8	*
296.33	100	38	62	*	65	20	12	28	27	38	*6
296.34	55	18	37	*	28	18	*8	18	11	19	*7
296.4	56	27	29	*	31	19	*5	19	12	19	*5
296.40	25	11	14	*	14	*7	*	*6	*6	11	*
296.43	*6	*	*	—	*	*	*	*	*	*	*
296.44	23	13	10	*	12	*8	*	11	*	*	*
296.5	54	26	28	*	36	13	*	14	16	21	*
296.50	29	14	15	*	18	*6	*	*6	*8	14	*
296.53	13	*6	*7	*	10	*	*	*5	*	*	*
296.54	10	*	*6	*	*6	*	*	*	*	*	*

See footnotes and notes at end of table.

Table 42. Number of first-listed diagnoses for discharges from short-stay hospitals, by ICD-9-CM code, sex, age, and geographic region: United States, 1999—Con.

[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table: see Methods section]

ICD-9-CM code	Sex			Age				Region			
	Total	Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	North-east	Midwest	South	West
Number of first-listed diagnoses in thousands											
296.6	59	25	34	*	35	16	*	17	14	25	*
296.60	34	16	19	*	21	10	*	*6	*7	20	*
296.63	10	*	*6	*	*7	*	*	*	*	*	*
296.64	14	*6	*9	*	*7	*	*	*6	*	*	*
296.7	45	22	23	*	28	10	*	10	9	20	*5
296.8	14	*6	*9	*	12	*	*	*7	*	*	*
296.89	10	*	*7	*	*8	*	*	*	*	*	*
296.9	22	14	*8	*	15	*	*	10	*	*7	*
296.90	20	13	*7	*	14	*	—	10	*	*7	*
297	*9	*	*5	—	*	*	*	*	*	*	*
297.9	*7	*	*	—	*	*	*	*	*	*	*
298	71	38	33	*	36	17	16	28	12	23	*8
298.9	67	36	31	*	34	16	15	27	11	22	*8
300	79	31	49	*	41	22	13	21	24	20	14
300.0	32	12	20	*	10	14	*8	*8	*9	10	*6
300.00	23	*9	15	*	*7	10	*6	*6	*7	*8	*
300.01	*7	*	*	—	*	*	*	*	*	*	*
300.1	11	*	*9	*	*7	*	*	*	*	*	*
300.11	*7	*	*5	*	*	*	*	*	*	*	*
300.4	23	*8	14	*	13	*	*	*9	*7	*	*
300.9	*6	*	*	*	*	*	*	*	*	*	*
301	10	*	*5	*	*7	*	*	*	*	*	*
303	164	119	45	*	92	60	11	28	40	75	21
303.0	33	24	*8	*	15	14	*	*7	*9	10	*7
303.00	11	*8	*	—	*6	*5	*	*	*	*	*
303.01	20	16	*	*	9	*9	*	*	*	*6	*5
303.9	131	94	37	*	77	46	*8	21	31	65	14
303.90	64	45	19	*	36	24	*	*	9	50	*
303.91	66	49	17	*	41	22	*	19	22	15	10
304	84	51	33	*	64	18	*	24	10	43	*7
304.0	34	20	14	—	25	*8	*	13	*	14	*
304.00	12	*6	*6	—	9	*	*	*	*	10	*
304.01	22	14	*8	—	16	*6	*	12	*	*	*
304.1	*5	*	*	—	*	*	*	*	*	*	*
304.2	21	16	*6	—	18	*	*	*5	*	13	*
304.20	11	*8	*	—	9	*	—	*	*	9	—
304.21	10	*8	*	—	*9	*	*	*5	*	*	*
304.7	*7	*	*	—	*6	*	*	*	*	*	*
304.71	*6	*	*	—	*5	*	*	*	*	*	*
304.9	*6	*	*	—	*	*	*	*	*	*6	*
304.90	*6	*	*	—	*	*	*	—	*	*5	*
305	82	55	26	*	52	23	*6	14	15	40	13
305.0	36	25	11	*	23	*9	*	*6	*7	17	*
305.00	26	19	*7	*	17	*6	*	*	*6	13	*
305.01	*8	*6	*	—	*	*	*	*	*	*	*
305.1	18	10	*8	—	9	*8	*	*	*	*7	*
305.6	10	*6	*	—	*9	*	—	*	*	*	*
305.60	*7	*	*	—	*6	*	—	*	*	*	*
305.9	*9	*7	*	*	*6	*	—	*	*	*6	*
305.90	*7	*5	*	*	*	*	—	*	*	*	*
307	11	*	*8	*	*6	*	*	*	*	*	*
308	*6	*	*	*	*	*	*	—	*	*	*

See footnotes and notes at end of table.

Table 42. Number of first-listed diagnoses for discharges from short-stay hospitals, by ICD-9-CM code, sex, age, and geographic region: United States, 1999—Con.[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table: see Methods section]

ICD-9-CM code	Sex			Age				Region			
	Total	Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	North-east	Midwest	South	West
Number of first-listed diagnoses in thousands											
309	85	39	46	*8	64	10	*	37	19	21	*8
309.0	34	15	19	*	27	*5	*	16	*7	*7	*
309.2	13	*7	*6	*	9	*	*	*	*	*5	*
309.28	10	*	*5	*	*9	*	*	*	*	*	*
309.4	15	*8	*7	*	11	*	*	*8	*	*	*
309.8	13	*	*9	*	*9	*	—	*5	*	*	*
309.81	12	*	*9	*	*9	*	—	*5	*	*	*
309.9	*7	*	*	*	*7	*	*	*	*	*	*
310	*8	*	*	*	*	*	*5	*	*	*	*
311	110	53	57	10	73	18	9	51	16	35	*8
312	29	22	*8	*8	18	*	*	15	*7	*5	*
312.3	21	15	*6	*	15	*	*	12	*	*	*
312.30	12	*8	*	*	9	*	*	*9	*	*	*
312.34	*8	*6	*	*	*5	*	*	*	*	*	*
312.9	*6	*5	*	*	*	*	*	*	*	*	—
313	12	*7	*	*6	*6	—	—	*7	*	*	*
313.8	11	*7	*	*6	*6	—	—	*7	*	*	*
313.81	11	*7	*	*5	*6	—	—	*7	*	*	*
314	21	15	*5	15	*5	—	—	*	13	*	*
314.0	21	15	*5	15	*5	—	—	*	13	*	*
314.01	19	14	*5	14	*	—	—	*	12	*	*
320-389	503	226	278	89	115	112	188	105	123	190	86
320	*8	*	*	*	*	*	*	*	*	*	*
322	*6	*	*	*	*	*	*	*	*	*	*
322.9	*6	*	*	*	*	*	*	*	*	*	*
324	*5	*	*	*	*	*	*	*	*	*	*
331	67	24	43	*	*	*	57	10	16	29	12
331.0	50	15	35	—	*	*	49	*7	13	23	*8
331.3	*7	*	*	*	*	*	*	*	*	*	*
331.4	*8	*	*	*	*	*	*	*	*	*	*
332	22	13	9	—	*	*	19	*6	*7	*7	*
332.0	22	12	9	—	*	*	19	*5	*7	*7	*
333	*8	*	*	*	*	*	*	*	*	*	*
335	*5	*	*	*	*	*	*	*	*	*	*
335.2	*5	*	*	—	*	*	*	*	*	*	*
340	22	*9	14	*	10	10	*	*	*8	*8	*
343	*7	*	*	*	*	*	*	*	*	*	*
344	*9	*	*	*	*	*	*	*	*	*	*
345	51	24	27	16	19	10	*7	18	*7	17	9
345.1	*8	*	*	*	*	*	*	*	*	*	*
345.10	*8	*	*	*	*	*	*	*	*	*	*
345.3	15	*7	*8	*	*5	*	*	*	*	*	*
345.5	*6	*	*	*	*	*	*	*	*	*	*
345.9	12	*7	*5	*	*	*	*	*	*	*	*
345.90	10	*5	*	*	*	*	*	*	*	*	*
346	39	10	29	*	20	15	*	*8	14	11	*6
346.8	*5	*	*	*	*	*	*	*	*	*	*
346.9	24	*	19	*	13	*9	*	*	*9	*7	*
346.90	17	*	13	*	*8	*7	*	*	*7	*6	*
346.91	*6	*	*6	*	*5	*	*	*	*	*	*
348	28	14	14	*6	*8	*8	*7	*5	*6	*9	*8
348.1	*6	*	*	*	*	*	*	*	*	*	*
348.3	*6	*	*	*	*	*	*	*	*	*	*

See footnotes and notes at end of table.

Table 42. Number of first-listed diagnoses for discharges from short-stay hospitals, by ICD-9-CM code, sex, age, and geographic region: United States, 1999—Con.

[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table: see Methods section]

ICD-9-CM code	Sex			Age				Region			
	Total	Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	North-east	Midwest	South	West
Number of first-listed diagnoses in thousands											
349	*9	*	*	*	*	*	*	*	*	*	*
350	*5	*	*	*	*	*	*	*	*	*	*
353	*6	*	*	*	*	*	*	*	*	*	*
356	*6	*	*	*	*	*	*	*	*	*	*
357	10	*5	*	*	*	*	*	*	*	*6	*
357.0	*5	*	*	*	*	*	*	*	*	*	*
358	*6	*	*	*	*	*	*	*	*	*	*
358.0	*5	*	*	—	*	*	*	*	*	*	*
361	*5	*	*	*	*	*	*	*	*	*	*
366	*5	*	*	—	*	*	*	*	*	*	*
368	*8	*	*	*	*	*	*	*	*	*	*
376	9	*	*	*5	*	*	*	*	*	*	*
376.0	*8	*	*	*5	*	*	*	*	*	*	*
376.01	*8	*	*	*	*	*	*	*	*	*	*
380	*8	*5	*	*	*	*	*	*	*	*	*
380.1	*6	*	*	*	*	*	*	*	*	*	*
380.10	*6	*	*	*	*	*	*	*	*	*	*
382	24	12	12	22	*	*	*	*	*7	11	*
382.9	23	11	12	21	*	*	*	*	*6	11	*
383	*6	*	*	*	*	*	*	*	*	*	*
386	39	15	23	*	*	*8	28	*7	12	13	*6
386.1	13	*	9	*	*	*	10	*	*	*	*
386.11	*6	*	*	*	*	*	*	*	*	*	*
386.3	21	*9	12	*	*	*	16	*	*8	*6	*
386.30	19	*7	11	—	*	*	14	*	*6	*6	*
390-459	6,344	3,161	3,183	25	417	1,797	4,105	1,485	1,523	2,328	1,009
394	*7	*	*6	*	*	*	*	*	*	*	*
396	14	*	10	—	*	*	11	*	*	*	*
396.2	*6	*	*	—	*	*	*5	*	*	*	*
396.3	*6	*	*	—	*	*	*	*	*	*	*
398	21	*8	14	—	*	*	19	*8	*	*9	*
398.9	21	*8	14	—	*	*	19	*8	*	*9	*
398.91	21	*8	14	—	*	*	19	*8	*	*9	*
401	224	83	141	*	37	86	100	55	44	93	32
401.0	26	*9	18	—	*6	*8	13	*5	*	15	*
401.1	*6	*	*	*	*	*	*	*	*	*	*
401.9	192	73	119	*	30	77	84	49	40	77	26
402	97	33	64	*	*	23	68	30	17	40	11
402.0	*6	*	*	—	*	*	*	*	*	*	—
402.9	89	29	60	*	*	21	63	27	16	36	10
402.90	16	*7	9	—	*	*8	*7	*5	*	*7	*
402.91	73	22	51	*	*	13	56	22	14	29	*9
403	88	42	46	*	20	31	36	11	14	45	18
403.9	82	40	42	*	18	29	35	10	13	42	18
403.91	79	40	40	*	17	28	33	10	13	39	18
404	30	14	16	*	*	*7	18	*6	*6	13	*
404.9	27	13	14	*	*	*6	17	*6	*6	12	*
404.93	21	9	12	*	*	*	14	*5	*	*9	*

See footnotes and notes at end of table.

Table 42. Number of first-listed diagnoses for discharges from short-stay hospitals, by ICD-9-CM code, sex, age, and geographic region: United States, 1999—Con.[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table: see Methods section]

ICD-9-CM code	Sex			Age				Region			
	Total	Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	North-east	Midwest	South	West
Number of first-listed diagnoses in thousands											
410	829	481	348	*	46	274	509	207	208	266	148
410.0	28	14	14	—	*	12	15	*7	*6	*7	*8
410.01	28	14	14	—	*	12	15	*7	*6	*7	*8
410.1	130	77	53	—	*9	49	72	26	29	48	28
410.11	127	76	51	—	*8	49	70	25	29	47	27
410.2	24	14	10	—	*	*6	16	*	*	10	*
410.21	24	14	10	—	*	*6	16	*	*	10	*
410.3	17	11	*5	—	*	*8	*8	*	*6	*	*
410.31	17	11	*5	—	*	*8	*8	*	*6	*	*
410.4	163	108	54	—	16	62	84	45	40	46	32
410.41	159	107	52	—	16	61	82	44	40	44	31
410.5	12	*8	*	—	*	*5	*6	*	*	*	*
410.51	12	*8	*	—	*	*5	*6	*	*	*	*
410.6	*8	*7	*	—	*	*	*	*	*	*	*
410.61	*8	*7	*	—	*	*	*	*	*	*	*
410.7	313	173	140	*	11	94	208	86	78	98	50
410.71	308	170	138	*	11	93	204	84	77	97	50
410.8	11	*7	*	—	*	*	*7	*	*	*	*
410.81	11	*7	*	—	*	*	*7	*	*	*	*
410.9	123	61	62	—	*	29	89	27	34	45	17
410.90	*8	*	*5	—	*	*	*6	*	*	*	*
410.91	114	58	56	—	*	28	82	23	32	43	16
411	166	80	86	*	12	64	91	41	33	72	21
411.1	153	73	80	*	10	61	83	37	30	67	20
411.8	12	*6	*6	—	*	*	*8	*	*	*	*
411.89	*9	*5	*	—	*	*	*5	*	*	*	*
412	*8	*5	*	—	*	*	*	*	*	*	*
413	82	35	47	*	11	27	44	18	20	35	10
413.9	82	35	47	*	11	27	44	17	20	35	10
414	1,177	716	461	*	50	464	660	278	273	426	200
414.0	1,153	703	450	*	49	455	648	274	268	415	197
414.00	193	106	87	—	*8	69	116	36	44	87	26
414.01	904	559	345	*	40	365	498	226	212	305	161
414.02	37	24	12	—	*	12	24	*8	9	11	*8
414.04	*6	*	*	—	*	*	*	*	*	*	*
414.05	13	*8	*5	—	*	*	*8	*	*	*8	*
414.8	12	*9	*	—	*	*5	*6	*	*	*5	*
414.9	10	*	*6	—	*	*	*6	*	*	*5	*
415	91	34	57	*	13	25	53	20	23	30	18
415.1	90	33	57	—	13	24	52	20	22	30	18
415.11	*6	*	*	—	*	*	*	*	*	*	*
415.19	84	30	53	—	12	21	50	17	21	28	17
416	12	*5	*7	*	*	*	*	*	*	*	*
416.0	*7	*	*	*	*	*	*	*	*	*	*
420	12	*	*7	—	*	*6	*	*	*	*	*
420.9	12	*	*7	—	*	*6	*	*	*	*	*
420.90	*9	*	*5	—	*	*	*	*	*	*	*
421	14	9	*	*	*5	*	*	*	*	*	*
421.0	13	*9	*	*	*	*	*	*	*	*	*
423	14	*5	*9	*	*	*	*7	*	*	*	*
423.9	12	*	*8	*	*	*	*6	*	*	*	*
424	97	43	54	*	10	25	62	27	20	32	18
424.0	43	18	25	*	*8	16	19	11	*8	17	*7
424.1	52	24	28	*	*	*8	42	16	12	14	11
425	35	15	19	*	*6	13	15	*7	*7	16	*
425.4	25	12	13	*	*	*9	12	*5	*7	11	*

See footnotes and notes at end of table.

Table 42. Number of first-listed diagnoses for discharges from short-stay hospitals, by ICD-9-CM code, sex, age, and geographic region: United States, 1999—Con.

[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table: see Methods section]

ICD-9-CM code	Sex			Age				Region			
	Total	Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	North-east	Midwest	South	West
Number of first-listed diagnoses in thousands											
426	57	32	24	*	*6	9	42	17	15	14	12
426.0	29	16	14	*	*	*	25	*9	*7	*8	*6
426.1	11	*6	*6	*	*	*	9	*	*	*	*
427	704	329	375	*5	53	156	491	186	166	246	107
427.0	28	10	18	*	*	10	14	*7	*6	12	*
427.1	64	42	22	*	*	17	42	17	15	19	13
427.3	384	174	210	*	22	84	277	107	98	122	57
427.31	347	150	197	*	20	73	254	97	88	111	51
427.32	37	24	13	*	*	11	23	10	10	11	*6
427.4	*9	*6	*	—	*	*	*	*	*	*	*
427.41	*9	*6	*	—	*	*	*	*	*	*	*
427.5	10	*7	*	*	*	*	*	*	*	*	*
427.6	18	*7	11	*	*	*	11	*	*	*8	*
427.69	12	*6	*7	*	*	*	*7	*	*	*6	*
427.8	176	74	102	*	15	31	128	42	38	72	25
427.81	68	24	44	*	*	*7	60	19	11	30	*8
427.89	108	50	58	*	13	24	68	23	27	41	17
427.9	16	*9	*7	*	*	*	10	*6	*	*6	*
428	975	434	541	*	31	175	767	215	260	371	129
428.0	962	430	532	*	30	173	757	214	258	364	126
428.1	*5	*	*	—	—	*	*	*	*	*	*
428.9	*7	*	*	*	*	*	*5	*	*	*6	*
429	19	10	10	*	*	*5	12	*	*5	11	*
429.2	*8	*	*	—	*	*	*6	—	*	*5	*
430	21	*7	14	*	*	*8	9	*5	*	*8	*
431	61	31	30	*	*5	17	39	11	13	21	16
432	18	12	*6	*	*	*	13	*6	*	*	*
432.1	15	11	*	*	*	*	11	*	*	*	*
433	170	86	84	—	*	41	128	33	46	65	26
433.1	126	64	62	—	*	29	96	22	34	51	19
433.10	104	54	50	—	*	23	81	19	27	40	17
433.11	22	10	12	—	*	*7	15	*	*6	10	*
433.3	40	19	20	—	*	10	29	*9	12	13	*7
433.30	32	15	17	—	*	*9	23	*8	10	10	*
433.31	*7	*	*	—	—	*	*6	*	*	*	*
434	289	127	162	*	12	67	209	65	70	107	47
434.0	21	9	11	*	*	*	16	*	*8	*8	*
434.01	16	*7	*8	*	*	*	12	*	*6	*5	*
434.1	35	17	18	—	*	*7	26	*9	*8	10	*8
434.11	32	15	17	—	*	*6	24	*7	*7	10	*8
434.9	233	101	132	*	*9	56	167	54	54	89	36
434.90	*9	*	*	*	*	*	*5	*	*	*	*
434.91	224	96	128	*	*8	54	162	53	52	85	35
435	196	79	118	*	*9	39	149	47	52	71	26
435.3	11	*	*7	—	*	*	10	*	*	*	*
435.9	180	73	107	*	*8	36	135	44	47	66	23
436	149	69	80	*	*	25	120	18	40	64	28
437	32	13	18	*	*	*9	18	*	12	10	*6
437.1	*6	*	*	—	*	*	*5	*	*	*	*
437.3	*7	*	*	*	*	*	*	*	*	*	*
437.9	*5	*	*	—	*	*	*	—	*	*	*
438	25	10	15	*	*	*5	19	*	*5	10	*6
438.2	*8	*	*	*	—	*	*6	*	*	*	*
438.20	*8	*	*	*	—	*	*6	*	*	*	*
438.8	11	*	*7	—	*	*	*8	*	*	*	*
438.89	10	*	*6	—	*	*	*6	*	*	*	*

See footnotes and notes at end of table.

Table 42. Number of first-listed diagnoses for discharges from short-stay hospitals, by ICD-9-CM code, sex, age, and geographic region: United States, 1999—Con.[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table: see Methods section]

ICD-9-CM code	Sex			Age				Region			
	Total	Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	North-east	Midwest	South	West
Number of first-listed diagnoses in thousands											
440	120	65	55	—	*	35	83	28	34	41	17
440.1	*5	*	*	—	*	*	*	*	*	*	*
440.2	99	53	46	—	*	27	71	22	31	33	13
440.20	10	*6	*	—	*	*	*	*	*	*5	*
440.21	25	16	*9	—	*	*8	16	*	*8	*6	*6
440.22	16	*7	*8	—	*	*	13	*	*	*8	*
440.23	14	*6	*9	—	*	*	12	*	*	*	*
440.24	33	18	16	—	*	*9	25	11	12	*8	*
440.3	*8	*6	*	—	*	*	*	*	*	*	*
441	63	42	20	—	*	13	47	17	14	21	10
441.0	*8	*5	*	—	*	*	*	*	*	*	*
441.4	43	30	13	—	*	*7	36	12	10	15	*6
442	*9	*6	*	*	*	*	*7	*	*	*	*
442.3	*5	*	*	*	*	*	*	*	*	*	*
443	23	10	13	*	*	9	13	*	*	12	*
443.9	20	9	11	—	*	*9	11	*	*	11	*
444	43	18	26	—	*	14	26	12	10	17	*
444.2	29	13	17	—	*	*7	20	*7	*8	12	*
444.22	25	11	14	—	*	*7	17	*6	*7	11	*
444.8	*7	*	*	—	*	*	*	*	*	*	*
444.81	*5	*	*	—	*	*	*	*	*	*	*
446	14	*8	*6	*	*	*	*5	*	*	*	*
447	13	*6	*7	*	*	*	*8	*	*	*	*
447.1	*8	*	*	—	*	*	*	*	*	*	*
451	31	10	21	*	*	*8	17	*9	*7	12	*
451.0	*6	*	*	—	*	*	*	*	*	*	*
451.1	13	*	*9	—	*	*	9	*	*	*6	*
451.19	11	*	*8	—	*	*	*8	*	*	*	*
451.2	*7	*	*	*	*	*	*	*	*	*	*
453	147	63	84	—	23	46	78	30	39	56	22
453.8	129	55	74	—	21	39	69	26	36	45	21
453.9	12	*	*8	—	*	*	*7	*	*	*5	*
454	9	*6	*	*	*	*	*6	*	*	*	*
455	24	11	13	—	*8	*7	9	*6	*	11	*
455.2	9	*	*5	—	*	*	*	*	*	*	*
456	*8	*5	*	*	*	*	*	*	*	*	*
456.0	*5	*	*	*	*	*	*	*	*	*	*
458	70	34	36	*	*	13	54	14	18	23	14
458.0	39	18	21	*	*	*5	31	10	12	*9	*8
458.9	27	13	14	*	*	*6	20	*	*6	13	*5
459	21	10	11	*	*	*6	13	*	*	12	*
459.8	*9	*	*6	*	*	*	*	*	*	*	*
459.81	*7	*	*	—	*	*	*	*	*	*	*
460-519	3,689	1,742	1,947	741	430	686	1,832	771	901	1,378	640
461	14	*7	*7	*	*6	*	*	*	*	*	*
462	13	*	9	*	*7	*	*	*	*	*5	*
463	10	*	*6	*	*7	*	—	*	*	*	*
464	47	31	16	40	*	*	*	*6	10	16	15
464.4	38	26	12	38	—	—	—	*	*6	14	13
465	44	19	25	18	*7	*6	12	10	*8	19	*8
465.9	42	18	24	17	*7	*6	12	10	*7	18	*7

See footnotes and notes at end of table.

Table 42. Number of first-listed diagnoses for discharges from short-stay hospitals, by ICD-9-CM code, sex, age, and geographic region: United States, 1999—Con.

[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table: see Methods section]

ICD-9-CM code	Sex			Age				Region			
	Total	Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	North-east	Midwest	South	West
Number of first-listed diagnoses in thousands											
466	298	153	145	204	19	20	56	52	65	123	58
466.0	108	45	63	16	18	19	55	20	27	51	11
466.1	190	108	82	188	*	*	*	32	38	72	48
466.11	106	62	45	106	*	*	*	16	20	47	23
466.19	83	47	37	82	*	*	*	16	17	26	25
473	17	9	*8	*	*	*	*	*	*	*7	*
473.9	*6	*	*	*	*	*	*	*	*	*	*
474	13	*7	*7	10	*	*	—	*5	*	*	*
474.0	*6	*	*	*	*	—	—	*	*	*	*
474.1	*7	*	*	*5	*	*	—	*	*	*	*
474.10	*5	*	*	*	*	—	—	*	*	*	*
475	11	*6	*	*	*7	*	*	*	*	*	*
478	20	12	*8	*7	*	*6	*	*6	*	*5	*7
478.2	*9	*6	*	*	*	*	*	*	*	*	*
478.7	*5	*	*	*	*	*	*	*	*	*	*
480	45	23	22	31	*	*	*8	*	11	16	14
480.1	22	12	10	20	*	*	*	*	*	10	*7
480.9	18	10	*8	*9	*	*	*	*	*	*5	*6
481	50	26	24	*5	*6	15	23	*9	13	19	*8
482	159	88	71	9	15	22	113	23	47	65	24
482.0	10	*5	*	*	*	*	*7	*	*	*6	*
482.1	36	20	16	*	*	*5	23	*	*9	14	10
482.2	16	*8	*9	*	*	*	14	*	*7	*5	*
482.3	*6	*	*	*	*	*	*	*	*	*	*
482.4	39	25	14	*	*	*	31	*6	*9	18	*6
482.41	34	22	13	*	*	*	26	*5	*8	15	*6
482.8	36	19	18	*	*	*	26	*	16	14	*
482.83	30	15	15	*	*	*	21	*	14	10	*
482.9	15	*8	*8	*	*	*	*9	*5	*	*6	*
483	12	*	*8	*	*	*	*6	*	*	*	*
483.0	11	*	*8	*	*	*	*5	*	*	*	*
485	35	17	18	14	*	*	13	*5	12	13	*
486	1,078	514	564	149	102	181	647	209	275	405	190
487	37	14	23	*6	*6	*6	18	*6	15	*8	*8
487.0	12	*	*7	*	*	*	*6	*	*	*	*
487.1	24	*8	16	*	*	*	12	*	11	*6	*
490	28	13	15	*	*5	*5	14	*	*	17	*
491	539	228	311	*	16	153	369	113	149	207	69
491.2	536	226	310	*	15	152	367	113	147	206	69
491.20	15	*6	9	*	*	*	9	*	*	*7	*
491.21	521	220	300	*	15	147	358	111	145	199	66
492	24	12	12	—	*	*7	16	*7	9	*5	*
492.8	22	11	12	—	*	*6	15	*6	9	*	*
493	478	190	288	190	122	94	73	136	107	154	81
493.0	40	23	17	33	*	*	*	9	*	*5	22
493.00	20	12	*9	17	*	*	*	*	*	*	11
493.01	20	11	*8	17	*	*	*	*5	*	*	11
493.2	69	22	47	*	*6	32	30	22	12	22	12
493.20	48	16	32	*	*	20	24	15	10	15	*9
493.21	21	*6	14	*	*	12	*6	*7	*	*8	*
493.9	368	144	224	156	110	60	42	104	92	127	46
493.90	228	79	148	73	73	47	35	66	48	87	28
493.91	141	65	76	83	38	13	*7	38	44	40	19
494	*6	*	*	*	*	*	*	*	*	*	*
496	116	50	66	*	*7	28	80	25	30	39	23

See footnotes and notes at end of table.

Table 42. Number of first-listed diagnoses for discharges from short-stay hospitals, by ICD-9-CM code, sex, age, and geographic region: United States, 1999—Con.[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table: see Methods section]

ICD-9-CM code	Sex		Age				Region				
	Total	Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	North-east	Midwest	South	West
Number of first-listed diagnoses in thousands											
507	170	97	73	*8	*9	17	137	40	36	61	33
507.0	167	95	72	*8	*8	17	135	40	36	60	31
510	9	*7	*	*	*	*	*	*	*	*	*
510.9	*9	*7	*	*	*	*	*	*	*	*	*
511	57	30	27	*	*8	15	32	10	15	24	*8
511.0	*9	*	*	*	*	*	*	*	*	*	*
511.8	*8	*	*	*	*	*	*	*	*	*	*
511.9	39	20	19	*	*5	10	23	*8	*8	18	*
512	41	22	19	*	17	*9	15	*9	10	14	*7
512.1	*7	*	*	—	*	*	*	*	*	*	*
512.8	31	16	15	*	15	*7	*8	*7	*8	10	*5
513	*5	*	*	*	*	*	*	*	*	*	*
513.0	*5	*	*	*	*	*	*	*	*	*	*
514	10	*	*6	*	*	*	*	*	*	*	*
515	17	*7	11	*	*	*	12	*	*5	*6	*
516	13	*6	*7	*	*	*	*8	*	*	*6	*
516.3	*5	*	*	—	*	*	*	*	*	*	*
516.8	*6	*	*	*	*	*	*	*	*	*	*
518	230	110	121	9	24	57	140	56	38	94	42
518.3	*6	*	*	—	*	*	*	*	*	*	*
518.4	*7	*	*	*	*	*	*5	*	*	*	*
518.8	206	98	108	*8	19	53	127	50	31	88	38
518.81	166	79	88	*	16	45	102	37	25	77	28
518.82	13	*7	*6	*	*	*	*6	*	*	*	*
518.84	17	*8	*9	*	*	*	14	*	*	*6	*
518.89	*9	*	*	*	*	*	*	*	*	*	*
519	28	15	13	9	*5	*	*9	*	*5	13	*6
519.1	11	*6	*	*	*	*	*	*	*	*	*
519.8	11	*6	*5	*	*	*	*6	*	*	*6	*
520-579	3,122	1,353	1,769	221	747	830	1,324	649	717	1,157	598
522	*8	*	*	*	*5	*	*	*	*	*	*
522.5	*6	*	*	*	*	*	*	*	*	*	*
524	16	*	12	*	14	*	*	*	*	*8	*
524.0	*8	*	*	*	*7	*	—	*	*	*	*
527	10	*	*	*	*	*	*	*	*	*	*
527.2	*5	*	*	*	*	*	*	*	*	*	*
528	12	*8	*	*	*5	*	*	*	*	*	*
528.3	*6	*	*	*	*	*	*	*	*	*	*
530	220	97	123	24	45	62	89	44	52	84	40
530.1	46	19	27	*	9	16	19	*8	14	17	*6
530.10	16	*7	9	*	*	*5	*7	*	*6	*	*
530.11	23	*8	15	*	*5	*8	9	*	*8	*9	*
530.19	*6	*	*	*	*	*	*	*	*	*	*
530.2	20	11	9	*	*	*	15	*	*	*7	*
530.3	*7	*	*	*	*	*	*	*	*	*	*
530.5	*7	*	*	*	*	*	*	*	*	*	*
530.7	16	*8	*8	*	*	*	*8	*	*	*5	*
530.8	118	50	69	22	27	32	38	24	27	46	22
530.81	112	46	66	22	27	30	34	23	25	44	20

See footnotes and notes at end of table.

Table 42. Number of first-listed diagnoses for discharges from short-stay hospitals, by ICD-9-CM code, sex, age, and geographic region: United States, 1999—Con.

[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table: see Methods section]

ICD-9-CM code	Sex			Age				Region			
	Total	Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	North-east	Midwest	South	West
Number of first-listed diagnoses in thousands											
531	98	43	54	*	14	24	59	18	28	33	18
531.0	16	*9	*7	*	*	*5	*8	*	*5	*5	*
531.00	16	*9	*7	*	*	*5	*8	*	*5	*5	*
531.4	51	24	27	*	*	12	34	10	14	17	10
531.40	50	23	27	*	*	12	34	10	14	17	10
531.5	*6	*	*	*	*	*	*	*	*	*	*
531.50	*6	*	*	*	*	*	*	*	*	*	*
531.9	15	*6	10	*	*	*	*8	*	*	*6	*
531.90	14	*5	*9	*	*	*	*7	*	*	*	*
532	70	42	28	*	9	15	45	16	18	23	13
532.0	14	*9	*5	—	*	*	9	*	*	*	*
532.00	14	*9	*5	—	*	*	*9	*	*	*	*
532.4	37	23	13	*	*	*7	25	*7	9	13	*7
532.40	36	23	13	*	*	*7	25	*7	9	13	*7
532.5	*6	*	*	—	*	*	*	*	*	*	*
532.50	*6	*	*	—	*	*	*	*	*	*	*
532.9	*7	*	*	—	*	*	*	*	*	*	*
532.90	*6	*	*	—	*	*	*	*	*	*	*
533	19	*9	10	*	*	*	10	*	*	*9	*
533.9	*9	*	*	*	*	*	*	*	*	*	*
533.90	*9	*	*	*	*	*	*	*	*	*	*
534	*5	*	*	—	*	*	*	*	*	*	*
535	156	66	91	*8	32	39	78	33	39	60	24
535.0	40	17	23	*	*8	11	19	*8	10	17	*6
535.00	25	10	16	*	*6	*6	11	*	*6	12	*
535.01	14	*7	*8	*	*	*	*9	*	*	*	*
535.1	15	*6	9	—	*	*5	*8	*	*	*7	*
535.10	*7	*	*	—	*	*	*	*	*	*	*
535.11	*8	*	*	—	*	*	*6	*	*	*	*
535.3	*6	*5	*	—	*	*	*	*	*	*	*
535.4	36	15	21	*	*7	*8	20	*6	12	11	*7
535.40	20	*7	13	*	*	*	10	*	*7	*6	*
535.41	16	*8	*8	*	*	*	10	*	*5	*5	*
535.5	50	18	32	*	10	11	25	12	10	22	*7
535.50	33	12	21	*	*8	*8	12	*8	*6	13	*5
535.51	18	*6	11	*	*	*	13	*	*	*8	*
535.6	10	*6	*	*	*	*	*	*	*	*	*
535.60	*6	*	*	*	*	*	*	*	*	*	*
536	46	18	28	*	*9	*9	25	*8	*9	22	*7
536.2	10	*	*7	*	*	*	*	*	*	*6	*
536.3	*6	*	*	—	*	*	*	*	*	*	*
536.4	17	*6	11	*	*	*	13	*	*	*7	*
536.41	*6	*	*	*	*	*	*	*	*	*	*
536.42	*7	*	*	*	*	*	*6	*	*	*	*
536.8	*9	*	*	*	*	*	*	*	*	*	*
537	21	10	11	*	*	*5	12	*	*	10	*
537.0	10	*	*	*	*	*	*	*	*	*	*
537.8	9	*	*6	—	*	*	*8	*	*	*	*
537.83	*7	*	*	—	—	*	*6	*	*	*	*
540	239	138	102	56	132	35	16	44	59	66	70
540.0	53	33	20	14	23	9	*6	9	12	14	17
540.1	24	13	10	*5	11	*	*	*	*6	*7	*6
540.9	162	91	71	37	98	22	*7	31	41	44	47
541	14	*7	*7	*	*7	*	*	*	*	*7	*

See footnotes and notes at end of table.

Table 42. Number of first-listed diagnoses for discharges from short-stay hospitals, by ICD-9-CM code, sex, age, and geographic region: United States, 1999—Con.[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table: see Methods section]

ICD-9-CM code	Sex			Age				Region			
	Total	Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	North-east	Midwest	South	West
Number of first-listed diagnoses in thousands											
550	40	36	*	*6	*8	*8	19	10	9	16	*5
550.1	17	15	*	*	*	*	*8	*5	*	*6	*
550.10	13	11	*	*	*	*	*5	*	*	*	*
550.9	24	21	*	*	*	*	11	*	*6	10	*
550.90	14	13	*	*	*	*	*7	*	*	*5	*
550.92	*6	*6	*	*	*	*	*	*	*	*	*
552	62	21	41	*	*8	22	32	14	12	23	12
552.0	*6	*	*	*	*	*	*5	*	*	*	*
552.00	*5	*	*	*	*	*	*	*	*	*	*
552.1	13	*	*8	*	*	*5	*5	*	*	*	*
552.2	37	12	25	*	*	15	17	*8	*5	16	*9
552.20	9	*	*6	—	*	*	*	*	*	*	*
552.21	24	*8	16	—	*	11	11	*5	*	11	*
553	79	29	50	*	14	36	28	16	16	29	18
553.1	*5	*	*	*	*	*	*	*	*	*	*
553.2	50	18	32	*	*7	25	18	12	11	17	11
553.20	*6	*	*	*	*	*	*	*	*	*	*
553.21	43	15	28	*	*6	21	16	10	9	14	9
553.3	21	*7	14	*	*	9	*8	*	*	*8	*5
555	41	16	24	*	20	10	*7	10	14	12	*5
555.0	10	*5	*	*	*6	*	*	*	*	*	*
555.1	*8	*	*6	*	*	*	*	*	*	*	*
555.9	18	*7	11	*	10	*	*	*	*5	*8	*
556	22	10	12	*	10	*5	*5	*6	*	*8	*
556.9	14	*6	*7	*	*6	*	*	*	*	*5	*
557	41	12	30	*	*	9	28	9	10	14	*8
557.0	20	*5	15	*	*	*	13	*	*6	*6	*
557.9	17	*	12	—	*	*	12	*	*	*	*
558	168	69	99	58	34	25	51	40	33	71	24
558.9	162	66	95	58	34	24	46	38	33	69	23
560	303	127	176	16	38	63	186	66	73	105	59
560.1	64	28	36	*	10	11	40	13	17	25	*9
560.2	14	*6	*8	*	*	*	10	*	*5	*5	*
560.3	35	13	22	*	*	*	27	*8	*6	16	*
560.39	33	13	20	*	*	*	25	*8	*6	14	*
560.8	77	28	50	*	14	18	43	16	18	22	22
560.81	63	24	39	*	11	14	36	13	13	18	18
560.89	15	*	11	*	*	*	*8	*	*	*	*
560.9	109	50	60	*	11	28	65	25	27	37	21
562	239	93	146	*	23	70	145	57	55	88	39
562.1	237	92	145	—	23	70	144	57	54	88	38
562.10	29	10	19	—	*	*7	19	*6	*8	9	*6
562.11	135	50	85	—	18	55	62	36	33	46	20
562.12	58	26	32	—	*	*6	50	11	12	25	10
562.13	15	*6	9	—	*	*	12	*	*	*7	*
564	58	17	41	*8	11	9	30	13	19	16	10
564.0	36	12	23	*6	*	*	22	10	13	*8	*5
564.1	13	*	11	*	*	*	*	*	*	*5	*
564.8	*6	*	*	*	*	*	*	*	*	*	*
564.89	*6	*	*	*	*	*	*	*	*	*	*
566	17	12	*6	*	*6	*8	*	*5	*	*7	*
567	18	*8	9	*	*	*8	*	*	*	*8	*
567.2	13	*7	*6	*	*	*7	*	*	*	*	*
568	12	*	11	*	*5	*	*	*	*	*	*
568.0	9	*	*8	*	*	*	*	*	*	*	*

See footnotes and notes at end of table.

Table 42. Number of first-listed diagnoses for discharges from short-stay hospitals, by ICD-9-CM code, sex, age, and geographic region: United States, 1999—Con.

[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table: see Methods section]

ICD-9-CM code	Sex			Age				Region			
	Total	Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	North-east	Midwest	South	West
Number of first-listed diagnoses in thousands											
569	93	40	53	*	12	21	57	14	21	41	17
569.1	*8	*	*6	—	*	*	*	*	*	*	*
569.3	15	*6	9	*	*	*	*9	*	*	*7	*
569.4	9	*6	*	*	*	*	*5	*	*	*	*
569.6	11	*7	*	*	*	*	*7	*	*	*	*
569.69	*7	*5	*	*	*	*	*	*	*	*	*
569.8	42	18	24	*	*	*9	29	*5	10	21	*6
569.81	*6	*	*	—	*	*	*	*	*	*	*
569.82	*5	*	*	—	*	*	*	*	*	*	*
569.83	12	*5	*7	*	*	*	*8	*	*	*6	*
569.85	12	*	*8	*	*	*	11	*	*	*5	*
569.89	*5	*	*	*	*	*	*	*	*	*	*
570	*7	*	*	—	*	*	*	*	*	*	*
571	87	50	37	*	18	45	23	15	13	42	17
571.1	*8	*6	*	—	*	*	*	*	*	*	*
571.2	41	30	11	*	*7	26	*7	*8	*6	19	*8
571.5	29	11	18	*	*5	12	12	*	*	17	*
572	33	18	14	*	*	18	10	*8	*	13	*7
572.2	21	11	10	—	*	14	*	*	*	*8	*5
573	10	*	*7	*	*	*	*	*	*	*	*
573.3	*5	*	*	*	*	*	*	*	*	*	*
574	357	110	247	*	114	109	133	82	70	129	77
574.0	122	43	79	*	37	41	44	28	26	40	29
574.00	114	42	73	*	35	38	41	26	24	37	27
574.01	*8	*	*6	—	*	*	*	*	*	*	*
574.1	136	37	99	*	46	41	48	34	24	51	27
574.10	133	36	97	*	45	40	47	33	24	50	26
574.2	40	11	29	*	14	11	15	11	*7	17	*
574.20	39	10	29	*	13	10	14	11	*6	16	*
574.5	21	*6	15	*	*5	*	11	*	*	*8	*5
574.50	14	*	11	—	*	*	*7	*	*	*5	*
574.51	*7	*	*	*	*	*	*	*	*	*	*
574.6	*6	*	*	—	*	*	*	*	*	*	*
574.7	12	*	*8	*	*	*	*5	*	*	*6	*
574.70	*7	*	*5	*	*	*	*	*	*	*	*
574.8	*9	*	*5	*	*	*	*	*	*	*	*
575	63	19	44	*	19	21	22	14	18	23	*8
575.0	21	*9	12	*	*	*6	10	*	*5	*9	*
575.1	36	9	27	*	13	13	10	10	10	12	*
575.10	*8	*	*	*	*	*	*	*	*	*	*
575.11	22	*	18	*	*6	9	*6	*	*6	9	*
575.12	*6	*	*	—	*	*	*	*	*	*	*
576	26	12	14	*	*	*6	15	*7	*6	*8	*5
576.1	14	*7	*6	*	*	*	*7	*	*	*	*
576.2	*6	*	*	*	*	*	*	*	*	*	*
576.8	*6	*	*	*	*	*	*	*	*	*	*
577	221	103	118	*	80	78	60	40	49	93	39
577.0	191	92	99	*	70	67	52	34	43	82	32
577.1	20	*8	12	*	*8	*7	*	*	*	*7	*
577.2	*7	*	*	*	*	*	*	*	*	*	*
578	163	76	88	*	13	38	110	30	45	56	32
578.0	15	*9	*7	*	*	*	*9	*	*	*7	*
578.1	32	15	17	*	*	*7	22	*6	*7	11	*7
578.9	116	52	64	*	*8	28	80	21	34	38	22
580-629	1,719	534	1,185	81	568	428	641	330	382	693	314
581	*7	*	*	*	*	*	*	*	*	*	*
581.9	*6	*	*	*	*	*	*	*	*	*	*

See footnotes and notes at end of table.

Table 42. Number of first-listed diagnoses for discharges from short-stay hospitals, by ICD-9-CM code, sex, age, and geographic region: United States, 1999—Con.[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table: see Methods section]

ICD-9-CM code	Sex			Age				Region			
	Total	Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	North-east	Midwest	South	West
Number of first-listed diagnoses in thousands											
584	94	50	44	*	*7	23	63	21	23	34	16
584.5	11	*8	*	*	*	*	*	*	*	*	*
584.9	81	41	40	*	*5	18	57	19	21	28	13
585	29	19	10	*	*	*8	15	*8	*	11	*6
586	*6	*	*	*	*	*	*	*	*	*	*
590	125	21	104	19	58	26	22	22	30	44	30
590.1	73	10	63	*8	34	18	13	13	18	25	17
590.10	72	10	63	*8	34	18	13	13	18	25	17
590.8	47	11	37	10	23	*6	*8	*8	10	18	12
590.80	47	11	37	10	23	*6	*8	*8	10	18	12
591	12	*7	*	*	*	*	*	*	*	*	*
592	192	118	73	*	82	75	34	34	52	79	27
592.0	58	32	27	*	26	21	10	12	15	22	*9
592.1	131	85	46	*	54	54	22	22	36	56	17
593	41	18	23	*9	*9	*9	15	10	10	15	*
593.4	12	*6	*7	*	*	*	*	*	*	*5	*
593.7	*7	*	*	*6	*	*	—	*	*	*	*
593.70	*5	*	*	*	*	*	—	*	*	*	*
593.9	14	*6	*8	*	*	*	*8	*	*	*7	*
594	*8	*7	*	*	*	*	*	*	*	*	*
594.1	*7	*6	*	*	*	*	*	*	*	*	*
595	18	10	*9	*	*	*	13	*6	*5	*	*
595.8	*5	*	*	—	*	*	*	*	*	*	*
596	14	10	*	*	*	*	*8	*	*	*	*
599	372	119	254	32	38	46	256	75	77	155	65
599.0	340	103	237	31	31	41	238	70	67	143	60
599.7	23	12	11	*	*	*	15	*	*8	*8	*
600	102	102	—	—	*	16	85	21	28	31	23
601	*7	*7	—	—	*	*	*	*	*	*	*
604	*8	*8	—	*	*	*	*	*	*	*5	*
604.9	*7	*7	—	*	*	*	*	*	*	*	*
607	*8	*8	—	*	*	*	*	*	*	*	*
607.8	*5	*5	—	*	*	*	*	*	*	*	*
608	*7	*7	—	*	*	*	*	*	*	*	*
610	*6	*	*5	—	*	*	*	*	*	*	*
611	20	*	19	*	13	*6	*	*9	*	*6	*
611.1	11	*	10	*	*8	*	*	*5	*	*	*
614	55	—	55	*	46	*7	*	12	10	23	10
614.1	*5	—	*5	—	*	*	*	*	*	*	*
614.3	*7	—	*7	*	*	*	*	*	*	*	*
614.6	18	—	18	—	15	*	*	*	*	9	*
614.9	14	—	14	*	13	*	—	*	*	*6	*
616	11	—	11	*	*7	*	*	*	*	*	*
616.0	*6	—	*6	—	*	*	*	*	*	*	*
617	86	—	86	—	63	21	*	12	17	35	21
617.0	51	—	51	—	35	15	*	*8	9	19	15
617.1	12	—	12	—	9	*	*	*	*	*	*
617.3	14	—	14	—	12	*	—	*	*	*7	*
617.9	*5	—	*5	—	*	*	—	*	*	*	*

See footnotes and notes at end of table.

Table 42. Number of first-listed diagnoses for discharges from short-stay hospitals, by ICD-9-CM code, sex, age, and geographic region: United States, 1999—Con.

[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table: see Methods section]

ICD-9-CM code	Sex			Age				Region			
	Total	Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	North-east	Midwest	South	West
Number of first-listed diagnoses in thousands											
618	159	—	159	—	34	66	60	24	35	65	35
618.0	37	—	37	—	*6	17	14	*6	*9	15	*8
618.1	15	—	15	—	*8	*	*	*	*	*6	*
618.2	19	—	19	—	*6	*9	*	*	*	*6	*
618.3	20	—	20	—	*	*8	*9	*	*	*7	*
618.4	23	—	23	—	*6	10	*7	*	*7	11	*
618.5	30	—	30	—	*	12	16	*	*	14	*6
618.6	*6	—	*6	—	*	*	*	*	*	*	*
618.8	11	—	11	—	*	*	*	*	*	*	*6
620	70	—	70	*	49	13	*5	12	15	30	14
620.0	*6	—	*6	—	*	*	—	*	*	*	*
620.1	14	—	14	*	11	*	—	*	*	*6	*
620.2	41	—	41	*	30	*6	*	*8	*7	18	*9
621	15	—	15	—	*	*9	*	*	*	*6	*
621.3	*9	—	*9	—	*	*7	*	*	*	*	*
622	9	—	9	—	*6	*	*	*	*	*	*
622.1	*8	—	*8	—	*6	*	*	*	*	*	*
625	104	—	104	—	59	33	12	14	16	58	16
625.3	23	—	23	—	22	*	—	*	*	15	*
625.6	45	—	45	—	*8	26	11	*7	*6	21	10
625.9	33	—	33	—	28	*	*	*	*6	20	*
626	89	—	89	*	59	27	*	*8	24	41	15
626.2	63	—	63	*	43	18	—	*6	16	31	10
626.8	19	—	19	*	13	*	*	*	*7	*7	*
627	13	—	13	—	*	*7	*6	*	*	*7	*
627.1	13	—	13	—	*	*6	*6	*	*	*6	*
630-677	495	*	495	*	491	*	—	89	90	220	96
632	15	—	15	—	15	*	—	*	*	*6	*
633	35	—	35	*	35	*	—	*7	*6	14	*8
633.1	33	—	33	*	32	*	—	*6	*5	14	*8
634	23	—	23	*	23	—	—	*7	*	*8	*5
634.9	17	—	17	*	17	—	—	*6	*	*5	*
634.91	14	—	14	*	14	—	—	*5	*	*	*
635	*7	—	*7	—	*7	—	—	*	*	*	*
640	*6	—	*6	—	*6	—	—	*	*	*	*
640.0	*6	—	*6	—	*6	—	—	*	*	*	*
640.03	*6	—	*6	—	*6	—	—	*	*	*	*
641	18	—	18	—	18	*	—	*	*	*8	*
641.1	*7	—	*7	—	*7	—	—	*	*	*	*
641.13	*7	—	*7	—	*7	—	—	*	*	*	*
641.9	*6	—	*6	—	*6	*	—	*	*	*	*
641.93	*6	—	*6	—	*6	*	—	*	*	*	*
642	29	—	29	*	28	*	—	*	*7	11	*6
642.3	10	—	10	—	9	*	—	*	*	*	*
642.33	10	—	10	—	9	*	—	*	*	*	*
642.4	10	—	10	*	10	—	—	*	*	*	*
642.43	*8	—	*8	*	*8	—	—	*	*	*	*
643	41	—	41	*	41	*	—	*6	10	20	*
643.0	18	—	18	—	18	—	—	*	*	12	*
643.03	18	—	18	—	18	—	—	*	*	12	*
643.1	19	—	19	—	19	*	—	*	*6	*7	*
643.13	19	—	19	—	19	*	—	*	*6	*7	*
644	116	—	116	*	116	*	—	19	25	51	21
644.0	98	—	98	*	98	*	—	18	22	41	17
644.03	98	—	98	*	98	*	—	18	22	41	17
644.1	18	—	18	—	18	—	—	*	*	9	*
644.13	18	—	18	—	18	—	—	*	*	9	*

See footnotes and notes at end of table.

Table 42. Number of first-listed diagnoses for discharges from short-stay hospitals, by ICD-9-CM code, sex, age, and geographic region: United States, 1999—Con.[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table: see Methods section]

ICD-9-CM code	Sex			Age				Region			
	Total	Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	North-east	Midwest	South	West
Number of first-listed diagnoses in thousands											
646	48	—	48	*	47	—	—	*5	*6	28	*9
646.6	34	—	34	*	33	—	—	*	*	19	*7
646.63	32	—	32	*	31	—	—	*	*	18	*7
646.8	10	—	10	—	10	—	—	*	*	*7	*
646.83	10	—	10	—	10	—	—	*	*	*7	*
647	10	—	10	*	9	—	—	*	*	*5	*
648	72	*	72	*	72	*	—	14	12	32	15
648.0	*8	—	*8	—	*8	—	—	*	*	*	*
648.03	*8	—	*8	—	*8	—	—	*	*	*	*
648.4	*6	*	*6	*	*6	*	—	*	*	*	*
648.43	*5	—	*5	*	*5	—	—	*	*	*	*
648.8	*7	—	*7	—	*7	—	—	*	*	*	*
648.83	*7	—	*7	—	*7	—	—	*	*	*	*
648.9	45	—	45	—	44	*	—	*7	*	23	10
648.93	44	—	44	—	43	*	—	*7	*	22	10
654	11	—	11	—	11	*	—	*	*	*	*
654.5	*6	—	*6	—	*6	*	—	*	*	*	*
654.53	*6	—	*6	—	*6	*	—	*	*	*	*
658	*8	—	*8	—	*8	—	—	*	*	*	*
659	*6	—	*6	*	*6	—	—	*	*	*	*
674	10	—	10	—	10	—	—	*	*	*	*
680-709	518	255	263	69	133	134	182	113	119	209	77
681	18	11	*7	*	*	*7	*	*	*	*7	*
681.0	*7	*	*	*	*	*	*	*	*	*	*
681.00	*6	*	*	*	*	*	*	*	*	*	*
681.1	10	*6	*	*	*	*	*	*	*	*	*
681.10	10	*6	*	*	*	*	*	*	*	*	*
682	340	172	169	30	90	98	122	85	85	115	56
682.0	20	*9	11	*	*7	*5	*	*6	*5	*6	*
682.1	*7	*	*	*	*	*	*	*	*	*	*
682.2	31	15	16	*	*8	15	*7	*	11	12	*
682.3	37	22	15	*	15	10	10	*9	*6	12	11
682.4	20	13	*7	*	*8	*	*5	*5	*	*	*5
682.5	*7	*	*	*	*	*	*	*	*	*	*
682.6	168	84	84	*7	39	52	69	49	43	54	22
682.7	44	20	24	*6	*6	*9	23	*9	10	19	*6
683	*6	*	*	*	*	*	*	*	*	*	*
686	*6	*	*	*	*	*	*	*	*	*	*
693	*8	*	*6	*	*	*	*	*	*	*	*
693.0	*7	*	*6	*	*	*	*	*	*	*	*
695	*6	*	*	*	*	*	*	*	*	*	*
707	68	31	38	*	9	15	44	11	17	32	*8
707.0	35	16	20	*	*6	*7	22	*	*8	19	*
707.1	29	14	15	*	*	*8	18	*6	*7	11	*5
709	38	21	17	23	11	*	*	*	*	34	*
709.2	31	18	14	20	10	*	*	*	*	30	*
710-739	1,543	675	868	39	326	481	697	310	369	556	308
710	22	*	20	*	*8	*8	*	*	*5	*8	*5
710.0	16	*	14	*	*7	*6	*	*	*	*6	*
711	21	13	*8	*	*6	*	*8	*	*	*8	*5
711.0	19	13	*7	*	*	*	*7	*	*	*8	*5
711.06	*9	*6	*	*	*	*	*	*	*	*	*
714	23	*7	15	*	*	*9	11	*	*7	11	*
714.0	20	*5	14	—	*	*9	10	*	*6	10	*

See footnotes and notes at end of table.

Table 42. Number of first-listed diagnoses for discharges from short-stay hospitals, by ICD-9-CM code, sex, age, and geographic region: United States, 1999—Con.

[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table: see Methods section]

ICD-9-CM code	Sex			Age				Region			
	Total	Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	North-east	Midwest	South	West
Number of first-listed diagnoses in thousands											
715	434	169	265	*	13	118	303	95	120	130	90
715.1	39	15	24	—	*	*8	29	*6	*7	11	15
715.15	15	*7	*8	—	*	*	13	*	*	*5	*6
715.16	22	*7	15	—	*	*6	15	*	*	*	*8
715.3	202	78	124	—	*7	59	137	47	59	62	35
715.35	68	27	41	—	*	19	46	17	21	16	14
715.36	127	49	79	—	*	39	86	29	36	43	20
715.8	*6	*	*	—	*	*	*5	*	*	*	*
715.89	*6	*	*	—	*	*	*5	*	*	*	*
715.9	182	72	110	*	*	48	130	39	51	55	37
715.90	9	*	*8	—	*	*	*7	*	*	*5	*
715.91	*5	*	*	—	*	*	*	*	*	*	*
715.95	55	25	31	*	*	19	34	16	16	14	9
715.96	109	42	67	—	*	28	81	21	32	31	26
716	21	11	11	*	*	*9	*8	*	*	*6	*7
716.1	10	*7	*	—	*	*6	*	*	*	*	*
716.9	*8	*	*6	—	*	*	*6	*	*	*	*
717	13	*6	*6	*	*9	*	*	*	*	*	*
717.8	*8	*	*	*	*7	*	*	*	*	*	*
717.83	*8	*	*	*	*7	*	—	*	*	*	*
718	24	14	10	*	14	*	*	*6	*5	*	*9
718.3	11	*7	*	*	*7	*	*	*	*	*	*
718.8	*5	*	*	—	*	*	*	*	*	*	*
719	30	11	19	*	*5	*8	16	*	*6	13	*7
719.4	20	*8	12	*	*	*	11	*	*	*9	*
719.45	10	*	*6	*	*	*	*8	*	*	*6	*
721	76	34	42	*	*8	28	40	10	16	31	18
721.0	24	13	11	—	*	13	*7	*	*6	10	*
721.1	15	*7	*8	—	*	*6	*7	*	*	*7	*
721.3	28	12	17	*	*	*7	18	*	*6	10	*9
722	341	179	161	*	144	136	60	61	82	138	59
722.0	56	32	25	—	25	29	*	*9	11	28	*8
722.1	196	103	93	—	89	73	34	38	50	72	36
722.10	194	102	92	—	88	72	33	38	50	71	35
722.4	10	*	*5	—	*	*7	*	*	*	*	*
722.5	37	17	20	—	9	12	15	*5	11	14	*7
722.52	36	17	19	—	9	12	15	*	10	14	*7
722.7	17	*9	*8	—	*7	*6	*	*	*	*7	*
722.71	9	*	*	—	*	*	*	*	*	*	*
722.73	*6	*	*	—	*	*	*	*	*	*	*
722.8	*7	*	*	—	*	*	*	*	*	*	*
722.83	*7	*	*	—	*	*	*	*	*	*	*
722.9	11	*6	*5	*	*	*6	*	*	*	*6	*
722.93	*8	*	*	*	*	*	*	*	*	*	*
723	25	15	10	*	*7	10	*7	*6	*6	9	*
723.0	*8	*	*	—	*	*	*	*	*	*	*
723.1	*6	*	*	*	*	*	*	*	*	*	*
723.4	*8	*5	*	—	*	*	*	*	*	*	*
724	134	52	82	*	21	43	70	28	32	52	22
724.0	71	28	43	—	*	24	44	17	15	26	12
724.02	68	27	41	—	*	23	42	16	15	26	12
724.2	32	13	19	*	9	10	12	*7	*9	10	*6
724.4	13	*	9	—	*	*	*	*	*	*8	*
724.5	13	*6	*8	*	*	*	*7	*	*5	*	*
726	33	18	16	*	*8	16	9	*8	*6	*9	10
726.1	11	*6	*5	*	*	*5	*	*	*	*	*
726.10	11	*6	*	*	*	*5	*	*	*	*	*
726.2	*6	*	*	—	*	*	*	*	*	*	*

See footnotes and notes at end of table.

Table 42. Number of first-listed diagnoses for discharges from short-stay hospitals, by ICD-9-CM code, sex, age, and geographic region: United States, 1999—Con.[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table: see Methods section]

ICD-9-CM code	Sex		Age				Region				
	Total	Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	North-east	Midwest	South	West
Number of first-listed diagnoses in thousands											
727	26	10	15	*	*7	*9	*8	*5	*5	12	*
727.0	9	*	*	*	*	*	*	*	*	*	*
727.05	*6	*	*	*	*	*	*	*	*	*	*
728	28	12	15	*	*5	*7	14	*7	*8	9	*
728.8	16	9	*7	*	*5	*	*6	*	*5	*5	*
728.89	12	*6	*6	*	*	*	*6	*	*	*	*
729	39	14	26	*	10	12	16	*7	*8	17	*7
729.1	9	*	*6	*	*	*	*	*	*	*	*
729.5	17	*6	11	*	*	*5	*8	*	*	*6	*
729.8	10	*	*7	*	*	*	*	*	*	*	*
730	50	32	18	*	10	12	24	11	10	22	*7
730.0	13	*6	*7	*	*	*	*6	*	*	*	*
730.1	11	*7	*	*	*	*	*5	*	*	*	*
730.2	26	19	*7	*	*	*6	13	*7	*	13	*
730.27	12	*8	*	*	*	*	*8	*	*	*	*
730.28	*5	*	*	*	*	*	*	*	*	*	*
732	*9	*7	*	*6	*	*	*	*	*	*	*
733	141	49	93	*	24	35	80	29	31	50	32
733.1	69	15	54	*	*	*9	59	13	16	22	19
733.13	49	10	39	—	*	*	44	*8	11	16	14
733.14	11	*	*8	*	*	*	*7	*	*	*	*
733.4	19	11	*8	*	*5	*9	*	*	*	*7	*
733.42	16	9	*7	*	*	*8	*	*	*	*6	*
733.6	14	*	9	—	*	*7	*5	*	*	*	*
733.8	30	16	14	*	13	*9	*7	*7	*	11	*6
733.81	*6	*	*	*	*	*	*	*	*	*	*
733.82	23	13	10	*	11	*7	*	*5	*	*9	*
736	10	*7	*	*	*	*	*	*	*	*	*
737	12	*	*9	*6	*	*	*	*	*	*	*
737.3	11	*	*9	*6	*	*	*	*	*	*	*
737.30	*7	*	*6	*	*	*	*	*	*	*	*
738	22	*8	14	*	*9	*7	*6	*	*	*7	*5
738.4	18	*7	11	—	*7	*6	*6	*	*	*6	*
740-759	182	94	88	130	34	11	*8	53	25	65	39
745	25	10	15	19	*	*	*	10	*	*7	*6
745.4	*7	*	*	*6	*	*	*	*	*	*	*
745.5	*8	*	*6	*	*	*	*	*	*	*	*
746	13	*7	*6	*8	*	*	*	*	*	*	*
747	16	*7	9	*9	*	*	*	*	*	*5	*
749	19	12	*7	18	*	*	*	*	*	10	*
749.2	12	*9	*	12	*	*	*	*	*	*7	*
749.20	*7	*	*	*7	*	*	—	*	—	*7	*
750	11	*8	*	11	*	—	—	*	*	*	*
750.5	*8	*6	*	*8	—	—	—	*	*	*	*
751	13	*8	*	10	*	*	*	*	*	*	*
752	*5	*	*	*	*	*	*	*	*	*	*
753	*6	*	*	*	*	*	*	*	*	*	*
754	13	*7	*6	10	*	—	*	*	*	*	*
755	*5	*	*	*	*	*	—	*	*	*	*
756	17	10	*7	*8	*	*	*	*	*	*6	*
756.1	*6	*	*	*	*	*	*	*	*	*	*
756.12	*5	*	*	—	*	*	*	*	*	*	*

See footnotes and notes at end of table.

Table 42. Number of first-listed diagnoses for discharges from short-stay hospitals, by ICD-9-CM code, sex, age, and geographic region: United States, 1999—Con.

[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table: see Methods section]

ICD-9-CM code	Sex			Age				Region			
	Total	Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	North-east	Midwest	South	West
Number of first-listed diagnoses in thousands											
757	13	*5	*8	*8	*	—	*	*	*	13	—
757.3	12	*5	*7	*8	*	—	—	*	—	12	—
757.32	12	*5	*7	*8	*	—	—	*	—	12	—
759	*6	*	*	*	*	*	*	*	*	*	*
760-779	170	102	68	170	*	—	—	27	34	50	59
765	23	16	*8	23	—	—	—	*	*	11	*7
765.1	20	14	*6	20	—	—	—	*	*	*9	*6
765.18	*6	*	*	*6	—	—	—	*	*	*	*
769	17	10	*7	17	—	—	—	*	*	*5	*5
770	32	21	12	32	—	—	—	*5	*7	*8	13
770.6	*6	*	*	*6	—	—	—	*	*	*	*
770.8	15	*9	*6	15	—	—	—	*	*	*	*6
771	13	*8	*	13	—	—	—	*	*	*	*
771.8	11	*7	*	11	—	—	—	*	*	*	*
774	39	24	15	39	—	—	—	*	*7	11	16
774.6	34	21	13	34	—	—	—	*	*6	*9	15
775	*7	*	*	*7	—	—	—	*	*	*	*
778	10	*5	*	10	—	—	—	*	*	*	*
778.4	*9	*	*	*9	—	—	—	*	*	*	*
779	11	*6	*	11	—	—	—	*	*	*	*
779.3	*6	*	*	*6	—	—	—	*	*	*	*
780-799	313	146	168	58	110	77	69	57	50	148	57
780	60	29	31	26	16	11	*6	17	12	24	*7
780.2	11	*5	*6	*	*	*	*	*	*	*6	—
780.3	26	12	14	13	*9	*	*	*9	*	10	*
780.39	23	10	13	10	*9	*	*	*7	*	9	*
780.6	12	*7	*6	11	*	*	*	*	*	*5	*
784	*6	*	*	*	*	*	*	*	*	*	*
786	87	47	39	*5	30	35	16	16	17	40	14
786.0	*6	*	*	*	*	*	*	*	*	*	*
786.5	76	41	35	—	29	34	14	15	15	35	11
786.50	51	28	22	—	17	24	11	9	*8	28	*6
786.59	18	9	9	—	*9	*7	*	*	*	*6	*
787	*8	*	*	*	*	*	*	*	*	*	*
789	51	19	32	10	28	*9	*	*9	*8	22	12
789.0	50	19	31	10	28	*9	*	*9	*8	22	12
789.00	15	*6	*9	*	*8	*	*	*	*	*8	*
789.03	18	*7	12	*	12	*	—	*	*	*6	*6
789.06	*5	*	*	*	*	*	*	*	*	*	*
799	87	38	49	*9	24	17	36	*9	10	47	21
799.9	86	38	48	*9	24	17	36	*9	10	47	21
800-999	2,565	1,248	1,317	228	787	526	1,024	541	587	938	500
800	10	*6	*	*5	*	*	*	*	*	*	*
801	25	18	*7	*	13	*	*	*6	*6	*9	*
801.0	*8	*6	*	*	*	*	*	*	*	*	*
801.2	*8	*	*	*	*	*	*	*	*	*	*
802	32	23	*8	*	19	*5	*6	*6	*8	12	*6
802.2	10	*8	*	*	*8	*	*	*	*	*	*
802.4	*7	*6	*	*	*	*	*	*	*	*	*
803	*7	*	*	*	*	*	*	*	*	*	*
805	69	30	38	*	19	13	35	12	22	19	16
805.0	13	*8	*5	*	*6	*	*	*	*	*	*
805.2	18	*8	10	*	*5	*	11	*	*5	*6	*
805.4	34	14	20	*	*7	*8	19	*6	12	9	*7

See footnotes and notes at end of table.

Table 42. Number of first-listed diagnoses for discharges from short-stay hospitals, by ICD-9-CM code, sex, age, and geographic region: United States, 1999—Con.

[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table: see Methods section]

ICD-9-CM code	Sex			Age				Region			
	Total	Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	North-east	Midwest	South	West
Number of first-listed diagnoses in thousands											
806	*6	*	*	—	*	*	*	*	*	*	*
807	35	21	14	*	*8	11	15	*7	*9	12	*8
807.0	29	18	11	*	*6	11	12	*6	*8	9	*6
807.00	*5	*	*	—	*	*	*	*	*	*	*
808	63	21	41	*	14	*	43	10	20	22	11
808.0	13	*8	*5	*	*5	*	*6	*	*	*	*
808.2	39	10	29	*	*7	*	29	*6	12	13	*8
810	*7	*5	*	*	*	*	*	*	*	*	*
810.0	*6	*5	*	*	*	*	*	*	*	*	*
812	67	28	40	15	10	11	31	15	16	22	15
812.0	32	10	22	*	*	*5	22	*7	*8	11	*
812.00	*8	*	*	*	*	*	*6	*	*	*	*
812.01	13	*	10	—	*	*	*9	*	*	*	*
812.09	*8	*	*	*	*	*	*6	*	*	*	*
812.2	*8	*	*	*	*	*	*	*	*	*	*
812.21	*6	*	*	*	*	*	*	*	*	*	*
812.4	23	11	11	13	*	*	*	*6	*	*5	*7
812.41	15	*8	*7	11	*	*	*	*	*	*	*
813	61	29	32	*7	20	13	20	13	12	23	12
813.0	11	*	*7	*	*	*	*5	*	*	*	*
813.01	*6	*	*	*	*	*	*	*	*	*	*
813.4	30	13	17	*	*9	*8	11	*7	*6	11	*7
813.41	*7	*	*6	*	*	*	*5	*	*	*	*
813.42	15	*6	*8	*	*5	*	*	*	*	*6	*
813.44	*8	*5	*	*	*	*	*	*	*	*	*
816	*8	*	*	*	*	*	*	*	*	*	*
816.1	*6	*	*	*	*	*	*	*	*	*	*
820	323	81	243	*	9	21	292	67	75	124	58
820.0	84	19	65	*	*	*7	75	19	19	29	18
820.00	*7	*	*	—	*	*	*6	*	*	*	*
820.02	10	*	*6	*	*	*	*9	*	*	*	*
820.03	*8	*	*6	—	*	*	*7	*	*	*	*
820.09	58	10	48	*	*	*	53	13	14	21	11
820.2	158	38	120	*	*5	*8	144	36	35	59	28
820.20	*7	*	*5	—	*	*	*6	*	*	*	*
820.21	138	33	105	*	*	*6	129	31	32	50	25
820.22	13	*	9	*	*	*	10	*	*	*6	*
820.8	78	22	56	*	*	*5	70	11	20	35	11
821	56	27	30	13	12	*9	23	12	15	20	9
821.0	30	19	12	10	*8	*	10	*7	9	11	*
821.00	*9	*	*	*	*	*	*	*	*	*	*
821.01	22	14	*7	*7	*7	*	*5	*5	*7	*6	*
821.2	21	*5	16	*	*	*5	13	*5	*5	*7	*
821.20	*7	*	*	*	*	*	*	*	*	*	*
821.23	9	*	*8	*	*	*	*7	*	*	*	*
822	16	*	11	*	*	*	*7	*	*	*8	*
822.0	14	*	10	*	*	*	*7	*	*	*7	*
823	79	46	32	*6	46	14	12	18	16	26	18
823.0	24	11	13	*	13	*5	*	*	*6	*9	*6
823.00	19	*9	10	*	10	*	*	*	*	*7	*
823.2	21	15	*6	*	15	*	*	*6	*	*	*6
823.22	17	12	*	*	12*	*	*	*	*	*	*
823.3	*8	*7	*	*	*5	*	*	*	*	*	*
823.32	*6	*5	*	*	*	*	*	*	*	*	*
823.8	17	9	*8	*	*9	*	*	*5	*	*7	*
823.82	13	*7	*5	*	*6	*	*	*	*	*5	*
823.9	*6	*	*	*	*	*	*	*	*	*	*
823.92	*5	*	*	*	*	*	*	*	*	*	*

See footnotes and notes at end of table.

Table 42. Number of first-listed diagnoses for discharges from short-stay hospitals, by ICD-9-CM code, sex, age, and geographic region: United States, 1999—Con.

[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table: see Methods section]

ICD-9-CM code	Sex			Age				Region			
	Total	Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	North-east	Midwest	South	West
Number of first-listed diagnoses in thousands											
824	105	42	63	*	40	33	30	21	25	38	20
824.0	*7	*	*	*	*	*	*	*	*	*	*
824.2	*7	*	*	*	*	*	*	*	*	*	*
824.4	30	11	18	*	10	*9	11	*7	*8	*9	*6
824.6	26	*	21	*	*9	*8	*8	*	*8	*9	*6
824.8	25	15	11	*	11	*7	*5	*6	*5	9	*
825	17	13	*	*	*8	*6	*	*	*	*5	*
825.0	*8	*7	*	*	*	*	*	*	*	*	*
825.2	*6	*	*	*	*	*	*	*	*	*	*
840	24	10	14	*	*	12	*7	*	*	11	*5
840.4	22	*9	13	*	*	11	*6	*	*	11	*
844	18	10	*8	*	11	*	*	*	*	*6	*
844.2	12	*6	*6	*	9	*	—	*	*	*	*
845	*6	*	*	—	*	*	*	*	*	*	*
845.0	*6	*	*	—	*	*	*	*	*	*	*
847	15	*5	10	*	*9	*	*	*	*	*6	*
847.0	*7	*	*	*	*	*	*	*	*	*	*
847.2	*7	*	*	*	*	*	*	*	*	*	*
850	37	23	14	*6	20	*	*7	13	*8	11	*
850.0	*8	*	*	*	*	*	*	*	*	*	*
850.1	14	10	*	*	10	*	*	*6	*	*	*
850.5	*6	*	*	*	*	*	*	*	*	*	*
850.9	*8	*5	*	*	*	*	*	*	*	*	*
851	21	14	*7	*	*9	*	*8	*	*5	*6	*6
851.8	15	9	*6	*	*6	*	*6	*	*	*	*
852	39	19	20	*	12	*7	17	*5	*8	13	12
852.0	*9	*	*6	*	*	*	*	*	*	*	*
852.2	27	15	12	*	*6	*5	14	*	*5	*9	*9
852.20	11	*7	*	*	*	*	*5	*	*	*5	*
852.21	*8	*	*	*	*	*	*5	*	*	*	*
853	*9	*	*	*	*	*	*	*	*	*	*
853.0	*9	*	*	*	*	*	*	*	*	*	*
854	14	*9	*6	*	10	*	*	*	*5	*	*
854.0	13	*8	*6	*	9	*	*	*	*5	*	*
860	29	19	10	*	16	*	*8	*	9	12	*
860.0	13	*8	*	*	*6	*	*	*	*	*	*
860.4	*6	*	*	*	*	*	*	*	*	*	*
861	*8	*6	*	*	*	*	*	*	*	*	*
863	12	11	*	*	*9	*	*	*	*	*5	*
864	10	*5	*	*	*6	*	*	*	*	*	*
864.0	*8	*	*	*	*	*	*	*	*	*	*
865	13	10	*	*	*6	*	*	*	*	*	*
865.0	12	*9	*	*	*5	*	*	*	*	*	*
868	*5	*	*	*	*	*	*	*	*	*	*
873	20	11	9	*	*8	*	*5	*5	*	*6	*5
873.4	*9	*	*	*	*	*	*	*	*	*	*
879	13	10	*	*	*6	*	*	*	*	10	*
879.8	*5	*	*	*	*	*	*	—	*	*	—
881	*8	*6	*	*	*6	*	*	*	*	*	*
882	*6	*	*	*	*	*	*	*	*	*	*
883	*7	*	*	*	*	*	*	*	*	*	*
886	*5	*	*	*	*	*	*	*	*	*	*

See footnotes and notes at end of table.

Table 42. Number of first-listed diagnoses for discharges from short-stay hospitals, by ICD-9-CM code, sex, age, and geographic region: United States, 1999—Con.

[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table: see Methods section]

ICD-9-CM code	Sex		Age				Region				
	Total	Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	North-east	Midwest	South	West
Number of first-listed diagnoses in thousands											
891	18	10	*7	*	*7	*	*	*	*	*8	*
891.0	*8	*5	*	*	*	*	*	*	*	*	*
891.1	*7	*	*	*	*	*	*	*	*	*	*
892	*6	*	*	*	*	*	*	*	*	*	*
920	*8	*	*	*	*	*	*	*	*	*	*
922	21	11	10	*	*	*	12	*	*8	*6	*
922.1	12	*7	*5	—	*	*	*7	*	*	*	*
924	19	*7	12	*	*	*	13	*	*6	*	*6
924.0	*9	*	*6	*	*	*	*7	*	*	*	*
924.01	*6	*	*	—	*	*	*5	*	*	*	*
935	9	*5	*	*	*	*	*	*	*	*	*
935.1	*8	*	*	*	*	*	*	*	*	*	*
941	*5	*	*	*	*	*	*	*	*	*	*
942	*7	*	*	*	*	*	*	*	*	*	*
943	*6	*	*	*	*	*	*	*	*	*	*
944	*7	*6	*	*	*	*	—	*	*	*	*
945	12	*8	*	*	*	*	*	*	*	*	*
945.2	*5	*	*	*	*	*	*	*	*	*	*
945.3	*6	*	*	*	*	*	*	*	*	*	*
948	22	15	*7	19	*	*	*	*	—	21	—
948.0	*6	*	*	*6	—	*	*	*	—	*5	—
948.00	*6	*	*	*6	—	*	*	*	—	*5	—
952	*6	*	*	*	*	*	*	*	*	*	*
958	*9	*6	*	*	*	*	*	*	*	*	*
959	37	18	19	*6	14	*	12	*7	9	11	10
959.0	26	13	13	*	10	*	*8	*	*8	*6	*8
959.01	25	13	12	*	10	*	*8	*	*7	*6	*8
959.1	*5	*	*	*	*	*	*	*	*	*	*
962	*7	*5	*	*	*	*	*	*	*	*	*
962.3	*6	*5	*	*	*	*	*	*	*	*	*
963	*5	*	*	*	*	*	—	*	*	*	*
963.0	*5	*	*	*	*	*	—	*	*	*	*
965	38	11	26	*	23	*6	*	*7	10	13	*7
965.0	11	*	*6	*	*6	*	*	*	*	*5	*
965.09	*6	*	*	*	*	*	*	*	*	*	*
965.4	18	*	14	*	12	*	*	*	*6	*6	*
966	*9	*	*7	*	*	*	*	*	*	*	*
966.3	*6	*	*	*	*	*	*	*	*	*	*
968	9	*7	*	—	*7	*	—	*	*	*	*
968.5	*5	*	*	—	*	*	—	*	*	*	*
969	53	22	30	*	33	14	*	*9	10	21	13
969.0	19	10	9	*	13	*	*	*	*	*7	*
969.4	23	*8	15	*	13	*8	*	*	*	10	*6
972	*9	*	*	*	*	*	*	*	*	*	*
977	10	*5	*5	*	*7	*	*	*	*	*6	*
977.9	*7	*	*	*	*	*	*	*	*	*	*
989	*7	*	*	*	*	*	*	*	*	*	*
995	26	11	15	*	*7	*8	*8	*	*6	11	*
995.0	*5	*	*	*	*	*	*	*	*	*	*
995.1	*8	*	*	*	*	*	*	*	*	*	*
995.2	*8	*	*	*	*	*	*	*	*	*	*

See footnotes and notes at end of table.

Table 42. Number of first-listed diagnoses for discharges from short-stay hospitals, by ICD-9-CM code, sex, age, and geographic region: United States, 1999—Con.

[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table: see Methods section]

ICD-9-CM code	Sex			Age				Region			
	Total	Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	North-east	Midwest	South	West
Number of first-listed diagnoses in thousands											
996	412	201	211	26	79	117	190	97	86	150	79
996.0	24	13	11	*	*	*6	16	*6	*5	*8	*
996.01	15	*7	*8	*	*	*	10	*	*	*	*
996.04	*6	*	*	—	*	*	*	*	*	*	*
996.1	12	*6	*6	*	*	*	*	*	*	*	*
996.2	26	13	13	10	12	*	*	*	10	*7	*
996.3	*5	*	*	*	*	*	*	*	*	*	*
996.4	77	32	45	*	*9	18	50	17	20	22	17
996.5	13	*6	*8	*	*	*	*6	*	*	*	*
996.59	*7	*	*	*	*	*	*	*	*	*	*
996.6	113	58	55	*8	25	36	44	29	22	46	16
996.61	*6	*	*	—	*	*	*	*	*	*	*
996.62	59	29	30	*6	12	21	20	16	10	25	*9
996.63	*6	*	*	*	*	*	*	*	*	*	*
996.64	*8	*	*	—	*	*	*	*	*	*	*
996.66	9	*	*	—	*	*	*6	*	*	*	*
996.67	9	*	*	*	*	*	*	*	*	*	*
996.68	*5	*	*	*	*	*	*	*	*	*	*
996.69	*7	*	*	*	*	*	*	*	*	*	*
996.7	120	61	59	*	18	37	64	27	18	48	28
996.72	*7	*	*	*	*	*	*	*	*	*	*
996.73	49	25	24	*	*8	15	24	*7	*6	20	14
996.74	31	17	13	*	*	*8	20	12	*5	11	*
996.77	*9	*	*	—	*	*	*6	*	*	*	*
996.78	13	*7	*6	*	*	*	*	*	*	*7	*
996.8	19	*9	10	*	*8	9	*	*5	*	*8	*
996.81	12	*6	*6	*	*6	*5	*	*	*	*6	*
997	103	54	50	*	20	35	45	24	20	37	21
997.0	*6	*	*	*	*	*	*	*	*	*	*
997.1	11	*6	*	*	*	*	*8	*	*	*	*
997.2	*8	*	*	—	*	*	*	*	*	*	*
997.3	10	*7	*	*	*	*	*	*	*	*	*
997.4	37	14	22	*	12	11	12	*6	*7	13	10
997.5	13	*9	*	*	*	*	*8	*	*	*	*
997.6	16	9	*7	*	*	*8	*7	*	*	*7	*
997.62	10	*6	*	*	*	*	*	*	*	*	*
997.69	*5	*	*	—	*	*	*	*	*	*	*
998	200	96	104	12	53	62	72	48	41	71	41
998.1	53	27	26	*5	16	10	21	14	12	16	11
998.11	34	19	15	*5	*9	*5	15	11	*8	*7	*8
998.12	15	*6	*9	*	*6	*	*5	*	*	*6	*
998.2	9	*	*7	*	*	*	*	*	*	*	*
998.3	10	*	*6	*	*	*	*	*	*	*	*
998.5	106	52	54	*	27	40	36	24	21	41	20
998.59	103	51	53	*	26	39	35	24	19	40	20
998.6	*5	*	*	*	*	*	*	*	*	*	*
998.8	16	*7	*8	*	*	*	*6	*	*	*6	*
998.89	11	*	*6	*	*	*	*	*	*	*	*
999	*8	*	*	*	*	*	*	*	*	*	*
999.3	*5	*	*	*	*	*	*	*	*	*	*
V01-V82	4,676	386	4,290	77	3,939	207	453	809	1,016	1,786	1,065
V10	10	*	*6	*	*	*	*6	*	*	*	*
V12	15	*6	9	*	*	*6	*7	*	*	*8	*
V12.5	9	*	*5	—	*	*	*5	*	*	*6	*
V12.59	*5	*	*	—	*	*	*	*	*	*	*
V15	*9	*6	*	*	*	*	*	*	*	*	*
V15.8	*7	*	*	*	*	*	*	*	*	*	*

See footnotes and notes at end of table.

Table 42. Number of first-listed diagnoses for discharges from short-stay hospitals, by ICD-9-CM code, sex, age, and geographic region: United States, 1999—Con.[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table: see Methods section]

ICD-9-CM code	Sex			Age				Region			
	Total	Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	North-east	Midwest	South	West
Number of first-listed diagnoses in thousands											
V17	*5	*	*	—	*	*	*	*	*	*	*
V22	*8	—	*8	—	*8	—	—	*	*	*	*
V27	3,810	—	3,810	*7	3,797	*6	—	636	809	1,454	912
V27.0	3,583	—	3,583	*7	3,572	*	—	601	694	1,403	884
V27.1	18	—	18	—	18	—	—	*	*	*8	*
V27.2	38	—	38	—	38	*	—	*8	*7	15	*9
V27.9	169	—	169	*	167	*	—	23	104	27	15
V30	12	9	*	12	—	—	—	*	*	*	*
V30.1	12	9	*	12	—	—	—	*	*	*	*
V42	*6	*	*	*	*	*	*	*	*	*	*
V43	*6	*	*	*	*	*	*	*	*	*	*
V45	18	12	*6	*	*	*6	*8	*5	*	*6	*
V45.8	13	10	*	*	*	*	*6	*	*	*	*
V45.89	*7	*	*	*	*	*	*	*	*	*	*
V53	11	*7	*	*	*	*	*8	*	*	*	*
V53.3	9	*6	*	*	*	*	*8	*	*	*	*
V53.31	*7	*	*	*	*	*	*6	*	*	*	*
V54	10	*	*6	*	*	*	*	*	*	*	*
V55	37	22	15	*	*8	13	13	10	*6	13	*8
V55.2	*6	*	*	*	*	*	*	*	*	*	*
V55.3	26	17	9	*	*5	11	*9	*6	*	10	*5
V57	427	168	259	*	30	79	315	65	125	178	59
V57.1	100	38	62	*	*5	22	72	26	12	53	10
V57.8	313	125	188	*	24	57	231	34	112	122	45
V57.89	313	125	188	*	24	57	230	34	112	122	45
V57.9	12	*	*8	—	*	*	11	*	*	*	*
V58	198	106	92	36	42	55	65	55	38	67	38
V58.0	16	*8	*8	*	*	*6	*7	*	*	11	*
V58.1	148	84	63	33	31	41	42	46	29	45	28
V58.4	21	10	11	*	*7	*	*8	*6	*	*6	*
V58.49	19	*9	11	*	*6	*	*7	*6	*	*6	*
V58.8	11	*	*7	*	*	*	*8	*	*	*	*
V58.89	10	*	*7	*	*	*	*8	*	*	*	*
V59	*6	*	*	*	*	*	*	*	*	*	*
V66	*6	*	*	—	*	*	*	*	*	*	*
V70	12	*7	*5	*	*7	*	*	*	*	*	*
V70.7	10	*6	*	*	*6	*	*	*	*	*	*
V71	17	*7	11	*	*6	*	*	*	*	*	*7
V71.7	*7	*	*5	*	*	*	*	*	*	*	*

* Figure does not meet standard of reliability or precision.

— Quantity zero.

NOTE: Estimates of 5,000-9,000 are to be used with caution; see Methods section.

Table 43. Number of days of care for discharges from short-stay hospitals, by ICD-9-CM code of first-listed diagnosis, sex, age, and geographic region: United States, 1999

[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table: see Methods section]

ICD-9-CM code	Sex			Age				Region			
	Total	Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	North-east	Midwest	South	West
Number of days of care in thousands											
All codes	160,128	68,719	91,409	11,026	37,236	34,595	77,270	39,814	33,988	59,635	26,692
001-139	5,297	2,523	2,774	522	1,072	1,016	2,687	1,583	896	2,004	814
003	*33	*	*	*	*	*	*	*	*	*	*
008	422	148	274	119	64	61	179	116	88	140	78
008.4	171	*42	129	*	*	*37	109	*51	*45	*44	*
008.45	153	*33	120	*	*	*34	96	*41	*42	*39	*
008.6	68	36	32	63	*	*	*	*	19	26	*14
008.61	65	36	29	62	*	*	-	*	19	24	*13
008.8	176	66	109	45	40	*22	69	56	23	65	31
009	43	*24	*19	*	*	*	*17	*	*	*18	*
009.0	31	*17	*14	*	*	*	*	*	*	*	*
011	*72	*56	*	*	*	*	*	*	*	*	*
011.9	*49	*	*	*	*	*	*	*	*	*	*
034	24	*10	*14	*10	*	*	*	*	*11	*	*
034.0	20	*9	*11	*9	*	*	*	*	*	*	*
038	2,880	1,289	1,591	80	201	483	2,116	821	512	1,135	411
038.0	235	84	151	*	*	*	173	*75	*56	80	*
038.1	571	342	229	*	*44	132	385	186	114	194	*77
038.10	118	*75	*	*	*	*	*86	*	*	*	*
038.11	354	231	123	*	*	*100	219	*103	*65	122	*64
038.19	99	*	*63	*	*	*	*80	*	*	*	*
038.2	88	*47	*41	*	*	*	*49	*	*	*	*
038.4	721	280	441	*	*44	161	508	225	131	238	128
038.40	*81	*	*62	*	*	*	*	*	*	*	*
038.42	388	138	250	*	*	*67	294	117	89	141	*41
038.49	173	66	107	*	*	*55	100	*	*	*62	*41
038.8	91	*	*73	*	*	*	*72	*	*	*	*
038.9	1,140	515	625	40	75	130	896	249	170	571	150
041	60	*28	*32	*	*	*	*	*	*	*	*
041.1	*37	*	*	*	*	*	*	*	-	*	*
042	602	444	158	*	378	205	*	251	*63	170	118
047	103	56	47	43	45	*	*	*	*	48	*26
047.9	93	54	40	40	40	*	*	*	*	46	*25
052	*23	*	*	*15	*	-	*	*	*	*	*
053	68	*	55	*	*	*	*44	*	*	*	*
053.9	*41	*	*33	*	*	*	*	*	*	*	*
054	65	*	*52	*36	*	*	*	*	*	*22	*
070	90	*45	*46	*	*37	*43	*	*	*	*36	*
070.5	*29	*	*	-	*	*	*	*	*	*	*
075	*21	*	*	*	*14	*	-	*	*	*	*
079	166	71	95	84	*21	*18	43	36	32	69	30
079.9	156	66	90	77	*20	*16	42	35	30	65	26
079.99	156	66	90	77	*20	*16	42	35	30	65	26
112	114	*61	*52	*	*	*	*73	*55	*	*30	*
117	*92	*	*	*	*	*	*	*	*	*	*
135	*38	*	*	-	*	*	*	*	*	*	*
136	94	*49	*	*	*58	*	*	*	*	*	*
136.3	*68	*	*	*	*	*	*	*	*	*	*
140-239	10,502	4,569	5,933	265	1,218	3,264	5,755	2,769	2,178	3,613	1,942
150	153	104	*	-	*	*72	*81	*	*	*56	*
150.5	*48	*	*	-	*	*	*	*	*	*	*

See footnotes and notes at end of table.

Table 43. Number of days of care for discharges from short-stay hospitals, by ICD-9-CM code of first-listed diagnosis, sex, age, and geographic region: United States, 1999—Con.[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table: see Methods section]

ICD-9-CM code	Sex		Age				Region				
	Total	Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	North-east	Midwest	South	West
Number of days of care in thousands											
151	330	220	110	—	*	*99	221	*46	*82	*86	*116
151.0	*119	*104	*	—	*	*	*94	*	*	*	*
153	1,099	501	597	*	*47	221	830	265	325	333	175
153.1	113	*67	*	—	*	*	91	*	*	*	*
153.2	90	*	*63	—	*	*	*53	*	*	*	*
153.3	275	126	149	—	*	*52	205	*98	*83	*61	*34
153.4	196	*79	118	—	*	*	155	*53	*	*59	*
153.6	152	*71	81	—	*	*	120	*	*71	*	*
153.9	133	*57	*75	*	*	*	106	*	*	*65	*
154	383	200	184	*	*	101	266	113	72	128	*71
154.0	154	79	*75	—	*	*	119	*50	*	*60	*
154.1	201	113	87	—	*	*57	132	*52	*50	*56	*
155	109	78	*31	*	*	*37	*61	*	*	*28	*
155.0	74	*50	*	*	*	*	*	*	*	*	*
156	107	*	*71	—	*	*	85	*	*	*	*
156.1	*48	*	*	—	—	*	*46	*	*	*	*
157	234	91	143	—	*	*60	160	*68	*50	68	*49
157.0	115	*51	*64	—	*	*32	77	*	*	*37	*
157.9	80	*	*55	—	*	*	*63	*	*	*	*
161	*85	*59	*	—	*	*	*	*	*	*	*
162	1,100	594	506	*	*	345	737	260	214	401	226
162.2	*52	*	*	—	*	*	*32	*	*	*	*
162.3	360	189	170	—	*	102	254	108	*62	116	74
162.4	*74	*	*	—	—	*	*	*	*	*	*
162.5	157	94	*62	*	*	*47	108	*	*	*46	*
162.8	133	63	*70	—	*	*44	87	*78	*30	*	*
162.9	322	164	158	*	*	99	214	*	59	166	*70
170	*29	*	*	*	*	*	*	*	*	*	*
171	*42	*	*	*	*	*	*	*	*	*	*
173	*28	*	*	*	*	*	*	*	*	*	*
174	289	—	289	—	34	111	144	84	58	98	48
174.2	*14	—	*14	—	*	*	*	*	*	*	*
174.4	52	—	52	—	*	*19	28	*21	*	*12	*
174.8	83	—	83	—	*	37	34	32	*16	*22	*
174.9	108	—	108	—	*	38	60	*19	*19	52	*
180	70	—	70	*	*23	*33	*	*	*	*24	*
180.8	*25	—	*25	—	*	*	*	*	*	*	*
180.9	*24	—	*24	*	*	*	*	*	*	*	*
182	125	—	125	—	*	42	75	*34	*29	38	*25
182.0	124	—	124	—	*	42	74	*34	*29	38	*
183	178	—	178	*	*	*79	*84	*	*	*65	*
183.0	177	—	177	*	*	*79	*84	*	*	*65	*
184	*24	—	*24	*	*	*	*	*	*	*	*
185	349	349	—	—	*	125	220	72	70	142	65
188	205	133	71	—	*	*31	171	68	50	65	*
188.2	*29	*24	*	—	*	*	*26	*	*	*	*
188.8	50	*30	*	—	—	*	*45	*	*	*	*
188.9	71	*33	*38	—	*	*	60	*	*	*27	*
189	218	123	95	*	*	*51	148	*47	*52	*64	*56
189.0	176	100	76	*	*	*46	111	*41	*40	*54	*41
191	189	123	67	*	*35	*58	*68	*57	*48	*57	*27
191.1	*36	*	*	*	*	*	*	*	*	*	*
191.2	*54	*	*	*	*	*	*	*	*	*	*
193	52	*	33	*	*19	*12	*	*	*	*19	*

See footnotes and notes at end of table.

Table 43. Number of days of care for discharges from short-stay hospitals, by ICD-9-CM code of first-listed diagnosis, sex, age, and geographic region: United States, 1999—Con.

[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table: see Methods section]

ICD-9-CM code	Sex			Age				Region			
	Total	Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	North-east	Midwest	South	West
Number of days of care in thousands											
195	*45	*	*	*	*	*	*	*	*	*	*
196	56	42	*	—	*	*	*17	*	*	*	*
196.0	*27	*	*	—	*	*	*	*	*	*	*
197	811	274	537	*	*53	286	465	200	181	292	138
197.0	101	*39	62	*	*	*37	*50	*	*	*	*
197.2	249	*61	188	*	*	110	133	*60	*	87	*
197.4	*46	*	*	—	*	*	*	*	*	*	*
197.6	123	*47	*76	—	*	*	77	*	*	*	*
197.7	188	92	96	*	*	*60	114	*45	*36	89	*
198	735	336	400	*	*	279	424	269	164	213	90
198.3	240	124	116	*	*	105	126	104	*37	71	*
198.5	291	145	146	*	*	111	168	105	100	*43	*43
198.8	161	*	*111	—	*	*	*105	*	*	*	*
198.89	156	*	*106	—	*	*	*101	*	*	*	*
199	232	93	139	*	*	73	128	*57	*39	57	*79
199.0	131	*47	85	*	*	*48	*64	*	*	*35	*
199.1	101	*47	*54	—	*	*	64	*	*	*	*
200	82	*	*57	*	*	*	*56	*	*	*	*
200.0	*40	*	*	—	*	*	*	*	*	*	*
202	376	161	215	*	*	188	151	*168	*44	96	*68
202.8	333	146	187	*	*	172	140	*164	*40	77	*51
202.80	118	77	*41	*	*	*31	73	*	*	*32	*
202.83	*41	*	*	—	*	*	*	*	*	*	*
203	155	88	67	—	*	*	109	*	*54	*47	*
203.0	150	87	*63	—	*	*	104	*	*54	*	*
203.00	148	86	*62	—	*	*	103	*	*53	*	*
204	160	*82	*78	*82	*	*	*36	*55	*	*61	*
204.0	121	*	*62	*82	*	*	*	*	*	*	*
204.00	*85	*	*	*54	*	*	*	*	*	*	*
204.1	*39	*	*	—	*	*	*34	*	*	*	*
204.10	*39	*	*	—	*	*	*33	*	*	*	*
205	378	209	170	*	*	*	207	*83	*	*185	*
205.0	330	*182	*148	*	*	*	199	*	*	*176	*
205.00	324	*178	*146	*	*	*	197	*	*	*175	*
210	*11	*	*	*	*	*	*	*	*	*	*
210.2	*9	*	*	—	*	*	*	*	*	*	*
211	188	90	98	*	*	46	127	*45	*41	61	*41
211.3	142	71	71	*	*	*32	104	*	*	42	*34
218	653	—	653	—	377	264	*	109	123	249	171
218.0	80	—	80	—	40	40	*	*22	*15	28	*
218.1	180	—	180	—	95	81	*	39	32	74	34
218.2	62	—	62	—	33	*24	*	*	*	23	*15
218.9	331	—	331	—	208	118	*	35	65	123	108
220	110	—	110	*	39	47	*19	*26	*23	34	27
225	173	*48	125	*	*35	*57	*81	*78	*	*49	*
225.2	136	*	103	—	*	*44	*76	*	*	*41	*
226	*10	*	*	*	*	*	*	*	*	*	*
227	47	*23	*24	*	*	*22	*	*	*	*	*
227.1	*13	*	*	—	*	*	*	*	*	*	*
227.3	*28	*	*	*	*	*	*	*	*	*	*
228	*25	*	*16	*11	*	*	*	*	*	*14	*
228.0	*22	*	*15	*8	*	*	*	*	*	*13	*

See footnotes and notes at end of table.

Table 43. Number of days of care for discharges from short-stay hospitals, by ICD-9-CM code of first-listed diagnosis, sex, age, and geographic region: United States, 1999—Con.

[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table: see Methods section]

ICD-9-CM code	Sex		Age				Region				
	Total	Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	North-east	Midwest	South	West
Number of days of care in thousands											
233	67	*	66	—	34	22	*11	*13	*11	30	*13
233.0	21	—	21	—	*	*	*9	*	*	*9	*
233.1	44	—	44	—	32	*	*	*	*	*20	*
235	*67	*	*	*	*	*	*33	*	*	*	*
235.2	*56	*	*	—	*	*	*	*	*	*	*
237	*39	*	*	*	*	*	*	*	*	*	*
238	61	*22	*39	*	*	*	39	*	*	*27	*
238.7	*46	*	*29	*	*	*	*33	*	*	*	*
239	60	*	51	*	*	*24	*28	*	*	36	*
239.6	*19	*	*	*	*	*	*	*	*	*	*
240-279	6,591	2,742	3,849	463	997	1,706	3,425	1,772	1,416	2,264	1,140
241	44	*	42	*	*	*16	*	*	*	*	*
241.1	*28	*	*27	*	*	*	*	*	*	*	*
242	33	*	*20	*	*11	*	*	*	*	*17	*
242.9	*19	*	*	*	*	*	*	*	*	*	*
242.90	*18	*	*	*	*	*	*	*	*	*	*
244	96	*	83	*	*	*	*32	*	*	*	*
244.9	90	*	79	*	*	*	*27	*	*	*	*
250	2,938	1,439	1,499	58	576	1,052	1,253	766	532	1,055	584
250.0	569	229	340	24	120	211	213	116	76	270	106
250.00	188	81	107	*	*25	88	74	37	24	108	*19
250.01	128	51	77	*	*55	*33	*30	*18	*	56	*
250.02	158	69	89	*	24	61	70	35	*23	74	27
250.03	95	27	67	*	*17	*30	*39	*26	*19	32	*
250.1	394	201	193	29	204	119	*42	108	94	142	51
250.10	66	*37	*30	*	*31	*	*	*	*	*28	*
250.11	106	52	54	*16	62	*23	*	*27	29	*27	*23
250.12	59	*26	*32	*	*19	*25	*	*	*	*24	*
250.13	163	86	77	*	92	*47	*	*48	35	62	*18
250.2	70	*39	*30	*	*	*25	*35	*	*	*	*
250.22	*37	*	*	*	*	*	*	*	*	*	*
250.3	*44	*	*	*	*	*	*	*	*	*	*
250.4	226	102	124	—	*	85	108	*35	*	71	*85
250.40	120	*53	*67	—	*	*45	*69	*	*	*33	*
250.41	64	*	*41	—	*	*	*	*	*	*	*
250.42	*31	*	*	—	*	*	*	*	*	*	*
250.5	*28	*	*	—	*	*	*	*	*	*	*
250.6	310	113	197	*	55	142	112	67	55	100	89
250.60	162	53	109	—	*20	69	73	*33	*	*51	*60
250.61	67	*30	*37	*	*	*37	*	*	*	*	*
250.62	*39	*	*	—	*	*	*	*	*	*	*
250.63	*42	*	*	—	*	*	*	*	*	*	*
250.7	677	387	290	—	*	256	380	220	117	232	108
250.70	318	165	152	—	*	*97	211	*83	*54	118	*
250.71	195	*110	*85	—	*	*94	87	*	*	*63	*
250.72	*91	*73	*	—	*	*	*52	*	*	*	*
250.73	*73	*	*	—	*	*	*	*	*	*	*
250.8	573	315	258	*	56	191	323	156	128	181	106
250.80	281	176	104	—	*	87	173	65	52	83	*81
250.81	116	57	60	*	*	39	55	*26	*40	*38	*
250.82	83	*41	43	—	*	*36	45	*	*	*39	*
250.83	93	*41	*51	*	*	*	*	*	*	*	*
250.9	47	*30	*	*	*22	*	*	*	*	*25	*
251	46	*21	*26	*	*	*	*	*	*	*17	*
251.2	*21	*	*	*	*	*	*	*	*	*	*
252	*21	*	*	—	*	*	*	*	*	*	*

See footnotes and notes at end of table.

Table 43. Number of days of care for discharges from short-stay hospitals, by ICD-9-CM code of first-listed diagnosis, sex, age, and geographic region: United States, 1999—Con.

[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table: see Methods section]

ICD-9-CM code	Sex		Age				Region				
	Total	Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	North-east	Midwest	South	West
Number of days of care in thousands											
253	80	*46	*34	*	*	*	*58	*	*	*	*
253.6	61	*	*25	*	*	*	*56	*	*	*	*
255	*26	*	*	*	*	*	*	*	*	*	*
261	*27	*	*	*	*	*	*	*	*	*	*
263	141	*69	72	*	*	*	118	*	*	*57	*
263.9	135	*66	69	*	*	*	115	*	*	*55	*
272	60	22	38	*	*14	25	*21	25	*	16	*
272.0	26	*11	*16	—	*	*13	*	*15	*	*	*
272.4	26	*9	*	—	*	*	*	*	*	*	*
274	43	*23	*20	—	*	*	*33	*	*	*	*
274.0	*25	*	*	—	*	*	*	*	*	*	*
275	95	*40	55	*	*	*	62	*	*	*29	*
275.4	87	*37	*50	*	*	*	59	*	*	*26	*
275.42	66	*27	*39	*	*	*	*47	*	*	*22	*
276	2,513	850	1,663	283	189	384	1,657	631	636	888	358
276.0	*59	*	*	*	*	*	*47	*	*	*	*
276.1	282	81	201	*	*	60	201	58	54	102	68
276.2	*28	*	*	*	*	*	*	*	*	*	*
276.5	1,886	646	1,240	265	144	233	1,243	501	516	634	235
276.7	87	*40	47	*	*	*	65	*	*	42	*
276.8	130	29	101	*	*13	51	65	*22	*26	58	*
277	147	*61	86	*54	*70	*	*	*65	*	*	*
277.0	120	*53	*67	*52	*67	*	—	*	*	*	*
277.00	118	*53	*65	*50	*67	*	—	*	*	*	*
278	134	*40	94	*	59	69	*	*47	*27	33	*26
278.0	125	*39	86	*	56	63	*	*46	*25	*31	*23
278.00	*10	*	*	*	*	*	*	*	*	*	*
278.01	115	*36	79	—	53	58	*	*44	*23	*28	*20
280-289	1,773	749	1,024	243	487	362	680	430	362	684	296
280	207	72	135	*	*22	*29	153	58	48	78	*23
280.0	119	*43	76	—	*	*	91	*36	*	*47	*
280.9	87	*29	58	*	*	*13	60	*	*26	30	*
281	*25	*	*	*	*	*	*	*	*	*	*
282	368	124	244	61	278	*	*	97	51	198	*
282.6	347	115	232	55	265	*	*	90	48	191	*
282.62	333	108	225	48	260	*	*	87	44	185	*
283	*37	*	*	*	*	*	*	*	*	*	*
284	219	112	107	*50	*	*71	69	*51	*52	*52	*63
284.8	197	103	94	*	*	*69	64	*46	*50	*40	*62
285	275	106	169	*	*23	66	174	54	72	119	30
285.1	57	*19	*37	—	*	*	*43	*	*	*	*
285.9	199	83	116	*	*17	50	121	41	44	97	*17
286	94	*44	51	*	*	*	44	*	*	*30	*
286.9	*27	*	*	*	*	*	*	*	*	*	*
287	169	85	84	*13	*49	44	62	36	*16	58	59
287.3	70	*33	*37	*	*	*	*	*	*	*	*
287.4	*25	*	*	*	*	*	*	*	*	*	*
287.5	47	*20	*27	*	*	*	*	*	*	*	*
288	279	136	143	59	*30	65	125	69	73	93	45
288.0	243	113	130	49	*25	61	108	61	62	78	*41
288.8	*25	*	*	*	*	*	*	*	*	*	*

See footnotes and notes at end of table.

Table 43. Number of days of care for discharges from short-stay hospitals, by ICD-9-CM code of first-listed diagnosis, sex, age, and geographic region: United States, 1999—Con.[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table: see Methods section]

ICD-9-CM code	Sex			Age				Region			
	Total	Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	North-east	Midwest	South	West
Number of days of care in thousands											
289	100	44	56	*24	*37	*	*	*	*	39	*24
289.2	*23	*	*	*	*	*	*	*	*	*	*
289.8	*32	*	*	*	*	*	*	*	*	*	*
290-319	15,090	7,835	7,254	1,198	7,764	3,512	2,615	5,625	3,099	4,874	1,492
290	281	111	170	*	—	*	255	*82	90	86	*
290.0	*36	*	*	—	—	—	*36	*	*	*	—
290.2	*81	*	*63	*	—	—	*80	*	*	*	*
290.4	136	*70	*66	—	—	*	114	*	*	*65	*
290.40	*65	*	*	—	—	*	*	*	*	*	*
291	553	413	139	—	250	247	56	131	238	106	78
291.0	68	56	*	—	*31	*31	*	*31	*	*	*
291.2	*39	*	*	—	*	*	*	*	*	*	*
291.8	413	300	113	—	207	175	*32	96	190	67	60
291.81	386	273	112	—	191	164	*31	90	176	67	54
292	333	182	151	*	209	68	*56	166	58	77	*33
292.0	168	92	76	*	113	*45	*	96	*34	*28	*
292.8	140	77	63	—	75	*	*45	*67	*	*36	*
292.81	56	*	*40	—	*	*	*45	*	*	*	*
292.84	*71	*56	*	—	*62	*	—	*	*	*	*
293	265	163	102	*	66	*58	141	109	81	*51	*
293.0	137	*80	56	*	*	*	117	*	*55	*	*
293.8	116	76	*40	—	63	*	*	82	*	*	*
293.83	101	68	*	—	*55	*	*	78	*	*	*
294	248	81	168	*	*	*	163	97	*41	*55	*
294.8	172	*51	121	*	*	*	135	*83	*33	*43	*
295	3,152	1,858	1,294	*	1,842	932	372	1,363	471	903	415
295.3	1,081	616	465	*	631	289	161	505	129	327	121
295.30	266	125	141	—	198	*53	*	*	*60	*81	*
295.32	227	154	*73	—	*101	*	*	*120	*	*58	*
295.34	573	333	239	*	328	175	*69	273	*59	180	*61
295.6	338	236	*102	*	227	*86	*	178	*	*81	*
295.60	*65	*	*	*	*	*	*	*	*	*	*
295.62	*85	*61	*	—	*	*	*	*	*	*	—
295.64	183	*148	*	—	*121	*	*	*	*	*	*
295.7	1,335	692	643	*	784	382	166	546	206	390	193
295.70	849	411	437	*	456	278	*112	271	174	250	*155
295.74	456	268	188	—	307	95	*	257	*	132	*
295.9	328	257	*71	*	155	*152	*	*97	*72	*89	*
295.90	230	188	*42	*	*83	*132	*	*	*69	*65	*
295.94	*70	*47	*	—	*59	*	*	*57	*	*	*
296	5,445	2,204	3,241	396	2,755	1,313	981	1,846	1,147	2,068	383
296.2	1,439	558	881	181	737	220	301	360	300	685	94
296.20	665	249	416	*65	347	98	154	167	163	273	*62
296.22	157	68	89	*	84	*33	*	*	*	135	*
296.23	311	110	201	*38	165	*37	*71	*53	65	182	*
296.24	302	130	173	*	140	*52	*56	134	61	96	*
296.3	1,909	686	1,223	*114	902	457	437	731	388	669	121
296.30	510	169	341	*	228	120	111	205	116	169	*
296.32	88	*25	63	*	41	*	*	*	*20	*45	*
296.33	745	302	443	*	382	146	170	289	153	273	*30
296.34	558	184	374	*	244	174	*128	213	99	179	*67
296.4	540	246	294	*	258	210	*67	215	122	164	*39
296.40	207	80	127	*	100	*61	*	*56	*60	79	*
296.43	*67	*	*	—	*	*	*	*	*	*	*
296.44	249	126	123	*	117	*109	*	135	*	*	*
296.5	420	183	237	*	223	136	*	133	112	152	*
296.50	212	95	116	*	108	*58	*	*43	*65	98	*

See footnotes and notes at end of table.

Table 43. Number of days of care for discharges from short-stay hospitals, by ICD-9-CM code of first-listed diagnosis, sex, age, and geographic region: United States, 1999—Con.

[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table: see Methods section]

ICD-9-CM code	Sex		Age				Region				
	Total	Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	North-east	Midwest	South	West
Number of days of care in thousands											
296.53	106	*50	*56	*	76	*	*	*54	*	*	*
296.54	88	*	*60	*	*34	*	*	*	*	*	*
296.6	507	206	301	*	286	159	*	181	105	190	*
296.60	277	134	144	*	178	75	*	*75	*42	150	*
296.63	89	*	*66	*	*42	*	*	*	*	*	*
296.64	137	*48	*88	*	*63	*	*	*49	*	*	*
296.7	383	190	193	*	191	79	*	123	70	132	*57
296.8	107	*43	*64	*	79	*	*	*52	*	*	*
296.89	78	*	*53	*	*61	*	*	*	*	*	*
296.9	115	75	*40	*	71	*	*	47	*	*46	*
296.90	111	71	*39	*	67	*	—	46	*	*44	*
297	*93	*	*63	—	*	*	*	*	*	*	*
297.9	*63	*	*	—	*	*	*	*	*	*	*
298	609	340	268	*	298	135	133	268	98	168	*76
298.9	582	330	251	*	290	122	127	260	92	158	*72
300	340	143	197	*	178	86	58	102	101	84	53
300.0	95	45	50	*	27	45	*23	*26	*25	25	*19
300.00	63	*34	29	*	*20	26	*16	*17	*20	*16	*
300.01	*19	*	*	—	*	*	*	*	*	*	*
300.1	48	*	*42	*	*33	*	*	*	*	*	*
300.11	*22	*	*18	*	*	*	*	*	*	*	*
300.4	106	*38	68	*	50	*	*	*41	*25	*	*
300.9	*20	*	*	*	*	*	*	*	*	*	*
301	89	*	*67	*	*70	*	*	*	*	*	*
303	868	608	260	*	505	286	71	171	246	322	129
303.0	108	68	*40	*	51	41	*	*22	*26	33	*27
303.00	32	*20	*	—	*14	*16	*	*	*	*	*
303.01	71	47	*	*	37	*24	*	*	*	*18	*24
303.9	760	540	220	*	454	245	*55	149	220	289	101
303.90	293	201	91	*	156	107	*	*	40	228	*
303.91	462	337	124	*	294	136	*	142	176	60	83
304	472	296	176	*	369	88	*	116	58	260	*38
304.0	159	100	60	—	119	*39	*	58	*	74	*
304.00	64	*35	*29	—	51	*	*	*	*	58	*
304.01	95	64	*31	—	67	*27	*	55	*	*	*
304.1	*37	*	*	—	*	*	*	*	*	*	*
304.2	124	94	*30	—	108	*	*	*26	*	79	*
304.20	65	*45	*	—	58	*	—	*	*	58	—
304.21	59	*48	*	—	*50	*	*	*26	*	*	*
304.7	*43	*	*	—	*38	*	*	*	*	*	*
304.71	*39	*	*	—	*34	*	*	*	*	*	*
304.9	*47	*	*	—	*	*	*	*	*	*46	*
304.90	*45	*	*	—	*	*	*	—	*	*44	*
305	294	209	85	*	197	75	*20	32	29	198	34
305.0	118	88	31	*	65	*36	*	*14	*12	86	*
305.00	94	74	*20	*	52	*27	*	*	*10	74	*
305.01	*20	*11	*	—	*	*	*	*	*	*	*
305.1	44	32	*13	—	32	*11	*	*	*	*27	*
305.6	33	*15	*	—	*29	*	—	*	*	*	*
305.60	*21	*	*	—	*18	*	—	*	*	*	*
305.9	*57	*43	*	*	*46	*	—	*	*	*40	*
305.90	*42	*29	*	*	*	*	—	*	*	*	*
307	77	*	*68	*	*46	*	*	*	*	*	*
308	*23	*	*	*	*	*	*	—	*	*	*

See footnotes and notes at end of table.

Table 43. Number of days of care for discharges from short-stay hospitals, by ICD-9-CM code of first-listed diagnosis, sex, age, and geographic region: United States, 1999—Con.

[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table: see Methods section]

ICD-9-CM code	Sex		Age				Region				
	Total	Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	North-east	Midwest	South	West
Number of days of care in thousands											
309	517	271	246	*214	252	42	*	328	57	108	*24
309.0	111	46	65	*	80	*18	*	73	*15	*18	*
309.2	77	*30	*47	*	32	*	*	*	*	*46	*
309.28	56	*	*45	*	*29	*	*	*	*	*	*
309.4	49	*29	*20	*	35	*	*	*28	*	*	*
309.8	228	*	*77	*	*66	*	—	*177	*	*	*
309.81	227	*	*77	*	*65	*	—	*176	*	*	*
309.9	*32	*	*	*	*30	*	*	*	*	*	*
310	*186	*	*	*	*	*	*175	*	*	*	*
311	587	265	321	79	383	75	50	331	73	149	*34
312	288	220	*69	*106	163	*	*	157	*47	*47	*
312.3	168	136	*32	*	116	*	*	105	*	*	*
312.30	102	*82	*	*	86	*	*	*88	*	*	*
312.34	*58	*51	*	*	*26	*	*	*	*	*	*
312.9	*54	*48	*	*	*	*	*	*	*	*	—
313	184	*161	*	*155	*28	—	—	*133	*	*	*
313.8	184	*161	*	*155	*28	—	—	*133	*	*	*
313.81	182	*159	*	*154	*28	—	—	*133	*	*	*
314	138	109	*29	110	*28	—	—	*	55	*	*
314.0	138	109	*29	110	*28	—	—	*	55	*	*
314.01	133	104	*29	106	*	—	—	*	55	*	*
320-389	2,573	1,184	1,388	341	574	520	1,138	544	604	1,022	404
320	*90	*	*	*	*	*	*	*	*	*	*
322	*38	*	*	*	*	*	*	*	*	*	*
322.9	*30	*	*	*	*	*	*	*	*	*	*
324	*92	*	*	*	*	*	*	*	*	*	*
331	496	179	316	*	*	*	449	80	116	238	61
331.0	415	136	279	—	*	*	403	*62	103	211	*38
331.3	*42	*	*	*	*	*	*	*	*	*	*
331.4	*32	*	*	*	*	*	*	*	*	*	*
332	125	72	53	—	*	*	116	*29	*44	*44	*
332.0	122	69	53	—	*	*	113	*28	*43	*44	*
333	*45	*	*	*	*	*	*	*	*	*	*
335	*30	*	*	*	*	*	*	*	*	*	*
335.2	*29	*	*	—	*	*	*	*	*	*	*
340	104	*32	72	*	49	49	*	*	*34	*38	*
343	*21	*	*	*	*	*	*	*	*	*	*
344	*63	*	*	*	*	*	*	*	*	*	*
345	231	97	134	44	103	46	*38	89	*27	90	25
345.1	*35	*	*	*	*	*	*	*	*	*	*
345.10	*33	*	*	*	*	*	*	*	*	*	*
345.3	73	*38	*36	*	*35	*	*	*	*	*	*
345.5	*22	*	*	*	*	*	*	*	*	*	*
345.9	43	*25	*	*	*	*	*	*	*	*	*
345.90	33	*19	*	*	*	*	*	*	*	*	*
346	134	41	93	*	65	60	*	*25	52	31	*26
346.8	*25	*	*	*	*	*	*	*	*	*	*
346.9	63	*	51	*	39	*20	*	*	*23	*19	*
346.90	44	*	33	*	*24	*17	*	*	*18	*15	*
346.91	*19	*	*18	*	*15	*	*	*	*	*	*
348	180	113	67	*49	*38	*34	*59	*22	*39	*57	*62
348.1	*51	*	*	*	*	*	*	*	*	*	*
348.3	*52	*	*	*	*	*	*	*	*	*	*
349	*56	*	*	*	*	*	*	*	*	*	*

See footnotes and notes at end of table.

Table 43. Number of days of care for discharges from short-stay hospitals, by ICD-9-CM code of first-listed diagnosis, sex, age, and geographic region: United States, 1999—Con.

[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table: see Methods section]

ICD-9-CM code	Sex			Age				Region			
	Total	Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	North-east	Midwest	South	West
Number of days of care in thousands											
350	*17	*	*	*	*	*	*	*	*	*	*
353	*22	*	*	*	*	*	*	*	*	*	*
356	*30	*	*	*	*	*	*	*	*	*	*
357	90	*52	*	*	*	*	*	*	*	*35	*
357.0	*53	*	*	*	*	*	*	*	*	*	*
358	*35	*	*	*	*	*	*	*	*	*	*
358.0	*35	*	*	—	*	*	*	*	*	*	*
361	*9	*	*	*	*	*	*	*	*	*	*
366	*11	*	*	—	*	*	*	*	*	*	*
368	*17	*	*	*	*	*	*	*	*	*	*
376	26	*	*	*16	*	*	*	*	*	*	*
376.0	*23	*	*	*16	*	*	*	*	*	*	*
376.01	*23	*	*	*	*	*	*	*	*	*	*
380	*40	*14	*	*	*	*	*	*	*	*	*
380.1	*15	*	*	*	*	*	*	*	*	*	*
380.10	*13	*	*	*	*	*	*	*	*	*	*
382	57	28	30	53	*	*	*	*	*16	30	*
382.9	53	24	29	49	*	*	*	*	*13	29	*
383	*20	*	*	*	*	*	*	*	*	*	*
386	109	41	67	*	*	*18	88	*25	32	38	*14
386.1	37	*	24	*	*	*	31	*	*	*	*
386.11	*19	*	*	*	*	*	*	*	*	*	*
386.3	59	*24	36	*	*	*	48	*	*21	*21	*
386.30	54	*20	34	—	*	*	43	*	*17	*20	*
390-459	30,820	14,858	15,961	150	1,683	7,735	21,252	7,960	6,977	11,403	4,480
394	*49	*	*42	*	*	*	*	*	*	*	*
396	107	*	85	—	*	*	77	*	*	*	*
396.2	*56	*	*	—	*	*	*52	*	*	*	*
396.3	*35	*	*	—	*	*	*	*	*	*	*
398	136	*57	80	—	*	*	119	*55	*	*50	*
398.9	136	*57	80	—	*	*	119	*55	*	*50	*
398.91	136	*56	80	—	*	*	119	*55	*	*50	*
401	512	189	323	*	76	190	244	148	91	217	56
401.0	77	*27	50	—	*16	*24	37	*14	*	48	*
401.1	*12	*	*	*	*	*	*	*	*	*	*
401.9	423	160	263	*	58	164	200	130	80	168	46
402	459	140	319	*	*	76	360	154	75	187	44
402.0	*28	*	*	—	*	*	*	*	*	*	—
402.9	422	120	302	*	*	71	331	142	71	168	42
402.90	47	*18	29	—	*	*25	*20	*19	*	*20	*
402.91	376	102	274	*	*	45	311	123	65	148	*39
403	523	238	286	*	95	166	249	85	82	243	113
403.9	485	227	258	*	77	153	242	78	66	230	111
403.91	469	223	246	*	74	148	236	78	65	216	111
404	201	91	111	*	*	*42	130	*43	*45	87	*
404.9	180	80	100	*	*	*35	119	*41	*43	78	*
404.93	154	65	90	*	*	*	108	*38	*	*64	*

See footnotes and notes at end of table.

Table 43. Number of days of care for discharges from short-stay hospitals, by ICD-9-CM code of first-listed diagnosis, sex, age, and geographic region: United States, 1999—Con.[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table: see Methods section]

ICD-9-CM code	Sex			Age				Region			
	Total	Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	North-east	Midwest	South	West
	Number of days of care in thousands										
410	4,608	2,530	2,078	*	171	1,275	3,161	1,247	1,040	1,599	721
410.0	154	69	85	—	*	51	98	*52	*25	*38	*39
410.01	152	68	84	—	*	51	96	*50	*25	*38	*39
410.1	756	406	349	—	*44	250	461	150	159	317	130
410.11	741	400	341	—	*43	247	450	149	158	307	127
410.2	118	67	50	—	*	*24	82	*	*	56	*
410.21	116	67	49	—	*	*24	80	*	*	56	*
410.3	106	76	*30	—	*	*67	*37	*	*28	*	*
410.31	104	76	*29	—	*	*65	*37	*	*28	*	*
410.4	810	543	267	—	56	249	505	235	193	245	138
410.41	795	537	258	—	56	245	494	230	192	236	137
410.5	73	*32	*	—	*	*19	*52	*	*	*	*
410.51	72	*32	*	—	*	*19	*51	*	*	*	*
410.6	*43	*39	*	—	*	*	*	*	*	*	*
410.61	*43	*39	*	—	*	*	*	*	*	*	*
410.7	1,828	944	884	*	34	446	1,347	568	415	581	264
410.71	1,801	935	867	*	34	443	1,324	557	409	576	259
410.8	63	*38	*	—	*	*	*42	*	*	*	*
410.81	63	*38	*	—	*	*	*42	*	*	*	*
410.9	657	316	341	—	*	125	520	137	161	267	93
410.90	*48	*	*29	—	*	*	*42	*	*	*	*
410.91	602	297	305	—	*	120	472	110	154	251	87
411	490	218	271	*	22	161	307	143	76	206	64
411.1	437	195	242	*	17	149	271	129	69	187	52
411.8	48	*22	*26	—	*	*	*35	*	*	*	*
411.89	*30	*19	*	—	*	*	*20	*	*	*	*
412	*22	*13	*	—	*	*	*	*	*	*	*
413	203	88	114	*	18	63	121	47	50	82	24
413.9	201	88	113	*	18	62	121	46	50	82	24
414	4,597	2,684	1,913	*	169	1,618	2,800	1,159	1,000	1,692	746
414.0	4,481	2,621	1,860	*	154	1,567	2,752	1,135	979	1,630	737
414.00	658	338	320	—	*28	194	437	110	126	325	98
414.01	3,569	2,132	1,437	*	123	1,286	2,152	967	806	1,199	597
414.02	174	97	77	—	*	45	127	*31	37	66	*39
414.04	*31	*	*	—	*	*	*	*	*	*	*
414.05	45	*27	*18	—	*	*	*30	*	*	*20	*
414.8	80	*49	*	—	*	*39	*28	*	*	*40	*
414.9	31	*	*20	—	*	*	*20	*	*	*19	*
415	687	237	451	*	82	169	432	162	163	247	115
415.1	673	229	444	—	79	166	427	154	160	245	114
415.11	*44	*	*	—	*	*	*	*	*	*	*
415.19	628	209	419	—	75	144	410	135	152	233	109
416	60	*24	*36	*	*	*	*	*	*	*	*
416.0	*37	*	*	*	*	*	*	*	*	*	*
420	72	*	*47	—	*	*30	*	*	*	*	*
420.9	72	*	*47	—	*	*30	*	*	*	*	*
420.90	*54	*	*39	—	*	*	*	*	*	*	*
421	176	108	*	*	*61	*	*	*	*	*	*
421.0	169	*104	*	*	*	*	*	*	*	*	*
423	90	*40	*50	*	*	*	*41	*	*	*	*
423.9	66	*	*44	*	*	*	*36	*	*	*	*
424	614	311	303	*	30	135	446	215	117	187	96
424.0	228	108	120	*	*19	76	133	71	*39	93	*25
424.1	377	198	179	*	*	*56	311	142	76	91	68
425	198	88	111	*	*30	88	73	*64	*35	85	*
425.4	155	62	93	*	*	*70	65	*49	*33	63	*

See footnotes and notes at end of table.

Table 43. Number of days of care for discharges from short-stay hospitals, by ICD-9-CM code of first-listed diagnosis, sex, age, and geographic region: United States, 1999—Con.

[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table: see Methods section]

ICD-9-CM code	Sex			Age				Region			
	Total	Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	North-east	Midwest	South	West
Number of days of care in thousands											
426	178	91	87	*	*9	18	149	59	52	38	29
426.0	100	46	54	*	*	*	91	*33	*31	*22	*14
426.1	36	*19	*17	*	*	*	30	*	*	*	*
427	2,541	1,132	1,409	*28	124	479	1,911	768	562	877	334
427.0	75	27	48	*	*	23	42	*22	*14	32	*
427.1	296	171	125	*	*	65	213	105	53	90	47
427.3	1,390	589	802	*	57	260	1,068	433	340	450	168
427.31	1,249	504	745	*	47	219	980	392	306	400	151
427.32	142	85	57	*	*	41	88	41	33	50	*17
427.4	*43	*33	*	—	*	*	*	*	*	*	*
427.41	*43	*33	*	—	*	*	*	*	*	*	*
427.5	48	*36	*	*	*	*	*	*	*	*	*
427.6	32	*11	21	*	*	*	21	*	*	*11	*
427.69	23	*8	*15	*	*	*	*13	*	*	*7	*
427.8	597	226	371	*	30	89	473	160	131	239	68
427.81	285	93	191	*	*	*41	240	85	55	118	*27
427.89	313	133	180	*	27	47	233	75	76	121	40
427.9	58	*40	*18	*	*	*	45	*28	*	*24	*
428	5,339	2,345	2,993	*	175	902	4,251	1,373	1,362	1,969	635
428.0	5,296	2,337	2,959	*	173	898	4,215	1,368	1,355	1,949	625
428.1	*22	*	*	—	—	*	*	*	*	*	*
428.9	*20	*	*	*	*	*	*16	*	*	*15	*
429	89	36	53	*	*	*28	51	*	*17	53	*
429.2	*33	*	*	—	*	*	*18	—	*	*25	*
430	265	*85	180	*	*	*77	151	*64	*	*128	*
431	563	335	228	*	*36	254	269	118	105	150	190
432	104	76	*28	*	*	*	79	*32	*	*	*
432.1	88	68	*	*	*	*	70	*	*	*	*
433	636	302	334	—	*	161	468	111	182	261	83
433.1	443	201	242	—	*	124	316	73	127	190	54
433.10	281	139	142	—	*	50	229	54	82	106	38
433.11	163	63	100	—	*	*74	87	*	*45	83	*
433.3	172	90	82	—	*	31	137	*28	50	66	*28
433.30	110	48	62	—	*	*23	84	*25	34	35	*
433.31	*62	*	*	—	—	*	*54	*	*	*	*
434	1,774	789	985	*	68	362	1,340	475	404	653	243
434.0	136	74	62	*	*	*	106	*	*44	*55	*
434.01	105	*56	*60	*	*	*	78	*	*33	*39	*
434.1	233	113	120	—	*	*47	174	*75	*49	73	*35
434.11	218	104	115	—	*	*45	161	*68	*44	72	*35
434.9	1,405	603	803	*	*49	293	1,059	383	310	525	188
434.90	*27	*	*	*	*	*	*19	*	*	*	*
434.91	1,378	593	786	*	*48	286	1,041	380	303	511	183
435	640	274	366	*	*22	116	501	191	175	209	65
435.3	37	*	*24	—	*	*	31	*	*	*	*
435.9	584	251	333	*	*22	102	460	177	157	192	58
436	918	362	556	*	*	137	761	117	209	445	147
437	133	56	77	*	*	*35	64	*	38	48	*23
437.1	*21	*	*	—	*	*	*15	*	*	*	*
437.3	*41	*	*	*	*	*	*	*	*	*	*
437.9	*13	*	*	—	*	*	*	—	*	*	*
438	186	55	130	*	*	*25	159	*	*27	77	*59
438.2	*72	*	*	*	—	*	*63	*	*	*	*
438.20	*66	*	*	*	—	*	*57	*	*	*	*
438.8	72	*	*49	—	*	*	*61	*	*	*	*
438.89	62	*	*42	—	*	*	*51	*	*	*	*

See footnotes and notes at end of table.

Table 43. Number of days of care for discharges from short-stay hospitals, by ICD-9-CM code of first-listed diagnosis, sex, age, and geographic region: United States, 1999—Con.[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table: see Methods section]

ICD-9-CM code	Sex			Age				Region			
	Total	Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	North-east	Midwest	South	West
Number of days of care in thousands											
440	768	381	387	—	*	224	534	190	224	259	96
440.1	*18	*	*	—	*	*	*	*	*	*	*
440.2	672	326	347	—	*	183	480	162	210	221	80
440.20	60	*31	*	—	*	*	*	*	*	*29	*
440.21	104	66	*39	—	*	*40	62	*	*34	*26	*32
440.22	82	*37	*45	—	*	*	61	*	*	*36	*
440.23	106	*36	*71	—	*	*	88	*	*	*	*
440.24	315	153	162	—	*	*82	229	97	101	*90	*
440.3	*52	*37	*	—	*	*	*	*	*	*	*
441	543	376	168	—	*	109	413	167	114	189	73
441.0	*79	*54	*	—	*	*	*	*	*	*	*
441.4	347	238	109	—	*	*63	283	102	84	118	*42
442	*67	*45	*	*	*	*	*54	*	*	*	*
442.3	*34	*	*	*	*	*	*	*	*	*	*
443	99	41	58	*	*	30	68	*	*	52	*
443.9	87	37	50	—	*	*28	57	*	*	51	*
444	301	116	185	—	*	83	195	92	68	121	*
444.2	219	83	136	—	*	*42	163	*63	*53	92	*
444.22	190	79	112	—	*	*40	141	*52	*49	80	*
444.8	*32	*	*	—	*	*	*	*	*	*	*
444.81	*26	*	*	—	*	*	*	*	*	*	*
446	78	*45	*34	*	*	*	*29	*	*	*	*
447	79	*31	*48	*	*	*	*57	*	*	*	*
447.1	*34	*	*	—	*	*	*	*	*	*	*
451	165	45	120	*	*	*36	102	*38	*34	74	*
451.0	*26	*	*	—	*	*	*	*	*	*	*
451.1	87	*	*66	—	*	*	60	*	*	*50	*
451.19	74	*	*56	—	*	*	*53	*	*	*	*
451.2	*27	*	*	*	*	*	*	*	*	*	*
453	842	365	477	—	125	245	472	169	222	340	112
453.8	732	317	414	—	109	202	421	147	209	271	105
453.9	70	*	*40	—	*	*	*38	*	*	*33	*
454	82	*59	*	*	*	*	*69	*	*	*	*
455	76	35	40	—	*22	*17	37	*21	*	37	*
455.2	32	*	*18	—	*	*	*	*	*	*	*
456	*39	*28	*	*	*	*	*	*	*	*	*
456.0	*30	*	*	*	*	*	*	*	*	*	*
458	309	141	168	*	*	58	243	58	89	111	51
458.0	180	87	94	*	*	*34	140	40	56	*51	*33
458.9	120	49	70	*	*	*21	96	*	*31	56	*17
459	114	45	69	*	*	*30	79	*	*	66	*
459.8	*43	*	*24	*	*	*	*	*	*	*	*
459.81	*38	*	*	—	*	*	*	*	*	*	*
460-519	20,054	9,225	10,829	2,328	1,966	3,662	12,097	4,838	4,154	7,729	3,333
461	50	*23	*27	*	*18	*	*	*	*	*	*
462	28	*	20	*	*15	*	*	*	*	*12	*
463	18	*	*11	*	*12	*	—	*	*	*	*
464	92	58	34	73	*	*	*	*19	20	25	28
464.4	58	39	19	58	—	—	—	*	*9	21	20
465	124	44	80	40	*15	*16	52	33	*20	51	*19
465.9	121	43	78	39	*14	*16	52	33	*20	49	*19

See footnotes and notes at end of table.

Table 43. Number of days of care for discharges from short-stay hospitals, by ICD-9-CM code of first-listed diagnosis, sex, age, and geographic region: United States, 1999—Con.

[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table: see Methods section]

ICD-9-CM code	Sex			Age				Region			
	Total	Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	North-east	Midwest	South	West
Number of days of care in thousands											
466	983	479	504	599	59	69	255	194	185	441	162
466.0	405	158	247	36	56	69	245	87	87	203	29
466.1	577	320	257	563	*	*	*	107	98	238	133
466.11	347	199	148	347	*	*	*	60	58	155	74
466.19	230	121	109	217	*	*	*	48	40	83	59
473	55	36	*19	*	*	*	*	*	*	*19	*
473.9	*14	*	*	*	*	*	*	*	*	*	*
474	24	*11	*13	18	*	*	—	*10	*	*	*
474.0	*10	*	*	*	*	—	—	*	*	*	*
474.1	*14	*	*	*10	*	*	—	*	*	*	*
474.10	*11	*	*	*	*	—	—	*	*	*	*
475	26	*15	*	*	*19	*	*	*	*	*	*
478	71	34	*38	*34	*	*13	*	*32	*	*14	*15
478.2	*23	*17	*	*	*	*	*	*	*	*	*
478.7	*18	*	*	*	*	*	*	*	*	*	*
480	245	129	115	182	*	*	*28	*	40	102	84
480.1	152	91	61	146	*	*	*	*	*	82	*47
480.9	71	32	*39	*26	*	*	*	*	*	*18	*25
481	366	179	187	*33	*29	116	189	*83	93	134	*57
482	1,445	739	706	77	253	171	943	262	331	618	233
482.0	76	*41	*	*	*	*	*59	*	*	*41	*
482.1	363	205	158	*	*	*37	233	*	*82	149	81
482.2	95	*43	*52	*	*	*	84	*	*35	*27	*
482.3	*73	*	*	*	*	*	*	*	*	*	*
482.4	398	215	183	*	*	*	314	*70	*75	166	*86
482.41	370	196	175	*	*	*	288	*63	*66	155	*86
482.8	343	124	219	*	*	*	169	*	96	185	*
482.83	269	88	181	*	*	*	129	*	77	154	*
482.9	97	*54	*43	*	*	*	*62	*35	*	*36	*
483	82	*	*66	*	*	*	*56	*	*	*	*
483.0	75	*	*64	*	*	*	*53	*	*	*	*
485	173	81	92	42	*	*	104	*44	40	70	*
486	5,985	2,807	3,177	487	471	975	4,051	1,377	1,338	2,313	957
487	150	55	94	*20	*21	*26	82	*33	59	*33	*25
487.0	66	*	*38	*	*	*	*33	*	*	*	*
487.1	81	*25	56	*	*	*	49	*	39	*21	*
490	98	41	57	*	*10	*16	60	*	*	69	*
491	2,816	1,111	1,705	*	66	704	2,035	664	699	1,110	343
491.2	2,798	1,102	1,696	*	66	701	2,026	662	689	1,105	343
491.20	70	*30	41	*	*	*	46	*	*	*34	*
491.21	2,728	1,072	1,656	*	64	678	1,980	651	678	1,071	328
492	123	59	64	—	*	*40	78	*46	46	*25	*
492.8	110	49	60	—	*	*33	75	*39	44	*	*
493	1,540	531	1,008	436	388	352	364	492	289	533	226
493.0	96	50	46	74	*	*	*	22	*	*14	50
493.00	47	25	*22	35	*	*	*	*	*	*	22
493.01	49	24	*24	40	*	*	*	*12	*	*	28
493.2	328	92	236	*	*27	142	157	134	48	98	48
493.20	223	66	156	*	*	84	120	81	38	68	*35
493.21	105	*26	79	*	*	58	*37	*53	*	*30	*
493.9	1,112	387	725	359	346	202	205	335	230	419	128
493.90	717	224	493	164	229	152	172	221	128	301	68
493.91	395	163	231	195	117	51	*32	114	102	118	60

See footnotes and notes at end of table.

Table 43. Number of days of care for discharges from short-stay hospitals, by ICD-9-CM code of first-listed diagnosis, sex, age, and geographic region: United States, 1999—Con.[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table: see Methods section]

ICD-9-CM code	Sex			Age				Region			
	Total	Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	North-east	Midwest	South	West
Number of days of care in thousands											
494	*38	*	*	*	*	*	*	*	*	*	*
496	539	238	301	*	*21	113	400	140	101	204	95
507	1,533	912	621	*74	*64	142	1,253	467	273	547	246
507.0	1,519	902	618	*74	*59	142	1,244	465	271	544	240
510	114	*81	*	*	*	*	*	*	*	*	*
510.9	*105	*80	*	*	*	*	*	*	*	*	*
511	362	206	156	*	*28	81	243	76	72	170	*43
511.0	*23	*	*	*	*	*	*	*	*	*	*
511.8	*56	*	*	*	*	*	*	*	*	*	*
511.9	274	151	122	*	*22	58	187	*66	*37	140	*
512	245	124	121	*	109	*45	86	*79	56	71	*39
512.1	*25	*	*	—	*	*	*	*	*	*	*
512.8	189	89	101	*	103	*37	*48	*72	*40	46	*31
513	*74	*	*	*	*	*	*	*	*	*	*
513.0	*74	*	*	*	*	*	*	*	*	*	*
514	53	*	*42	*	*	*	*	*	*	*	*
515	117	*52	65	*	*	*	85	*	*35	*35	*
516	83	*42	*41	*	*	*	*56	*	*	*49	*
516.3	*39	*	*	—	*	*	*	*	*	*	*
516.8	*39	*	*	*	*	*	*	*	*	*	*
518	2,257	955	1,302	80	180	540	1,456	566	289	839	562
518.3	*36	*	*	—	*	*	*	*	*	*	*
518.4	*25	*	*	*	*	*	*21	*	*	*	*
518.8	2,118	896	1,223	*67	156	525	1,371	535	249	803	531
518.81	1,828	780	1,048	*	129	479	1,178	422	200	720	486
518.82	69	*33	*36	*	*	*	*25	*	*	*	*
518.84	174	*66	*108	*	*	*	143	*	*	*54	*
518.89	*38	*	*	*	*	*	*	*	*	*	*
519	102	56	46	21	*18	*	*44	*	*22	54	*12
519.1	32	*16	*	*	*	*	*	*	*	*	*
519.8	46	*30	*16	*	*	*	*34	*	*	*32	*
520-579	15,060	6,343	8,717	795	2,800	3,912	7,553	3,474	3,266	5,628	2,692
522	*16	*	*	*	*8	*	*	*	*	*	*
522.5	*13	*	*	*	*	*	*	*	*	*	*
524	35	*	24	*	29	*	*	*	*	*16	*
524.0	*19	*	*	*	*17	*	—	*	*	*	*
527	35	*	*	*	*	*	*	*	*	*	*
527.2	*25	*	*	*	*	*	*	*	*	*	*
528	40	*28	*	*	*22	*	*	*	*	*	*
528.3	*25	*	*	*	*	*	*	*	*	*	*
530	734	333	401	91	108	189	345	160	164	283	127
530.1	144	55	89	*	24	55	61	*28	46	53	*17
530.10	51	*19	31	*	*	*18	*23	*	*12	*	*
530.11	68	*24	44	*	*13	*26	28	*	*30	*22	*
530.19	*25	*	*	*	*	*	*	*	*	*	*
530.2	92	51	40	*	*	*	67	*	*	*30	*
530.3	*41	*	*	*	*	*	*	*	*	*	*
530.5	*18	*	*	*	*	*	*	*	*	*	*
530.7	59	*27	*32	*	*	*	*32	*	*	*23	*
530.8	327	127	201	84	57	78	108	69	76	117	65
530.81	301	110	191	83	56	66	96	65	71	107	58

See footnotes and notes at end of table.

Table 43. Number of days of care for discharges from short-stay hospitals, by ICD-9-CM code of first-listed diagnosis, sex, age, and geographic region: United States, 1999—Con.

[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table: see Methods section]

ICD-9-CM code	Sex		Age				Region				
	Total	Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	North-east	Midwest	South	West
Number of days of care in thousands											
531	561	242	319	*	89	112	341	127	152	207	76
531.0	79	*38	*41	*	*	*20	*36	*	*28	*27	*
531.00	79	*38	*41	*	*	*20	*36	*	*28	*27	*
531.4	273	142	131	*	*	46	181	77	65	101	31
531.40	269	138	131	*	*	44	179	77	65	98	29
531.5	*67	*	*	*	*	*	*	*	*	*	*
531.50	*67	*	*	*	*	*	*	*	*	*	*
531.9	74	*18	56	*	*	*	*40	*	*	*25	*
531.90	55	*16	*39	*	*	*	*31	*	*	*	*
532	459	287	173	*	39	70	350	161	105	142	51
532.0	73	*44	*30	—	*	*	58	*	*	*	*
532.00	72	*42	*30	—	*	*	*57	*	*	*	*
532.4	237	170	67	*	*	*31	190	*88	53	71	*25
532.40	235	169	67	*	*	*31	189	*88	52	71	*25
532.5	*61	*	*	—	*	*	*	*	*	*	*
532.50	*61	*	*	—	*	*	*	*	*	*	*
532.9	*28	*	*	—	*	*	*	*	*	*	*
532.90	*24	*	*	—	*	*	*	*	*	*	*
533	59	*22	37	*	*	*	36	*	*	*32	*
533.9	*21	*	*	*	*	*	*	*	*	*	*
533.90	*21	*	*	*	*	*	*	*	*	*	*
534	*27	*	*	—	*	*	*	*	*	*	*
535	575	242	333	*19	93	130	333	150	136	217	72
535.0	123	53	70	*	*18	34	67	*24	26	53	*20
535.00	72	31	41	*	*14	*20	34	*	*15	34	*
535.01	51	*23	*29	*	*	*	*33	*	*	*	*
535.1	78	*29	48	—	*	*26	*45	*	*	*29	*
535.10	*30	*	*	—	*	*	*	*	*	*	*
535.11	*48	*	*	—	*	*	*32	*	*	*	*
535.3	*15	*14	*	—	*	*	*	*	*	*	*
535.4	130	53	77	*	*26	*23	78	*26	44	41	*18
535.40	62	*23	39	*	*	*	30	*	*20	*20	*
535.41	68	*30	*38	*	*	*	48	*	*24	*22	*
535.5	183	63	120	*	29	33	112	53	36	80	*15
535.50	94	27	67	*	*23	*20	42	*28	*17	40	*9
535.51	89	*36	53	*	*	*	70	*	*	*40	*
535.6	45	*30	*	*	*	*	*	*	*	*	*
535.60	*23	*	*	*	*	*	*	*	*	*	*
536	261	102	159	*	*41	*32	164	*57	*38	125	*40
536.2	48	*	*27	*	*	*	*	*	*	*29	*
536.3	*35	*	*	—	*	*	*	*	*	*	*
536.4	147	*53	94	*	*	*	117	*	*	*61	*
536.41	*66	*	*	*	*	*	*	*	*	*	*
536.42	*52	*	*	*	*	*	*40	*	*	*	*
536.8	*21	*	*	*	*	*	*	*	*	*	*
537	142	67	75	*	*	*34	77	*	*	74	*
537.0	82	*	*	*	*	*	*	*	*	*	*
537.8	44	*	*22	—	*	*	*37	*	*	*	*
537.83	*33	*	*	—	—	*	*28	*	*	*	*
540	834	502	332	197	392	134	111	169	191	246	229
540.0	298	194	104	85	115	49	*49	59	61	89	89
540.1	160	93	68	*35	67	*	*	*	*44	*56	*29
540.9	376	216	161	77	211	54	*35	79	85	102	110
541	43	*27	*16	*	*21	*	*	*	*	*26	*

See footnotes and notes at end of table.

Table 43. Number of days of care for discharges from short-stay hospitals, by ICD-9-CM code of first-listed diagnosis, sex, age, and geographic region: United States, 1999—Con.[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table: see Methods section]

ICD-9-CM code	Sex		Age				Region				
	Total	Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	North-east	Midwest	South	West
Number of days of care in thousands											
550	101	77	*	*10	*13	*16	62	35	21	34	*10
550.1	57	38	*	*	*	*	*40	*25	*	*19	*
550.10	32	25	*	*	*	*	*18	*	*	*	*
550.9	43	39	*	*	*	*	22	*	*13	15	*
550.90	26	23	*	*	*	*	*14	*	*	*9	*
550.92	*12	*11	*	*	*	*	*	*	*	*	*
552	343	90	254	*	*28	118	195	79	66	136	64
552.0	*27	*	*	*	*	*	*27	*	*	*	*
552.00	*26	*	*	*	*	*	*	*	*	*	*
552.1	56	*	*46	*	*	*17	*35	*	*	*	*
552.2	211	57	154	*	*	89	107	*50	*25	87	*48
552.20	45	*	*36	—	*	*	*	*	*	*	*
552.21	146	*42	105	—	*	71	67	*30	*	64	*
553	292	100	192	*	39	124	125	59	54	106	74
553.1	*13	*	*	*	*	*	*	*	*	*	*
553.2	175	63	112	*	*20	86	69	37	39	64	36
553.20	*24	*	*	*	*	*	*	*	*	*	*
553.21	146	53	93	*	*17	72	57	28	34	52	32
553.3	97	*30	67	*	*	33	*50	*	*	*32	*36
555	223	101	122	*	111	56	*33	55	82	60	*27
555.0	66	*36	*	*	*37	*	*	*	*	*	*
555.1	*49	*	*30	*	*	*	*	*	*	*	*
555.9	83	*36	47	*	45	*	*	*	*20	*36	*
556	133	50	83	*	59	*35	*32	*34	*	*50	*
556.9	81	*35	*45	*	*34	*	*	*	*	*28	*
557	259	82	177	*	*	63	167	49	60	97	*54
557.0	138	*43	95	*	*	*	82	*	*35	*58	*
557.9	90	*	54	—	*	*	64	*	*	*	*
558	489	180	309	116	91	90	192	137	90	213	49
558.9	453	162	291	116	89	86	162	120	86	201	45
560	1,936	762	1,173	89	212	365	1,269	429	414	698	395
560.1	330	153	176	*	50	48	224	71	69	149	*41
560.2	115	*53	*62	*	*	*	91	*	*47	*40	*
560.3	171	68	103	*	*	*	148	*44	*30	81	*
560.39	158	64	94	*	*	*	136	*42	*28	71	*
560.8	738	262	475	*	97	149	472	150	151	210	227
560.81	639	240	400	*	81	126	414	131	126	176	207
560.89	98	*	76	*	*	*	*58	*	*	*	*
560.9	568	216	352	*	53	142	333	151	112	215	90
562	1,347	494	853	*	110	354	883	361	295	495	197
562.1	1,336	486	850	—	110	352	873	359	292	491	194
562.10	171	69	102	—	*	*37	117	*41	*35	60	*35
562.11	784	266	518	—	83	280	420	223	175	279	107
562.12	298	122	176	—	*	*26	269	74	65	118	41
562.13	83	*29	53	—	*	*	66	*	*	*34	*
564	218	60	158	*28	32	39	118	53	66	63	36
564.0	119	33	86	*12	*	*	80	35	40	*31	*14
564.1	41	*	36	*	*	*	*	*	*	*18	*
564.8	*35	*	*	*	*	*	*	*	*	*	*
564.89	*34	*	*	*	*	*	*	*	*	*	*
566	57	31	*26	*	*17	*29	*	*22	*	*16	*
567	157	*55	102	*	*	*69	*	*	*	*80	*
567.2	107	*47	*60	*	*	*50	*	*	*	*	*
568	71	*	59	*	*27	*	*	*	*	*	*
568.0	48	*	*40	*	*	*	*	*	*	*	*

See footnotes and notes at end of table.

Table 43. Number of days of care for discharges from short-stay hospitals, by ICD-9-CM code of first-listed diagnosis, sex, age, and geographic region: United States, 1999—Con.

[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table: see Methods section]

ICD-9-CM code	Sex			Age				Region			
	Total	Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	North-east	Midwest	South	West
Number of days of care in thousands											
569	685	280	405	*	58	190	419	124	159	313	90
569.1	*44	*	*36	—	*	*	*	*	*	*	*
569.3	81	*27	54	*	*	*	*56	*	*	*45	*
569.4	33	*19	*	*	*	*	*21	*	*	*	*
569.6	72	*41	*	*	*	*	*44	*	*	*	*
569.69	*52	*33	*	*	*	*	*	*	*	*	*
569.8	403	165	239	*	*	*119	261	*67	101	201	*35
569.81	*83	*	*	—	*	*	*	*	*	*	*
569.82	*58	*	*	—	*	*	*	*	*	*	*
569.83	134	*50	*84	*	*	*	*87	*	*	*58	*
569.85	104	*	*81	*	*	*	91	*	*	*48	*
569.89	*20	*	*	*	*	*	*	*	*	*	*
570	*45	*	*	—	*	*	*	*	*	*	*
571	549	284	265	*	112	261	164	123	84	248	95
571.1	*45	*36	*	—	*	*	*	*	*	*	*
571.2	250	173	76	*	*46	150	*53	*60	*34	108	*48
571.5	182	65	117	*	*31	74	77	*	*	102	*
572	231	124	106	*	*	131	63	*50	*	90	*62
572.2	143	71	72	—	*	101	*	*	*	*49	*51
573	74	*	*51	*	*	*	*	*	*	*	*
573.3	*29	*	*	*	*	*	*	*	*	*	*
574	1,377	491	886	*	303	382	683	343	291	445	298
574.0	511	204	307	*	105	140	265	134	111	152	114
574.00	477	196	280	*	98	133	245	130	99	142	105
574.01	*34	*	*27	—	*	*	*	*	*	*	*
574.1	446	137	309	*	108	144	193	103	92	160	91
574.10	431	132	300	*	105	139	186	98	90	156	87
574.2	119	31	88	*	28	36	53	46	*22	39	*
574.20	108	26	82	*	28	30	49	46	*16	35	*
574.5	84	*25	59	*	*19	*	49	*	*	*26	*26
574.50	50	*	42	—	*	*	*29	*	*	*11	*
574.51	*34	*	*	*	*	*	*	*	*	*	*
574.6	*33	*	*	—	*	*	*	*	*	*	*
574.7	72	*	*47	*	*	*	*40	*	*	*25	*
574.70	*38	*	*25	*	*	*	*	*	*	*	*
574.8	*55	*	*20	*	*	*	*	*	*	*	*
575	258	86	172	*	50	71	135	84	54	87	*32
575.0	97	*38	59	*	*	*20	60	*	*17	*42	*
575.1	140	44	95	*	29	46	63	56	32	39	*
575.10	*35	*	*	*	*	*	*	*	*	*	*
575.11	67	*	56	*	*14	32	*21	*	*23	30	*
575.12	*37	*	*	—	*	*	*	*	*	*	*
576	158	88	70	*	*	*25	99	*38	*30	*57	*34
576.1	92	*56	*36	*	*	*	*55	*	*	*	*
576.2	*23	*	*	*	*	*	*	*	*	*	*
576.8	*40	*	*	*	*	*	*	*	*	*	*
577	1,400	576	824	*	430	522	427	273	307	550	271
577.0	1,184	503	681	*	367	438	363	223	260	480	220
577.1	143	*57	86	*	*45	*55	*	*	*	*47	*
577.2	*59	*	*	*	*	*	*	*	*	*	*
578	713	322	391	*	45	142	517	152	196	258	107
578.0	55	*31	*23	*	*	*	*34	*	*	*18	*
578.1	114	47	67	*	*	*21	87	*28	*29	36	*21
578.9	544	243	301	*	*35	111	397	114	151	204	75
580-629	6,618	2,370	4,248	262	1,619	1,510	3,228	1,395	1,308	2,627	1,289
581	*32	*	*	*	*	*	*	*	*	*	*
581.9	*28	*	*	*	*	*	*	*	*	*	*

See footnotes and notes at end of table.

Table 43. Number of days of care for discharges from short-stay hospitals, by ICD-9-CM code of first-listed diagnosis, sex, age, and geographic region: United States, 1999—Con.

[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table: see Methods section]

ICD-9-CM code	Sex			Age				Region			
	Total	Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	North-east	Midwest	South	West
Number of days of care in thousands											
584	804	406	398	*	*100	173	527	201	157	291	155
584.5	133	*107	*	*	*	*	*	*	*	*	*
584.9	647	294	354	*	*57	115	474	171	140	208	128
585	265	211	55	*	*	*145	90	*54	*	49	*136
586	*39	*	*	*	*	*	*	*	*	*	*
590	470	80	390	58	186	104	122	98	106	173	93
590.1	276	42	234	*26	113	69	67	52	67	104	52
590.10	275	42	233	*26	113	69	67	52	67	104	52
590.8	173	35	139	32	66	*26	*49	*38	33	65	37
590.80	173	35	139	32	66	*26	*49	*38	33	65	37
591	36	*19	*	*	*	*	*	*	*	*	*
592	443	248	195	*	175	163	100	85	99	197	61
592.0	173	89	85	*	67	65	38	37	34	77	*25
592.1	264	154	109	*	105	98	60	47	64	119	33
593	148	62	86	*27	*24	*43	54	43	30	55	*
593.4	33	*15	*18	*	*	*	*	*	*	*14	*
593.7	*22	*	*	*20	*	*	—	*	*	*	*
593.70	*18	*	*	*	*	*	—	*	*	*	*
593.9	55	*22	*33	*	*	*	*34	*	*	*27	*
594	*25	*22	*	*	*	*	*	*	*	*	*
594.1	*23	*20	*	*	*	*	*	*	*	*	*
595	90	35	*55	*	*	*	72	*21	*16	*	*
595.8	*38	*	*	—	*	*	*	*	*	*	*
596	56	32	*	*	*	*	*29	*	*	*	*
599	1,859	647	1,212	106	154	205	1,393	419	342	821	277
599.0	1,755	585	1,170	103	130	190	1,332	399	317	776	263
599.7	70	41	29	*	*	*	51	*	*22	*24	*
600	406	406	—	—	*	41	362	69	77	115	145
601	*21	*21	—	—	*	*	*	*	*	*	*
604	*34	*34	—	*	*	*	*	*	*	*23	*
604.9	*28	*28	—	*	*	*	*	*	*	*	*
607	*20	*20	—	*	*	*	*	*	*	*	*
607.8	*11	*11	—	*	*	*	*	*	*	*	*
608	*21	*21	—	*	*	*	*	*	*	*	*
610	*11	*	*10	—	*	*	*	*	*	*	*
611	48	*	44	*	23	*21	*	*20	*	*11	*
611.1	13	*	13	*	*10	*	*	*6	*	*	*
614	225	—	225	*	154	*45	*	63	54	68	40
614.1	*15	—	*15	—	*	*	*	*	*	*	*
614.3	*51	—	*51	*	*30	*	*	*	*	*	*
614.6	51	—	51	—	41	*	*	*	*	23	*
614.9	40	—	40	*	39	*	—	*	*	*18	*
616	35	—	35	*	*20	*	*	*	*	*	*
616.0	*17	—	*17	—	*	*	*	*	*	*	*
617	223	—	223	—	164	54	*	32	41	91	59
617.0	127	—	127	—	85	38	*	*21	23	46	37
617.1	34	—	34	—	29	*	*	*	*	*	*
617.3	37	—	37	—	33	*	—	*	*	*18	*
617.9	*12	—	*12	—	*	*	—	*	*	*	*

See footnotes and notes at end of table.

Table 43. Number of days of care for discharges from short-stay hospitals, by ICD-9-CM code of first-listed diagnosis, sex, age, and geographic region: United States, 1999—Con.

[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table: see Methods section]

ICD-9-CM code	Sex			Age				Region			
	Total	Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	North-east	Midwest	South	West
Number of days of care in thousands											
618	398	—	398	—	72	156	169	64	89	154	92
618.0	84	—	84	—	*11	36	37	*15	*19	33	*17
618.1	31	—	31	—	*15	*	*	*	*	*10	*
618.2	49	—	49	—	*15	*21	*	*	*	*14	*
618.3	54	—	54	—	*	*21	*27	*	*	*20	*
618.4	60	—	60	—	*14	26	*21	*	*18	26	*
618.5	73	—	73	—	*	30	38	*	*	34	*17
618.6	*15	—	*15	—	*	*	*	*	*	*	*
618.8	32	—	32	—	*	*	*	*	*	*	*20
620	183	—	183	*	122	36	*20	29	35	84	36
620.0	*14	—	*14	—	*	*	—	*	*	*	*
620.1	34	—	34	*	28	*	—	*	*	*14	*
620.2	114	—	114	*	77	*19	*	*17	*18	56	*24
621	45	—	45	—	*	*23	*	*	*	*17	*
621.3	*24	—	*24	—	*	*18	*	*	*	*	*
622	22	—	22	—	*15	*	*	*	*	*	*
622.1	*17	—	*17	—	*12	*	*	*	*	*	*
625	287	—	287	—	134	100	52	38	52	160	38
625.3	50	—	50	—	46	*	—	*	*	32	*
625.6	127	—	127	—	*19	59	49	*15	*15	73	24
625.9	97	—	97	—	65	*	*	*	*30	51	*
626	211	—	211	*	141	62	*	*20	61	98	32
626.2	148	—	148	*	104	43	—	*16	40	71	21
626.8	44	—	44	*	33	*	*	*	*16	*16	*
627	38	—	38	—	*	*15	*22	*	*	*20	*
627.1	36	—	36	—	*	*14	*22	*	*	*19	*
630-677	1,307	*	1,306	*	1,279	*	—	229	272	581	224
632	20	—	20	—	20	*	—	*	*	*9	*
633	77	—	77	*	76	*	—	*17	*15	30	*16
633.1	71	—	71	*	70	*	—	*15	*13	28	*15
634	28	—	28	*	27	—	—	*9	*	*9	*5
634.9	20	—	20	*	20	—	—	*6	*	*7	*
634.91	16	—	16	*	16	—	—	*6	*	*	*
635	*19	—	*19	—	*19	—	—	*	*	*	*
640	*12	—	*12	—	*12	—	—	*	*	*	*
640.0	*10	—	*10	—	*10	—	—	*	*	*	*
640.03	*10	—	*10	—	*10	—	—	*	*	*	*
641	59	—	59	—	58	*	—	*	*	*19	*
641.1	*25	—	*25	—	*25	—	—	*	*	*	*
641.13	*25	—	*25	—	*25	—	—	*	*	*	*
641.9	*15	—	*15	—	*14	*	—	*	*	*	*
641.93	*15	—	*15	—	*14	*	—	*	*	*	*
642	64	—	64	*	59	*	—	*	*16	21	*17
642.3	22	—	22	—	17	*	—	*	*	*	*
642.33	21	—	21	—	16	*	—	*	*	*	*
642.4	20	—	20	*	20	—	—	*	*	*	*
642.43	*18	—	*18	*	*18	—	—	*	*	*	*
643	122	—	122	*	121	*	—	*19	27	64	*
643.0	55	—	55	—	55	—	—	*	*	41	*
643.03	55	—	55	—	55	—	—	*	*	41	*
643.1	58	—	58	—	57	*	—	*	*20	*19	*
643.13	58	—	58	—	57	*	—	*	*20	*19	*

See footnotes and notes at end of table.

Table 43. Number of days of care for discharges from short-stay hospitals, by ICD-9-CM code of first-listed diagnosis, sex, age, and geographic region: United States, 1999—Con.[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table: see Methods section]

ICD-9-CM code	Sex		Age				Region				
	Total	Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	North-east	Midwest	South	West
Number of days of care in thousands											
644	307	—	307	*	301	*	—	49	75	136	47
644.0	278	—	278	*	272	*	—	48	71	117	42
644.03	278	—	278	*	272	*	—	48	71	117	42
644.1	29	—	29	—	29	—	—	*	*	19	*
644.13	29	—	29	—	29	—	—	*	*	19	*
646	117	—	117	*	116	—	—	*20	*14	62	*22
646.6	86	—	86	*	85	—	—	*	*	44	*19
646.63	80	—	80	*	79	—	—	*	*	43	*18
646.8	24	—	24	—	24	—	—	*	*	*15	*
646.83	23	—	23	—	23	—	—	*	*	*13	*
647	26	—	26	*	25	—	—	*	*	*15	*
648	234	*	233	*	224	*	—	44	36	120	33
648.0	*26	—	*26	—	*26	—	—	*	*	*	*
648.03	*26	—	*26	—	*26	—	—	*	*	*	*
648.4	*27	*	*26	*	*26	*	—	*	*	*	*
648.43	*24	—	*24	*	*23	—	—	*	*	*	*
648.8	*25	—	*25	—	*25	—	—	*	*	*	*
648.83	*25	—	*25	—	*25	—	—	*	*	*	*
648.9	134	—	134	—	125	*	—	*20	*	84	20
648.93	124	—	124	—	114	*	—	*20	*	75	20
654	35	—	35	—	35	*	—	*	*	*	*
654.5	*25	—	*25	—	*25	*	—	*	*	*	*
654.53	*25	—	*25	—	*25	*	—	*	*	*	*
658	*24	—	*24	—	*24	—	—	*	*	*	*
659	*8	—	*8	*	*8	—	—	*	*	*	*
674	30	—	30	—	30	—	—	*	*	*	*
680-709	2,973	1,488	1,486	323	615	772	1,263	651	593	1,356	373
681	69	39	*30	*	*	*27	*	*	*	*26	*
681.0	*23	*	*	*	*	*	*	*	*	*	*
681.00	*18	*	*	*	*	*	*	*	*	*	*
681.1	46	*25	*	*	*	*	*	*	*	*	*
681.10	44	*25	*	*	*	*	*	*	*	*	*
682	1,796	893	903	99	367	553	777	453	398	681	264
682.0	78	*30	48	*	*26	*22	*	*29	*21	*19	*
682.1	*32	*	*	*	*	*	*	*	*	*	*
682.2	172	75	97	*	*32	76	*59	*	51	70	*
682.3	140	79	62	*	45	34	54	*35	*21	51	34
682.4	78	49	*29	*	*25	*	*24	*16	*	*	*25
682.5	*36	*	*	*	*	*	*	*	*	*	*
682.6	991	515	476	*22	176	333	460	270	215	384	122
682.7	238	107	131	*15	*29	*51	142	*55	46	101	*36
683	*33	*	*	*	*	*	*	*	*	*	*
686	*31	*	*	*	*	*	*	*	*	*	*
693	*24	*	*18	*	*	*	*	*	*	*	*
693.0	*23	*	*18	*	*	*	*	*	*	*	*
695	*31	*	*	*	*	*	*	*	*	*	*
707	624	287	337	*	74	138	406	124	135	312	*53
707.0	394	195	199	*	*59	*83	247	*	*87	215	*
707.1	205	80	126	*	*	*51	140	*55	*41	83	*27
709	273	194	79	163	89	*	*	*	*	250	*
709.2	239	173	66	152	84	*	*	*	*	237	*
710-739	6,651	2,820	3,831	170	1,074	1,770	3,637	1,397	1,522	2,432	1,300
710	170	*	153	*	*72	*45	*	*	*30	*68	*45
710.0	90	*	75	*	*47	*32	*	*	*	*46	*

See footnotes and notes at end of table.

Table 43. Number of days of care for discharges from short-stay hospitals, by ICD-9-CM code of first-listed diagnosis, sex, age, and geographic region: United States, 1999—Con.

[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table: see Methods section]

ICD-9-CM code	Sex		Age				Region				
	Total	Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	North-east	Midwest	South	West
Number of days of care in thousands											
711	141	92	*49	*	*41	*	*47	*	*	*56	*33
711.0	134	91	*43	*	*	*	*43	*	*	*53	*33
711.06	*58	*36	*	*	*	*	*	*	*	*	*
714	91	*31	60	*	*	*32	47	*	*25	44	*
714.0	78	*23	54	—	*	*31	43	*	*21	41	*
715	1,962	769	1,193	*	44	493	1,423	470	509	597	386
715.1	197	77	120	—	*	*31	161	*26	*32	71	68
715.15	82	*36	*47	—	*	*	72	*	*	*44	*21
715.16	106	*39	68	—	*	*25	81	*	*	*	*47
715.3	932	376	555	—	*25	250	657	240	257	282	154
715.35	325	134	190	—	*	89	222	89	89	75	71
715.36	590	236	354	—	*	158	422	149	160	200	80
715.8	*27	*	*	—	*	*	*25	*	*	*	*
715.89	*27	*	*	—	*	*	*25	*	*	*	*
715.9	789	304	485	*	*	204	571	193	213	229	154
715.90	29	*	*26	—	*	*	*23	*	*	*19	*
715.91	*14	*	*	—	*	*	*	*	*	*	*
715.95	249	110	139	*	*	84	158	83	65	60	40
715.96	489	182	308	—	*	115	370	104	135	142	107
716	83	38	44	*	*	*29	*40	*	*	*28	*28
716.1	31	*23	*	—	*	*18	*	*	*	*	*
716.9	*35	*	*27	—	*	*	*30	*	*	*	*
717	19	*8	*11	*	*12	*	*	*	*	*	*
717.8	*10	*	*	*	*9	*	*	*	*	*	*
717.83	*10	*	*	*	*9	*	—	*	*	*	*
718	57	25	31	*	22	*	*	*12	*16	*	*19
718.3	24	*12	*	*	*12	*	*	*	*	*	*
718.8	*8	*	*	—	*	*	*	*	*	*	*
719	112	39	73	*	*16	*23	69	*	*19	43	*32
719.4	62	*24	37	*	*	*	39	*	*	*29	*
719.45	40	*	*25	*	*	*	*30	*	*	*20	*
721	317	140	177	*	*18	86	212	47	59	126	85
721.0	58	35	23	—	*	23	*29	*	*18	18	*
721.1	63	*30	*33	—	*	*32	*29	*	*	*26	*
721.3	154	66	88	*	*	*26	118	*	*27	54	*54
722	1,018	518	499	*	398	367	249	178	262	406	172
722.0	93	50	43	—	38	46	*	*16	18	43	*15
722.1	559	292	267	—	243	188	128	106	135	220	97
722.10	548	289	259	—	239	185	124	103	131	218	95
722.4	18	*	*10	—	*	*13	*	*	*	*	*
722.5	159	69	90	—	35	49	75	*26	52	49	*32
722.52	154	69	85	—	35	48	70	*	47	49	*32
722.7	61	*27	*34	—	*17	*28	*	*	*	*30	*
722.71	26	*	*	—	*	*	*	*	*	*	*
722.73	*18	*	*	—	*	*	*	*	*	*	*
722.8	*29	*	*	—	*	*	*	*	*	*	*
722.83	*29	*	*	—	*	*	*	*	*	*	*
722.9	75	*50	*26	*	*	*27	*	*	*	*24	*
722.93	*68	*	*	*	*	*	*	*	*	*	*
723	63	38	26	*	*12	32	*19	*16	*14	26	*
723.0	*34	*	*	—	*	*	*	*	*	*	*
723.1	*13	*	*	*	*	*	*	*	*	*	*
723.4	*13	*9	*	—	*	*	*	*	*	*	*
724	509	169	339	*	62	135	311	110	116	205	77
724.0	303	93	210	—	*	79	216	68	68	121	46
724.02	295	92	204	—	*	78	210	65	66	118	46
724.2	107	40	66	*	27	29	51	*24	*26	38	*19
724.4	34	*	26	—	*	*	*	*	*	*21	*
724.5	37	*17	*20	*	*	*	*20	*	*15	*	*

See footnotes and notes at end of table.

Table 43. Number of days of care for discharges from short-stay hospitals, by ICD-9-CM code of first-listed diagnosis, sex, age, and geographic region: United States, 1999—Con.[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table: see Methods section]

ICD-9-CM code	Sex		Age				Region				
	Total	Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	North-east	Midwest	South	West
Number of days of care in thousands											
726	83	51	32	*	*33	27	24	*20	*21	*21	21
726.1	20	*12	*8	*	*	*7	*	*	*	*	*
726.10	19	*12	*	*	*	*7	*	*	*	*	*
726.2	*7	*	*	—	*	*	*	*	*	*	*
727	69	23	46	*	*18	*18	*28	*12	*20	30	*
727.0	33	*	*	*	*	*	*	*	*	*	*
727.05	*17	*	*	*	*	*	*	*	*	*	*
728	247	93	154	*	*56	*54	134	*78	*61	79	*
728.8	137	59	*79	*	*55	*	*50	*	*33	*43	*
728.89	84	*41	*44	*	*	*	*43	*	*	*	*
729	122	42	80	*	24	34	61	*20	*15	63	*23
729.1	26	*	*17	*	*	*	*	*	*	*	*
729.5	57	*19	38	*	*	*15	*32	*	*	*30	*
729.8	28	*	*20	*	*	*	*	*	*	*	*
730	624	417	207	*	83	171	343	119	108	290	*107
730.0	200	*112	*88	*	*	*	*122	*	*	*	*
730.1	137	*96	*	*	*	*	*58	*	*	*	*
730.2	287	209	*78	*	*	*66	163	*80	*	142	*
730.27	123	*75	*	*	*	*	*87	*	*	*	*
730.28	*87	*	*	*	*	*	*	*	*	*	*
732	*16	*13	*	*8	*	*	*	*	*	*	*
733	707	221	487	*	67	128	502	137	147	251	173
733.1	472	122	349	*	*	*50	416	84	100	155	133
733.13	323	70	253	—	*	*	305	*51	68	98	107
733.14	67	*	*49	*	*	*	*48	*	*	*	*
733.4	80	46	*34	*	*20	*39	*	*	*	*30	*
733.42	70	41	*29	*	*	*34	*	*	*	*26	*
733.6	29	*	22	—	*	*10	*14	*	*	*	*
733.8	85	38	47	*	30	*23	*26	*19	*	36	*16
733.81	*18	*	*	*	*	*	*	*	*	*	*
733.82	66	28	39	*	25	*18	*	*15	*	*31	*
736	29	*17	*	*	*	*	*	*	*	*	*
737	74	*	*58	*41	*	*	*	*	*	*	*
737.3	71	*	*57	*40	*	*	*	*	*	*	*
737.30	*47	*	*39	*	*	*	*	*	*	*	*
738	101	*35	67	*	*34	*31	*35	*	*	*32	*24
738.4	88	*29	58	—	*27	*28	*33	*	*	*25	*
740-759	1,048	499	550	823	128	58	*39	295	152	332	269
745	242	83	159	215	*	*	*	64	*	*76	*77
745.4	*29	*	*	*27	*	*	*	*	*	*	*
745.5	*34	*	*25	*	*	*	*	*	*	*	*
746	120	*57	*64	*94	*	*	*	*	*	*	*
747	98	*45	52	*56	*	*	*	*	*	*38	*
749	45	28	*17	41	*	*	*	*	*	30	*
749.2	31	*21	*	29	*	*	*	*	*	*23	*
749.20	*22	*	*	*21	*	*	—	*	—	*21	*
750	57	*35	*	57	*	—	—	*	*	*	*
750.5	*19	*14	*	*19	—	—	—	*	*	*	*
751	134	*76	*	117	*	*	*	*	*	*	*
752	*19	*	*	*	*	*	*	*	*	*	*
753	*21	*	*	*	*	*	*	*	*	*	*
754	40	*19	*20	30	*	—	*	*	*	*	*
755	*18	*	*	*	*	*	—	*	*	*	*

See footnotes and notes at end of table.

Table 43. Number of days of care for discharges from short-stay hospitals, by ICD-9-CM code of first-listed diagnosis, sex, age, and geographic region: United States, 1999—Con.

[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table: see Methods section]

ICD-9-CM code	Sex			Age				Region			
	Total	Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	North-east	Midwest	South	West
Number of days of care in thousands											
756	127	69	*58	*79	*	*	*	*	*	*56	*
756.1	*41	*	*	*	*	*	*	*	*	*	*
756.12	*27	*	*	—	*	*	*	*	*	*	*
757	18	*6	*12	*10	*	—	*	*	*	17	—
757.3	14	*6	*8	*10	*	—	—	*	—	13	—
757.32	13	*5	*8	*9	*	—	—	*	—	13	—
759	*21	*	*	*	*	*	*	*	*	*	*
760-779	1,604	963	640	1,603	*	—	—	236	309	528	531
765	541	351	*190	541	—	—	—	*	*	261	*186
765.1	389	283	*106	389	—	—	—	*	*	*180	*130
765.18	*83	*	*	*83	—	—	—	*	*	*	*
769	385	195	*190	385	—	—	—	*	*	*85	*132
770	252	173	79	252	—	—	—	*39	*54	*68	91
770.6	*45	*	*	*45	—	—	—	*	*	*	*
770.8	83	*48	*35	83	—	—	—	*	*	*	*34
771	56	*36	*	56	—	—	—	*	*	*	*
771.8	49	*31	*	49	—	—	—	*	*	*	*
774	81	52	30	81	—	—	—	*	*15	26	31
774.6	71	46	25	71	—	—	—	*	*12	*22	29
775	*28	*	*	*28	—	—	—	*	*	*	*
778	28	*15	*	28	—	—	—	*	*	*	*
778.4	*22	*	*	*22	—	—	—	*	*	*	*
779	80	*64	*	80	—	—	—	*	*	*	*
779.3	*44	*	*	*44	—	—	—	*	*	*	*
780-799	998	456	541	188	255	204	351	144	130	506	218
780	130	63	68	53	35	32	*10	39	23	51	*19
780.2	20	*8	*11	*	*	*	*	*	*	*10	—
780.3	55	23	32	26	*19	*	*	*20	*	21	*
780.39	50	20	31	21	*19	*	*	*18	*	19	*
780.6	30	*15	*15	24	*	*	*	*	*	*14	*
784	*12	*	*	*	*	*	*	*	*	*	*
786	164	84	80	*10	48	59	48	25	22	101	17
786.0	*11	*	*	*	*	*	*	*	*	*	*
786.5	118	57	61	—	46	44	28	22	19	66	12
786.50	78	41	37	—	24	31	24	14	*10	47	*7
786.59	32	12	20	—	*18	*10	*	*	*	*16	*
787	*16	*	*	*	*	*	*	*	*	*	*
789	134	41	93	19	69	*20	*	*17	*12	69	36
789.0	133	41	92	19	68	*20	*	*17	*12	69	36
789.00	60	*19	*41	*	*28	*	*	*	*	*36	*
789.03	36	*10	26	*	27	*	—	*	*	*18	*7
789.06	*10	*	*	*	*	*	*	*	*	*	*
799	506	240	266	*82	80	84	259	*50	66	251	139
799.9	505	240	265	*82	80	84	258	*50	66	250	139
800-999	13,831	6,651	7,180	1,009	3,471	2,865	6,486	3,201	2,827	5,288	2,516
800	59	*29	*	*15	*	*	*	*	*	*	*
801	202	158	*44	*	80	*	*	*84	*31	*57	*
801.0	*27	*17	*	*	*	*	*	*	*	*	*
801.2	*74	*	*	*	*	*	*	*	*	*	*
802	103	78	*25	*	51	*19	*27	*24	*28	30	*21
802.2	27	*21	*	*	*18	*	*	*	*	*	*
802.4	*29	*24	*	*	*	*	*	*	*	*	*

See footnotes and notes at end of table.

Table 43. Number of days of care for discharges from short-stay hospitals, by ICD-9-CM code of first-listed diagnosis, sex, age, and geographic region: United States, 1999—Con.[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table: see Methods section]

ICD-9-CM code	Sex		Age				Region				
	Total	Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	North-east	Midwest	South	West
Number of days of care in thousands											
803	*75	*	*	*	*	*	*	*	*	*	*
805	426	183	243	*	103	79	242	92	113	120	101
805.0	121	*70	*52	*	*42	*	*	*	*	*	*
805.2	110	*35	75	*	*23	*	78	*	*27	*40	*
805.4	171	67	104	*	*31	*37	103	*36	48	51	*37
806	*95	*	*	—	*	*	*	*	*	*	*
807	166	70	96	*	*24	68	73	*52	*31	40	*44
807.0	143	53	89	*	*16	66	59	*50	*30	34	*28
807.00	*15	*	*	—	*	*	*	*	*	*	*
808	417	154	263	*	85	*	299	86	98	158	75
808.0	85	*55	*30	*	*40	*	*30	*	*	*	*
808.2	263	82	181	*	*32	*	220	*62	46	101	*54
810	*19	*13	*	*	*	*	*	*	*	*	*
810.0	*18	*12	*	*	*	*	*	*	*	*	*
812	255	103	153	21	33	37	165	54	58	81	62
812.0	154	60	94	*	*	*21	120	*33	*35	43	*
812.00	*53	*	*	*	*	*	*48	*	*	*	*
812.01	60	*	49	—	*	*	*42	*	*	*	*
812.09	*32	*	*	*	*	*	*25	*	*	*	*
812.2	*32	*	*	*	*	*	*	*	*	*	*
812.21	*25	*	*	*	*	*	*	*	*	*	*
812.4	49	20	29	17	*	*	*	*13	*	*16	*9
812.41	31	*14	*17	14	*	*	*	*	*	*	*
813	172	72	100	*11	52	29	81	41	31	71	29
813.0	35	*	*23	*	*	*	*22	*	*	*	*
813.01	*21	*	*	*	*	*	*	*	*	*	*
813.4	71	28	43	*	*18	*13	37	*19	*13	26	*14
813.41	*24	*	*21	*	*	*	*20	*	*	*	*
813.42	26	*12	*15	*	*9	*	*	*	*	*9	*
813.44	*18	*10	*	*	*	*	*	*	*	*	*
816	*20	*	*	*	*	*	*	*	*	*	*
816.1	*17	*	*	*	*	*	*	*	*	*	*
820	2,115	565	1,550	*	50	119	1,939	504	466	809	336
820.0	541	112	429	*	*	*47	485	132	117	195	97
820.00	*42	*	*	—	*	*	*31	*	*	*	*
820.02	65	*	*47	*	*	*	*62	*	*	*	*
820.03	*54	*	*42	—	*	*	*47	*	*	*	*
820.09	376	66	309	*	*	*	343	86	95	133	61
820.2	1,034	268	765	*	*32	*46	954	272	223	376	162
820.20	*42	*	*33	—	*	*	*36	*	*	*	*
820.21	910	239	672	*	*	*36	854	241	200	322	147
820.22	81	*	61	*	*	*	64	*	*	*40	*
820.8	523	174	349	*	*	*24	486	95	121	236	72
821	400	217	183	75	70	*92	163	80	97	144	79
821.0	211	151	60	58	*42	*	82	*40	51	101	*
821.00	*82	*	*	*	*	*	*	*	*	*	*
821.01	129	90	*39	*43	*39	*	*28	*33	*43	*39	*
821.2	148	*40	107	*	*	*53	80	*35	*39	*32	*
821.20	*65	*	*	*	*	*	*	*	*	*	*
821.23	53	*	*47	*	*	*	*35	*	*	*	*
822	70	*	50	*	*	*	*28	*	*	*38	*
822.0	59	*	43	*	*	*	*28	*	*	*30	*

See footnotes and notes at end of table.

Table 43. Number of days of care for discharges from short-stay hospitals, by ICD-9-CM code of first-listed diagnosis, sex, age, and geographic region: United States, 1999—Con

[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table: see Methods section]

ICD-9-CM code	Sex		Age				Region				
	Total	Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	North-east	Midwest	South	West
Number of days of care in thousands											
823	380	229	151	*12	214	92	62	89	67	153	70
823.0	108	47	61	*	53	*25	*	*	*24	*44	*25
823.00	82	*36	46	*	37	*	*	*	*	*33	*
823.2	65	42	*23	*	47	*	*	*15	*	*	*22
823.22	53	35	*	*	39	*	*	*	*	*	*
823.3	*65	*62	*	*	*47	*	*	*	*	*	*
823.32	*57	*54	*	*	*	*	*	*	*	*	*
823.8	81	51	*30	*	*40	*	*	*31	*	*33	*
823.82	61	*44	*17	*	*29	*	*	*	*	*27	*
823.9	*47	*	*	*	*	*	*	*	*	*	*
823.92	*38	*	*	*	*	*	*	*	*	*	*
824	383	123	260	*	119	94	162	80	73	139	91
824.0	*18	*	*	*	*	*	*	*	*	*	*
824.2	*19	*	*	*	*	*	*	*	*	*	*
824.4	119	27	92	*	25	*26	68	*25	*23	*25	*46
824.6	92	*	81	*	*21	*27	*43	*	*20	*39	*17
824.8	78	45	33	*	36	*19	*19	*20	*16	28	*
825	54	43	*	*	*28	*20	*	*	*	*18	*
825.0	*21	*19	*	*	*	*	*	*	*	*	*
825.2	*17	*	*	*	*	*	*	*	*	*	*
840	38	15	23	*	*	20	*10	*	*	16	*10
840.4	35	*13	22	*	*	19	*8	*	*	16	*
844	37	20	*17	*	18	*	*	*	*	*10	*
844.2	18	*11	*7	*	13	*	—	*	*	*	*
845	*21	*	*	—	*	*	*	*	*	*	*
845.0	*21	*	*	—	*	*	*	*	*	*	*
847	45	*13	32	*	*20	*	*	*	*	*21	*
847.0	*17	*	*	*	*	*	*	*	*	*	*
847.2	*26	*	*	*	*	*	*	*	*	*	*
850	96	64	32	*16	36	*	*34	37	*29	20	*
850.0	*14	*	*	*	*	*	*	*	*	*	*
850.1	27	17	*	*	18	*	*	*12	*	*	*
850.5	*22	*	*	*	*	*	*	*	*	*	*
850.9	*33	*25	*	*	*	*	*	*	*	*	*
851	177	126	*50	*	*89	*	*53	*	*39	*49	*53
851.8	140	102	*37	*	*76	*	*39	*	*	*	*
852	405	206	199	*	185	*64	145	*60	*69	179	97
852.0	*81	*	*58	*	*	*	*	*	*	*	*
852.2	225	100	125	*	*62	*37	120	*	*44	*78	*74
852.20	90	*56	*	*	*	*	*38	*	*	*42	*
852.21	*81	*	*	*	*	*	*54	*	*	*	*
853	*67	*	*	*	*	*	*	*	*	*	*
853.0	*66	*	*	*	*	*	*	*	*	*	*
854	76	*27	*48	*	38	*	*	*	*25	*	*
854.0	70	*22	*48	*	34	*	*	*	*25	*	*
860	246	145	100	*	108	*	*113	*	59	141	*
860.0	89	*58	*	*	*29	*	*	*	*	*	*
860.4	*79	*	*	*	*	*	*	*	*	*	*
861	*45	*36	*	*	*	*	*	*	*	*	*
863	99	91	*	*	*68	*	*	*	*	*48	*
864	94	*38	*	*	*72	*	*	*	*	*	*
864.0	*73	*	*	*	*	*	*	*	*	*	*
865	76	57	*	*	*32	*	*	*	*	*	*
865.0	70	*52	*	*	*27	*	*	*	*	*	*

See footnotes and notes at end of table.

Table 43. Number of days of care for discharges from short-stay hospitals, by ICD-9-CM code of first-listed diagnosis, sex, age, and geographic region: United States, 1999—Con.[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table: see Methods section]

ICD-9-CM code	Sex			Age				Region			
	Total	Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	North-east	Midwest	South	West
Number of days of care in thousands											
868	*45	*	*	*	*	*	*	*	*	*	*
873	55	27	28	*	*14	*	*18	*12	*	*22	*11
873.4	*20	*	*	*	*	*	*	*	*	*	*
879	48	32	*	*	*23	*	*	*	*	37	*
879.8	*19	*	*	*	*	*	*	—	*	*	—
881	*19	*16	*	*	*15	*	*	*	*	*	*
882	*17	*	*	*	*	*	*	*	*	*	*
883	*20	*	*	*	*	*	*	*	*	*	*
886	*9	*	*	*	*	*	*	*	*	*	*
891	74	38	*37	*	*25	*	*	*	*	*32	*
891.0	*31	*17	*	*	*	*	*	*	*	*	*
891.1	*33	*	*	*	*	*	*	*	*	*	*
892	*28	*	*	*	*	*	*	*	*	*	*
920	*19	*	*	*	*	*	*	*	*	*	*
922	68	36	32	*	*	*	47	*	*31	*17	*
922.1	31	*18	*13	—	*	*	*24	*	*	*	*
924	83	*21	61	*	*	*	63	*	*25	*	*23
924.0	*39	*	*29	*	*	*	*32	*	*	*	*
924.01	*28	*	*	—	*	*	*27	*	*	*	*
935	25	*17	*	*	*	*	*	*	*	*	*
935.1	*21	*	*	*	*	*	*	*	*	*	*
941	*17	*	*	*	*	*	*	*	*	*	*
942	*74	*	*	*	*	*	*	*	*	*	*
943	*45	*	*	*	*	*	*	*	*	*	*
944	*44	*38	*	*	*	*	—	*	*	*	*
945	88	*62	*	*	*	*	*	*	*	*	*
945.2	*32	*	*	*	*	*	*	*	*	*	*
945.3	*51	*	*	*	*	*	*	*	*	*	*
948	275	176	*99	223	*	*	*	*	—	266	—
948.0	*22	*	*	*17	—	*	*	*	—	*13	—
948.00	*22	*	*	*17	—	*	*	*	—	*13	—
952	*38	*	*	*	*	*	*	*	*	*	*
958	*38	*33	*	*	*	*	*	*	*	*	*
959	105	50	55	*10	31	*	54	*22	20	43	20
959.0	66	29	38	*	20	*	*34	*	*16	*23	*14
959.01	65	28	37	*	19	*	*34	*	*15	*22	*14
959.1	*12	*	*	*	*	*	*	*	*	*	*
962	*17	*14	*	*	*	*	*	*	*	*	*
962.3	*15	*14	*	*	*	*	*	*	*	*	*
963	*9	*	*	*	*	*	—	*	*	*	*
963.0	*8	*	*	*	*	*	—	*	*	*	*
965	102	39	64	*	54	*16	*	*26	21	30	*25
965.0	35	*	*13	*	*20	*	*	*	*	*13	*
965.09	*15	*	*	*	*	*	*	*	*	*	*
965.4	40	*	28	*	24	*	*	*	*11	*11	*
966	*29	*	*23	*	*	*	*	*	*	*	*
966.3	*16	*	*	*	*	*	*	*	*	*	*
968	19	*13	*	—	*14	*	—	*	*	*	*
968.5	*14	*	*	—	*	*	—	*	*	*	*
969	112	46	67	*	63	32	*	*21	20	42	30
969.0	35	18	16	*	21	*	*	*	*	*11	*
969.4	56	*19	37	*	25	*19	*	*	*	24	*13

See footnotes and notes at end of table.

Table 43. Number of days of care for discharges from short-stay hospitals, by ICD-9-CM code of first-listed diagnosis, sex, age, and geographic region: United States, 1999—Con.

[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table: see Methods section]

ICD-9-CM code	Sex			Age				Region			
	Total	Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	North-east	Midwest	South	West
Number of days of care in thousands											
972	*25	*	*	*	*	*	*	*	*	*	*
977	18	*7	*11	*	*11	*	*	*	*	*10	*
977.9	*10	*	*	*	*	*	*	*	*	*	*
989	*13	*	*	*	*	*	*	*	*	*	*
995	80	23	57	*	*15	*17	*18	*	*14	26	*
995.0	*8	*	*	*	*	*	*	*	*	*	*
995.1	*16	*	*	*	*	*	*	*	*	*	*
995.2	*23	*	*	*	*	*	*	*	*	*	*
996	2,605	1,306	1,299	150	467	741	1,247	688	471	960	487
996.0	163	88	75	*	*	*47	74	*26	*18	*68	*
996.01	81	*23	*58	*	*	*	29	*	*	*	*
996.04	*20	*	*	—	*	*	*	*	*	*	*
996.1	51	*30	*20	*	*	*	*	*	*	*	*
996.2	121	60	61	36	48	*	*	*	39	*38	*
996.3	*23	*	*	*	*	*	*	*	*	*	*
996.4	362	156	206	*	*40	100	221	79	103	91	89
996.5	63	*23	*40	*	*	*	*28	*	*	*	*
996.59	*37	*	*	*	*	*	*	*	*	*	*
996.6	1,096	564	531	*79	205	303	509	291	184	441	179
996.61	*74	*	*	—	*	*	*	*	*	*	*
996.62	536	240	296	*45	103	157	232	165	76	202	*94
996.63	*96	*	*	*	*	*	*	*	*	*	*
996.64	*74	*	*	—	*	*	*	*	*	*	*
996.66	126	*	*	—	*	*	*104	*	*	*	*
996.67	80	*	*	*	*	*	*	*	*	*	*
996.68	*33	*	*	*	*	*	*	*	*	*	*
996.69	*53	*	*	*	*	*	*	*	*	*	*
996.7	589	303	287	*	59	166	352	173	81	233	102
996.72	*37	*	*	*	*	*	*	*	*	*	*
996.73	180	80	100	*	*21	63	95	*33	*21	79	47
996.74	218	136	83	*	*	*52	155	86	*36	86	*
996.77	*31	*	*	—	*	*	*20	*	*	*	*
996.78	50	*26	*25	*	*	*	*	*	*	*17	*
996.8	138	*64	74	*	*60	66	*	*45	*	*52	*
996.81	82	*42	*40	*	*44	*32	*	*	*	*33	*
997	554	275	279	*	84	208	253	164	98	181	112
997.0	*32	*	*	*	*	*	*	*	*	*	*
997.1	33	*15	*	*	*	*	*26	*	*	*	*
997.2	*42	*	*	—	*	*	*	*	*	*	*
997.3	41	*33	*	*	*	*	*	*	*	*	*
997.4	190	67	123	*	43	65	77	*35	*34	55	66
997.5	48	*22	*	*	*	*	*23	*	*	*	*
997.6	145	86	*59	*	*	*71	*62	*	*	*55	*
997.62	104	*62	*	*	*	*	*	*	*	*	*
997.69	*36	*	*	—	*	*	*	*	*	*	*
998	1,317	600	718	48	264	425	581	340	249	473	257
998.1	187	89	98	*9	54	32	91	47	52	53	35
998.11	104	59	44	*9	*25	*16	54	32	*29	*21	*21
998.12	72	*25	*47	*	*26	*	*33	*	*	*26	*
998.2	57	*	*45	*	*	*	*	*	*	*	*
998.3	81	*	*42	*	*	*	*	*	*	*	*
998.5	889	416	473	*	161	327	377	218	156	344	170
998.59	876	410	467	*	156	324	373	218	150	341	167
998.6	*36	*	*	*	*	*	*	*	*	*	*
998.8	62	*28	*34	*	*	*	*23	*	*	*24	*
998.89	31	*	*16	*	*	*	*	*	*	*	*
999	*56	*	*	*	*	*	*	*	*	*	*
999.3	*49	*	*	*	*	*	*	*	*	*	*

See footnotes and notes at end of table.

Table 43. Number of days of care for discharges from short-stay hospitals, by ICD-9-CM code of first-listed diagnosis, sex, age, and geographic region: United States, 1999—Con.

[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table: see Methods section]

ICD-9-CM code	Sex			Age				Region			
	Total	Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	North-east	Midwest	South	West
Number of days of care in thousands											
V01-V82	17,340	3,444	13,896	332	10,233	1,712	5,064	3,271	3,924	6,764	3,380
V10	30	*	*13	*	*	*	*20	*	*	*	*
V12	36	*11	25	*	*	*9	*20	*	*	*19	*
V12.5	22	*	*16	—	*	*	*16	*	*	*13	*
V12.59	*12	*	*	—	*	*	*	*	*	*	*
V15	*18	*12	*	*	*	*	*	*	*	*	*
V15.8	*14	*	*	*	*	*	*	*	*	*	*
V17	*7	*	*	—	*	*	*	*	*	*	*
V22	*19	—	*19	—	*19	—	—	*	*	*	*
V27	9,421	—	9,421	*18	9,379	*23	—	1,736	1,995	3,581	2,109
V27.0	8,635	—	8,635	*18	8,601	*	—	1,587	1,652	3,388	2,008
V27.1	43	—	43	—	43	—	—	*	*	*20	*
V27.2	144	—	144	—	143	*	—	*32	*26	53	*33
V27.9	584	—	584	*	578	*	—	103	306	115	60
V30	49	39	*	49	—	—	—	*	*	*	*
V30.1	49	39	*	49	—	—	—	*	*	*	*
V42	*64	*	*	*	*	*	*	*	*	*	*
V43	*29	*	*	*	*	*	*	*	*	*	*
V45	59	41	*17	*	*	*11	*39	*11	*	*27	*
V45.8	46	35	*	*	*	*	*33	*	*	*	*
V45.89	*33	*	*	*	*	*	*	*	*	*	*
V53	18	*12	*	*	*	*	*13	*	*	*	*
V53.3	16	*11	*	*	*	*	*13	*	*	*	*
V53.31	*11	*	*	*	*	*	*10	*	*	*	*
V54	69	*	*50	*	*	*	*	*	*	*	*
V55	224	135	89	*	*43	79	88	62	*41	83	*39
V55.2	*36	*	*	*	*	*	*	*	*	*	*
V55.3	176	113	64	*	*30	70	*70	*46	*	70	*29
V57	5,872	2,395	3,478	*	455	1,136	4,232	1,029	1,573	2,461	809
V57.1	1,533	605	929	*	*95	317	1,099	446	167	820	99
V57.8	4,200	1,746	2,454	*	358	805	3,008	534	1,401	1,594	670
V57.89	4,197	1,746	2,451	*	358	805	3,005	534	1,399	1,593	670
V57.9	118	*	*81	—	*	*	114	*	*	*	*
V58	1,029	532	497	162	189	277	402	236	203	340	250
V58.0	103	*59	*45	*	*	*37	*54	*	*	69	*
V58.1	739	406	333	155	152	202	231	211	116	214	199
V58.4	81	36	44	*	*22	*	*44	*13	*	*24	*
V58.49	74	*31	43	*	*17	*	*43	*13	*	*23	*
V58.8	101	*	*70	*	*	*	*72	*	*	*	*
V58.89	100	*	*70	*	*	*	*72	*	*	*	*
V59	*22	*	*	*	*	*	*	*	*	*	*
V66	*120	*	*	—	*	*	*	*	*	*	*
V70	26	*18	*8	*	*15	*	*	*	*	*	*
V70.7	22	*15	*	*	*12	*	*	*	*	*	*
V71	30	*10	20	*	*9	*	*	*	*	*	*14
V71.7	*14	*	*11	*	*	*	*	*	*	*	*

* Figure does not meet standard of reliability or precision.

— Quantity zero.

NOTE: If estimate is preceded by asterisk, use according to size of corresponding estimate in table 42; see Methods section.

Table 44. Number of all-listed diagnoses for discharges from short-stay hospitals, by ICD-9-CM code, sex, age, and geographic region: United States, 1999

[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table: see Methods section]

ICD-9-CM code	Sex			Age				Region			
	Total	Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	North-east	Midwest	South	West
Number of all-listed diagnoses in thousands											
All codes	135,765	54,901	80,864	6,393	33,320	30,008	66,044	29,665	33,022	48,432	24,646
001-139	3,554	1,552	2,002	465	884	701	1,504	788	778	1,340	647
003	*8	*	*	*	*	*	*	*	*	*	*
003.0	*5	*	*	*	*	*	*	*	*	*	*
008	263	116	147	89	38	37	98	67	63	85	48
008.4	108	40	68	*7	13	21	68	35	26	29	18
008.45	102	38	64	*6	12	20	65	32	25	27	17
008.6	44	27	18	42	*	*	*	*7	15	14	*9
008.61	43	26	17	42	*	*	-	*6	15	13	*8
008.8	103	46	58	37	23	15	28	24	21	39	19
009	21	10	11	*	*	*5	*8	*	*5	*8	*
009.0	16	*8	*8	*	*	*	*7	*	*	*6	*
011	15	11	*	*	*6	*	*	*	*	*7	*
011.9	12	10	*	*	*	*	*	*	*	*6	*
011.90	*6	*	*	*	*	*	*	*	*	*	*
031	12	*7	*	*	*	*	*	*	*	*	*
031.0	*6	*	*	*	*	*	*	*	*	*	*
034	29	14	15	15	10	*	*	*6	10	10	*
034.0	27	14	13	14	10	*	*	*6	*8	*9	*
038	692	329	363	50	78	134	429	165	141	268	118
038.0	56	27	29	*5	*6	11	34	12	14	22	*8
038.1	132	73	59	*6	18	33	75	31	28	50	23
038.10	24	11	12	*	*	*6	13	*8	*5	*8	*
038.11	75	44	31	*	12	21	40	18	14	28	15
038.19	34	18	16	*	*	*6	22	*	*9	14	*6
038.2	27	14	13	*	*6	*6	12	*7	*5	11	*
038.4	140	56	84	*6	15	32	87	38	32	43	28
038.40	18	*7	11	*	*	*	10	*5	*	*6	*
038.42	71	23	48	*	*7	13	49	18	18	22	13
038.43	13	*8	*	*	*	*	*8	*	*	*	*
038.49	35	15	20	*	*6	10	18	*9	*7	11	*8
038.8	27	10	16	*	*	*	18	*8	*	*9	*
038.9	305	146	159	27	29	50	199	66	57	132	50
041	1,253	456	797	104	282	222	644	227	314	475	236
041.0	250	82	168	14	89	40	107	46	62	98	44
041.00	24	11	13	*	*	*6	13	*	*	9	*6
041.01	12	*6	*6	*	*5	*	*	*	*	*	*
041.02	81	12	70	*	63	*	13	19	18	32	11
041.04	102	39	63	*	11	21	66	13	27	43	19
041.09	28	12	16	*	*5	*6	13	*7	*8	10	*
041.1	244	131	113	21	59	59	105	47	55	96	46
041.10	35	16	19	*	*6	10	16	9	*8	13	*5
041.11	167	91	76	15	44	38	70	30	38	67	33
041.19	42	24	18	*	*9	11	19	*8	9	16	*8
041.3	79	20	59	*6	*9	10	55	15	17	32	16
041.4	327	68	259	32	63	46	186	60	81	114	72
041.5	12	*8	*	*	*	*	*6	*	*	*	*
041.6	44	12	32	*	*5	*5	32	*7	12	19	*6
041.7	84	43	41	9	12	17	46	14	22	35	13
041.8	190	83	107	14	37	39	100	31	57	66	35
041.84	16	*6	9	*	*	*5	*6	*	*5	*	*
041.85	59	23	37	*7	10	*9	33	13	17	18	12
041.86	77	38	39	*	15	18	44	12	23	30	13
041.89	31	13	18	*	*7	*7	14	*	*9	14	*5
041.9	18	*6	12	*	*7	*	*6	*	*	12	*

See footnotes and notes at end of table.

Table 44. Number of all-listed diagnoses for discharges from short-stay hospitals, by ICD-9-CM code, sex, age, and geographic region: United States, 1999—Con.[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table: see Methods section]

ICD-9-CM code	Sex			Age				Region			
	Total	Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	North-east	Midwest	South	West
Number of all-listed diagnoses in thousands											
042	135	91	44	*	98	33	*	49	15	51	20
047	31	15	16	13	15	*	*	*	*5	13	*8
047.9	29	15	14	12	14	*	*	*	*	13	*7
052	12	*5	*7	*8	*	*	*	*	*	*	*
052.9	*7	*	*	*	*	*	*	*	*	*	*
053	42	14	28	*	*	*7	31	10	10	13	*9
053.1	12	*	*9	*	*	*	10	*	*	*	*
053.19	9	*	*7	*	*	*	*8	*	*	*	*
053.9	24	*9	15	*	*	*	16	*5	*6	*8	*5
054	55	12	43	10	26	*	15	10	12	22	11
054.1	14	*	12	*	12	*	*	*	*	*7	*
054.10	10	*	*8	—	*8	*	*	*	*	*	*
054.2	10	*	*8	*5	*	*	*	*	*	*	*
054.7	*5	*	*	*	*	*	*	*	*	*	*
054.9	19	*	15	*	10	*	*7	*	*	*7	*
057	9	*5	*	*7	*	*	*	*	*	*	*5
057.9	*7	*	*	*6	*	*	*	*	*	*	*
070	204	117	86	*	87	88	27	52	27	82	43
070.1	*8	*	*	*	*	*	*	*	*	*	*
070.3	33	20	13	*	14	11	*7	9	*5	14	*
070.30	20	11	*9	*	10	*5	*	*6	*	*8	*
070.32	12	*8	*	*	*	*	*	*	*	*5	*
070.4	*6	*	*	*	*	*	*	*	*	*	*
070.5	149	86	63	*	66	68	15	39	19	58	33
070.51	73	43	30	—	33	33	*7	16	10	29	18
070.54	75	43	32	*	33	35	*8	23	9	29	15
070.9	*6	*	*	*	*	*	*	*	*	*	*
075	11	*	*6	*	*7	*	—	*	*	*	*
078	31	13	18	*5	18	*5	*	*6	*6	15	*
078.1	17	*6	11	*	12	*	*	*	*	*9	*
078.10	11	*	*7	*	*8	*	*	*	*	*7	*
078.5	*8	*	*	*	*	*	*	*	*	*	*
079	178	75	103	85	42	18	32	29	38	70	40
079.6	10	*6	*	10	*	*	—	*	*	*5	*
079.8	*7	*	*	*	*	*	*	*	*	*	*
079.89	*5	*	*	*	*	*	*	*	*	*	*
079.9	153	65	89	69	36	17	31	26	34	60	34
079.99	151	65	86	69	33	17	31	26	33	59	32
094	*5	*	*	—	*	*	*	*	*	*	*
097	*8	*	*6	—	*5	*	*	*	*	*	*
098	*6	*	*	*	*6	*	*	*	*	*	*
110	58	33	26	*	11	14	28	17	16	18	*7
110.1	28	14	14	*	*	*8	18	*8	*8	*8	*
110.3	*6	*	*	*	*	*	*	*	*	*	*
110.4	15	11	*	*	*	*	*6	*	*	*	*
110.5	*6	*	*	*	*	*	*	*	*	*	*
111	*7	*	*	*	*	*	*	*	*	*	*
111.9	*6	*	*	*	*	*	*	*	*	*	*
112	266	112	154	37	60	59	110	66	55	99	47
112.0	138	62	76	22	32	34	51	32	30	51	25
112.1	17	—	17	*	*9	*	*	*6	*	*7	*
112.2	25	12	12	*	*	*6	13	*6	*	11	*
112.3	21	*7	13	*9	*	*	*9	*	*5	*8	*
112.4	12	*6	*6	*	*	*	*9	*	*	*	*

See footnotes and notes at end of table.

Table 44. Number of all-listed diagnoses for discharges from short-stay hospitals, by ICD-9-CM code, sex, age, and geographic region: United States, 1999—Con.

[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table: see Methods section]

ICD-9-CM code	Sex			Age				Region			
	Total	Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	North-east	Midwest	South	West
Number of all-listed diagnoses in thousands											
112.5	*9	*	*	*	*	*	*	*	*	*	*
112.8	36	17	19	*	9	*8	18	10	*7	14	*6
112.84	28	15	13	*	*6	*6	15	*8	*	11	*
112.89	*7	*	*5	*	*	*	*	*	*	*	*
112.9	*9	*	*7	*	*	*	*	*	*	*	*
117	24	11	13	*	*9	*5	*7	*8	*	*9	*
117.3	*7	*	*	*	*	*	*	*	*	*	*
117.5	*7	*	*	*	*6	*	*	*	*	*	*
117.9	10	*	*6	*	*	*	*	*	*	*	*
131	12	*	11	*	9	*	*	*	*	*6	*
131.0	9	*	*9	—	*8	*	*	*	*	*	*
131.01	*6	—	*6	—	*5	*	*	*	*	*	*
135	37	12	25	—	15	15	*7	9	9	15	*
136	31	20	11	*	20	*	*5	9	*6	11	*5
136.3	20	14	*5	*	14	*	*	*6	*	*7	*
136.9	10	*5	*	*	*	*	*	*	*	*	*
138	23	14	9	*	*	*6	16	*7	*6	*5	*
140-239	4,432	1,888	2,544	126	623	1,378	2,305	1,073	1,042	1,531	785
141	9	*6	*	*	*	*5	*	*	*	*	*
146	*6	*5	*	—	*	*	*	*	*	*	—
150	34	27	*7	*	*	10	23	13	*8	10	*
150.5	9	*8	*	—	*	*	*7	*	*	*	*
150.9	16	13	*	—	*	*	11	*7	*	*	*
151	46	31	15	—	*	14	30	10	12	11	13
151.0	12	10	*	—	*	*	*8	*	*	*	*
151.8	*6	*	*	—	*	*	*5	*	*	*	*
151.9	14	*9	*6	—	*	*	*8	*	*	*	*
152	*8	*6	*	—	*	*	*6	*	*	*	*
153	186	87	99	*	10	47	128	40	51	67	27
153.0	*8	*	*5	—	*	*	*	*	*	*	*
153.1	15	*9	*6	—	*	*	11	*	*	*	*
153.2	12	*	*8	—	*	*	*7	*	*5	*	*
153.3	34	15	19	—	*	9	23	9	9	10	*6
153.4	28	11	17	—	*	*	22	*7	*6	9	*6
153.6	24	10	13	—	*	*	18	*	*8	*7	*
153.8	*7	*	*	—	*	*	*	*	*	*	*
153.9	51	27	24	*	*	13	32	10	12	24	*
154	73	41	32	*	*	19	49	20	15	24	13
154.0	23	13	10	—	*	*7	16	*7	*	*8	*
154.1	43	25	19	—	*	10	31	11	11	14	*8
155	30	20	11	*	*	9	17	*8	*	12	*6
155.0	21	14	*7	*	*	*6	11	*	*	*8	*
155.2	*6	*	*	*	*	*	*	*	*	*	*
156	18	*6	12	—	*	*6	13	*6	*	*6	*
156.0	*7	*	*6	—	—	*	*	*	*	*	*
156.1	*7	*	*	—	—	*	*7	*	*	*	*
157	56	26	30	—	*	18	34	13	13	20	*9
157.0	19	*9	11	—	*	*7	11	*	*5	*8	*
157.9	28	14	14	—	*	*8	18	*6	*7	10	*
158	*9	*	*6	*	*	*	*	*	*	*	*
161	15	9	*6	—	*	*7	*6	*	*	*6	*
161.9	*5	*	*	—	*	*	*	*	*	*	—

See footnotes and notes at end of table.

Table 44. Number of all-listed diagnoses for discharges from short-stay hospitals, by ICD-9-CM code, sex, age, and geographic region: United States, 1999—Con.[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table: see Methods section]

ICD-9-CM code	Sex			Age				Region			
	Total	Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	North-east	Midwest	South	West
Number of all-listed diagnoses in thousands											
162	370	213	157	*	*6	114	250	82	91	149	48
162.2	16	*8	*8	—	*	*6	11	*	*	*7	*
162.3	80	43	37	*	*	22	57	24	16	27	14
162.4	9	*6	*	—	*	*	*5	*	*	*	*
162.5	38	21	17	*	*	11	26	*7	11	12	*8
162.8	33	20	13	—	*	11	22	16	9	*	*
162.9	193	114	78	*	*	60	129	30	50	95	18
163	*7	*	*	—	*	*	*7	*	*	*	*
170	30	22	*9	12	14	*	*	10	*	*	13
170.7	18	14	*	*7	11	*	*	*6	*	*	*9
171	17	*8	9	*	*	*	*5	*	*	*6	*
172	15	*7	*8	*	*	*5	*8	*	*	*	*
172.9	*5	*	*	—	*	*	*	*	*	*	*
173	23	12	11	*	*	*	16	*7	*	*7	*
173.3	*7	*	*	—	*	*	*6	*	*	*	*
174	170	—	170	—	20	65	85	46	42	59	23
174.2	*7	—	*7	—	*	*	*	*	*	*	*
174.4	24	—	24	—	*	10	12	*8	*6	*7	*
174.8	41	—	41	—	*7	19	15	16	*9	9	*6
174.9	84	—	84	—	10	28	46	17	23	37	*8
180	29	—	29	*	10	13	*5	*6	*5	12	*6
180.8	*7	—	*7	—	*	*	*	*	*	*	*
180.9	17	—	17	*	*5	*9	*	*	*	*8	*
182	45	—	45	—	*	14	27	10	10	18	*7
182.0	44	—	44	—	*	14	27	10	10	18	*7
183	49	—	49	*	*5	19	24	11	10	17	10
183.0	47	—	47	*	*5	19	23	11	10	17	10
184	*9	—	*9	*	*	*	*6	*	*	*	*
185	223	223	—	—	*	43	178	49	50	87	37
186	*5	*5	—	*	*	*	*	*	*	*	*
188	69	51	18	—	*	*8	61	20	19	20	10
188.2	10	*8	*	—	*	*	*9	*	*	*	*
188.4	*5	*5	*	—	—	*	*	*	*	*	*
188.8	14	9	*	—	—	*	13	*6	*	*	*
188.9	32	22	10	—	*	*	28	*8	*9	12	*
189	47	27	20	*	*	12	28	11	12	14	10
189.0	39	22	16	*	*	11	21	10	10	12	*7
191	48	26	22	10	10	15	13	10	13	16	*8
191.1	*8	*	*	*	*	*	*	*	*	*	*
191.2	*7	*	*	*	*	*	*	*	*	*	*
191.8	*7	*	*	*	*	*	*	*	*	*	*
191.9	12	*6	*6	*	*	*	*	*	*	*6	*
192	*7	*	*5	*	*	*	*	*	*	*	*
193	26	*5	20	*	12	*9	*6	*	*	10	*7
194	11	*	*6	*6	*	*	*	*	*	*5	*
194.0	*7	*	*	*5	*	*	*	*	*	*	*
195	17	10	*8	*	*	*5	*8	*8	*	*6	*
195.0	*7	*7	*	—	*	*	*	*	*	*	*
196	179	75	104	*	20	68	90	42	46	53	38
196.0	30	17	13	*	*6	14	9	*5	*9	10	*6
196.1	22	12	10	*	*	*8	13	*6	*	*6	*6
196.2	63	29	34	*	*	23	37	13	17	19	14
196.3	37	*	33	*	*6	17	14	11	*8	12	*6
196.5	*6	*	*	*	*	*	*	*	*	*	*
196.6	10	*6	*	—	*	*	*7	*	*	*	*
196.9	*8	*	*	—	*	*	*5	*	*	*	*

See footnotes and notes at end of table.

Table 44. Number of all-listed diagnoses for discharges from short-stay hospitals, by ICD-9-CM code, sex, age, and geographic region: United States, 1999—Con.

[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table: see Methods section]

ICD-9-CM code	Sex			Age				Region			
	Total	Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	North-east	Midwest	South	West
Number of all-listed diagnoses in thousands											
197	501	216	285	*6	42	167	286	131	125	156	88
197.0	108	49	59	*	19	37	48	31	29	24	24
197.1	10	*6	*	*	*	*	*	*	*	*	*
197.2	60	18	42	*	*	19	39	15	11	23	11
197.4	15	*	11	*	*	*	10	*	*	*5	*
197.5	15	*	10	*	*	*	10	*	*	*	*
197.6	63	26	37	*	*6	19	38	18	14	20	11
197.7	207	97	110	*	12	73	121	50	56	70	30
197.8	20	*9	12	—	*	*5	14	*6	*5	*7	*
198	420	206	214	*6	26	147	240	121	111	113	75
198.1	10	*	*6	*	*	*	*5	*	*	*	*
198.2	*9	*	*6	—	*	*	*	*	*	*	*
198.3	94	46	48	*	*6	40	46	30	23	27	15
198.4	9	*	*	*	*	*	*	*	*	*	*
198.5	188	101	87	*	*8	56	120	54	52	45	37
198.6	*5	—	*5	—	*	*	*	*	*	*	*
198.7	12	*8	*	—	*	*5	*6	*	*	*	*
198.8	89	38	51	*	*9	31	48	25	22	25	16
198.82	13	*	10	—	*	*	9	*	*	*	*
198.89	75	34	40	*	*8	28	39	21	18	22	14
199	151	62	88	*	12	47	91	29	37	56	29
199.0	51	18	33	*	*	18	29	*7	12	19	13
199.1	100	44	55	*	*8	29	62	22	24	37	16
200	30	15	15	*	*	*8	18	*6	*5	*8	10
200.0	18	9	9	*	*	*5	11	*	*	*	*6
200.00	*7	*	*	*	*	*	*	*	*	*	*
201	19	*8	12	*	*6	*5	*7	*	*8	*6	*
201.9	17	*6	10	*	*	*5	*7	*	*6	*	*
201.90	12	*	*7	*	*	*	*6	*	*	*	*
202	116	68	48	*	14	39	62	28	27	36	25
202.8	106	61	44	*	12	34	59	26	26	32	22
202.80	69	42	27	*	*8	22	38	16	18	22	13
202.81	*7	*	*	*	*	*	*	*	*	*	*
202.83	12	*5	*6	*	*	*	*7	*	*	*	*
202.88	*7	*	*	—	*	*	*	*	*	*	*
203	71	33	38	—	*	20	48	19	16	23	13
203.0	70	33	37	—	*	20	47	19	16	22	13
203.00	67	31	36	—	*	19	44	18	15	22	12
204	99	52	47	29	*8	9	53	27	22	31	19
204.0	40	21	19	28	*8	*	*	10	*8	12	10
204.00	27	15	12	17	*6	*	*	*8	*6	9	*
204.01	13	*6	*7	11	*	*	*	*	*	*	*6
204.1	57	30	28	—	*	*6	51	17	14	18	*9
204.10	55	28	27	—	*	*6	49	16	13	17	*9
205	60	35	25	9	9	12	29	17	12	20	11
205.0	41	23	18	*7	*7	*9	19	13	*7	14	*6
205.00	35	20	16	*	*6	*7	18	11	*7	13	*
205.01	*6	*	*	*	*	*	*	*	*	*	*
205.1	17	11	*6	*	*	*	9	*	*	*5	*
205.10	15	9	*6	*	*	*	9	*	*	*	*
208	10	*6	*	*	*	*	*6	*	*	*	*
210	9	*	*	*	*	*	*	*	*	*	*
210.2	*7	*	*	—	*	*	*	*	*	*	*
211	165	79	86	*	9	44	112	37	37	64	27
211.1	17	*8	10	—	*	*6	10	*	*	*7	*
211.3	131	64	67	*	*	33	93	29	29	51	22
211.4	*8	*	*	—	*	*	*6	*	*	*	*

See footnotes and notes at end of table.

Table 44. Number of all-listed diagnoses for discharges from short-stay hospitals, by ICD-9-CM code, sex, age, and geographic region: United States, 1999—Con.[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table: see Methods section]

ICD-9-CM code	Sex		Age				Region				
	Total	Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	North-east	Midwest	South	West
Number of all-listed diagnoses in thousands											
212	*9	*	*5	*	*	*	*	*	*	*	*
213	*5	*	*	*	*	*	*	*	*	*	*
214	16	*8	*8	*	*	*6	*7	*5	*	*	*
216	17	*	12	*	*8	*	*	*	*	*6	*
216.5	*9	*	*7	*	*	*	*	*	*	*	*
218	437	—	437	—	224	192	21	83	94	172	88
218.0	50	—	50	—	24	24	*	12	10	18	9
218.1	123	—	123	—	63	54	*5	26	26	44	26
218.2	61	—	61	—	32	26	*	13	12	21	15
218.9	203	—	203	—	104	88	10	31	45	89	38
219	*7	—	*7	—	*	*	*	*	*	*	*
220	58	—	58	*	26	21	10	11	12	23	13
221	*6	—	*6	—	*	*	*	*	*	*	*
223	*6	*	*	—	*	*	*	*	*	*	*
225	38	12	27	*	*7	13	17	11	*8	14	*6
225.1	*6	*	*	—	*	*	*	*	*	*	*
225.2	26	*6	19	—	*	10	13	*7	*	10	*
226	10	*	*7	*	*	*	*	*	*	*	*
227	19	*8	11	*	*	*8	*7	*5	*	*7	*
227.1	*8	*	*5	—	*	*	*	*	*	*	*
227.3	*8	*	*	*	*	*	*	*	*	*	*
228	21	*7	14	*8	*	*	*6	*	*	10	*
228.0	20	*7	14	*7	*	*	*6	*	*	10	*
228.04	*6	*	*	*	*	*	*	*	*	*	*
230	*6	*	*	—	*	*	*	*	*	*	*
233	42	*	40	—	18	13	11	*9	*8	18	*7
233.0	18	*	18	—	*	*7	*9	*	*	*7	*
233.1	20	—	20	—	16	*	*	*	*	9	*
235	18	*8	10	*	*	*	11	*	*	*6	*
235.2	12	*	*7	—	*	*	*7	*	*	*5	*
236	*7	*	*6	*	*	*	*	*	*	*	*
237	12	*	*7	*	*	*	*	*	*	*	*
237.7	*6	*	*	*	*	*	*	*	*	*	*
238	70	32	38	*	*	*8	55	18	19	22	11
238.4	18	*8	10	*	*	*	13	*	*7	*5	*
238.7	45	20	24	*	*	*	39	11	11	14	*8
239	35	12	24	*	*	11	19	*7	*6	18	*
239.0	*7	*	*6	—	*	*	*6	*	*	*	*
239.5	*7	*	*	*	*	*	*	*	*	*5	*
239.6	*8	*	*	*	*	*	*	*	*	*	*
240-279	14,053	5,801	8,252	536	1,702	3,916	7,899	3,112	3,559	4,961	2,421
240	11	*	10	—	*	*	*6	*	*	*	*
240.9	11	*	10	—	*	*	*6	*	*	*	*
241	27	*	24	*	*	12	11	*8	*7	*8	*
241.0	*7	*	*7	*	*	*	*	*	*	*	*
241.1	14	*	12	*	*	*6	*6	*	*	*6	*
241.9	*5	*	*	*	*	*	*	*	*	*	*
242	62	14	48	*	18	17	27	19	12	22	10
242.0	16	*	13	*	*7	*	*	*	*	*	*
242.00	15	*	13	*	*6	*	*	*	*	*	*
242.9	43	11	32	*	10	11	21	13	*8	17	*
242.90	41	10	31	*	*9	11	20	12	*8	16	*

See footnotes and notes at end of table.

Table 44. Number of all-listed diagnoses for discharges from short-stay hospitals, by ICD-9-CM code, sex, age, and geographic region: United States, 1999—Con.

[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table: see Methods section]

ICD-9-CM code	Sex			Age				Region			
	Total	Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	North-east	Midwest	South	West
Number of all-listed diagnoses in thousands											
244	1,224	234	989	*	120	284	815	283	347	361	232
244.0	45	*5	39	*	*5	14	26	*9	12	11	13
244.1	12	*	10	-	*	*6	*	*	*	*	*
244.8	*5	*	*	*	*	*	*	*	*	*	*
244.9	1,159	225	934	*	112	262	782	271	330	343	215
245	13	*	11	*	*5	*	*	*	*	*	*
245.2	10	*	*8	*	*	*	*	*	*	*	*
246	17	*	15	*	*	*	10	*	*	*8	*
246.8	*5	*	*	*	*	*	*	*	*	*	*
246.9	11	*	10	-	*	*	*6	*	*	*	*
250	4,571	2,063	2,508	30	461	1,494	2,586	1,063	1,051	1,722	735
250.0	3,236	1,428	1,808	18	253	1,009	1,956	744	745	1,251	496
250.00	2,345	1,063	1,282	*	145	727	1,469	538	554	879	375
250.01	530	213	318	*8	61	164	297	130	118	213	69
250.02	267	119	148	*	28	88	151	55	57	113	41
250.03	93	33	60	*	20	30	39	21	16	45	11
250.1	128	65	62	10	72	30	16	29	33	45	20
250.10	21	12	10	*	10	*6	*5	*7	*	*9	*
250.11	43	21	21	*5	28	*8	*	*9	13	11	10
250.12	17	*7	10	*	*6	*6	*	*	*	*7	*
250.13	46	25	21	*	28	10	*	9	11	19	*7
250.2	19	*9	11	*	*	*7	9	*6	*	*7	*
250.20	*5	*	*	-	*	*	*	*	*	*	*
250.22	10	*	*6	*	*	*	*	*	*	*	*
250.3	*9	*	*	*	*	*	*	*	*	*	*
250.4	294	137	157	-	34	121	139	62	57	111	64
250.40	157	71	86	-	*6	67	84	28	28	61	40
250.41	98	49	48	-	21	40	36	28	20	31	18
250.42	20	*9	11	-	*	*6	14	*	*	10	*
250.43	20	*8	11	-	*6	*7	*6	*	*	*9	*
250.5	139	61	78	-	22	58	58	40	31	39	29
250.50	58	27	31	-	*	24	32	15	12	16	15
250.51	54	25	29	-	15	24	15	16	13	13	11
250.52	12	*	*9	-	*	*	*8	*	*	*5	*
250.53	15	*6	*9	-	*	*6	*	*6	*	*	*
250.6	342	154	188	*	37	139	166	82	87	113	60
250.60	177	82	94	-	10	65	101	41	45	57	34
250.61	98	42	56	*	18	43	37	28	27	27	16
250.62	39	16	23	-	*	18	19	*8	*8	18	*6
250.63	29	14	15	-	*7	13	*8	*6	*7	12	*
250.7	158	83	76	-	*7	49	102	43	35	55	26
250.70	84	44	40	-	*	23	59	23	18	29	14
250.71	41	21	21	-	*	16	23	12	*8	14	*7
250.72	23	13	10	-	*	*6	16	*	*6	*9	*
250.73	10	*5	*	-	*	*	*	*	*	*	*
250.8	204	104	100	*	19	68	116	49	52	74	30
250.80	94	49	46	-	*6	28	60	24	23	30	17
250.81	49	26	23	*	*6	19	23	12	12	18	*7
250.82	38	16	21	*	*	13	24	*7	10	18	*
250.83	23	13	10	*	*6	*9	*8	*6	*7	*8	*
250.9	41	18	23	*	10	12	19	*6	*8	24	*
250.90	9	*	*5	*	*	*	*5	*	*	*7	*
250.92	19	*6	13	*	*	*5	10	*	*	10	*
250.93	10	*6	*	*	*	*	*	*	*	*6	*
251	63	26	37	*8	10	15	30	17	18	18	10
251.2	27	12	16	*6	*	*	12	*7	*8	*9	*
251.8	28	10	18	*	*	9	14	*7	*8	*8	*

See footnotes and notes at end of table.

Table 44. Number of all-listed diagnoses for discharges from short-stay hospitals, by ICD-9-CM code, sex, age, and geographic region: United States, 1999—Con.[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table: see Methods section]

ICD-9-CM code	Sex			Age				Region			
	Total	Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	North-east	Midwest	South	West
Number of all-listed diagnoses in thousands											
252	30	10	20	*	*	*9	17	*6	*5	13	*6
252.0	22	*8	14	—	*	*7	14	*	*	10	*
252.1	*8	*	*6	*	*	*	*	*	*	*	*
253	73	35	38	*6	13	19	35	15	21	21	17
253.2	17	11	*6	*	*	*7	*6	*	*6	*	*
253.5	12	*	*7	*	*	*	*	*	*	*	*
253.6	36	15	21	*	*	*6	26	*8	*9	11	*8
255	54	19	35	*	12	15	24	11	15	19	10
255.0	11	*	*8	*	*5	*	*	*	*	*6	*
255.4	27	11	16	*	*	*7	14	*6	10	*7	*
255.9	*6	*	*	—	*	*	*	*	*	*	*
256	*6	—	*6	—	*6	*	*	*	*	*	*
259	*8	*	*	*	*	*	*	*	*	*	*
260	11	*	*7	*	*	*	*6	*	*	*6	*
261	27	13	14	*	*	*	20	*9	*6	9	*
262	*6	*	*	*	*	*	*	*	*	*	*
263	365	162	203	*7	26	66	266	67	91	149	59
263.0	*7	*	*	*	*	*	*5	*	*	*	*
263.1	10	*	*7	*	*	*	*9	*	*	*5	*
263.8	27	12	15	*	*	*	18	*	*9	9	*6
263.9	320	143	178	*5	22	60	234	57	80	132	52
266	34	14	21	*	*	*	28	10	10	*9	*5
266.2	31	12	19	*	*	*	25	*9	9	*8	*
269	*9	*	*	*	*	*	*6	*	*	*	*
269.9	*5	*	*	*	*	*	*	*	*	*	*
270	14	*7	*8	*	*	*	*	*	*	*	*
270.6	*6	*	*	*	*	*	*	*	*	*	*
271	22	*7	14	*	*	*5	*9	*9	*	*8	*
271.3	18	*5	13	*	*	*	*9	*7	*	*6	*
272	1,649	850	799	*	113	644	890	421	443	507	277
272.0	826	423	403	*	56	324	444	262	215	234	115
272.1	29	15	14	*	*8	13	*8	*7	11	*8	*
272.2	14	*6	*9	—	*	*7	*6	*	*	*5	*
272.4	753	390	363	*	43	288	422	144	212	248	149
272.9	11	*8	*	*	*	*	*5	*	*	*	*
273	53	26	27	*	*5	10	35	10	15	17	11
273.1	*6	*	*	—	—	*	*	*	*	*	*
273.3	*5	*	*	—	*	*	*	*	*	*	*
273.8	38	17	21	*	*5	*6	24	*6	11	12	*9
274	197	134	63	*	12	54	131	51	49	61	36
274.0	54	36	18	—	*5	16	33	15	13	20	*6
274.9	140	96	44	*	*6	38	97	36	35	41	29
275	188	81	107	*8	38	50	92	36	53	69	30
275.0	*7	*	*	*	*	*	*	*	*	*	*
275.2	70	30	40	*	17	15	37	12	28	20	10
275.3	21	*8	13	*	*	*7	*8	*	*	*7	*
275.4	88	38	50	*	13	25	45	17	20	37	14
275.41	32	14	18	*	*9	10	10	*	*9	11	*7
275.42	47	20	27	*	*	14	29	10	*8	23	*6
275.49	*8	*	*	*	*	*	*6	*	*	*	*
276	4,458	1,771	2,688	416	588	830	2,625	893	1,173	1,609	783
276.0	104	47	57	*6	*8	15	74	23	27	36	18
276.1	670	258	412	17	56	137	460	117	185	231	137
276.2	208	96	112	27	31	50	100	40	48	75	45
276.3	43	15	28	*	*8	9	24	9	11	14	*8
276.4	14	*6	*8	*	*	*	*8	*	*	*	*

See footnotes and notes at end of table.

Table 44. Number of all-listed diagnoses for discharges from short-stay hospitals, by ICD-9-CM code, sex, age, and geographic region: United States, 1999—Con.[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table: see Methods section]

ICD-9-CM code	Sex			Age				Region			
	Total	Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	North-east	Midwest	South	West
Number of all-listed diagnoses in thousands											
276.5	2,171	877	1,294	330	304	317	1,220	459	563	783	366
276.6	70	32	37	*	*6	21	41	*8	16	31	15
276.7	276	129	147	*7	25	58	186	56	74	102	44
276.8	817	271	545	21	137	202	456	171	216	299	131
276.9	86	39	47	*	11	17	55	9	31	32	15
277	48	24	24	18	20	*	*7	13	13	14	*8
277.0	26	12	14	13	13	*	*	*7	*8	*7	*
277.00	26	12	14	13	13	*	*	*7	*8	*7	*
277.3	*6	*	*	—	*	*	*	*	*	*	*
277.4	*5	*	*	*	*	*	*	*	*	*	*
278	774	266	508	10	216	351	198	143	200	271	160
278.0	761	263	497	9	212	342	197	140	198	265	158
278.00	529	197	332	*8	138	234	149	100	141	173	115
278.01	232	66	166	*	74	108	48	40	56	92	43
278.1	9	*	*7	*	*	*5	*	*	*	*	*
279	24	13	11	*6	*7	*	*7	*	*	*9	*7
279.0	9	*6	*	*	*	*	*	*	*	*	*
280-289	4,170	1,683	2,487	176	849	861	2,284	908	981	1,483	798
280	635	216	419	15	106	129	384	141	169	223	102
280.0	380	133	248	*	58	83	237	86	99	137	59
280.8	*5	*	*	*	*	*	*	*	*	*	*
280.9	248	80	168	12	48	45	143	53	69	84	42
281	101	36	65	*	9	15	75	19	29	37	16
281.0	36	10	26	*	*	*5	28	*7	12	12	*
281.1	*9	*	*	—	*	*	*7	*	*	*	*
281.9	53	20	33	*	*5	*8	39	9	14	20	10
282	124	49	74	27	80	10	*8	40	22	50	12
282.4	21	*6	15	*	11	*	*5	*8	*	*7	*
282.5	16	*	13	*	12	*	*	*	*	*7	*
282.6	80	37	44	20	54	*	*	25	16	35	*
282.60	14	*7	*7	*	*9	*	*	*	*	*6	*
282.62	60	26	34	12	45	*	*	18	10	28	*
283	23	*7	16	*	*	*5	11	*8	*	*7	*
283.0	10	*	*7	*	*	*	*7	*	*	*	*
283.9	*8	*	*6	*	*	*	*	*	*	*	*
284	183	100	84	14	26	52	91	43	38	58	44
284.8	166	91	74	11	23	49	82	37	35	51	42
284.9	17	*8	*9	*	*	*	9	*6	*	*6	*
285	2,186	829	1,357	43	434	413	1,295	444	510	797	434
285.1	670	268	402	*5	122	128	414	125	180	215	150
285.8	31	14	17	*	*	*7	20	*	*9	9	*8
285.9	1,481	545	936	37	308	278	857	314	321	571	276
286	156	80	76	*8	30	41	76	30	32	55	39
286.0	*7	*5	*	*	*	*	*	*	*	*	*
286.3	*8	*	*	*	*	*	*	*	*	*	*
286.4	10	*	*6	*	*	*	*	*	*	*	*
286.5	11	*	*6	*	*	*	*5	*	*	*5	*
286.6	24	15	9	*	*7	*5	10	*	*6	*8	*6
286.7	30	14	16	*	*	*8	19	*6	*6	*8	10
286.9	65	35	30	*	14	19	31	10	13	27	15
287	387	188	199	21	78	104	184	100	83	125	79
287.0	*6	*	*	*	*	*	*	*	*	*	*
287.3	39	15	24	*6	11	*8	15	11	*7	12	*8
287.4	70	39	31	*	14	24	29	20	12	19	18
287.5	263	129	134	9	50	69	134	64	61	89	49

See footnotes and notes at end of table.

Table 44. Number of all-listed diagnoses for discharges from short-stay hospitals, by ICD-9-CM code, sex, age, and geographic region: United States, 1999—Con.[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table: see Methods section]

ICD-9-CM code	Sex			Age				Region			
	Total	Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	North-east	Midwest	South	West
Number of all-listed diagnoses in thousands											
288	271	128	143	32	54	62	123	63	65	95	47
288.0	168	80	88	23	28	42	75	43	40	57	29
288.8	91	40	50	*8	23	18	42	17	23	35	16
289	106	51	55	13	26	30	38	20	29	36	21
289.0	*9	*7	*	*	*	*	*	*	*	*	*
289.2	10	*	*6	*5	*	*	*	*	*	*	*
289.3	9	*	*5	*	*	*	*	*	*	*	*
289.4	*6	*	*	*	*	*	*	*	*	*	*
289.5	12	*6	*6	*	*	*	*	*	*	*	*
289.59	9	*	*	*	*	*	*	*	*	*	*
289.8	31	15	16	*	*7	12	11	*6	*9	10	*6
289.9	27	10	17	*	*	*5	18	*5	*8	10	*
290-319	8,927	4,367	4,560	284	3,846	2,312	2,486	2,241	2,193	3,088	1,405
290	246	85	161	*	*	*7	238	56	61	97	31
290.0	132	39	94	*	—	*	131	36	28	52	16
290.2	18	*	13	*	*	—	18	*	*7	*7	*
290.20	*7	*	*	*	*	—	*7	*	*	*	—
290.21	11	*	*9	—	—	—	11	*	*	*	*
290.3	18	*5	13	—	—	*	18	*	*7	*	*
290.4	74	36	38	—	*	*5	68	13	19	32	10
290.40	57	27	30	—	*	*	52	11	13	23	10
290.41	*8	*	*	—	—	*	*8	*	*	*	*
291	228	181	47	—	108	85	35	62	67	57	42
291.0	38	33	*5	—	17	15	*7	13	*6	13	*6
291.2	19	16	*	—	*	*9	10	*	*7	*	*5
291.3	*6	*	*	—	*	*	*	*	*	*	*
291.8	158	123	35	—	86	57	16	44	48	38	28
291.81	151	117	34	—	81	54	15	43	45	36	27
291.89	*8	*7	*	—	*	*	*	*	*	*	*
292	152	81	71	*	81	32	38	50	34	43	25
292.0	69	41	28	*	48	17	*	31	13	16	*9
292.1	12	*	*7	*	*6	*	*	*	*	*	*
292.12	10	*	*6	*	*	*	*	*	*	*	*
292.8	65	32	33	*	22	13	29	16	15	22	12
292.81	39	16	23	*	*	*8	27	*7	*9	14	10
292.84	19	12	*7	*	15	*	*	*7	*	*6	*
292.9	*6	*	*	—	*	*	*	*	*	*	*
293	120	57	63	*	19	19	81	34	34	31	21
293.0	79	35	44	*	*5	10	63	17	23	23	16
293.8	29	15	14	*	13	*7	*8	15	*7	*	*
293.83	22	12	10	*	11	*5	*5	13	*5	*	*
293.9	10	*6	*	—	*	*	*7	*	*	*	*
294	620	202	418	*	9	24	586	147	151	216	105
294.1	241	79	162	—	*	*7	232	63	57	81	40
294.8	357	113	245	*	*	14	340	80	89	128	60
294.9	19	*9	10	*	*	*	12	*	*	*6	*5
295	416	225	191	*	206	128	80	132	79	150	55
295.3	115	65	50	*	59	37	19	37	18	42	17
295.30	40	22	18	—	21	12	*7	*8	11	14	*7
295.32	24	13	11	—	9	*9	*6	10	*	*9	*
295.34	50	29	20	*	29	15	*6	19	*6	19	*7
295.6	40	25	15	*	17	14	*9	16	*7	13	*
295.60	11	*6	*	*	*	*	*	*	*	*	*
295.62	15	9	*6	—	*5	*5	*	*7	*	*5	*
295.64	14	*9	*	—	*8	*5	*	*	*	*	*
295.7	152	72	79	*	93	41	16	48	32	54	17
295.70	104	48	56	*	63	29	12	27	26	40	12
295.72	*5	*	*	—	*	*	*	*	*	*	*

See footnotes and notes at end of table.

Table 44. Number of all-listed diagnoses for discharges from short-stay hospitals, by ICD-9-CM code, sex, age, and geographic region: United States, 1999—Con.

[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table: see Methods section]

ICD-9-CM code	Sex			Age				Region			
	Total	Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	North-east	Midwest	South	West
Number of all-listed diagnoses in thousands											
295.74	41	22	19	*	27	10	*	19	*5	12	*
295.9	98	56	43	*	32	32	34	26	20	36	16
295.90	83	45	38	*	24	28	31	17	19	33	14
295.92	*6	*	*	—	*	*	*	*	*	*	*
295.94	*8	*6	*	—	*6	*	*	*6	*	*	*
296	1,043	421	622	51	568	264	159	265	258	415	105
296.0	*6	*	*	*	*	*	*	*	*	*	*
296.00	*5	*	*	*	*	*	*	*	*	*	*
296.2	299	117	182	23	156	61	57	57	77	142	23
296.20	169	62	107	11	86	37	35	36	44	75	14
296.22	30	13	17	*	17	*6	*	*	*	23	*
296.23	59	26	33	*7	31	11	11	10	17	29	*
296.24	38	15	23	*	21	*7	*7	10	12	14	*
296.3	313	113	200	10	173	78	52	88	82	114	29
296.30	102	35	67	*	50	28	20	31	26	35	*9
296.32	23	*9	14	*	13	*	*	*	*8	10	*
296.33	121	47	74	*	76	25	16	32	33	47	10
296.34	62	20	42	*	30	20	10	20	13	20	*8
296.4	62	30	32	*	34	21	*6	21	14	21	*7
296.40	28	12	16	*	16	*8	*	*6	*7	12	*
296.43	*6	*	*	—	*	*	*	*	*	*	*
296.44	24	13	11	*	13	9	*	12	*5	*	*
296.5	76	31	45	*	46	20	9	20	22	28	*5
296.50	46	16	29	*	26	12	*7	9	13	20	*
296.53	16	*8	*9	*	12	*	*	*7	*	*	*
296.54	11	*5	*6	*	*7	*	*	*	*	*	*
296.6	68	29	39	*	40	19	*	19	16	28	*5
296.60	41	19	23	*	24	12	*	*7	9	23	*
296.63	11	*	*6	*	*7	*	*	*5	*	*	*
296.64	15	*6	10	*	*8	*	*	*7	*	*	*
296.7	147	64	83	*7	73	45	22	36	31	56	25
296.8	32	13	19	*	19	10	*	10	*7	*9	*6
296.80	16	*7	9	*	*7	*7	*	*	*	*6	*
296.89	12	*	*8	*	9	*	*	*	*	*	*
296.9	38	22	16	*	23	*9	*	14	*7	14	*
296.90	34	19	15	*	21	*7	*	13	*6	13	*
297	26	10	17	*	*6	*	15	*7	*7	11	*
297.1	*7	*	*	—	*	*	*	*	*	*	*
297.9	19	*6	12	*	*	*	13	*	*	9	*
298	165	77	89	*	51	30	81	41	39	60	25
298.9	158	74	84	*	48	27	79	39	38	58	23
299	11	*9	*	*5	*6	*	—	*	*	*	*
299.0	*8	*5	*	*	*	*	—	*	*	*	*
299.00	*7	*5	*	*	*	*	—	*	*	*	*
300	756	238	518	16	248	216	276	184	204	262	106
300.0	415	127	288	*	102	131	177	95	114	151	56
300.00	340	100	240	*	68	110	159	79	93	123	45
300.01	58	20	38	*	31	15	13	13	16	21	*8
300.02	16	*6	10	*	*	*6	*5	*	*	*7	*
300.1	25	*	20	*	17	*	*	*	*6	*9	*5
300.11	12	*	10	*	*7	*	*	*	*	*	*
300.15	*7	*	*6	*	*7	*	—	*	*	*	*
300.2	19	*8	11	*	11	*6	*	*5	*	*	*
300.21	11	*	*6	*	*7	*	*	*	*	*	*
300.3	29	14	14	*	18	*6	*	*8	*9	*8	*
300.4	198	55	143	*	54	55	84	60	51	63	23
300.8	*8	*	*7	*	*	*	*	*	*	*	*
300.81	*5	*	*	*	*	*	*	*	*	*	*
300.9	62	28	34	*	42	12	*5	9	16	23	13

See footnotes and notes at end of table.

Table 44. Number of all-listed diagnoses for discharges from short-stay hospitals, by ICD-9-CM code, sex, age, and geographic region: United States, 1999—Con.[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table: see Methods section]

ICD-9-CM code	Sex			Age				Region			
	Total	Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	North-east	Midwest	South	West
Number of all-listed diagnoses in thousands											
301	309	136	174	*9	213	66	21	122	81	79	27
301.1	*5	*	*	*	*	*	*	*	*	*	*
301.2	9	*5	*	*	*5	*	*	*	*	*	*
301.20	*8	*	*	*	*	*	*	*	*	*	*
301.3	*7	*6	*	*	*	*	*	*	*	*	*
301.4	*5	*	*	*	*	*	*	*	*	*	*
301.5	*7	*	*6	*	*	*	*	*	*	*	*
301.50	*6	*	*5	*	*	*	*	*	*	*	*
301.6	27	10	17	*	13	11	*	*6	12	*8	*
301.7	26	20	*6	*	21	*	*	*7	12	*	*
301.8	142	51	92	*	108	24	*6	64	35	31	12
301.81	11	*7	*	*	*5	*	*	*	*	*	*
301.83	82	14	68	*	66	13	*	29	23	22	*7
301.84	*5	*	*	*	*	*	*	*	*	*	*
301.89	44	26	18	*	34	*7	*	32	*	*	*
301.9	76	35	42	*	51	19	*5	39	11	20	*7
303	724	521	203	*	362	276	85	195	180	227	121
303.0	105	76	29	*	55	40	9	20	38	24	24
303.00	34	24	11	*	21	11	*	*5	11	*9	9
303.01	67	50	17	*	32	28	*7	14	26	14	13
303.9	619	445	175	*	306	236	76	175	143	204	98
303.90	278	196	82	*	128	107	42	36	69	133	40
303.91	288	211	77	*	159	104	25	124	61	59	44
303.92	11	*7	*	*	*6	*	*	*	*	*	*
303.93	42	30	12	*	13	22	*7	12	*9	9	12
304	435	251	184	*	329	92	11	155	75	144	62
304.0	146	84	62	*	98	43	*	67	18	34	27
304.00	42	24	18	*	27	12	*	*	*6	21	10
304.01	98	57	41	*	68	28	*	59	11	12	16
304.03	*5	*	*	—	*	*	—	*	*	*	*
304.1	28	10	17	—	15	9	*	*	*5	14	*5
304.10	15	*	12	—	*7	*6	*	*	*	*9	*
304.11	12	*7	*	—	*8	*	*	*	*	*5	*
304.2	105	69	37	—	90	15	*	37	18	44	*6
304.20	45	29	16	—	37	*8	*	*	*8	30	*
304.21	54	36	18	—	47	*7	*	32	*7	13	*
304.3	51	34	18	*	45	*	—	*8	19	20	*5
304.30	22	13	10	*	19	*	—	*	*	15	*
304.31	25	18	*7	*	22	*	—	*5	13	*	*
304.4	11	*6	*	—	9	*	*	*	*	*	*5
304.40	*7	*	*	—	*6	*	*	—	*	*	*
304.6	11	*6	*5	—	*7	*	*	*	*	*	*
304.60	*6	*	*	—	*	*	*	*	*	*	*
304.7	28	16	12	—	22	*5	*	15	*	*5	*
304.70	*5	*	*	—	*	*	—	*	*	*	*
304.71	22	13	*9	—	17	*	*	14	*	*	*
304.8	24	12	12	*	20	*	*	12	*	*5	*
304.80	*8	*5	*	*	*6	*	—	*	*	*	*
304.81	11	*5	*6	*	10	*	*	*7	*	*	*
304.9	30	15	15	—	20	*8	*	*6	*7	12	*5
304.90	20	10	10	—	12	*6	*	*	*6	9	*
304.91	*7	*	*	—	*6	*	*	*	*	*	*
305	2,110	1,198	912	10	1,033	737	330	375	526	776	433
305.0	409	292	116	*	226	127	53	92	84	160	72
305.00	293	214	79	*	164	90	37	46	69	125	54
305.01	75	51	23	—	41	25	9	29	10	25	11
305.02	14	9	*	*	11	*	*	*9	*	*	*
305.03	26	18	*9	*	11	9	*6	*9	*	*8	*5
305.1	1,242	637	605	*	447	528	265	168	341	454	279

See footnotes and notes at end of table.

Table 44. Number of all-listed diagnoses for discharges from short-stay hospitals, by ICD-9-CM code, sex, age, and geographic region: United States, 1999—Con.

[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table: see Methods section]

ICD-9-CM code	Sex			Age				Region			
	Total	Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	North-east	Midwest	South	West
Number of all-listed diagnoses in thousands											
305.2	106	64	42	*	93	10	*	30	29	36	12
305.20	71	42	29	*	63	*6	*	12	23	29	*7
305.21	21	13	*8	*	19	*	—	12	*	*	*
305.22	*8	*5	*	—	*7	*	*	*	*	*	*
305.23	*6	*	*	—	*	*	*	*	*	*	*
305.4	15	*7	*8	—	*9	*5	*	*	*	*7	*
305.40	12	*6	*6	—	*6	*	*	*	*	*6	*
305.5	51	30	21	*	36	15	*	15	12	16	*7
305.50	32	19	13	*	22	10	*	*6	*9	12	*5
305.51	15	*8	*7	—	10	*	*	*8	*	*	*
305.6	158	94	64	*	128	28	*	41	35	58	25
305.60	109	63	47	*	90	17	*	17	29	43	20
305.61	38	24	14	—	28	*9	*	19	*	13	*
305.62	*6	*	*	*	*5	*	—	*	*	*	*
305.63	*6	*	*	—	*	*	*	*	*	*	*
305.7	25	15	10	*	22	*	*	*	*	*6	16
305.70	19	10	*8	*	16	*	*	*	*	*	13
305.9	99	57	43	*	69	21	*7	25	20	37	18
305.90	69	40	29	*	47	14	*6	13	16	28	12
305.91	13	*7	*6	*	10	*	*	*7	*	*	*
305.93	14	*7	*7	*	9	*	*	*	*	*5	*
306	11	*	*8	*	*	*	*	*	*	*	*
306.1	*5	*	*	—	*	*	*	*	*	*	*
307	74	25	49	*7	36	14	17	17	18	29	10
307.1	*8	*	*8	*	*6	*	*	*	*	*	*
307.2	*7	*6	*	*	*	*	*	*	*	*	*
307.23	*6	*	*	*	*	*	*	*	*	*	*
307.5	18	*	15	*	14	*	*	*	*5	*6	*
307.50	*6	*	*	*	*	*	*	*	*	*	*
307.51	*8	*	*8	*	*8	*	*	*	*	*	*
307.8	22	*6	16	*	10	*8	*	*	*	12	*8
307.81	18	*5	13	*	*8	*6	*	*	*	10	*
307.9	12	*7	*5	*	*	*	9	*	*	*	*
308	20	*7	13	*	11	*	*	*	*	11	*
308.0	*6	*	*5	*	*	*	*	*	*	*	*
308.9	9	*	*5	*	*6	*	*	*	*	*	*
309	229	92	138	20	145	41	24	79	59	60	31
309.0	68	28	40	*	41	14	10	23	19	17	9
309.2	31	14	17	*	18	*	*5	*8	*8	11	*
309.28	23	9	14	*	15	*	*	*5	*7	*8	*
309.4	25	12	12	*	16	*	*	10	*7	*5	*
309.8	79	26	53	*8	52	16	*	30	20	19	11
309.81	75	23	52	*8	51	14	*	29	19	19	9
309.9	18	9	9	*	12	*	*	*5	*5	*5	*
310	77	31	46	*	*6	*6	64	18	15	35	*9
310.1	*6	*	*	*	*	*	*	*	*	*	*
310.2	*5	*	*	*	*	*	*	*	*	*	*
310.9	64	23	41	*	*	*	59	15	11	33	*6
311	795	285	510	18	253	212	312	202	179	265	149
312	67	47	21	21	37	*	*	30	19	13	*5
312.3	41	29	12	11	28	*	*	20	11	*6	*
312.30	23	16	*7	*	17	*	*	15	*5	*	*
312.34	13	11	*	*5	*8	*	*	*	*	*	*
312.8	*7	*	*	*	*	*	*	*	*	*	*
312.9	16	11	*5	*6	*6	*	*	*6	*6	*	*
313	41	23	17	24	16	*	*	13	13	13	*
313.8	40	23	17	24	16	*	*	13	13	12	*
313.81	38	22	17	22	15	*	*	13	12	11	*

See footnotes and notes at end of table.

Table 44. Number of all-listed diagnoses for discharges from short-stay hospitals, by ICD-9-CM code, sex, age, and geographic region: United States, 1999—Con.[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table: see Methods section]

ICD-9-CM code	Sex			Age				Region			
	Total	Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	North-east	Midwest	South	West
Number of all-listed diagnoses in thousands											
314	75	56	19	44	29	*	*	13	29	26	*6
314.0	74	55	19	44	28	*	*	13	29	26	*6
314.00	14	11	*	*	*9	*	*	*	*5	*	*
314.01	60	44	15	39	20	*	*	10	24	22	*
315	31	18	13	19	10	*	*	*6	12	*7	*7
315.2	*6	*	*	*	*	*	—	*	*	*	*
315.3	*6	*	*	*	*	—	*	*	*	*	*
315.8	*8	*	*	*6	*	*	—	*	*	*	*
315.9	*9	*6	*	*	*	*	*	*	*	*	*
317	29	16	14	*	18	*5	*	*8	*7	11	*
318	29	20	10	*	14	*8	*	*5	13	*8	*
318.0	12	*8	*	*	*6	*	*	*	*6	*	*
318.1	12	*8	*	*	*5	*	*	*	*	*	*
318.2	*6	*	*	*	*	*	*	*	*	*	*
319	80	49	30	14	30	24	11	18	20	33	9
320-389	3,519	1,533	1,986	403	574	717	1,825	765	871	1,222	661
320	14	*8	*6	*	*	*	*	*	*	*	*
321	*5	*	*	—	*	*	*	*	*	*	*
321.0	*5	*	*	—	*	*	*	*	*	*	*
322	11	*6	*5	*	*	*	*	*	*	*	*
322.9	*8	*	*	*	*	*	*	*	*	*	*
323	11	*	*7	*	*5	*	*	*	*	*	*
323.9	*6	*	*	*	*	*	*	*	*	*	*
324	11	*8	*	*	*	*	*	*	*	*5	*
324.1	*6	*	*	*	*	*	*	*	*	*	*
331	453	158	295	20	17	25	391	106	106	173	68
331.0	358	108	249	—	*	*8	350	86	82	141	50
331.3	15	*8	*6	*	*	*	9	*	*	*	*
331.4	61	32	29	18	14	10	19	13	18	18	12
331.9	14	*6	*8	*	*	*	11	*	*	*6	*
332	245	124	121	—	*	18	227	56	58	90	41
332.0	243	123	120	—	*	17	225	55	58	89	41
333	77	32	45	*	*9	15	50	13	26	20	18
333.0	*8	*	*	*	—	*	*8	*	*	*	*
333.1	27	12	15	—	*	*	22	*	10	*7	*6
333.8	*7	*	*6	—	*	*	*	*	*	*	*
333.82	*7	*	*6	—	*	*	*	*	*	*	*
333.9	20	*8	11	*	*	*7	*9	*	*8	*	*
333.90	*5	*	*	*	*	*	*	*	*	*	*
333.99	14	*5	*8	*	*	*	*7	*	*6	*	*
334	13	*5	*8	*	*	*	*7	*	*	*	*
334.3	*6	*	*	*	*	*	*	*	*	*	*
335	17	*8	*9	*	*	*5	*8	*	*6	*	*
335.2	15	*8	*7	—	*	*5	*7	*	*	*	*
335.20	12	*6	*5	—	*	*	*5	*	*	*	*
336	20	9	10	*	*	*7	9	*5	*	*7	*
336.3	*7	*	*	—	*	*	*	*	*	*	*
337	74	32	42	*	19	31	23	18	18	26	12
337.1	53	23	30	*	11	23	19	13	12	20	*8
337.2	14	*5	*9	*	*6	*6	*	*	*	*	*
337.22	*5	*	*	*	*	*	*	*	*	*	*
340	83	24	59	*	23	40	20	19	29	25	9

See footnotes and notes at end of table.

Table 44. Number of all-listed diagnoses for discharges from short-stay hospitals, by ICD-9-CM code, sex, age, and geographic region: United States, 1999—Con.

[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table: see Methods section]

ICD-9-CM code	Sex			Age				Region			
	Total	Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	North-east	Midwest	South	West
Number of all-listed diagnoses in thousands											
342	215	95	120	*	20	47	146	32	60	69	53
342.0	9	*	*5	—	*	*	*6	*	*	*	*
342.00	*7	*	*	—	*	*	*	*	*	*	*
342.8	*9	*	*	*	*	*	*	*	*	*	*
342.9	196	85	111	*	18	41	135	30	53	63	50
342.90	165	72	93	*	14	34	115	25	44	56	40
342.91	19	*7	11	*	*	*	12	*	*5	*	*6
342.92	13	*6	*7	*	*	*	*8	*	*	*	*
343	73	36	36	32	25	10	*5	17	17	22	17
343.2	15	*6	9	*7	*6	*	*	*	*	*6	*
343.9	49	26	23	21	16	*9	*	12	11	15	11
344	142	87	55	*	54	42	42	23	33	60	26
344.0	47	31	16	*	20	12	13	*8	12	19	*8
344.00	36	25	11	*	14	10	10	*7	10	16	*
344.1	52	32	20	*	23	19	10	9	10	22	11
344.3	*6	*	*	*	*	*	*	*	*	*	*
344.30	*6	*	*	*	*	*	*	*	*	*	*
344.6	9	*5	*	*	*	*	*	*	*	*	*
344.8	*6	*	*	*	*	*	*	*	*	*	*
344.89	*6	*	*	*	*	*	*	*	*	*	*
344.9	15	*7	*8	*	*	*	*8	*	*5	*6	*
345	100	47	53	25	36	21	18	29	19	34	18
345.1	18	*9	10	*	*6	*	*5	*6	*	*	*
345.10	16	*7	*9	*	*6	*	*5	*	*	*	*
345.3	19	*9	10	*5	*6	*	*	*	*	*7	*
345.4	*7	*	*	*	*	*	*	*	*	*	*
345.5	11	*	*6	*	*	*	*	*	*	*	*
345.50	*8	*	*	*	*	*	*	*	*	*	*
345.9	35	17	19	*8	14	*7	*6	*8	*7	14	*7
345.90	31	15	16	*7	11	*7	*6	*7	*6	11	*7
346	157	29	129	*	74	65	15	35	40	59	23
346.0	*5	*	*	*	*	*	*	*	*	*	*
346.2	*8	*	*	*	*	*	*	*	*	*	*
346.20	*5	*	*	*	*	*	*	*	*	*	*
346.8	10	*	*9	*	*6	*	*	*	*	*	*
346.80	*7	*	*6	*	*	*	*	*	*	*	*
346.9	133	21	112	*	64	54	13	29	33	51	20
346.90	125	21	105	*	58	52	13	27	31	49	18
346.91	*7	*	*7	*	*6	*	*	*	*	*	*
347	*8	*	*5	*	*	*	*	*	*	*	*
348	151	74	77	25	28	36	61	28	39	49	35
348.0	9	*	*5	*	*	*	*	*	*	*	*
348.1	56	31	25	9	*6	15	26	*9	14	18	16
348.2	*8	*	*7	*	*	*	*	*	*	*	*
348.3	38	19	19	*	*	*9	21	*9	10	15	*
348.4	*8	*	*	*	*	*	*	*	*	*	*
348.5	11	*5	*6	*	*	*	*	*	*	*	*
348.8	14	*8	*6	*	*	*	*	*	*5	*	*
348.9	*7	*	*	*	—	*	*	*	*	*	*
349	43	14	29	*	20	*9	11	10	*9	14	10
349.0	20	*	16	*	16	*	*	*	*	*7	*
349.2	*6	*	*	*	*	*	*	*	*	*	*
349.8	14	*7	*8	*	*	*	*8	*	*	*	*
349.82	*8	*	*	*	*	*	*5	*	*	*	*
350	17	*	12	*	*	*	9	*	*5	*5	*
350.1	16	*	11	*	*	*	9	*	*5	*5	*
351	24	11	13	*	*	*6	12	*6	*	*7	*7
351.0	21	10	11	*	*	*5	11	*5	*	*7	*6

See footnotes and notes at end of table.

Table 44. Number of all-listed diagnoses for discharges from short-stay hospitals, by ICD-9-CM code, sex, age, and geographic region: United States, 1999—Con.[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table: see Methods section]

ICD-9-CM code	Sex			Age				Region			
	Total	Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	North-east	Midwest	South	West
Number of all-listed diagnoses in thousands											
353	16	*7	*9	*	*6	*	*5	*	*	*	*
353.0	*9	*	*6	*	*	*	*	*	*	*	*
354	34	15	19	*	9	*9	15	*8	*9	11	*6
354.0	20	*8	12	—	*7	*7	*6	*	*	*8	*
355	38	14	24	*	*8	*9	21	*8	12	13	*5
355.8	19	*8	11	*	*	*	11	*	*6	*7	*
355.9	14	*	10	*	*	*	*7	*	*	*	*
356	62	32	29	*	*7	14	40	9	16	26	11
356.9	52	27	25	*	*5	11	35	*6	14	23	*9
357	223	105	117	*	21	84	116	49	57	78	38
357.0	10	*7	*	*	*	*	*	*	*	*	*
357.2	195	89	106	—	15	74	106	45	51	64	34
358	21	12	9	*	*	*	13	*7	*6	*6	*
358.0	17	9	*8	*	*	*	11	*6	*	*	*
359	25	14	11	*	*8	*	*8	*	*	*9	*7
359.1	*9	*6	*	*	*	*	*	*	*	*	*
359.4	*5	*	*	—	*	*	*	*	*	*	*
360	*8	*	*	*	*	*	*	*	*	*	*
360.0	*5	*	*	*	*	*	*	*	*	*	*
361	*8	*	*	*	*	*	*	*	*	*	*
362	166	68	98	*6	20	46	95	40	37	51	38
362.0	98	42	57	—	17	39	42	28	19	30	21
362.01	93	39	54	—	15	38	40	27	17	28	20
362.02	*6	*	*	—	*	*	*	*	*	*	*
362.2	*6	*	*	*	*	*	*	*	*	*	*
362.3	10	*	*6	—	*	*	*7	*	*	*	*
362.34	*7	*	*	—	*	*	*5	*	*	*	*
362.5	41	14	27	*	*	*	39	*	12	13	12
362.50	38	12	26	—	*	*	36	*	11	11	12
364	*5	*	*	*	*	*	*	*	*	*	*
365	179	66	113	*	*6	22	151	47	49	52	31
365.9	169	62	107	*	*	20	144	44	47	49	29
366	30	11	20	*	*	*	23	*7	*9	10	*5
366.1	*5	*	*	—	*	*	*	*	*	*	*
366.9	22	*8	14	—	*	*	18	*5	*8	*6	*
368	55	26	29	*	11	18	25	11	17	17	11
368.1	10	*	*7	*	*	*	*	*	*	*	*
368.16	*6	*	*	*	*	*	*	*	*	*	*
368.2	17	*8	9	*	*	*7	*5	*	*5	*	*
368.4	14	*7	*7	*	*	*	*9	*	*	*	*
368.46	12	*6	*6	*	*	*	*8	*	*	*	*
368.8	*9	*6	*	*	*	*	*6	*	*	*	*
369	80	30	49	*	10	13	54	21	20	27	12
369.0	30	11	20	*	*	*6	21	*7	*7	12	*
369.00	28	10	18	*	*	*6	20	*6	*6	11	*
369.4	19	*8	12	*	*	*	14	*7	*6	*	*
369.6	20	9	11	*	*	*	13	*5	*	*7	*
369.60	18	*9	10	*	*	*	11	*	*	*7	*
369.9	*6	*	*	*	*	*	*	*	*	*	*
370	*6	*	*	*	*	*	*	*	*	*	*
372	53	26	27	17	10	*5	21	12	12	21	*8
372.0	11	*6	*5	*	*	*	*	*	*	*5	*
372.00	11	*5	*5	*	*	*	*	*	*	*	*
372.3	31	14	17	12	*	*	12	*8	*8	12	*
372.30	30	13	17	12	*	*	12	*7	*8	12	*
372.7	*7	*	*	*	*	*	*	*	*	*	*
372.72	*7	*	*	*	*	*	*	*	*	*	*

See footnotes and notes at end of table.

Table 44. Number of all-listed diagnoses for discharges from short-stay hospitals, by ICD-9-CM code, sex, age, and geographic region: United States, 1999—Con.

[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table: see Methods section]

ICD-9-CM code	Sex			Age				Region			
	Total	Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	North-east	Midwest	South	West
Number of all-listed diagnoses in thousands											
374	14	*	10	*	*	*8	*	*	*	*9	*
374.3	*8	*	*6	*	*	*	*	*	*	*5	*
374.30	*7	*	*	*	*	*	*	*	*	*	*
376	20	9	11	*8	*	*6	*	*	*	*6	*6
376.0	13	*8	*5	*8	*	*	*	*	*	*	*
376.01	13	*7	*5	*8	*	*	*	*	*	*	*
377	12	*	*8	*	*5	*	*	*	*	*	*
378	17	9	*8	*	*	*	*7	*	*5	*	*
378.5	13	*7	*6	*	*	*	*6	*	*	*	*
378.51	*5	*	*	—	*	*	*	*	*	*	*
378.54	*7	*	*	*	*	*	*	*	*	*	*
379	16	*7	*9	*	*6	*	*5	*	*	*6	*
380	35	19	16	*8	10	*6	11	*6	11	10	*7
380.1	18	*9	9	*5	*7	*	*	*	*	*7	*
380.10	17	*8	*9	*5	*6	*	*	*	*	*7	*
380.4	13	*8	*6	*	*	*	*8	*	*7	*	*
381	17	*9	*8	10	*	*	*	*6	*	*5	*
382	197	108	88	174	14	*	*6	31	45	79	41
382.9	191	105	87	169	13	*	*5	30	43	78	40
383	11	*6	*	*	*	*	*	*	*	*	*
384	*5	*	*	*	*	*	*	*	*	*	*
385	*7	*5	*	*	*	*	*	*	*	*	*
386	68	23	44	*	*5	15	47	15	17	23	13
386.0	11	*	*8	—	*	*	*5	*	*	*	*
386.00	10	*	*7	—	*	*	*	*	*	*	*
386.1	22	*7	15	*	*	*	16	*	*	*7	*6
386.11	11	*	*7	*	*	*	*9	*	*	*	*
386.12	*7	*	*	—	*	*	*5	*	*	*	*
386.3	32	11	20	*	*	*5	24	*6	10	11	*
386.30	29	10	20	—	*	*5	22	*5	*8	11	*
388	18	*7	11	*	*	*	*8	*	*	*6	*
388.3	*6	*	*	*	*	*	*	*	*	*	*
388.30	*5	*	*	*	*	*	*	*	*	*	*
389	81	40	41	*7	13	10	50	16	20	24	20
389.1	*8	*6	*	*	*	*	*6	*	*	*	*
389.10	*7	*	*	*	*	*	*	*	*	*	*
389.9	62	28	34	*	*9	*8	41	13	16	19	15
390-459	29,558	13,999	15,560	119	1,527	7,134	20,778	6,933	7,432	10,439	4,755
394	33	*7	26	*	*	*7	24	10	*7	13	*
394.0	14	*	13	*	*	*	11	*	*	*8	*
394.1	10	*	*7	*	*	*	*8	*	*	*	*
394.2	*6	*	*	—	*	*	*	*	*	*	*
396	148	59	89	*	*	13	130	41	34	51	22
396.2	43	17	26	—	*	*	40	14	11	13	*
396.3	72	31	41	*	*	*7	63	17	18	29	*9
396.8	15	*	11	—	*	*	13	*5	*	*	*
396.9	10	*	*6	—	*	*	*8	*	*	*	*
397	110	39	71	*	*7	15	87	33	30	34	13
397.0	107	38	69	*	*7	14	85	33	30	33	11
398	61	20	42	—	*	*9	49	22	11	21	*7
398.9	61	20	42	—	*	*9	49	22	11	21	*7
398.90	10	*	*8	—	*	*	*7	*	*	*	*
398.91	50	18	32	—	*	*6	43	19	*9	17	*5

See footnotes and notes at end of table.

Table 44. Number of all-listed diagnoses for discharges from short-stay hospitals, by ICD-9-CM code, sex, age, and geographic region: United States, 1999—Con.[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table: see Methods section]

ICD-9-CM code	Sex			Age				Region			
	Total	Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	North-east	Midwest	South	West
Number of all-listed diagnoses in thousands											
401	6,532	2,713	3,818	12	422	1,870	4,227	1,493	1,644	2,329	1,066
401.0	65	21	43	*	*9	20	36	14	9	34	*7
401.1	192	86	106	*	10	53	129	62	17	16	97
401.9	6,275	2,606	3,669	12	403	1,797	4,063	1,416	1,617	2,279	962
402	388	139	249	*	12	77	297	97	71	163	55
402.0	*9	*	*	—	*	*	*6	*	*	*	*
402.01	*7	*	*	—	*	*	*5	*	*	*	*
402.1	*9	*	*	*	*	*	*6	*	*	*	*
402.11	*7	*	*	*	*	*	*	*	*	*	*
402.9	370	129	241	*	11	74	285	91	69	158	52
402.90	181	67	114	—	*6	44	132	45	39	74	24
402.91	189	62	127	*	*6	30	153	47	31	84	28
403	625	307	318	*	87	192	344	115	127	267	117
403.0	16	*5	10	—	*	*5	*7	*	*	*8	*
403.01	14	*	9	—	*	*	*6	*	*	*7	*
403.1	15	*9	*6	*	*	*	10	*	*	*	10
403.11	14	*8	*6	*	*	*	*8	*	*	*	*9
403.9	594	293	301	*	81	184	328	108	123	258	105
403.90	47	24	23	*	*7	12	27	9	10	23	*
403.91	547	269	278	*	73	172	300	99	113	236	100
404	84	39	45	*	*7	20	56	20	19	31	14
404.0	*5	*	*	—	*	*	*	*	*	*	*
404.9	77	37	41	*	*6	17	53	19	17	29	12
404.90	*6	*	*	—	*	*	*	*	*	*	*
404.91	*7	*	*	—	*	*	*5	*	*	*	*
404.92	18	10	*8	—	*	*	11	*5	*	*6	*
404.93	47	23	24	*	*	10	32	12	12	17	*7
405	11	*	*6	*	*	*	*	*	*	*	*
405.9	10	*	*6	*	*	*	*	*	*	*	*
405.91	*7	*	*	*	*	*	*	*	*	*	*
410	981	556	425	*	53	307	620	240	248	316	177
410.0	32	15	17	—	*	14	17	*8	*6	*8	10
410.01	30	14	16	—	*	14	16	*7	*6	*8	9
410.1	153	90	64	—	10	54	89	30	34	55	34
410.11	136	80	56	—	*9	50	77	27	30	49	29
410.12	15	*9	*6	—	*	*	9	*	*	*	*
410.2	27	15	12	—	*	*7	18	*	*5	11	*6
410.21	26	15	11	—	*	*6	17	*	*5	11	*
410.3	18	12	*6	—	*	*9	*9	*	*6	*5	*
410.31	18	12	*6	—	*	*8	*9	*	*6	*5	*
410.4	181	120	61	*	19	67	94	49	44	51	36
410.41	167	113	54	—	18	61	88	45	43	47	32
410.42	11	*6	*	*	*	*5	*	*	*	*	*
410.5	16	9	*7	—	*	*6	10	*	*7	*	*
410.51	14	*9	*6	—	*	*6	*8	*	*5	*	*
410.6	*8	*7	*	—	*	*	*	*	*	*	*
410.61	*8	*7	*	—	*	*	*	*	*	*	*
410.7	374	202	172	*	12	107	255	102	96	119	58
410.71	352	191	161	*	11	102	238	95	91	112	54
410.72	19	10	*9	—	*	*	14	*6	*	*5	*
410.8	14	*9	*6	—	*	*	10	*	*	*	*
410.81	13	*8	*	—	*	*	*8	*	*	*	*
410.9	156	76	80	—	*6	35	115	33	42	58	24
410.90	9	*	*5	—	*	*	*8	*	*	*	*
410.91	132	66	66	—	*	30	97	26	35	51	20
410.92	15	*6	*9	—	*	*	10	*	*	*5	*

See footnotes and notes at end of table.

Table 44. Number of all-listed diagnoses for discharges from short-stay hospitals, by ICD-9-CM code, sex, age, and geographic region: United States, 1999—Con.

[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table: see Methods section]

ICD-9-CM code	Sex			Age				Region			
	Total	Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	North-east	Midwest	South	West
Number of all-listed diagnoses in thousands											
411	909	503	405	*	45	330	533	240	183	334	152
411.0	*7	*	*	—	*	*	*	*	*	*	*
411.1	859	477	382	*	41	316	502	223	174	317	145
411.8	43	23	20	—	*	11	28	15	*8	15	*
411.81	13	*8	*	—	*	*6	*7	*	*	*5	*
411.89	30	14	15	—	*	*5	22	12	*6	10	*
412	694	416	278	*	27	217	448	174	165	220	135
413	559	270	290	*	22	155	380	133	153	198	74
413.9	558	269	289	*	22	154	380	132	153	198	74
414	4,492	2,477	2,014	10	123	1,217	3,142	1,118	1,133	1,543	698
414.0	4,200	2,311	1,889	*8	116	1,147	2,929	1,042	1,058	1,442	658
414.00	1,235	634	601	*	21	269	942	257	410	434	135
414.01	2,777	1,553	1,224	*	92	827	1,853	735	612	934	497
414.02	102	72	30	*	*	31	69	23	27	30	22
414.04	14	11	*	—	*	*6	*7	*	*	*5	*
414.05	70	40	29	—	*	12	56	22	*6	39	*
414.1	14	*7	*7	*	*	*	*8	*	*	*	*
414.10	9	*	*	*	*	*	*6	*	*	*	*
414.8	198	120	78	—	*	47	147	51	56	63	29
414.9	79	39	41	—	*	20	58	23	15	34	*7
415	142	58	85	*	19	40	84	32	36	48	26
415.1	138	55	83	*	19	38	81	31	35	47	25
415.11	14	*7	*8	*	*	*7	*6	*	*	*	*
415.19	124	49	75	*	17	32	75	27	32	43	23
416	258	111	146	*	18	59	176	55	57	92	53
416.0	154	62	92	*	14	35	101	35	33	61	25
416.8	29	12	17	*	*	*8	20	*	*7	*	14
416.9	74	37	37	*	*	16	55	16	17	26	15
420	25	13	12	*	*	11	10	*7	*5	*9	*
420.9	22	12	10	*	*	10	*9	*6	*	*8	*
420.90	17	9	*8	*	*	*8	*6	*	*	*6	*
421	29	17	11	*	*8	*9	9	*	*	12	*8
421.0	24	16	*8	*	*6	*8	*9	*	*	12	*
423	45	21	24	*	*7	15	21	*8	13	14	10
423.9	35	15	20	*	*5	13	16	*5	9	12	*8
424	912	358	554	*	85	163	660	236	221	315	140
424.0	519	187	332	*	72	114	332	132	126	193	69
424.1	316	140	176	*	*7	35	272	90	82	86	58
424.2	44	21	23	*	*	10	32	*7	*6	23	*8
424.3	*7	*	*	*	*	*	*	*	*	*	*
424.9	26	*8	19	*	*	*	20	*6	*	11	*5
424.90	25	*7	18	*	*	*	19	*6	*	10	*
425	518	286	233	*	41	140	335	112	126	193	88
425.1	19	*	14	*	*	*	14	*	*	*9	*
425.4	457	260	197	*	33	120	302	100	112	167	78
425.5	14	9	*5	—	*	*8	*	*	*	*6	*
425.8	17	*9	*9	—	*	*6	11	*	*	*8	*
425.9	*6	*	*	*	*	*	*	*	*	*	*
426	429	235	194	*	20	76	330	99	126	123	81
426.0	73	38	35	*	*	13	55	20	16	22	15
426.1	106	61	45	*	*	16	86	24	35	29	19
426.10	*8	*	*	*	*	*	*7	*	*	*	*
426.11	60	35	26	*	*	*8	50	13	22	15	10
426.12	*9	*6	*	*	*	*	*8	*	*	*	*
426.13	29	17	12	*	*	*5	22	*6	*7	*8	*7
426.2	20	11	*9	—	*	*	14	*	*7	*5	*
426.3	78	33	45	*	*	13	65	15	21	23	20

See footnotes and notes at end of table.

Table 44. Number of all-listed diagnoses for discharges from short-stay hospitals, by ICD-9-CM code, sex, age, and geographic region: United States, 1999—Con.[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table: see Methods section]

ICD-9-CM code	Sex			Age				Region			
	Total	Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	North-east	Midwest	South	West
Number of all-listed diagnoses in thousands											
426.4	77	50	28	*	*6	15	56	19	20	25	14
426.5	35	21	15	—	*	*	31	10	10	9	*6
426.52	10	*	*6	—	—	*	10	*	*	*	*
426.53	19	13	*6	—	*	*	17	*6	*5	*	*
426.6	*6	*	*	—	*	*	*5	*	*	*	*
426.7	10	*6	*	*	*	*	*	*	*	*	*
426.9	18	9	9	*	*	*6	12	*	11	*	*
427	3,773	1,795	1,978	21	152	605	2,994	909	976	1,243	645
427.0	113	45	68	*	10	27	76	25	21	47	20
427.1	239	150	89	*	16	64	159	57	58	78	45
427.2	*6	*	*	—	*	*	*	*	*	*	*
427.3	2,402	1,103	1,298	*	48	301	2,049	614	642	753	393
427.31	2,219	1,001	1,218	*	42	266	1,909	567	595	699	358
427.32	182	102	80	*	*7	34	140	47	47	53	35
427.4	53	33	19	*	*	19	32	10	12	18	13
427.41	52	33	19	*	*	18	31	10	12	17	13
427.5	98	54	45	*	*9	22	64	16	20	39	23
427.6	130	67	63	*	*8	28	93	27	34	42	27
427.60	10	*	*5	*	*	*	*8	*	*	*	*
427.61	30	14	16	*	*	*	25	*	9	*7	*9
427.69	91	48	42	*	*7	23	61	21	24	31	15
427.8	637	296	342	11	51	125	451	137	162	226	112
427.81	153	63	90	*	*	15	136	40	29	68	16
427.89	484	232	252	11	49	110	315	97	134	158	96
427.9	94	46	48	*	*8	18	66	22	26	35	11
428	3,135	1,359	1,776	11	78	509	2,537	698	865	1,159	413
428.0	3,105	1,347	1,759	11	75	504	2,515	694	859	1,145	408
428.1	11	*	*7	*	*	*	*9	*	*	*	*
428.9	19	9	10	*	*	*	13	*	*	11	*
429	268	119	149	*	15	48	201	56	73	104	35
429.2	119	53	66	—	*	13	104	27	24	57	11
429.3	83	34	49	*	*8	18	53	15	24	26	17
429.4	*9	*	*	*	*	*	*6	*	*	*	*
429.7	*6	*	*	—	*	*	*	*	*	*	*
429.9	41	21	20	*	*	10	29	*8	17	12	*
430	26	*8	17	*	*5	9	11	*6	*	9	*6
431	78	39	39	*	*8	23	47	13	17	28	20
432	29	18	12	*	*	*6	20	*8	*7	*9	*5
432.1	25	16	*9	*	*	*	18	*7	*6	*8	*
433	324	162	161	—	*	75	245	66	91	118	48
433.1	234	118	116	—	*	53	179	47	66	85	36
433.10	201	103	98	—	*	41	158	41	57	71	32
433.11	33	16	17	—	*	11	21	*6	*8	14	*
433.2	*5	*	*	—	*	*	*	*	*	*	*
433.3	76	36	41	—	*	20	56	17	23	27	10
433.30	66	31	36	—	*	17	49	15	21	23	*7
433.31	10	*	*	—	*	*	*7	*	*	*	*
434	365	155	210	*	15	81	267	82	85	133	64
434.0	25	11	14	*	*	*	19	*	*9	*8	*
434.00	*5	*	*	—	*	*	*	*	*	*	—
434.01	20	*9	11	*	*	*	15	*	*6	*5	*
434.1	48	23	25	—	*	9	36	13	9	15	12
434.11	44	22	22	—	*	*9	33	11	*8	14	11
434.9	292	121	171	*	11	67	211	65	67	111	49
434.90	14	*6	*8	*	*	*	*9	*	*	*7	*
434.91	278	115	163	*	10	63	202	64	65	104	45

See footnotes and notes at end of table.

Table 44. Number of all-listed diagnoses for discharges from short-stay hospitals, by ICD-9-CM code, sex, age, and geographic region: United States, 1999—Con.

[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table: see Methods section]

ICD-9-CM code	Sex			Age				Region			
	Total	Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	North-east	Midwest	South	West
Number of all-listed diagnoses in thousands											
435	265	107	158	*	12	54	199	59	65	104	37
435.3	16	*6	10	*	*	*	14	*	*5	*	*
435.8	*7	*	*5	*	*	*	*	*	*	*	*
435.9	234	96	138	*	10	45	178	51	56	94	32
436	203	89	114	*	*	32	166	25	52	86	39
437	159	67	92	*	*9	25	124	22	47	66	24
437.0	62	29	33	—	*	*5	57	10	20	25	*8
437.1	14	*6	*8	—	*	*	12	*	*6	*6	*
437.2	10	*	*7	*	*	*	*	*	*	*	*
437.3	15	*	11	*	*	*	*7	*	*	*8	*
437.7	*6	*	*	—	*	*	*	*	*	*	*
437.9	44	18	26	*	*	*7	36	*6	14	19	*5
438	528	248	281	*	14	94	419	118	136	179	95
438.0	11	*5	*5	*	*	*	*8	*	*	*	*
438.1	90	41	49	*	*	13	74	26	21	27	16
438.10	*5	*	*	—	*	*	*	*	*	*	*
438.11	63	28	35	—	*	9	53	18	14	19	12
438.12	13	*6	*7	*	*	*	11	*5	*	*	*
438.19	*8	*	*	—	*	*	*6	*	*	*	*
438.2	238	111	127	*	*6	46	185	58	66	73	42
438.20	210	95	115	*	*	41	164	54	57	68	31
438.21	14	*8	*5	*	*	*	10	*	*	*	*
438.22	14	*7	*7	—	*	*	10	*	*	*	*7
438.5	*8	*	*	*	*	*	*6	*	*	*	*
438.50	*5	*	*	—	*	*	*	*	*	*	*
438.8	130	62	69	*	*	25	103	21	33	50	27
438.82	32	18	13	—	*	*	29	*	9	13	*6
438.89	97	42	55	*	*	21	74	17	23	37	20
438.9	45	21	24	*	*	*5	38	10	10	21	*5
440	423	216	207	*	*8	98	317	98	121	151	54
440.0	18	*8	11	—	*	*5	13	*9	*	*	*
440.1	34	15	19	*	*	9	24	*9	*9	14	*
440.2	251	136	114	*	*5	61	185	54	78	82	38
440.20	70	35	35	*	*	16	52	9	24	26	11
440.21	49	32	17	—	*	13	35	11	15	13	11
440.22	23	12	11	—	*	*	18	*	*7	10	*
440.23	31	15	16	—	*	*7	23	*	10	11	*
440.24	75	40	34	—	*	19	54	24	22	20	9
440.3	11	*7	*	—	*	*	*6	*	*	*	*
440.8	*7	*	*	—	—	*	*	*	*	*	*
440.9	101	46	55	—	*	15	86	22	25	47	*8
441	157	93	65	*	*	27	127	45	42	49	22
441.0	13	*7	*6	—	*	*6	*5	*	*	*6	*
441.01	*7	*	*	—	*	*	*	*	*	*	*
441.2	10	*5	*5	*	*	*	*7	*	*	*	*
441.3	*6	*	*	—	—	*	*6	*	*	*	*
441.4	117	70	47	—	*	15	101	35	33	36	14
442	27	17	10	*	*	*7	19	*8	*6	9	*
442.2	*9	*8	*	—	—	*	*7	*	*	*	*
442.3	12	*6	*5	*	*	*	*9	*	*	*	*
443	428	219	209	*	10	100	317	106	97	168	56
443.0	11	*	*8	*	*	*	*5	*	*	*	*
443.8	77	39	39	*	*	20	54	23	13	29	13
443.81	66	34	33	—	*	18	45	18	11	27	10
443.89	11	*5	*6	*	*	*	*9	*	*	*	*
443.9	339	177	162	*	*	75	258	81	82	136	40

See footnotes and notes at end of table.

Table 44. Number of all-listed diagnoses for discharges from short-stay hospitals, by ICD-9-CM code, sex, age, and geographic region: United States, 1999—Con.[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table: see Methods section]

ICD-9-CM code	Sex			Age				Region			
	Total	Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	North-east	Midwest	South	West
Number of all-listed diagnoses in thousands											
444	105	51	53	*	*7	34	63	30	22	39	14
444.0	12	*7	*5	—	*	*	*6	*	*	*6	*
444.2	69	34	35	—	*	20	44	20	14	25	10
444.21	9	*	*6	—	*	*	*5	*	*	*	*
444.22	59	30	29	—	*	17	39	18	12	23	*7
444.8	17	*8	9	*	*	*7	*9	*	*	*6	*
444.81	12	*6	*5	—	*	*5	*6	*	*	*	*
444.89	*6	*	*	*	*	*	*	*	*	*	*
446	49	21	28	*6	*5	*8	30	14	12	13	11
446.1	*6	*	*	*5	*	*	—	*	*	*	*
446.2	*8	*	*7	*	*	*	*	*	*	*	*
446.4	*7	*	*	—	*	*	*	*	*	*	*
446.5	22	*9	14	*	*	*	20	*8	*6	*6	*
447	56	24	32	*	*	16	34	15	12	22	*7
447.1	26	12	14	*	*	*7	16	*7	*6	10	*
447.6	12	*	*8	*	*	*	*	*	*	*	*
447.9	*8	*	*	—	—	*	*6	*	*	*5	*
448	*7	*	*	*	*	*	*	*	*	*	*
448.0	*5	*	*	*	*	*	*	*	*	*	*
451	70	28	43	*	11	22	35	19	18	23	10
451.0	*9	*	*6	*	*	*	*	*	*	*	*
451.1	21	*8	13	—	*	*	14	*6	*	*7	*
451.19	17	*7	10	—	*	*	12	*	*	*6	*
451.2	16	*7	10	*	*	*6	*7	*	*5	*5	*
451.8	19	*8	11	*	*	*6	*8	*	*6	*6	*
451.82	*9	*	*	*	*	*	*	*	*	*	*
451.9	*5	*	*	—	*	*	*	*	*	*	*
453	309	133	176	*	43	98	168	77	77	108	47
453.2	11	*6	*	*	*	*6	*	*	*	*7	*
453.8	253	109	144	*	36	79	138	59	69	85	40
453.9	42	16	26	—	*	13	25	16	*5	15	*6
454	33	16	17	*	*	9	21	9	11	*8	*
454.0	*6	*	*	—	*	*	*	*	*	*	*
454.1	*5	*	*	—	*	*	*	*	*	*	*
454.2	*6	*	*	—	*	*	*	*	*	*	*
454.9	16	*7	*9	*	*	*	9	*	*5	*	*
455	150	60	91	*	36	37	77	31	27	66	27
455.0	54	21	33	—	*9	11	34	10	10	24	10
455.2	24	11	13	—	*	*9	11	*5	*	*8	*6
455.3	15	*	10	—	*	*	*8	*	*	*5	*
455.5	10	*5	*	—	*	*	*	*	*	*	*
455.6	25	*8	18	*	10	*6	*9	*	*	13	*
455.8	11	*	*8	—	*	*	*6	*	*	*	*
456	62	35	28	*	12	29	21	11	14	22	15
456.0	*9	*5	*	*	*	*	*5	*	*	*	*
456.1	*8	*	*	—	*	*	*	*	*	*	*
456.2	35	21	15	*	*7	19	10	*6	*7	14	*8
456.20	21	12	*9	*	*	11	*	*	*	9	*
456.21	15	*9	*6	—	*	*8	*	*	*	*5	*
456.8	*7	*	*	—	*	*	*	*	*	*	*
457	24	11	14	*	*	11	9	*	*7	*9	*
457.1	16	*6	9	*	*	*6	*7	*	*	*5	*
458	362	182	180	*6	28	84	244	69	95	122	76
458.0	105	49	56	*	*8	18	77	19	30	35	21
458.2	31	16	15	*	*	*9	20	*6	*6	*8	12
458.8	22	11	11	*	*	*6	13	*6	*6	*5	*
458.9	203	106	97	*	16	50	133	38	53	74	39

See footnotes and notes at end of table.

Table 44. Number of all-listed diagnoses for discharges from short-stay hospitals, by ICD-9-CM code, sex, age, and geographic region: United States, 1999—Con.

[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table: see Methods section]

ICD-9-CM code	Sex			Age				Region			
	Total	Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	North-east	Midwest	South	West
Number of all-listed diagnoses in thousands											
459	144	68	75	*	13	42	88	35	37	51	21
459.0	14	*8	*6	*	*	*	*8	*	*	*5	*
459.2	14	*8	*6	*	*	*5	*6	*	*	*7	*
459.8	94	42	52	*	*6	28	59	23	27	28	16
459.81	87	39	48	—	*5	26	56	22	25	25	15
459.89	*7	*	*	*	*	*	*	*	*	*	*
459.9	19	*9	10	*	*	*	13	*	*6	9	*
460-519	10,508	5,030	5,478	1,235	1,274	2,156	5,844	2,280	2,563	3,844	1,821
460	*8	*	*	*	*	*	*	*	*	*	*
461	72	31	41	11	25	15	22	14	16	29	13
461.0	17	*6	10	*	*6	*	*6	*	*6	*	*
461.2	*6	*	*	*	*	*	*	*	*	*	*
461.8	11	*	*6	*	*	*	*	*	*	*7	*
461.9	33	13	20	*	12	*9	10	*6	*6	15	*6
462	52	17	35	15	21	*8	*7	*9	13	20	10
463	20	*9	11	*6	13	*	*	*	*7	*6	*
464	60	38	22	48	*	*	*	*9	13	22	16
464.1	*8	*	*	*	*	*	*	*	*	*	*
464.10	*7	*	*	*	*	*	*	*	*	*	*
464.3	*5	*	*	*	*	*	*	*	*	*	*
464.4	42	28	14	42	*	—	—	*	*7	17	13
465	183	80	103	61	49	25	48	39	50	66	28
465.9	180	79	101	60	48	24	48	38	49	65	28
466	460	230	231	249	48	46	117	85	104	182	90
466.0	231	100	132	23	46	45	116	48	59	94	29
466.1	229	130	99	225	*	*	*	36	44	88	61
466.11	124	70	55	124	*	*	*	18	23	54	29
466.19	105	60	44	101	*	*	*	19	21	33	32
470	17	13	*	*	*6	*5	*5	*	*	*5	*6
471	11	*6	*	*	*	*	*	*	*	*	*
472	13	*6	*7	*	*	*	*	*	*5	*	*
472.0	13	*6	*7	*	*	*	*	*	*5	*	*
473	162	78	84	24	48	40	49	33	40	69	20
473.0	30	16	14	*	*8	*6	10	*7	*6	13	*
473.1	*7	*	*	*	*	*	*	*	*	*	*
473.2	11	*7	*	*	*	*	*	*	*	*	*
473.8	15	*8	*7	*	*	*	*5	*	*	*	*
473.9	97	42	54	13	29	26	28	18	25	45	10
474	26	14	11	15	*8	*	*	*9	*	*7	*7
474.0	*9	*	*	*	*	*	*	*	*	*	*
474.00	*7	*	*	*	*	*	*	*	*	*	*
474.1	16	10	*6	9	*	*	—	*7	*	*	*
474.10	*7	*	*	*6	*	—	—	*	*	*	*
474.11	*7	*5	*	*	*	*	—	*	*	*	*
475	12	*7	*5	*	*9	*	*	*	*	*	*
477	39	17	22	*	11	11	14	*7	*9	16	*8
477.9	38	16	21	*	11	11	14	*6	*9	16	*7
478	83	47	35	14	17	28	23	20	14	26	23
478.0	*6	*	*	*	*	*	*	*	*	*	*
478.1	19	9	9	*	*5	*7	*	*	*	*7	*6
478.2	23	15	*8	*	*6	10	*	*7	*	*6	*7
478.29	16	11	*	*	*	*6	*	*	*	*5	*6
478.3	13	*6	*7	*	*	*	*8	*	*	*	*
478.7	12	*7	*6	*6	*	*	*	*	*	*	*
478.74	*5	*	*	*	*	*	*	*	*	*	*

See footnotes and notes at end of table.

Table 44. Number of all-listed diagnoses for discharges from short-stay hospitals, by ICD-9-CM code, sex, age, and geographic region: United States, 1999—Con.[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table: see Methods section]

ICD-9-CM code	Sex			Age				Region			
	Total	Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	North-east	Midwest	South	West
Number of all-listed diagnoses in thousands											
480	52	27	25	35	*	*	10	*	13	19	16
480.1	24	14	11	23	*	*	*	*	*	12	*8
480.9	22	12	10	11	*	*	*6	*	*6	*7	*6
481	75	38	37	*8	12	21	35	14	19	27	15
482	266	154	111	16	29	43	178	48	69	101	48
482.0	18	9	*9	*	*	*	13	*	*	*9	*
482.1	57	30	26	*5	*7	10	35	*8	14	20	15
482.2	26	13	13	*	*	*	19	*	9	*9	*
482.3	13	*8	*	*	*	*	*8	*	*	*	*
482.30	*6	*	*	*	*	*	*	*	*	*	*
482.4	71	48	23	*	*7	10	52	15	14	30	13
482.41	63	42	21	*	*6	9	45	13	13	26	11
482.8	59	34	26	*	*7	*9	41	*8	21	22	*9
482.82	*6	*	*	*	*	*	*	*	*	*	*
482.83	47	26	21	*	*5	*6	33	*7	17	16	*7
482.9	21	11	10	*	*	*	11	*7	*	*8	*
483	16	*5	11	*	*	*	*7	*	*6	*6	*
483.0	13	*	10	*	*	*	*6	*	*	*	*
485	49	25	24	18	*	*7	21	*7	15	21	*5
486	1,673	801	872	191	162	291	1,029	342	404	642	285
487	61	22	39	10	13	*8	31	*8	22	16	14
487.0	13	*6	*8	*	*	*	*7	*	*	*	*
487.1	45	15	30	*6	11	*6	23	*	18	13	9
490	103	46	57	*8	22	26	47	19	21	48	15
491	1,064	492	572	*	36	269	755	220	269	420	155
491.2	1,017	472	545	*	27	255	734	213	254	403	147
491.20	118	62	56	*	*	26	87	24	22	51	21
491.21	899	410	489	*	22	228	647	189	231	352	126
491.9	35	15	20	*	*8	11	16	*	11	15	*
492	209	120	89	*	9	52	147	40	64	75	30
492.0	14	12	*	*	*	*	*5	*	*	*	*
492.8	194	107	87	*	*6	47	142	36	61	71	27
493	1,347	487	860	315	377	310	345	369	321	410	248
493.0	70	37	32	50	11	*6	*	15	*6	13	35
493.00	44	24	20	29	*7	*5	*	9	*	10	21
493.01	25	13	12	21	*	*	*	*6	*	*	14
493.2	229	84	145	*	15	78	135	62	56	66	44
493.20	200	75	125	*	12	62	126	52	53	56	39
493.21	29	9	20	*	*	16	10	10	*	10	*6
493.9	1,048	365	682	264	350	227	207	290	258	331	168
493.90	872	285	587	162	305	208	196	246	206	276	144
493.91	176	80	95	102	45	18	11	44	52	55	25
494	36	13	23	*	*	*	29	*8	*8	15	*6
496	1,844	986	858	*	45	394	1,403	410	505	650	278
500	*8	*7	*	—	*	*	*6	*6	*	*	*
501	14	14	*	—	*	*	13	*	*	*7	*
507	319	188	131	12	26	40	241	79	66	111	62
507.0	315	185	130	12	24	40	239	78	66	111	60
508	*9	*	*	—	*	*	*7	*	*	*	*
508.0	*7	*	*	—	*	*	*6	*	*	*	*
510	26	18	*8	*	*8	*6	10	*	*	10	*6
510.9	23	16	*7	*	*7	*	9	*	*	*9	*6

See footnotes and notes at end of table.

Table 44. Number of all-listed diagnoses for discharges from short-stay hospitals, by ICD-9-CM code, sex, age, and geographic region: United States, 1999—Con.

[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table: see Methods section]

ICD-9-CM code	Sex			Age				Region			
	Total	Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	North-east	Midwest	South	West
Number of all-listed diagnoses in thousands											
511	395	180	215	15	44	91	246	89	93	153	60
511.0	28	14	14	*	*6	*9	12	*	*8	13	*
511.8	34	17	16	*	*	10	18	*	*7	15	*8
511.9	332	148	184	13	33	72	215	81	78	125	48
512	109	56	53	*	27	27	51	20	25	42	21
512.0	*6	*	*	*	*	*	*	*	*	*	*
512.1	51	26	25	*	*7	13	28	*9	11	18	13
512.8	53	26	26	*	19	13	20	11	12	22	*8
513	11	*6	*	*	*	*	*	*	*	*	*
513.0	10	*6	*	*	*	*	*	*	*	*	*
514	30	12	18	*	*	*8	15	*5	*7	15	*
515	147	58	88	*	*5	26	114	23	41	60	23
516	29	13	16	*	*	*7	19	*6	*5	11	*7
516.3	17	*9	*9	—	*	*	13	*	*	*6	*
516.8	10	*	*6	*	*	*	*	*	*	*	*
517	15	*	10	*	*5	*5	*	*	*	*6	*
517.8	14	*	9	*	*5	*5	*	*	*	*6	*
518	1,279	602	677	103	145	295	735	289	260	473	257
518.0	311	132	180	24	48	75	164	62	65	125	60
518.1	*7	*	*	*	*	*	*	*	*	*	*
518.3	21	*9	13	*	*	*7	*8	*	*5	*7	*
518.4	23	10	12	*	*	*	13	*	*	9	*
518.5	94	50	44	*6	12	23	53	22	19	33	20
518.8	822	398	425	69	73	185	495	194	164	298	167
518.81	549	271	279	14	51	131	353	132	106	216	96
518.82	138	67	71	47	12	24	55	27	21	36	55
518.83	27	11	16	*	*	*	20	*8	*9	9	*
518.84	29	13	16	*	*	*7	19	*6	*6	12	*
518.89	79	35	44	*7	*5	18	48	21	22	25	10
519	90	44	45	25	14	16	36	14	26	31	19
519.0	10	*5	*5	*	*	*	*	*	*	*	*
519.09	*6	*	*	*	*	*	*	*	*	*	*
519.1	40	18	23	12	*5	9	13	*6	13	12	*8
519.8	27	14	13	*6	*	*	14	*	*7	12	*6
519.9	*6	*	*	*	*	*	*	*	*	*	*
520-579	8,898	3,754	5,144	478	1,699	2,413	4,309	1,929	2,114	3,271	1,584
521	23	14	*9	*5	*8	*5	*	*6	*	*9	*
521.0	22	14	*9	*5	*8	*5	*	*6	*	*8	*
522	24	12	13	*	14	*	*	*6	*5	*8	*5
522.5	19	*9	10	*	10	*	*	*	*	*7	*
523	16	*9	*7	*	*6	*	*	*	*	*9	*
523.1	*6	*	*	*	*	*	*	*	*	*	*
524	42	16	26	*	33	*	*	*9	*6	21	*7
524.0	16	*8	*9	*	14	*	*	*	*	9	*
524.1	*9	*	*	*	*7	*	—	*	*	*	*
524.10	*7	*	*	*	*6	*	—	*	*	*	*
524.6	11	*	*9	*	*6	*	*	*	*	*	*
524.69	*5	*	*	*	*	*	*	*	*	*	*
525	*6	*	*	*	*	*	*	*	*	*	*
526	*6	*	*	*	*	*	*	*	*	*	*
527	20	*9	11	*	*7	*	*7	*	*6	*	*
527.2	12	*	*8	*	*	*	*	*	*	*	*
528	46	27	19	9	15	10	12	10	12	13	12
528.0	22	12	10	*	*	*	*9	*5	*6	*	*6
528.3	11	*7	*	*	*7	*	*	*	*	*	*
528.9	10	*6	*	*	*	*	*	*	*	*	*

See footnotes and notes at end of table.

Table 44. Number of all-listed diagnoses for discharges from short-stay hospitals, by ICD-9-CM code, sex, age, and geographic region: United States, 1999—Con.[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table: see Methods section]

ICD-9-CM code	Sex			Age				Region			
	Total	Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	North-east	Midwest	South	West
Number of all-listed diagnoses in thousands											
529	*5	*	*	*	*	*	*	*	*	*	*
530	1,310	542	767	86	177	389	658	301	304	491	214
530.0	*5	*	*	—	*	*	*	*	*	*	*
530.1	228	99	128	*	39	67	117	47	60	85	36
530.10	95	40	55	*	17	29	47	23	25	36	11
530.11	92	35	57	*	17	26	48	19	26	32	15
530.19	40	24	16	*	*	12	21	*	*9	17	10
530.2	63	37	26	*	9	16	37	13	15	21	14
530.3	54	24	29	*	*	10	41	10	13	21	9
530.5	20	*6	13	*	*	*7	10	*	*	*9	*
530.7	26	14	12	*	*6	*7	12	*	*6	10	*7
530.8	905	353	552	79	114	279	433	219	203	341	142
530.81	875	338	537	78	111	272	414	215	195	330	135
530.82	18	10	*9	*	*	*	12	*	*	*7	*
530.89	11	*	*6	*	*	*	*7	*	*	*	*
531	175	76	100	*	19	45	110	33	50	60	32
531.0	20	12	*9	*	*	*7	10	*	*6	*7	*
531.00	20	11	*9	*	*	*7	10	*	*6	*7	*
531.3	*6	*	*	*	*	*	*	*	*	*	*
531.30	*6	*	*	*	*	*	*	*	*	*	*
531.4	74	34	39	*	*5	17	51	15	19	24	16
531.40	74	34	39	*	*5	17	51	15	19	23	16
531.5	*8	*	*5	*	*	*	*5	*	*	*	*
531.50	*8	*	*5	*	*	*	*5	*	*	*	*
531.7	*5	*	*	—	*	*	*	*	*	*	*
531.9	57	21	35	*	*7	16	33	12	17	22	*6
531.90	55	21	34	*	*6	16	32	12	16	21	*6
532	117	67	50	*	15	26	74	27	27	43	20
532.0	18	11	*7	—	*	*	11	*	*	*	*
532.00	18	11	*7	—	*	*	10	*	*	*	*
532.4	46	29	17	*	*	9	31	10	11	17	*9
532.40	45	28	17	*	*	9	31	10	11	16	*9
532.5	*7	*	*	*	*	*	*	*	*	*	*
532.50	*7	*	*	*	*	*	*	*	*	*	*
532.9	33	18	15	*	*5	*9	18	*7	*8	14	*
532.90	31	16	15	*	*	*9	17	*7	*8	13	*
533	167	71	96	*	17	51	99	41	39	64	24
533.4	*9	*5	*	—	*	*	*6	*	*	*	*
533.40	*9	*5	*	—	*	*	*6	*	*	*	*
533.7	11	*	*6	—	*	*5	*	*	*	*	*
533.70	11	*	*6	—	*	*5	*	*	*	*	*
533.9	140	57	82	*	13	42	85	34	32	51	22
533.90	139	57	82	*	13	42	85	34	32	51	21
534	9	*	*	—	*	*	*	*	*	*	*
535	609	261	348	16	122	153	317	125	156	229	99
535.0	85	38	47	*	15	24	42	16	21	33	14
535.00	63	26	37	*	12	19	28	12	17	25	9
535.01	22	12	10	*	*	*5	14	*	*	*8	*
535.1	71	30	41	*	*8	19	45	20	13	27	10
535.10	55	23	32	*	*7	16	32	17	11	19	*8
535.11	16	*7	9	—	*	*	12	*	*	*8	*
535.3	19	14	*5	—	12	*6	*	*	*	*7	*
535.30	14	10	*	—	*9	*	*	*	*	*6	*
535.31	*6	*	*	—	*	*	*	*	*	*	*
535.4	133	56	77	*	22	31	77	22	41	41	28
535.40	97	38	59	*	18	24	52	15	30	31	21
535.41	36	17	18	*	*	*7	24	*7	11	10	*7

See footnotes and notes at end of table.

Table 44. Number of all-listed diagnoses for discharges from short-stay hospitals, by ICD-9-CM code, sex, age, and geographic region: United States, 1999—Con.

[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table: see Methods section]

ICD-9-CM code	Sex			Age				Region			
	Total	Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	North-east	Midwest	South	West
Number of all-listed diagnoses in thousands											
535.5	229	84	145	*7	47	57	118	44	55	97	33
535.50	193	71	122	*7	44	48	95	37	46	81	29
535.51	35	13	23	*	*	*9	24	*7	9	15	*
535.6	70	38	32	*	19	17	33	17	20	24	10
535.60	59	31	29	*	18	14	26	14	19	19	*8
535.61	11	*7	*	*	*	*	*7	*	*	*5	*
536	254	99	155	12	46	72	124	39	69	93	53
536.2	27	9	18	*	*8	*6	*8	*	*6	14	*
536.3	76	29	46	*	19	33	23	15	13	30	18
536.4	42	14	28	*	*	*	31	10	*7	17	*8
536.41	15	*	11	*	*	*	12	*	*	*7	*
536.42	17	*5	12	*	*	*	13	*	*	*6	*
536.49	*8	*	*	*	*	*	*5	*	*	*	*
536.8	88	36	52	*	14	23	48	*9	38	21	20
536.9	19	*8	10	*	*	*5	11	*	*	9	*
537	56	23	33	*	*	15	35	10	15	20	12
537.0	24	11	13	*	*	*8	12	*	*6	*8	*
537.8	26	*9	17	—	*	*5	19	*	*7	*9	*6
537.83	10	*	*7	—	—	*	9	*	*	*	*
537.89	10	*	*6	—	*	*	*6	*	*	*	*
540	258	145	113	58	140	40	21	49	64	72	73
540.0	57	34	22	15	24	9	*9	10	14	15	18
540.1	27	16	12	*5	12	*6	*	*	*7	*9	*7
540.9	174	95	79	37	104	24	*8	34	44	48	48
541	17	*9	*8	*	*8	*	*	*	*	*9	*
542	*7	*	*	*	*	*	*	*	*	*	*
543	14	*5	9	*	*9	*	*	*	*	*	*
543.9	12	*	*7	*	*7	*	*	*	*	*	*
550	84	71	13	*9	15	18	41	22	21	28	12
550.1	22	19	*	*	*	*	11	*7	*	*7	*
550.10	16	14	*	*	*	*	*7	*6	*	*	*
550.9	62	52	10	*6	11	14	30	15	17	21	9
550.90	43	35	*8	*	*7	11	22	9	12	15	*7
550.92	13	11	*	*	*	*	*	*	*	*	*
552	81	28	53	*	11	27	43	18	18	29	15
552.0	*6	*	*	*	*	*	*5	*	*	*	*
552.00	*5	*	*	*	*	*	*5	*	*	*	*
552.1	19	*8	11	*	*	*7	*8	*	*	*7	*
552.2	45	14	31	*	*5	18	21	10	*8	18	10
552.20	11	*	*8	—	*	*	*6	*	*	*	*
552.21	29	*9	20	*	*	12	14	*6	*5	11	*6
552.29	*5	*	*	*	*	*	*	*	*	*	*
552.3	*7	*	*	*	*	*	*6	*	*	*	*
553	538	196	342	*6	66	154	311	120	136	191	91
553.1	46	19	27	*	11	17	16	*9	10	16	11
553.2	94	36	58	*	12	39	42	18	25	30	20
553.20	24	*8	16	*	*	10	11	*5	*6	9	*
553.21	65	26	39	*	9	27	29	13	18	20	14
553.3	388	137	251	*	42	97	246	92	98	142	56
555	82	32	50	*5	33	23	20	17	26	29	9
555.0	15	*6	9	*	9	*	*	*	*6	*	*
555.1	11	*	*7	*	*	*	*	*	*	*	*
555.2	*7	*	*	*	*	*	*	*	*	*	*
555.9	50	20	29	*	18	15	14	*9	15	20	*6
556	44	19	26	*	17	11	15	15	9	13	*7
556.9	32	14	18	*	11	*8	11	11	*6	10	*

See footnotes and notes at end of table.

Table 44. Number of all-listed diagnoses for discharges from short-stay hospitals, by ICD-9-CM code, sex, age, and geographic region: United States, 1999—Con.[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table: see Methods section]

ICD-9-CM code	Sex			Age				Region			
	Total	Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	North-east	Midwest	South	West
Number of all-listed diagnoses in thousands											
557	73	22	50	*	*7	16	48	16	15	26	15
557.0	39	12	27	*	*	10	23	*8	*9	14	*9
557.1	*7	*	*5	*	*	*	*5	*	*	*	*
557.9	26	*9	18	*	*	*	19	*7	*5	*9	*6
558	378	151	227	130	73	55	120	81	76	166	54
558.1	13	*5	*8	—	*	*	10	*	*	*	*
558.9	362	145	217	130	72	51	108	77	73	162	50
560	707	288	420	30	103	161	414	137	164	271	134
560.1	304	129	175	9	59	76	160	52	74	120	58
560.2	19	*8	11	*	*	*	14	*	*6	*7	*
560.3	94	31	63	*7	*	*8	75	20	17	48	*9
560.30	*8	*	*7	*	*	*	*8	*	*	*6	*
560.39	85	29	56	*6	*	*7	67	18	17	42	*8
560.8	123	45	78	*	20	30	70	25	30	37	31
560.81	80	29	51	*	13	20	44	17	18	23	22
560.89	43	16	27	*	*6	10	26	*8	12	13	10
560.9	163	72	90	*6	18	44	95	37	36	58	31
562	526	197	328	*	30	123	372	125	134	185	81
562.0	*7	*	*	*	*	*	*5	*	*	*	*
562.1	519	195	324	—	30	122	367	123	133	183	80
562.10	245	91	154	—	*8	42	195	59	69	82	36
562.11	183	64	118	—	20	69	94	45	48	63	27
562.12	73	33	40	—	*	*9	63	15	14	30	14
562.13	18	*7	11	—	*	*	15	*	*	*8	*
564	401	121	280	22	74	82	223	78	112	139	72
564.0	282	100	182	17	40	52	174	55	79	100	48
564.1	84	9	75	*	25	23	34	18	25	28	13
564.8	18	*7	11	*	*	*	*7	*	*	*5	*6
564.89	14	*	9	*	*	*	*6	*	*	*5	*
565	15	*7	*8	*	*6	*	*	*	*	*5	*
565.0	10	*	*6	*	*	*	*	*	*	*	*
565.1	*5	*	*	*	*	*	*	*	*	*	*
566	25	16	*9	*	*9	10	*	*7	*5	*8	*5
567	93	45	48	*	17	34	39	18	22	33	21
567.2	55	30	26	*	10	25	19	12	12	18	14
567.8	12	*	*8	*	*	*	*6	*	*	*	*
567.9	25	11	15	*	*	*5	14	*	*6	10	*
568	147	35	112	*	51	43	49	27	31	61	28
568.0	122	29	93	*	38	39	43	24	26	50	23
568.8	23	*	18	*	12	*	*6	*	*5	10	*5
568.81	15	*	12	*	10	*	*	*	*	*7	*
568.89	*8	*	*6	*	*	*	*	*	*	*	*
569	265	110	155	*7	33	64	161	59	58	104	43
569.0	13	*6	*7	*	*	*	10	*	*	*7	*
569.1	13	*	10	*	*	*	*6	*	*	*	*
569.3	50	20	30	*	*6	12	31	15	*8	19	*8
569.4	30	11	18	*	*7	*6	17	*7	*	13	*6
569.41	11	*	*7	*	*	*	*7	*	*	*	*
569.49	17	*6	10	*	*	*	9	*	*	*7	*
569.5	19	*7	12	—	*	*8	*8	*5	*	*8	*
569.6	23	14	*9	*	*	*5	13	*	*6	*6	*6
569.69	17	12	*5	*	*	*	10	*	*5	*	*
569.8	105	44	61	*	*9	24	71	21	27	40	18
569.81	14	*7	*7	*	*	*7	*5	*	*	*6	*
569.82	12	*5	*7	*	*	*	*9	*	*	*5	*
569.83	29	14	16	*	*	*7	19	*	*8	11	*6
569.84	*7	*	*5	*	*	*	*6	*	*	*	*

See footnotes and notes at end of table.

Table 44. Number of all-listed diagnoses for discharges from short-stay hospitals, by ICD-9-CM code, sex, age, and geographic region: United States, 1999—Con.

[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table: see Methods section]

ICD-9-CM code	Sex			Age				Region			
	Total	Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	North-east	Midwest	South	West
Number of all-listed diagnoses in thousands											
569.85	22	*7	15	*	*	*	20	*	*7	*9	*
569.89	20	*8	11	*	*	*	11	*	*5	*7	*
569.9	*8	*	*6	*	*	*	*	*	*	*6	*
570	15	*9	*6	*	*6	*5	*	*	*	*7	*
571	353	201	151	*	71	175	103	79	66	138	71
571.1	34	24	9	—	14	15	*	10	*	11	*8
571.2	133	96	37	*	24	81	27	28	26	49	30
571.3	22	15	*6	—	*6	12	*	*	*6	*	*8
571.4	14	*	9	*	*	*5	*	*	*	*6	*
571.40	*7	*	*	*	*	*	*	*	*	*	*
571.49	*7	*	*6	*	*	*	*	*	*	*	*
571.5	117	50	66	*	15	49	52	27	21	54	15
571.6	*7	*	*6	*	*	*	*	*	*	*	*
571.8	19	*6	13	*	*	*7	*6	*	*	*8	*
571.9	*5	*	*	*	*	*	*	*	*	*	*
572	131	81	50	*	19	74	37	27	28	48	28
572.0	*6	*	*	*	*	*	*	*	*	*	*
572.2	59	34	25	*	*8	35	17	14	10	23	13
572.3	41	26	14	*	*5	24	11	*8	14	12	*7
572.4	*8	*6	*	—	*	*5	*	*	*	*	*
572.8	16	10	*6	*	*	*8	*5	*	*	*7	*
573	68	29	39	*	14	22	29	19	14	25	10
573.0	*5	*	*	*	*	*	*	*	*	*	*
573.3	32	14	18	*	*6	12	11	11	*6	10	*
573.8	16	*7	10	*	*	*6	*7	*	*	*	*
573.9	13	*6	*7	*	*	*	*8	*	*	*5	*
574	624	202	422	*	165	181	275	145	135	219	126
574.0	139	49	90	*	40	45	54	31	31	44	33
574.00	130	47	84	*	38	42	50	30	29	42	30
574.01	*9	*	*7	—	*	*	*	*	*	*	*
574.1	245	76	169	*	72	72	99	60	51	85	49
574.10	238	73	165	*	70	70	96	59	49	84	47
574.11	*7	*	*	*	*	*	*	*	*	*	*
574.2	143	47	96	*	28	39	75	38	34	52	20
574.20	141	46	95	*	27	38	75	37	34	51	19
574.3	*6	*	*6	—	*	*	*	*	*	*	*
574.4	11	*	*7	—	*	*	*6	*	*	*6	*
574.41	*7	*	*	—	*	*	*	*	*	*	*
574.5	39	11	28	*	*8	10	21	*8	*6	16	9
574.50	23	*6	18	*	*	*7	11	*5	*	9	*
574.51	16	*6	10	*	*	*	9	*	*	*7	*
574.6	*7	*	*	—	*	*	*	*	*	*	*
574.7	16	*5	11	*	*	*	*7	*	*	*7	*
574.70	10	*	*7	*	*	*	*	*	*	*5	*
574.71	*6	*	*	—	*	*	*	*	*	*	*
574.8	9	*	*6	*	*	*	*	*	*	*	*
574.9	*7	*	*5	*	*	*	*	*	*	*	*
575	126	41	84	*	38	36	50	27	38	43	17
575.0	28	12	16	*	*7	*7	14	*	*8	11	*
575.1	59	20	39	*	18	20	20	14	17	20	*8
575.10	14	*6	*8	*	*	*5	*	*	*	*	*
575.11	35	10	25	*	10	14	12	*6	10	13	*6
575.12	9	*	*6	—	*	*	*	*	*	*	*
575.3	*9	*	*7	*	*	*	*5	*	*	*	*
575.8	15	*	11	*	*7	*	*	*	*6	*	*
575.9	*6	*	*6	*	*	*	*	*	*	*	*

See footnotes and notes at end of table.

Table 44. Number of all-listed diagnoses for discharges from short-stay hospitals, by ICD-9-CM code, sex, age, and geographic region: United States, 1999—Con.[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table: see Methods section]

ICD-9-CM code	Sex			Age				Region			
	Total	Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	North-east	Midwest	South	West
Number of all-listed diagnoses in thousands											
576	90	41	50	*	17	19	51	21	20	31	19
576.1	34	18	16	*	*6	*8	19	*9	*7	10	*9
576.2	30	14	16	*	*5	*5	19	*6	*6	14	*
576.8	23	*7	15	*	*	*5	12	*5	*6	*6	*5
577	372	176	196	*6	126	132	108	78	81	146	68
577.0	270	128	142	*	95	91	80	49	61	112	48
577.1	65	32	33	*	22	27	16	18	13	21	13
577.2	18	*8	10	*	*6	*6	*5	*	*	*6	*
577.8	13	*	*8	*	*	*	*	*	*	*	*
577.9	*7	*	*	—	*	*	*	*	*	*	*
578	436	214	221	*9	40	94	292	98	114	152	71
578.0	49	29	20	*	11	10	25	9	*9	20	10
578.1	106	49	57	*	*9	23	69	25	27	34	20
578.9	281	136	145	*	21	61	198	64	79	98	40
579	37	15	22	13	*	*7	11	12	10	9	*6
579.0	*6	*	*	*	*	*	*	*	*	*	*
579.3	10	*	*6	*	*	*	*	*	*	*	*
579.8	14	*6	*8	*6	*	*	*	*	*	*	*
579.9	*6	*	*	*	*	*	*	*	*	*	*
580-629	7,198	2,429	4,769	178	1,917	1,703	3,399	1,498	1,702	2,708	1,291
580	*7	*	*	*	*	*	*	*	*	*	*
581	38	18	20	*	*8	12	13	*8	9	12	*8
581.8	16	*6	10	*	*	*6	*6	*	*	*	*5
581.81	15	*6	9	*	*	*6	*6	*	*	*	*5
581.9	18	10	*8	*	*	*	*7	*	*5	*7	*
582	31	14	17	*	*7	*6	17	9	*7	10	*5
582.8	*7	*	*7	*	*	*	*	*	*	*	*
582.81	*5	*	*5	*	*	*	*	*	*	*	*
582.9	20	12	*8	*	*	*	14	*8	*	*6	*
583	136	62	74	*	27	51	56	34	25	50	27
583.8	123	56	67	*	22	48	53	29	23	46	25
583.81	115	51	65	*	19	45	51	28	21	43	24
583.89	*8	*6	*	*	*	*	*	*	*	*	*
583.9	10	*	*6	*	*	*	*	*	*	*	*
584	417	225	192	*6	37	88	286	106	93	151	68
584.5	51	31	20	*	*7	15	27	9	10	19	12
584.8	*6	*	*	*	*	*	*	*	*	*	*
584.9	361	192	169	*	30	72	256	95	81	129	55
585	323	177	146	*	26	90	204	89	75	109	49
586	47	24	23	*	*	*7	36	12	10	20	*
588	13	*6	*6	*	*	*	*5	*	*	*	*
588.8	*9	*	*	*	*	*	*	*	*	*	*
590	209	34	176	24	98	41	48	36	44	81	48
590.0	*6	*	*5	*	*	*	*	*	*	*	*
590.00	*6	*	*5	*	*	*	*	*	*	*	*
590.1	104	14	90	10	47	23	23	20	24	35	24
590.10	103	13	90	9	47	23	23	20	24	35	24
590.8	93	18	75	13	47	13	20	13	16	42	22
590.80	93	18	75	13	47	13	20	13	16	42	22
591	155	72	83	*9	43	37	66	39	32	51	33

See footnotes and notes at end of table.

Table 44. Number of all-listed diagnoses for discharges from short-stay hospitals, by ICD-9-CM code, sex, age, and geographic region: United States, 1999—Con.

[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table: see Methods section]

ICD-9-CM code	Sex			Age				Region			
	Total	Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	North-east	Midwest	South	West
Number of all-listed diagnoses in thousands											
592	273	156	117	*	111	98	60	57	75	102	39
592.0	115	60	55	*	44	38	32	29	31	38	17
592.1	151	92	59	*	64	60	26	28	39	63	20
592.9	*7	*	*	*	*	*	*	*	*	*	*
593	514	256	258	23	41	97	353	114	141	176	83
593.2	15	*8	*8	*	*	*	10	*6	*	*5	*
593.3	*8	*	*	*	*	*	*	*	*	*	*
593.4	28	13	16	*	*7	*7	13	*6	*8	11	*
593.5	9	*	*6	*	*	*	*	*	*	*	*
593.7	14	*	9	13	*	*	*	*5	*	*	*
593.70	12	*	*8	11	*	*	*	*	*	*	*
593.8	17	*8	9	*	*	*6	*5	*	*6	*	*
593.81	*6	*	*	*	*	*	*	*	*	*	*
593.89	11	*5	*6	*	*	*	*	*	*	*	*
593.9	422	217	205	*	23	77	318	86	117	150	69
594	17	14	*	*	*	*	11	*7	*	*	*
594.1	14	12	*	*	*	*	10	*6	*	*	*
595	70	30	39	*	12	14	42	16	22	24	*7
595.0	16	*8	*8	*	*	*	10	*	*5	*7	*
595.1	*6	*	*6	—	*	*	*	*	*	*	*
595.2	11	*5	*6	*	*	*	*8	*	*	*	*
595.8	12	*7	*	*	*	*	*8	*	*	*	*
595.82	*8	*	*	—	*	*	*7	*	*	*	*
595.9	25	11	14	*	*	*	15	*	9	*9	*
596	169	118	51	*	21	28	115	38	48	54	29
596.0	62	60	*	*	*	*8	54	10	19	20	13
596.3	*5	*	*	*	—	*	*	*	*	*	*
596.4	*8	*	*	*	*	*	*5	*	*	*	*
596.5	63	32	31	*	13	12	34	17	17	20	9
596.54	57	30	27	*	13	12	29	16	16	16	*9
596.59	*5	*	*	*	*	*	*	*	*	*	*
596.8	19	11	*8	*	*	*	12	*5	*6	*6	*
597	*7	*	*	*	*	*	*	*	*	*	*
597.8	*6	*	*	*	*	*	*	*	*	*	*
598	31	26	*5	*	*	*6	20	11	*5	*9	*6
598.8	*7	*6	*	—	*	*	*	*	*	*	*
598.9	23	19	*	*	*	*	15	9	*	*7	*
599	1,961	647	1,313	66	236	287	1,372	424	483	730	324
599.0	1,658	454	1,203	54	187	237	1,179	367	402	617	272
599.6	29	23	*6	*	*	*	21	*	*6	11	*7
599.7	250	165	86	10	42	40	159	48	69	94	39
599.8	12	*	11	*	*	*	*7	*	*	*	*
600	388	388	—	—	*	58	328	104	99	112	74
601	54	54	—	—	*	16	36	13	14	13	14
601.0	16	16	—	—	*	*6	*8	*	*5	*	*5
601.1	23	23	—	—	*	*6	17	*	*6	*6	*7
601.9	11	11	—	—	*	*	*8	*	*	*	*
602	*6	*6	—	—	—	*	*	*	*	*	*
603	15	15	—	*	*	*7	*5	*	*	*5	*
603.9	13	13	—	*	*	*5	*	*	*	*	*
604	17	17	—	*	*	*6	*7	*	*	9	*
604.9	15	15	—	*	*	*6	*6	*	*	*8	*
604.90	11	11	—	*	*	*	*	*	*	*7	*
605	11	11	—	*	*	*	*5	*	*	*	*
607	23	23	—	*	*	*8	*8	*	*	13	*
607.8	15	15	—	*	*	*5	*7	*	*	*8	*
607.84	9	9	—	—	*	*	*	*	*	*	*

See footnotes and notes at end of table.

Table 44. Number of all-listed diagnoses for discharges from short-stay hospitals, by ICD-9-CM code, sex, age, and geographic region: United States, 1999—Con.[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table: see Methods section]

ICD-9-CM code	Sex			Age				Region			
	Total	Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	North-east	Midwest	South	West
Number of all-listed diagnoses in thousands											
608	21	21	—	*	*6	*5	*7	*	*6	*8	*
608.4	*6	*6	—	*	*	*	*	*	*	*	*
608.8	*8	*8	—	*	*	*	*	*	*	*	*
610	29	*	28	—	*5	14	10	*8	*7	*9	*6
610.1	21	*	21	—	*	11	*7	*7	*	*6	*
611	51	*	48	*	21	17	12	17	*8	15	11
611.0	10	*	9	*	*	*	*	*	*	*	*
611.1	14	*	13	*	*8	*	*	*6	*	*5	*
611.7	15	*	15	*	*	*	*7	*	*	*	*
611.72	12	*	12	—	*	*	*6	*	*	*	*
611.8	*8	*	*8	—	*	*	*	*	*	*	*
614	254	—	254	*	180	58	15	46	50	102	55
614.1	28	—	28	—	18	*8	*	*	*6	10	*7
614.2	11	—	11	*	*8	*	*	*	*	*	*
614.3	12	—	12	*	*9	*	*	*	*	*	*
614.4	*8	—	*8	*	*5	*	*	*	*	*	*
614.6	166	—	166	*	115	41	10	27	32	71	37
614.9	21	—	21	*	19	*	—	*	*5	*9	*
615	20	—	20	*	18	*	*	*	*5	9	*
615.1	*5	—	*5	—	*	*	*	*	*	*	*
615.9	10	—	10	*	10	*	*	*	*	*5	*
616	159	—	159	*	92	51	15	29	39	63	27
616.0	123	—	123	—	69	45	9	21	32	49	21
616.1	31	—	31	*	20	*5	*5	*7	*7	12	*
616.10	30	—	30	*	19	*5	*5	*7	*7	12	*
617	295	—	295	—	187	99	*8	41	61	130	63
617.0	171	—	171	—	97	66	*7	25	35	72	39
617.1	43	—	43	—	31	11	*	*	*9	21	*9
617.2	10	—	10	—	*8	*	*	*	*	*	*
617.3	43	—	43	—	32	11	—	*6	11	20	*7
617.5	*5	—	*5	—	*	*	—	*	*	*	*
617.9	16	—	16	—	11	*5	*	*	*	*9	*
618	290	—	290	—	71	116	103	39	58	128	65
618.0	80	—	80	—	14	38	28	11	13	36	19
618.1	32	—	32	—	17	*8	*7	*7	*7	15	*
618.2	27	—	27	—	10	12	*	*	*8	9	*5
618.3	22	—	22	—	*	9	10	*	*	*8	*5
618.4	31	—	31	—	*8	13	9	*	*9	14	*
618.5	35	—	35	—	*	14	19	*	*6	17	*7
618.6	39	—	39	—	*7	11	21	*	*7	18	12
618.8	23	—	23	—	*8	11	*	*	*	10	*9
619	10	—	10	*	*	*	*5	*	*	*	*
619.1	*7	—	*7	*	*	*	*	*	*	*	*
620	296	—	296	*7	182	88	20	51	68	117	60
620.0	51	—	51	*	32	18	*	*8	16	23	*6
620.1	63	—	63	*	42	19	*	9	17	24	13
620.2	129	—	129	*	82	32	11	23	26	52	28
620.5	*8	—	*8	*	*5	*	*	*	*	*	*
620.8	36	—	36	*	17	14	*	*8	*8	11	9
621	94	—	94	—	32	45	17	19	23	34	18
621.0	30	—	30	—	9	15	*	*7	*7	*8	*7
621.2	12	—	12	—	*7	*	*	*	*	*6	*
621.3	27	—	27	—	*	18	*5	*5	*6	11	*
621.5	*5	—	*5	—	*	*	*	*	*	*	*
621.8	14	—	14	—	*7	*	*	*	*	*	*

See footnotes and notes at end of table.

Table 44. Number of all-listed diagnoses for discharges from short-stay hospitals, by ICD-9-CM code, sex, age, and geographic region: United States, 1999—Con.

[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table: see Methods section]

ICD-9-CM code	Sex			Age				Region			
	Total	Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	North-east	Midwest	South	West
Number of all-listed diagnoses in thousands											
622	32	—	32	*	20	*9	*	*	*7	14	*6
622.1	20	—	20	—	15	*	*	*	*	*9	*
622.7	*7	—	*7	*	*	*	*	*	*	*	*
623	22	—	22	*	10	*	*7	*	*	10	*
623.8	15	—	15	*	*7	*	*	*	*	*6	*
624	*5	—	*5	*	*	*	*	*	*	*	*
625	360	—	360	*	208	106	47	40	76	175	69
625.0	25	—	25	—	23	*	—	*	*6	12	*
625.3	88	—	88	*	76	12	—	*	16	52	16
625.6	157	—	157	*	38	77	43	19	33	69	35
625.9	79	—	79	—	63	13	*	13	19	37	11
626	268	—	268	*	175	88	*	36	66	115	51
626.2	188	—	188	*	126	60	*	25	45	81	37
626.6	*6	—	*6	—	*	*	*	*	*	*	*
626.8	54	—	54	*	34	18	*	*7	14	23	10
626.9	11	—	11	*	*7	*	*	*	*	*5	*
627	51	—	51	—	*5	28	18	9	15	19	*8
627.1	30	—	30	—	*	17	12	*7	*7	11	*
627.2	*8	—	*8	—	*	*5	*	*	*	*	*
627.9	*5	—	*5	—	—	*	*	*	*	*	*
630-677	8,350	*	8,350	17	8,312	22	—	1,422	1,822	3,082	2,024
632	17	—	17	—	16	*	—	*	*	*7	*
633	36	—	36	*	35	*	—	*7	*6	15	*9
633.1	33	—	33	*	32	*	—	*6	*5	14	*8
634	26	—	26	*	25	—	—	*8	*	*9	*5
634.9	19	—	19	*	18	—	—	*6	*	*6	*
634.91	14	—	14	*	14	—	—	*5	*	*	*
635	9	—	9	—	9	—	—	*	*	*	*
635.9	*5	—	*5	—	*5	—	—	*	*	*	*
639	10	—	10	—	10	—	—	*	*	*	*
640	10	—	10	—	10	—	—	*	*	*	*
640.0	*7	—	*7	—	*7	—	—	*	*	*	*
640.03	*7	—	*7	—	*7	—	—	*	*	*	*
641	87	—	87	*	87	*	—	15	18	33	22
641.0	*7	—	*7	*	*7	—	—	*	*	*	*
641.01	*5	—	*5	*	*5	—	—	*	*	*	*
641.1	17	—	17	—	17	—	—	*	*	*7	*
641.11	10	—	10	—	10	—	—	*	*	*	*
641.13	*7	—	*7	—	*7	—	—	*	*	*	*
641.2	47	—	47	*	47	—	—	*7	9	17	13
641.21	41	—	41	*	41	—	—	*6	*8	16	12
641.23	*5	—	*5	—	*5	—	—	*	*	*	*
641.9	12	—	12	*	12	*	—	*	*	*	*
641.91	*5	—	*5	*	*	—	—	*	*	*	*
641.93	*7	—	*7	—	*7	*	—	*	*	*	*
642	326	—	326	*	323	*	—	54	67	141	65
642.0	23	—	23	—	23	*	—	*	*	12	*
642.01	18	—	18	—	18	*	—	*	*	10	*
642.3	112	—	112	—	111	*	—	18	25	43	26
642.31	99	—	99	—	99	*	—	16	23	37	23
642.33	11	—	11	—	10	*	—	*	*	*5	*
642.4	97	—	97	*	96	—	—	14	19	47	17
642.41	85	—	85	*	84	—	—	13	15	42	15
642.43	*9	—	*9	*	*9	—	—	*	*	*	*

See footnotes and notes at end of table.

Table 44. Number of all-listed diagnoses for discharges from short-stay hospitals, by ICD-9-CM code, sex, age, and geographic region: United States, 1999—Con.[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table: see Methods section]

ICD-9-CM code	Sex			Age				Region			
	Total	Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	North-east	Midwest	South	West
Number of all-listed diagnoses in thousands											
642.5	34	—	34	*	33	*	—	*6	*7	14	*7
642.51	31	—	31	*	30	*	—	*6	*6	13	*7
642.7	*9	—	*9	—	*9	*	—	*	*	*	*
642.71	*7	—	*7	—	*7	*	—	*	*	*	*
642.9	42	—	42	*	42	*	—	*8	*7	19	*8
642.91	36	—	36	*	36	*	—	*7	*6	15	*8
643	50	—	50	*	50	*	—	*8	11	24	*7
643.0	20	—	20	—	20	—	—	*	*	12	*
643.03	19	—	19	—	19	—	—	*	*	12	*
643.1	21	—	21	—	21	*	—	*	*6	*7	*
643.13	21	—	21	—	21	*	—	*	*6	*7	*
644	364	—	364	*	361	*	—	62	76	154	72
644.0	111	—	111	*	110	*	—	20	25	47	20
644.03	111	—	111	*	110	*	—	20	25	47	20
644.1	18	—	18	—	18	—	—	*	*	10	*
644.13	18	—	18	—	18	—	—	*	*	10	*
644.2	234	—	234	*	233	*	—	41	48	97	48
644.21	234	—	234	*	232	*	—	41	48	97	48
645	131	—	131	*	130	*	—	22	30	39	40
645.0	131	—	131	*	130	*	—	22	30	39	40
645.01	128	—	128	*	127	*	—	22	30	39	38
646	233	—	233	*	232	*	—	43	47	100	44
646.1	17	—	17	—	17	—	—	*	*	*7	*7
646.11	14	—	14	—	14	—	—	*	*	*5	*6
646.2	12	—	12	—	12	—	—	*	*	*6	*
646.21	*6	—	*6	—	*6	—	—	*	*	*	*
646.23	*5	—	*5	—	*5	—	—	*	*	*	*
646.6	151	—	151	*	151	*	—	30	31	67	24
646.61	91	—	91	*	91	*	—	23	20	35	13
646.62	*5	—	*5	*	*5	—	—	*	*	*	*
646.63	51	—	51	*	51	—	—	*6	*9	27	10
646.8	39	—	39	*	39	—	—	*6	10	15	*9
646.81	22	—	22	*	22	—	—	*	*5	*7	*6
646.83	14	—	14	*	14	—	—	*	*	*8	*
647	119	—	119	*	118	*	—	16	26	54	24
647.6	43	—	43	*	42	—	—	*5	*6	21	10
647.61	35	—	35	*	35	—	—	*	*5	17	9
647.63	*7	—	*7	*	*6	—	—	*	*	*	*
647.8	68	—	68	*	67	*	—	*8	16	30	13
647.81	55	—	55	*	55	*	—	*7	13	24	10
647.83	11	—	11	*	11	—	—	*	*	*5	*
648	964	*	964	*	960	*	—	176	228	344	216
648.0	35	—	35	*	35	*	—	*7	*6	13	*9
648.01	22	—	22	*	22	*	—	*	*	*8	*7
648.03	12	—	12	—	12	—	—	*	*	*	*
648.1	42	—	42	—	42	*	—	13	10	9	10
648.11	35	—	35	—	35	*	—	10	*9	*8	*9
648.13	*7	—	*7	—	*7	—	—	*	*	*	*
648.2	275	—	275	*	275	*	—	44	70	96	65
648.21	164	—	164	*	164	*	—	23	47	59	35
648.22	86	—	86	*	86	—	—	16	17	27	25
648.23	21	—	21	*	21	—	—	*	*5	*7	*
648.3	10	—	10	—	10	—	—	*	*	*	*
648.31	*7	—	*7	—	*7	—	—	*	*	*	*
648.4	103	*	103	*	103	*	—	18	28	36	21
648.41	80	—	80	*	80	*	—	13	23	25	19
648.43	19	—	19	*	19	*	—	*	*	9	*

See footnotes and notes at end of table.

Table 44. Number of all-listed diagnoses for discharges from short-stay hospitals, by ICD-9-CM code, sex, age, and geographic region: United States, 1999—Con.

[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table: see Methods section]

ICD-9-CM code	Sex			Age				Region			
	Total	Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	North-east	Midwest	South	West
Number of all-listed diagnoses in thousands											
648.6	37	—	37	*	36	*	—	*8	*8	17	*
648.61	32	—	32	*	31	*	—	*7	*6	16	*
648.8	134	—	134	—	133	*	—	31	32	42	30
648.81	118	—	118	—	117	*	—	29	28	36	26
648.83	16	—	16	—	16	*	—	*	*	*6	*
648.9	319	—	319	*	318	*	—	48	71	125	75
648.91	239	—	239	*	239	*	—	34	57	91	57
648.92	*9	—	*9	—	*9	—	—	*	*	*	*
648.93	69	—	69	*	68	*	—	12	11	31	15
650	572	—	572	*	570	*	—	88	105	259	119
651	56	—	56	—	55	*	—	11	12	22	11
651.0	50	—	50	—	50	*	—	10	10	21	11
651.01	38	—	38	—	37	*	—	*7	*7	15	*8
651.03	13	—	13	—	13	*	—	*	*	*5	*
652	261	—	261	*	261	*	—	50	62	95	55
652.1	*8	—	*8	—	*8	—	—	*	*	*	*
652.11	*6	—	*6	—	*6	—	—	*	*	*	*
652.2	117	—	117	*	117	*	—	22	25	45	26
652.21	115	—	115	*	114	*	—	21	25	44	25
652.3	30	—	30	—	30	*	—	*6	*7	11	*6
652.31	30	—	30	—	30	*	—	*6	*6	11	*6
652.4	*6	—	*6	—	*6	*	—	*	*	*	*
652.41	*6	—	*6	—	*6	*	—	*	*	*	*
652.5	33	—	33	—	33	*	—	*6	*8	15	*
652.51	33	—	33	—	33	*	—	*6	*8	15	*
652.6	13	—	13	—	13	*	—	*	*	*	*
652.61	13	—	13	—	12	*	—	*	*	*	*
652.8	47	—	47	—	47	*	—	9	14	12	12
652.81	47	—	47	—	46	*	—	*9	14	12	12
653	115	—	115	*	115	—	—	14	23	48	30
653.4	91	—	91	*	91	—	—	10	17	40	23
653.41	91	—	91	*	91	—	—	10	17	40	23
653.5	15	—	15	—	15	—	—	*	*	*	*
653.51	15	—	15	—	15	—	—	*	*	*	*
654	546	—	546	—	544	*	—	97	113	208	128
654.0	9	—	9	—	9	—	—	*	*	*	*
654.01	*7	—	*7	—	*7	—	—	*	*	*	*
654.1	24	—	24	—	23	*	—	*	*	*9	*8
654.11	21	—	21	—	21	*	—	*	*	*8	*7
654.2	448	—	448	—	448	*	—	79	96	166	106
654.21	438	—	438	—	437	*	—	78	95	164	101
654.23	10	—	10	—	10	—	—	*	*	*	*5
654.4	16	—	16	—	16	*	—	*	*	*5	*
654.41	13	—	13	—	13	*	—	*	*	*	*
654.5	25	—	25	—	25	*	—	*	*	13	*
654.51	15	—	15	—	15	—	—	*	*	*9	*
654.53	10	—	10	—	10	*	—	*	*	*	*
654.6	11	—	11	—	11	*	—	*	*	*5	*
654.61	*8	—	*8	—	*8	—	—	*	*	*	*
654.9	*7	—	*7	—	*7	—	—	*	*	*	*
654.91	*7	—	*7	—	*7	—	—	*	*	*	*
655	31	—	31	—	31	*	—	*	*7	10	10
655.7	13	—	13	—	13	—	—	*	*	*	*6
655.71	11	—	11	—	11	—	—	*	*	*	*
655.8	11	—	11	—	11	—	—	*	*	*	*
655.81	10	—	10	—	10	—	—	*	*	*	*

See footnotes and notes at end of table.

Table 44. Number of all-listed diagnoses for discharges from short-stay hospitals, by ICD-9-CM code, sex, age, and geographic region: United States, 1999—Con.[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table: see Methods section]

ICD-9-CM code	Sex			Age				Region			
	Total	Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	North-east	Midwest	South	West
Number of all-listed diagnoses in thousands											
656	398	—	398	*	397	*	—	60	94	150	94
656.1	69	—	69	—	69	—	—	*9	16	31	14
656.11	66	—	66	—	66	—	—	*9	16	28	13
656.3	43	—	43	*	43	*	—	13	10	13	*8
656.31	43	—	43	*	43	*	—	13	10	13	*8
656.4	16	—	16	—	16	—	—	*	*	*9	*
656.41	16	—	16	—	16	—	—	*	*	*9	*
656.5	61	—	61	*	61	*	—	11	11	22	17
656.51	58	—	58	*	58	*	—	11	10	22	16
656.6	92	—	92	*	92	—	—	10	22	35	24
656.61	92	—	92	*	91	—	—	10	22	35	24
656.7	23	—	23	—	23	—	—	*	13	*	*
656.71	23	—	23	—	23	—	—	*	13	*	*
656.8	88	—	88	*	88	—	—	11	19	34	24
656.81	87	—	87	*	87	—	—	10	19	34	24
657	21	—	21	—	21	—	—	*5	*7	*6	*
657.0	21	—	21	—	21	—	—	*5	*7	*6	*
657.01	19	—	19	—	19	—	—	*5	*6	*	*
658	360	—	360	*	358	*	—	72	70	123	95
658.0	88	—	88	*	87	*	—	22	15	29	22
658.01	82	—	82	*	81	*	—	21	14	28	19
658.03	*6	—	*6	—	*5	*	—	*	*	*	*
658.1	146	—	146	*	145	*	—	25	30	54	37
658.11	140	—	140	*	139	*	—	24	29	51	36
658.13	*6	—	*6	—	*6	—	—	*	*	*	*
658.2	68	—	68	*	68	*	—	14	16	22	16
658.21	66	—	66	*	66	*	—	14	16	21	15
658.4	53	—	53	*	52	—	—	10	*7	18	18
658.41	52	—	52	*	51	—	—	10	*7	17	18
659	605	—	605	*	600	*	—	91	114	205	196
659.1	37	—	37	—	36	*	—	*7	*7	11	12
659.11	30	—	30	—	29	*	—	*6	*6	9	*9
659.13	*7	—	*7	—	*7	—	—	*	*	*	*
659.2	39	—	39	—	39	—	—	*6	*6	16	11
659.21	39	—	39	—	39	—	—	*6	*6	16	11
659.4	17	—	17	—	17	*	—	*	*	*5	*5
659.41	17	—	17	—	17	*	—	*	*	*5	*5
659.5	19	—	19	—	18	*	—	*	*	*	*9
659.51	18	—	18	—	18	*	—	*	*	*	*8
659.6	118	—	118	—	117	*	—	18	18	38	44
659.61	113	—	113	—	112	*	—	18	17	37	41
659.7	303	—	303	*	303	*	—	46	57	116	84
659.71	300	—	300	*	300	*	—	45	57	115	83
659.8	61	—	61	*	59	—	—	*7	17	*9	28
659.81	58	—	58	*	57	—	—	*6	17	*8	27
659.9	*9	—	*9	*	*9	—	—	*	*	*	*
659.91	*9	—	*9	*	*9	—	—	*	*	*	*
660	192	—	192	*	191	*	—	27	49	67	50
660.0	36	—	36	*	36	—	—	*	*8	14	*8
660.01	36	—	36	*	36	—	—	*	*8	14	*8
660.1	38	—	38	*	38	—	—	*	*6	13	15
660.11	38	—	38	*	38	—	—	*	*6	13	15
660.2	*6	—	*6	—	*6	—	—	*	*	*	*
660.21	*6	—	*6	—	*6	—	—	*	*	*	*
660.3	36	—	36	*	35	*	—	*6	10	14	*7
660.31	36	—	36	*	35	*	—	*6	10	14	*7
660.4	57	—	57	—	57	*	—	11	18	16	12
660.41	57	—	57	—	57	*	—	11	18	16	12

See footnotes and notes at end of table.

Table 44. Number of all-listed diagnoses for discharges from short-stay hospitals, by ICD-9-CM code, sex, age, and geographic region: United States, 1999—Con.

[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table: see Methods section]

ICD-9-CM code	Sex			Age				Region			
	Total	Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	North-east	Midwest	South	West
Number of all-listed diagnoses in thousands											
660.6	*8	—	*8	—	*8	—	—	*	*	*	*
660.61	*8	—	*8	—	*8	—	—	*	*	*	*
660.7	10	—	10	—	10	—	—	*	*	*	*
660.71	10	—	10	—	10	—	—	*	*	*	*
661	338	—	338	*	337	—	—	57	77	120	84
661.0	75	—	75	*	75	—	—	12	20	26	17
661.01	74	—	74	*	74	—	—	11	19	26	17
661.1	67	—	67	*	67	—	—	13	13	26	16
661.11	67	—	67	*	67	—	—	13	13	25	16
661.2	92	—	92	*	91	—	—	17	23	34	19
661.21	92	—	92	*	91	—	—	17	22	34	19
661.3	89	—	89	—	89	—	—	12	20	26	31
661.31	89	—	89	—	89	—	—	12	20	26	31
661.4	*7	—	*7	—	*7	—	—	*	*	*	*
661.41	*7	—	*7	—	*7	—	—	*	*	*	*
661.9	*7	—	*7	—	*7	—	—	*	*	*	*
661.91	*6	—	*6	—	*6	—	—	*	*	*	*
662	43	—	43	*	43	—	—	10	*9	11	13
662.0	*8	—	*8	*	*8	—	—	*	*	*	*
662.01	*8	—	*8	*	*8	—	—	*	*	*	*
662.1	*6	—	*6	—	*6	—	—	*	*	*	*
662.11	*6	—	*6	—	*6	—	—	*	*	*	*
662.2	29	—	29	—	29	—	—	*7	*6	*7	*9
662.21	29	—	29	—	29	—	—	*7	*6	*7	*9
663	721	—	721	*	718	*	—	122	163	240	195
663.0	12	—	12	—	12	—	—	*	*	*	*
663.01	12	—	12	—	12	—	—	*	*	*	*
663.1	95	—	95	*	94	*	—	17	19	30	29
663.11	95	—	95	*	94	*	—	17	19	30	29
663.2	30	—	30	*	30	—	—	*6	*8	9	*7
663.21	30	—	30	*	30	—	—	*6	*8	9	*7
663.3	545	—	545	*	543	*	—	92	126	184	142
663.31	545	—	545	*	543	*	—	92	126	184	142
663.4	17	—	17	—	17	—	—	*	*	*	*7
663.41	17	—	17	—	17	—	—	*	*	*	*7
663.5	*5	—	*5	—	*5	—	—	*	*	*	*
663.8	13	—	13	—	13	—	—	*	*	*7	*
663.81	13	—	13	—	13	—	—	*	*	*7	*
664	1,108	—	1,108	*	1,105	*	—	210	262	344	291
664.0	472	—	472	*	471	*	—	93	111	149	120
664.01	471	—	471	*	470	*	—	92	111	148	120
664.1	445	—	445	*	444	*	—	83	102	131	130
664.11	444	—	444	*	443	*	—	83	102	129	130
664.2	111	—	111	*	111	—	—	22	29	38	22
664.21	111	—	111	*	111	—	—	22	29	38	22
664.3	55	—	55	*	54	*	—	9	15	20	11
664.31	54	—	54	*	53	*	—	9	15	20	11
664.4	11	—	11	—	11	—	—	*	*	*	*
664.41	11	—	11	—	11	—	—	*	*	*	*
664.9	*7	—	*7	—	*7	—	—	*	*	*	*
664.91	*7	—	*7	—	*7	—	—	*	*	*	*
665	164	—	164	*	164	*	—	26	38	68	33
665.3	11	—	11	—	11	—	—	*	*	*6	*
665.31	11	—	11	—	11	—	—	*	*	*6	*
665.4	50	—	50	*	50	—	—	*8	*8	24	9
665.41	50	—	50	*	50	—	—	*8	*8	24	9
665.5	93	—	93	*	92	*	—	16	24	35	18
665.51	92	—	92	*	92	*	—	16	24	35	18

See footnotes and notes at end of table.

Table 44. Number of all-listed diagnoses for discharges from short-stay hospitals, by ICD-9-CM code, sex, age, and geographic region: United States, 1999—Con.[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table: see Methods section]

ICD-9-CM code	Sex			Age				Region			
	Total	Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	North-east	Midwest	South	West
Number of all-listed diagnoses in thousands											
666	107	—	107	*	107	—	—	16	26	39	26
666.0	13	—	13	—	13	—	—	*	*	*	*
666.02	13	—	13	—	13	—	—	*	*	*	*
666.1	79	—	79	*	79	—	—	11	19	30	19
666.12	79	—	79	*	79	—	—	11	19	30	19
666.2	11	—	11	—	11	—	—	*	*	*	*
666.22	*7	—	*7	—	*7	—	—	*	*	*	*
667	24	—	24	—	24	*	—	*	*6	*8	*6
667.0	17	—	17	—	17	*	—	*	*	*6	*
667.02	16	—	16	—	16	*	—	*	*	*5	*
667.1	*7	—	*7	—	*7	—	—	*	*	*	*
667.12	*7	—	*7	—	*7	—	—	*	*	*	*
668	*6	—	*6	*	*6	—	—	*	*	*	*
668.8	*5	—	*5	*	*	—	—	*	*	*	*
669	170	—	170	*	169	*	—	17	39	71	42
669.5	96	—	96	*	95	—	—	*8	25	45	18
669.51	96	—	96	*	95	—	—	*8	25	45	18
669.7	16	—	16	—	16	—	—	*	*	10	*
669.71	16	—	16	—	16	—	—	*	*	10	*
669.8	47	—	47	—	47	—	—	*6	11	12	18
669.81	45	—	45	—	45	—	—	*6	10	12	16
670	24	—	24	—	24	—	—	*	*	11	*
670.0	24	—	24	—	24	—	—	*	*	11	*
670.02	19	—	19	—	19	—	—	*	*	*8	*
670.04	*5	—	*5	—	*5	—	—	*	*	*	*
671	33	—	33	—	33	—	—	*7	*8	13	*6
671.2	*6	—	*6	—	*6	—	—	*	*	*	*
671.8	14	—	14	—	14	—	—	*	*	*5	*
671.81	*7	—	*7	—	*7	—	—	*	*	*	*
671.82	*6	—	*6	—	*6	—	—	*	*	*	*
672	19	—	19	*	19	—	—	*	*	*6	*6
672.0	19	—	19	*	19	—	—	*	*	*6	*6
672.02	17	—	17	*	17	—	—	*	*	*5	*6
674	32	—	32	—	32	—	—	*	*8	12	*8
674.3	12	—	12	—	12	—	—	*	*	*	*
674.32	*6	—	*6	—	*6	—	—	*	*	*	*
674.34	*5	—	*5	—	*5	—	—	*	*	*	*
674.8	13	—	13	—	13	—	—	*	*	*6	*
674.82	*9	—	*9	—	*9	—	—	*	*	*	*
676	11	—	11	—	10	*	—	*	*	*8	*
680-709	1,833	885	948	147	354	440	892	429	428	679	296
680	*5	*	*	*	*	*	*	*	*	*	*
681	48	29	19	*5	10	16	17	12	13	16	*7
681.0	14	*8	*6	*	*6	*	*	*	*	*	*
681.00	11	*6	*	*	*5	*	*	*	*	*	*
681.1	33	21	12	*	*	11	16	*8	*9	11	*6
681.10	28	19	*9	*	*	9	14	*6	*8	9	*
681.11	*5	*	*	*	*	*	*	*	*	*	*
682	743	367	376	46	156	209	331	180	182	254	126
682.0	30	14	17	*7	12	*7	*6	*7	*7	11	*5
682.1	14	*7	*8	*	*	*	*	*	*	*	*
682.2	87	41	46	*	22	34	27	18	24	29	16
682.3	77	43	34	*	25	19	29	17	15	26	19

See footnotes and notes at end of table.

Table 44. Number of all-listed diagnoses for discharges from short-stay hospitals, by ICD-9-CM code, sex, age, and geographic region: United States, 1999—Con.

[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table: see Methods section]

ICD-9-CM code	Sex			Age				Region			
	Total	Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	North-east	Midwest	South	West
Number of all-listed diagnoses in thousands											
682.4	34	22	12	*	11	9	*9	*9	*8	*8	*9
682.5	17	*7	9	*	*5	*	*6	*	*	*6	*
682.6	340	161	179	11	57	98	174	94	85	117	44
682.7	118	61	56	*6	16	31	65	26	27	44	20
682.8	10	*5	*	*	*	*	*	*	*	*	*
682.9	16	*6	10	*	*	*	9	*	*6	*	*
683	*7	*	*5	*	*	*	*	*	*	*	*
684	*7	*	*	*	*	*	*	*	*	*	*
686	20	10	10	*	*7	*	*7	*	*6	*8	*
686.9	*9	*	*	*	*	*	*	*	*	*	*
690	12	*8	*	*	*	*	*	*	*	*	*
690.1	11	*8	*	*	*	*	*	*	*	*	*
690.10	*8	*6	*	*	*	*	*	*	*	*	*
691	30	14	16	26	*	*	*	*	*7	11	*9
691.0	24	9	14	22	*	—	*	*	*	*9	*7
691.8	*7	*	*	*	*	*	*	*	*	*	*
692	52	26	26	11	11	11	19	16	13	14	*9
692.9	45	23	22	10	*9	10	17	14	12	13	*6
693	45	15	29	*	14	11	17	12	9	16	*8
693.0	43	14	28	*	13	10	17	11	*9	15	*8
694	*7	*	*	—	*	*	*6	*	*	*	*
695	26	10	16	*	*7	*7	*9	*7	*5	9	*
695.1	*5	*	*	*	*	*	*	*	*	*	*
695.4	*7	*	*	—	*	*	*	*	*	*	*
695.8	*6	*	*	*	*	*	*	*	*	*	*
695.89	*5	*	*	*	*	*	*	*	*	*	*
696	48	23	24	—	13	12	23	13	12	15	*8
696.0	*7	*	*	—	*	*	*	*	*	*	*
696.1	40	19	21	—	10	11	19	11	9	14	*6
698	30	11	19	*	10	*8	11	*5	*7	11	*7
698.3	*6	*	*	*	*	*	*	*	*	*	*
698.9	20	*8	11	*	*6	*6	*7	*	*	*6	*5
701	20	*6	14	*	*6	*6	*6	*	*	*8	*
701.4	*5	*	*	*	*	*	*	*	*	*	*
702	*8	*	*	—	*	*	*	*	*	*	*
702.1	*5	*	*	—	*	*	*	*	*	*	*
702.19	*5	*	*	—	*	*	*	*	*	*	*
703	16	*7	*9	*	*	*	11	*	*	*6	*
703.0	9	*	*6	*	*	*	*6	*	*	*	*
703.8	*6	*	*	*	*	*	*5	*	*	*	*
704	15	*7	*7	*	*7	*	*	*	*	*9	*
704.0	9	*	*	*	*	*	*	*	*	*6	*
704.00	*9	*	*	*	*	*	*	*	*	*6	*
705	*8	*	*5	*	*	*	*	*	*	*	*
705.8	*5	*	*	*	*	*	*	*	*	*	*
706	20	11	*8	*	*7	*	*7	*5	*6	*6	*
706.1	*5	*	*	*	*	*	—	*	*	*	*
706.2	10	*5	*	*	*	*	*	*	*	*	*
707	567	276	291	*	53	119	391	135	126	224	82
707.0	316	148	169	*	26	49	240	69	64	138	45
707.1	230	121	109	*	25	66	139	61	53	79	36
707.8	15	*	10	*	*	*	9	*	*8	*	*
707.9	*6	*	*	—	*	*	*	*	*	*	*
708	18	*6	11	*	*5	*5	*	*	*5	*	*5
708.0	10	*	*6	*	*	*	*	*	*	*	*

See footnotes and notes at end of table.

Table 44. Number of all-listed diagnoses for discharges from short-stay hospitals, by ICD-9-CM code, sex, age, and geographic region: United States, 1999—Con.[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table: see Methods section]

ICD-9-CM code	Sex			Age				Region			
	Total	Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	North-east	Midwest	South	West
Number of all-listed diagnoses in thousands											
709	76	35	40	27	22	11	16	*8	10	51	*7
709.2	46	22	24	22	16	*5	*	*	*	38	*
709.8	10	*	*6	*	*	*	*5	*	*	*	*
709.9	13	*6	*7	*	*	*	*6	*	*	*7	*
710-739	5,054	1,810	3,244	72	755	1,279	2,949	1,049	1,340	1,744	920
710	143	17	125	*	47	52	41	30	30	54	28
710.0	98	12	86	*	39	35	23	20	15	39	23
710.1	19	*	17	-	*	*8	*7	*	*6	*6	*
710.2	10	*	9	-	*	*5	*	*	*	*	*
710.3	*6	*	*5	*	*	*	*	*	*	*	*
710.4	*6	*	*	-	*	*	*	*	*	*	*
711	39	26	13	*5	12	*8	15	*8	*8	16	*8
711.0	36	24	12	*5	10	*7	14	*7	*7	15	*7
711.06	15	11	*	*	*5	*	*6	*	*	*7	*
712	*6	*	*	-	*	*	*	*	*	*	*
713	*8	*	*5	*	*	*	*	*	*	*	*
713.5	*6	*	*	-	*	*	*	*	*	*	*
714	225	69	156	*	18	62	144	55	56	76	39
714.0	211	62	149	*	15	60	136	51	53	71	36
714.8	*7	*	*	-	*	*	*5	*	*	*	*
715	1,281	425	856	*	36	242	1,002	266	394	424	197
715.0	33	10	24	-	*	*	28	11	10	*6	*5
715.00	*7	*	*6	-	*	*	*6	*	*	*	*
715.09	26	*9	18	-	*	*	22	11	*6	*5	*
715.1	50	18	31	-	*	10	38	*7	11	15	17
715.15	17	*7	9	-	*	*	14	*	*	*6	*7
715.16	29	9	20	-	*	*8	21	*	*7	*7	10
715.2	*6	*	*	*	*	*	*	*	*	*	*
715.3	298	119	179	*	11	81	204	69	82	96	51
715.30	*7	*	*5	-	-	*	*6	*	*	*	*
715.31	14	*6	*8	*	*	*	*9	*	*	*6	*
715.35	89	35	54	-	*	24	60	21	26	22	19
715.36	177	72	105	*	*	50	122	41	49	60	26
715.37	*5	*	*	-	*	*	*	*	*	*	*
715.8	48	13	36	-	*	*7	41	*8	23	11	*6
715.89	48	13	35	-	*	*7	40	*8	22	11	*6
715.9	847	263	583	*	20	137	688	170	268	293	116
715.90	542	149	393	*	10	64	468	103	177	208	54
715.91	18	*7	10	-	*	*	13	*	*6	*	*
715.95	86	35	51	*	*	25	57	21	24	25	15
715.96	175	64	111	-	*	39	133	34	55	49	38
715.97	*7	*	*	-	*	*	*	*	*	*	*
715.98	15	*	12	-	*	*	12	*6	*	*	*
716	224	82	142	*	21	57	144	43	61	89	32
716.1	14	*9	*5	-	*	*7	*	*	*	*	*
716.16	*7	*	*	-	*	*	*	*	*	*	*
716.5	*8	*	*	*	*	*	*	*	*	*	*
716.59	*7	*	*	*	*	*	*	*	*	*	*
716.9	198	69	130	*	15	47	135	36	57	80	26
716.90	144	47	97	*	*8	34	102	27	40	63	15
716.91	*6	*	*	-	*	*	*	*	*	*	*
716.95	*8	*	*	*	*	*	*	*	*	*	*
716.96	17	*7	11	*	*	*	11	*	*	*6	*5
716.98	*7	*	*	-	*	*	*	*	*	*	*
716.99	*9	*	*6	*	*	*	*6	*	*	*	*

See footnotes and notes at end of table.

Table 44. Number of all-listed diagnoses for discharges from short-stay hospitals, by ICD-9-CM code, sex, age, and geographic region: United States, 1999—Con.

[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table: see Methods section]

ICD-9-CM code	Sex			Age				Region			
	Total	Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	North-east	Midwest	South	West
Number of all-listed diagnoses in thousands											
717	36	17	20	*	21	10	*6	*7	*9	10	11
717.4	*5	*	*	—	*	*	*	*	*	*	*
717.7	*6	*	*	*	*	*	*	*	*	*	*
717.8	13	*7	*6	*	10	*	*	*	*	*	*
717.83	11	*5	*6	*	10	*	*	*	*	*	*
718	57	27	30	*	22	16	15	11	11	15	19
718.3	13	*7	*6	*	*8	*	*	*	*	*	*6
718.36	*5	*	*	*	*	*	*	*	*	*	*
718.4	19	*9	10	*	*	*	*9	*	*	*7	*
718.46	*7	*	*	*	*	*	*	*	*	*	*
718.8	14	*6	*8	*	*7	*	*	*	*	*	*5
719	180	71	109	*	31	38	108	35	63	52	30
719.0	16	*9	*7	*	*	*	*9	*	*	*5	*
719.06	13	*7	*5	*	*	*	*7	*	*	*	*
719.4	96	37	59	*	22	22	51	15	29	37	16
719.40	*5	*	*	*	*	*	*	*	*	*	*
719.41	29	12	17	*	*6	11	11	*	*8	9	*7
719.45	30	11	19	*	*	*	20	*5	*9	13	*
719.46	16	*6	10	*	*5	*	*8	*	*7	*6	*
719.49	*7	*	*	*	*	*	*	*	*	*	*
719.7	55	21	33	—	*	10	43	13	28	*	10
719.70	19	*7	12	—	*	*	17	*	14	*	*
719.75	14	*5	9	—	*	*	11	*	*6	*	*5
719.76	13	*6	*7	—	*	*	10	*	*6	*	*
720	15	9	*6	*	*	*7	*	*	*	*7	*
720.0	10	*8	*	—	*	*6	*	*	*	*	*
721	219	92	128	*	22	66	131	37	59	78	46
721.0	58	30	28	—	*6	27	25	11	15	20	12
721.1	24	12	12	—	*	*9	13	*	*5	9	*
721.2	*6	*	*	—	*	*	*	*	*	*	*
721.3	83	30	53	*	*9	18	56	13	21	31	19
721.9	41	14	27	*	*	*7	31	*7	13	15	*6
721.90	40	13	26	*	*	*7	30	*6	13	14	*6
722	489	243	246	*	176	192	121	83	129	189	88
722.0	66	36	30	—	27	34	*	10	13	32	11
722.1	226	115	111	*	96	84	46	45	58	80	42
722.10	221	113	109	*	95	82	44	45	57	79	40
722.2	*5	*	*	—	*	*	*	*	*	*	—
722.4	25	14	11	—	*5	12	*8	*	*6	*9	*6
722.5	93	42	52	—	22	31	41	12	33	33	15
722.52	90	40	50	—	22	31	38	11	32	33	14
722.6	11	*	*7	—	*	*	*	*	*	*7	*
722.7	19	10	9	—	*8	*8	*	*	*	*8	*
722.71	11	*5	*5	—	*	*5	*	*	*	*	*
722.73	*6	*	*	—	*	*	*	*	*	*	*
722.8	16	*7	*9	—	*6	*7	*	*	*	*7	*
722.83	15	*6	*9	—	*6	*6	*	*	*	*7	*
722.9	27	12	15	*	*5	13	*8	*	*7	10	*6
722.91	*5	*	*	—	*	*	*	*	*	*	*
722.93	19	9	10	*	*	9	*6	*	*6	*6	*
723	67	34	34	*	18	23	26	14	18	25	9
723.0	22	11	11	—	*	*6	13	*	*6	*9	*
723.1	24	12	12	*	*8	*8	*8	*	*7	9	*
723.4	15	*8	*6	*	*	*8	*	*	*	*6	*
724	426	161	265	*	90	132	202	86	108	147	85
724.0	132	52	80	—	*6	39	87	33	30	42	27
724.00	21	10	11	—	*	*	17	*8	*	*	*5
724.02	107	40	67	—	*6	34	67	25	25	37	21

See footnotes and notes at end of table.

Table 44. Number of all-listed diagnoses for discharges from short-stay hospitals, by ICD-9-CM code, sex, age, and geographic region: United States, 1999—Con.[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table: see Methods section]

ICD-9-CM code	Sex			Age				Region			
	Total	Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	North-east	Midwest	South	West
Number of all-listed diagnoses in thousands											
724.2	124	46	78	*	36	39	49	22	34	42	26
724.3	19	*5	14	*	*	*	12	*	*	*	*6
724.4	37	13	24	-	9	11	16	*6	*8	15	*7
724.5	93	37	57	*	27	32	34	15	26	36	16
724.8	9	*	*5	-	*	*	*	*	*	*	*
725	49	*7	41	-	*	*	45	11	16	12	10
726	89	38	51	*	16	30	44	22	20	27	20
726.0	*6	*	*	-	*	*	*	*	*	*	*
726.1	31	11	20	*	*	9	18	*8	*8	*9	*7
726.10	28	11	17	*	*	*9	16	*7	*8	*7	*6
726.2	15	*5	9	-	*	*6	*6	*	*	*	*
726.3	*8	*7	*	-	*	*	*	*	*	*	*
726.33	*7	*6	*	-	*	*	*	*	*	*	*
726.5	*9	*	*6	*	*	*	*6	*	*	*	*
726.6	*7	*	*	*	*	*	*	*	*	*	*
726.7	9	*	*7	-	*	*	*	*	*	*	*
727	66	29	37	*	14	21	27	13	11	28	14
727.0	25	12	12	*	*7	*8	*8	*6	*	*8	*7
727.00	12	*6	*6	*	*	*	*6	*	*	*	*
727.05	*8	*	*	*	*	*	*	*	*	*	*
727.1	*8	*	*6	*	*	*	*6	*	*	*	*
727.4	9	*	*6	*	*	*5	*	*	*	*	*
727.40	*6	*	*	-	*	*	*	*	*	*	*
727.5	*7	*	*	-	*	*	*	*	*	*	*
727.51	*7	*	*	-	*	*	*	*	*	*	*
727.6	*6	*	*	-	*	*	*	*	*	*	*
727.8	*8	*	*	*	*	*	*	*	*	*	*
728	119	62	57	*	32	25	59	28	32	36	24
728.2	*8	*	*	*	*	*	*6	*	*	*	*
728.3	10	*	*8	*	*	*	*8	*	*	*7	*
728.8	78	45	33	*	27	17	31	19	19	21	19
728.85	20	11	*9	*	*7	*	*7	*	*5	*8	*
728.86	*7	*	*	*	*	*	*	*	*	*	*
728.89	47	30	17	*	17	*8	22	15	11	11	*9
728.9	16	*8	*8	*	*	*	10	*	*6	*	*
729	242	81	161	*	53	85	101	45	53	93	52
729.1	92	19	74	*	24	42	26	19	18	35	20
729.2	*8	*	*	-	*	*	*	*	*	*	*
729.5	56	21	35	*	14	16	25	13	15	18	10
729.8	78	35	43	*	12	22	44	9	17	34	18
729.81	14	*6	*8	*	*	*	*7	*	*	*6	*
729.82	*6	*	*	*	*	*	*	*	*	*	*
729.89	59	28	30	*	*8	16	34	*6	14	25	14
730	127	81	46	*5	24	39	59	32	30	46	20
730.0	35	21	14	*	*6	12	16	9	10	10	*7
730.07	19	13	*7	*	*	*5	10	*6	*6	*	*
730.08	*6	*	*	*	*	*	*	*	*	*	*
730.1	26	14	12	*	*	*7	14	*6	*6	9	*
730.15	*6	*	*	-	*	*	*	*	*	*	*
730.17	9	*	*5	*	*	*	*5	*	*	*	*
730.2	65	46	20	*	13	19	29	17	13	27	*9
730.25	*7	*6	*	*	*	*	*	*	*	*	*
730.26	*7	*	*	*	*	*	*	*	*	*	*
730.27	32	22	10	*	*	11	17	11	*6	11	*
730.28	12	9	*	*	*	*	*6	*	*	*5	*

See footnotes and notes at end of table.

Table 44. Number of all-listed diagnoses for discharges from short-stay hospitals, by ICD-9-CM code, sex, age, and geographic region: United States, 1999—Con.

[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table: see Methods section]

ICD-9-CM code	Sex			Age				Region			
	Total	Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	North-east	Midwest	South	West
Number of all-listed diagnoses in thousands											
731	48	30	19	*	*	13	33	15	15	11	*7
731.0	23	11	12	—	*	*	21	*6	10	*6	*
731.8	26	19	*7	—	*	11	12	9	*	*5	*6
732	12	*9	*	*6	*	*	*	*	*	*	*
733	692	130	562	*6	49	105	533	147	166	245	134
733.0	380	30	350	*	*	38	338	85	97	131	67
733.00	352	27	324	*	*	34	314	78	91	125	57
733.01	16	*	16	—	—	*	16	*	*	*	*7
733.09	11	*	9	—	*	*	*7	*	*	*	*
733.1	157	39	118	*	*	20	131	29	36	57	34
733.10	*7	*	*6	*	*	—	*6	*	*	*	*
733.13	115	28	87	*	*	10	103	18	26	43	27
733.14	15	*	11	*	*	*5	9	*	*	*	*
733.15	*6	*	*	*	*	*	*	*	*	*	*
733.19	*7	*	*6	*	*	*	*5	*	*	*	*
733.2	*5	*	*	*	*	*	*	*	*	*	*
733.4	33	18	16	*	11	13	9	*7	*8	13	*6
733.42	27	14	12	*	9	10	*7	*	*6	10	*6
733.6	31	9	22	*	*6	13	12	*7	*7	13	*
733.8	39	19	19	*	16	11	11	9	*7	14	*8
733.81	*8	*	*	*	*	*	*	*	*	*	*
733.82	31	16	15	*	14	*8	*8	*6	*7	12	*6
733.9	46	12	34	*	*	10	30	*9	10	16	11
733.90	38	*9	29	*	*	*8	27	*8	*9	13	*9
733.99	*6	*	*	*	*	*	*	*	*	*	*
735	13	*	11	*	*	*	*5	*	*	*	*
736	40	22	18	*6	*7	10	17	*8	11	10	10
736.4	*6	*	*	*	*	*	*	*	*	*	*
736.7	18	11	*8	*	*	*	*9	*	*7	*6	*
736.79	14	*8	*6	*	*	*	*7	*	*6	*5	*
736.8	*7	*	*	*	*	*	*	*	*	*	*
737	68	14	54	10	14	11	34	15	16	25	12
737.1	14	*	11	*	*	*	12	*	*	*5	*
737.10	14	*	11	*	*	*	11	*	*	*5	*
737.3	49	10	39	9	12	*9	19	12	11	17	*9
737.30	42	*8	34	*7	11	*8	17	11	*9	15	*7
738	50	18	32	*	12	15	21	*9	11	15	15
738.4	41	14	26	—	*8	13	19	*7	10	12	13
739	21	*7	14	—	9	*	*7	18	*	*	*
739.2	*7	*	*	—	*	*	*	*6	*	*	—
739.3	10	*	*9	—	*8	*	*	10	*	*	—
740-759	623	300	324	368	134	64	58	168	121	186	148
741	21	*9	12	10	*8	*	*	*8	*6	*5	*
741.0	*8	*	*	*5	*	*	—	*	*	*	*
741.9	12	*5	*7	*	*5	*	*	*5	*	*	*
741.90	*9	*	*	*	*	*	*	*	*	*	*
742	27	11	16	19	*	*	*	*7	*8	*6	*6
742.1	*6	*	*	*	*	*	—	*	*	*	*
742.3	*9	*	*	*7	*	*	*	*	*	*	*
744	*8	*5	*	*7	*	—	—	*	*	*5	*
745	91	38	52	66	16	*	*	28	12	22	28
745.1	*7	*	*	*6	*	*	—	*	*	*	*
745.2	*7	*	*	*6	*	*	—	*	*	*	*
745.4	29	15	15	24	*	*	*	10	*	*	11
745.5	35	14	21	21	*8	*	*	*8	*5	12	10
745.6	*6	*	*	*5	*	*	—	*	*	*	*

See footnotes and notes at end of table.

Table 44. Number of all-listed diagnoses for discharges from short-stay hospitals, by ICD-9-CM code, sex, age, and geographic region: United States, 1999—Con.[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table: see Methods section]

ICD-9-CM code	Sex			Age				Region			
	Total	Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	North-east	Midwest	South	West
Number of all-listed diagnoses in thousands											
746	50	25	25	27	14	*	*	14	*8	16	12
746.0	*5	*	*	*	*	—	—	*	*	*	*
746.4	*7	*	*	*	*	*	*	*	*	*	*
746.8	19	10	9	10	*	*	*	*5	*	*6	*
746.89	*8	*	*	*5	*	*	*	*	*	*	*
746.9	*6	*	*	*	*	*	*	*	*	*	*
747	65	33	31	41	*8	*7	9	16	13	17	18
747.0	21	10	11	20	*	—	*	*	*	*	11
747.3	10	*	*6	9	*	—	—	*	*	*	*
747.6	14	*9	*	*	*	*	*7	*	*6	*5	*
747.61	11	*7	*	*	*	*	*6	*	*6	*	*
747.8	9	*	*	*	*	*	*	*	*	*	*
748	20	12	*8	17	*	*	*	*	*6	*	*
748.3	10	*6	*	10	*	*	*	*	*	*	*
749	23	14	9	21	*	*	*	*6	*	12	*
749.0	*6	*	*	*5	*	—	*	*	*	*	*
749.2	14	10	*	13	*	*	*	*	*	*8	*
749.20	*8	*6	*	*7	*	*	*	*	*	*8	*
750	15	10	*	13	*	*	*	*	*	*	*
750.5	*9	*7	*	*8	—	—	*	*	*	*	*
751	34	17	16	19	*8	*	*	10	10	*5	9
751.0	*5	*	*	*	*	*	*	*	*	*	*
751.3	*7	*7	*	*6	*	—	—	*	*	*	*
751.5	*7	*	*	*	*	*	*	*	*	*	*
752	29	9	19	10	14	*	*	*7	*5	*9	*8
752.1	*9	—	*9	*	*5	*	*	*	*	*	*
752.11	*8	—	*8	*	*	*	*	*	*	*	*
752.6	*5	*5	—	*	*	*	*	*	*	*	*
753	58	26	32	20	*9	14	15	14	13	17	13
753.0	*8	*	*	*	*	*	*	*	*	*	*
753.1	27	12	15	*	*	9	11	*7	*5	9	*6
753.10	*6	*	*	*	*	*	*	*	*	*	*
753.12	10	*	*5	*	*	*	*	*	*	*	*
753.2	*7	*	*	*6	*	*	*	*	*	*	*
753.3	*7	*	*	*	*	*	*	*	*	*	*
754	25	12	13	16	*5	*	*	*7	*	*8	*5
754.7	*6	*	*	*	*	—	*	*	*	*	*
755	18	*7	11	9	*	*	*	*6	*	*	*6
755.6	10	*	*7	*	*	*	*	*	*	*	*
755.63	*8	*	*6	*	*	*	*	*	*	*	*
756	50	25	25	22	14	*7	*7	11	*9	18	12
756.0	11	*8	*	10	*	*	*	*	*	*	*
756.1	20	9	11	*	*6	*6	*6	*	*	*8	*
756.12	11	*	*7	—	*	*	*	*	*	*	*
756.5	*6	*	*	*	*	*	*	*	*	*	*
757	20	*8	12	12	*6	—	*	*	*	15	*
757.3	16	*7	*8	11	*	—	*	*	*	14	—
757.32	12	*5	*7	*8	*	—	—	*	—	12	—
758	37	19	19	23	*8	*5	*	10	*7	9	11
758.0	27	15	13	16	*7	*	*	*8	*6	*7	*7
759	30	17	13	13	10	*5	*	12	*6	*8	*
759.8	19	13	*6	*9	*6	*	*	*8	*	*	*
759.89	16	11	*	*8	*	*	*	*7	*	*	*
760-779	487	304	183	485	*	*	*	85	98	147	157
763	*5	*	*	*5	*	—	—	*	*	*	*
764	*6	*	*	*6	—	—	—	*	*	*	*

See footnotes and notes at end of table.

Table 44. Number of all-listed diagnoses for discharges from short-stay hospitals, by ICD-9-CM code, sex, age, and geographic region: United States, 1999—Con.

[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table: see Methods section]

ICD-9-CM code	Sex			Age				Region			
	Total	Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	North-east	Midwest	South	West
Number of all-listed diagnoses in thousands											
765	55	34	20	55	—	—	*	*8	11	18	18
765.0	*7	*	*	*7	—	—	—	*	*	*	*
765.1	48	31	17	48	—	—	*	*8	9	15	16
765.16	*8	*7	*	*8	—	—	—	*	*	*	*
765.17	*6	*	*	*6	—	—	—	*	*	*	*
765.18	14	10	*	14	—	—	—	*	*	*	*5
765.19	10	*	*	10	—	—	—	*	*	*	*
767	*8	*	*	*8	*	—	*	*	*	*	*
769	28	18	10	28	—	*	*	*	*6	11	*8
770	99	64	35	98	*	*	*	18	23	25	32
770.0	*6	*	*	*6	—	—	—	*	*	*	*
770.1	*6	*	*	*6	—	—	—	*	*	*	*
770.2	*7	*5	*	*7	—	*	—	*	*	*	*
770.6	13	*9	*	13	—	—	—	*	*	*	*5
770.7	17	*8	*8	17	*	—	—	*	*6	*	*
770.8	45	30	15	45	*	*	*	10	9	10	16
771	44	26	18	44	*	*	—	10	9	17	*8
771.8	34	19	15	33	—	*	—	*6	*7	13	*7
772	9	*5	*	9	—	*	—	*	*	*	*
773	*9	*	*5	*9	—	—	—	*	*	*	*
774	91	59	32	91	—	*	—	13	17	26	35
774.2	29	20	*9	29	—	—	—	*	*5	*8	12
774.6	60	38	22	60	—	*	—	9	11	17	23
775	32	21	11	32	—	—	—	*5	*5	10	11
775.5	14	9	*5	14	—	—	—	*	*	*	*6
775.6	*7	*	*	*7	—	—	—	*	*	*	*
776	20	12	*9	20	—	*	—	*	*	*9	*6
776.1	*6	*	*	*6	—	*	—	*	*	*	*
776.6	*9	*5	*	*9	—	—	—	*	*	*	*
777	9	*5	*	9	—	—	—	*	*	*	*
778	17	10	*7	17	—	—	—	*	*5	*	*5
778.4	11	*6	*5	11	—	—	—	*	*	*	*
779	44	29	15	44	*	—	*	11	*6	10	17
779.0	*7	*	*	*7	—	—	*	*	*	*	*
779.3	30	20	10	29	*	—	—	*6	*	*6	14
779.8	*5	*	*	*5	—	—	—	*	*	*	*
780-799	7,144	3,156	3,988	593	1,482	1,832	3,238	1,522	1,848	2,530	1,244
780	2,109	990	1,119	208	428	492	982	502	537	743	328
780.0	94	40	54	*	17	19	54	20	19	30	24
780.01	23	*8	14	*	*	*	16	*5	*	*7	*6
780.09	64	29	35	*	14	13	34	14	14	21	15
780.1	17	*8	*8	*	*	*	9	*	*	*9	*
780.2	379	163	215	*	43	82	250	102	87	134	56
780.3	798	395	404	91	218	194	295	212	210	269	107
780.31	23	12	11	22	*	*	*	*	*6	*8	*5
780.39	775	382	393	70	217	194	295	209	204	261	102
780.4	112	37	75	*	16	22	73	23	34	34	20
780.5	168	98	70	*8	43	74	42	42	41	57	28
780.50	*6	*	*	*	*	*	*	*	*	*	*
780.52	15	*6	9	—	*	*	*7	*	*	*	*
780.57	139	85	54	*8	36	64	31	34	32	48	25
780.6	285	144	142	89	56	53	88	56	65	120	45
780.7	170	69	100	*7	17	27	119	26	55	58	31
780.79	165	68	97	*6	16	25	117	26	54	56	29
780.8	*6	*	*	*	*	*	*	*	*	*	*
780.9	81	33	49	*	11	17	50	16	21	32	13

See footnotes and notes at end of table.

Table 44. Number of all-listed diagnoses for discharges from short-stay hospitals, by ICD-9-CM code, sex, age, and geographic region: United States, 1999—Con.[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table: see Methods section]

ICD-9-CM code	Sex			Age				Region			
	Total	Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	North-east	Midwest	South	West
Number of all-listed diagnoses in thousands											
781	177	75	102	9	18	30	120	44	58	52	22
781.0	21	10	11	*	*6	*5	*9	*7	*5	*8	*
781.2	93	35	58	*	*5	12	75	27	33	25	*8
781.3	47	22	25	*	*	11	28	*7	17	16	*7
781.9	11	*	*6	*	*	*	*5	*	*	*	*
782	203	89	114	20	39	58	86	39	60	60	44
782.0	43	17	26	*	12	17	13	*7	12	14	10
782.1	34	17	17	9	9	*5	10	*9	10	10	*6
782.3	94	40	55	*	12	29	52	16	31	27	20
782.4	18	9	*8	*	*	*5	*7	*	*	*5	*
782.7	*7	*	*	*	*	*	*	*	*	*	*
783	170	76	94	65	21	17	68	41	37	59	33
783.0	38	15	22	*	*	*	25	*5	11	18	*
783.2	42	18	24	*	*7	*8	24	*7	10	17	*7
783.3	14	*7	*7	10	*	—	*	*	*	*	*
783.4	69	35	34	46	*6	*	14	21	12	18	17
784	329	136	193	15	87	79	148	52	95	114	68
784.0	143	46	97	*6	65	38	34	22	44	50	28
784.2	12	*7	*	*	*	*	*	*	*	*	*
784.3	66	25	41	*	*7	13	45	10	18	24	14
784.4	*7	*	*	*	*	*	*5	*	*	*	*
784.49	*5	*	*	*	*	*	*	*	*	*	*
784.5	54	28	26	*	*	14	35	*7	20	17	10
784.7	40	21	19	*	*6	*8	23	*9	*8	14	*8
785	324	144	181	20	57	86	162	73	90	103	59
785.0	62	23	39	*	20	13	26	10	22	17	13
785.1	36	11	25	—	*6	13	17	*8	11	13	*
785.2	27	12	15	*8	*8	*	*7	*	*6	9	*7
785.4	85	39	45	*	*8	29	46	21	17	31	15
785.5	81	43	38	*	*8	17	53	21	24	22	14
785.50	*8	*	*	*	*	*	*	*	*	*	*
785.51	32	20	12	*	*	*6	23	*9	*9	*9	*6
785.59	41	20	21	*	*	*9	27	11	13	11	*5
785.6	25	12	13	*	*7	*7	*8	*8	*6	*8	*
785.9	*9	*	*5	*	*	*	*6	*	*	*	*
786	1,277	590	688	65	222	478	511	266	307	479	225
786.0	171	78	93	49	25	34	63	29	45	66	32
786.01	*5	*	*	*	*	*	*	*	*	*	*
786.03	20	11	9	15	*	*	*	*	*5	*6	*6
786.05	36	14	22	*	*7	*8	21	*7	9	14	*
786.06	*6	*	*	*	*	*	*	*	*	*	*
786.07	9	*	*5	*	*	*	*	*	*	*	*
786.09	88	40	48	26	11	20	31	15	24	35	13
786.1	*8	*	*	*	*	*	*	*	*	*	*
786.2	25	9	15	*7	*	*6	*7	*	*7	*9	*7
786.3	62	37	25	*	10	20	30	17	15	21	*8
786.5	965	434	531	*	177	406	380	209	231	360	165
786.50	543	246	298	*	92	230	220	108	120	220	95
786.51	27	14	14	*	*5	10	12	*5	*	16	*
786.52	51	21	30	*	11	19	21	*7	20	17	*8
786.59	343	154	189	*	69	146	128	88	87	108	61
786.6	38	20	18	*	*	*8	26	*6	*7	19	*6
787	691	255	436	84	156	164	287	119	180	260	132
787.0	356	113	243	50	101	90	116	56	93	134	74
787.01	206	57	148	15	66	57	68	31	49	86	39
787.02	85	27	59	*	24	24	35	13	28	23	22
787.03	65	29	36	33	11	*8	12	12	16	25	12
787.1	*6	*	*	—	*	*	*	*	*	*	*

See footnotes and notes at end of table.

Table 44. Number of all-listed diagnoses for discharges from short-stay hospitals, by ICD-9-CM code, sex, age, and geographic region: United States, 1999—Con.[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table: see Methods section]

ICD-9-CM code	Sex			Age				Region			
	Total	Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	North-east	Midwest	South	West
Number of all-listed diagnoses in thousands											
787.2	121	53	67	*	12	25	81	23	32	48	18
787.3	11	*7	*	*	*	*	*5	*	*	*	*
787.6	11	*	*7	*	*	*	*	*	*	*	*
787.9	185	76	109	28	36	44	78	35	48	68	34
787.91	183	76	108	28	36	44	76	34	48	68	34
788	434	242	192	*7	66	81	281	105	114	143	73
788.0	15	9	*6	*	*9	*	*	*6	*	*	*
788.1	*8	*	*6	*	*	*	*	*	*	*	*
788.2	260	174	86	*	39	46	173	70	62	83	45
788.20	217	143	73	*	34	36	145	59	50	69	37
788.29	39	27	11	*	*	*9	25	10	10	12	*6
788.3	97	34	63	*	11	17	66	18	31	34	15
788.30	73	27	47	*	*9	11	52	13	24	26	11
788.31	*8	*	*5	—	*	*	*6	*	*	*	*
788.33	*7	*	*7	—	*	*	*	*	*	*	*
788.39	*5	*	*	*	*	*	*	*	*	*	*
788.4	14	*7	*7	*	*	*6	*6	*	*6	*	*
788.41	*8	*	*	*	*	*	*	*	*	*	*
788.5	*6	*	*	*	*	*	*	*	*	*	*
788.9	31	12	19	*	*	*	24	*6	*7	12	*
789	519	190	328	40	160	149	169	105	125	185	104
789.0	373	122	251	36	139	94	104	68	94	134	77
789.00	140	45	95	17	49	33	42	23	28	66	23
789.01	33	10	23	*	13	12	*8	*7	*7	10	*8
789.02	10	*	*7	*	*	*	*	*	*	*	*
789.03	58	18	39	*8	29	11	10	12	17	17	11
789.04	21	*7	14	*	*8	*6	*7	*	*7	*6	*
789.05	*8	*	*	*	*	*	*	*	*	*	*
789.06	46	16	30	*	12	16	14	*7	12	18	9
789.07	19	*5	13	*	*7	*5	*	*	*	*	*8
789.09	39	14	25	*	16	*7	13	*9	13	*9	*9
789.1	*9	*	*7	*	*	*	*	*	*	*	*
789.2	12	*5	*7	*	*	*	*	*	*	*	*
789.3	14	*	11	*	*	*	*7	*	*	*	*
789.30	*6	*	*5	—	*	*	*	*	*	*	*
789.5	110	58	53	*	13	47	49	26	25	40	20
790	316	131	185	23	76	74	143	69	91	104	52
790.2	47	*6	41	*	33	*	11	9	12	14	12
790.4	*8	*	*	*	*	*	*	*	*	*	*
790.5	24	9	14	*	*8	*8	*7	*6	*6	*8	*
790.6	113	48	66	*	17	30	63	26	31	41	15
790.7	71	36	35	13	11	17	30	16	21	23	12
790.9	44	24	20	*	*	10	27	*8	17	12	*8
790.92	25	*8	18	*	*	*7	15	*5	*8	*7	*6
790.93	11	11	—	—	—	*	10	*	*	*	*
790.99	*8	*5	*	*	*	*	*	*	*	*	*
791	26	13	13	*	*9	*6	*8	*7	*6	10	*
791.0	11	*6	*	*	*	*	*	*	*	*	*
791.9	11	*	*7	*	*	*	*	*	*	*	*
792	63	12	51	*	39	*6	17	12	21	16	14
792.1	27	11	16	*	*	*6	16	*7	*8	*9	*
792.3	35	—	35	—	35	*	—	*	13	*7	10
793	21	9	12	*	*	*	13	*	*8	*7	*
793.1	10	*	*5	*	*	*	*7	*	*	*	*
794	91	43	48	*	23	34	31	20	16	42	13
794.3	41	21	20	*	*7	15	18	*7	10	21	*
794.31	32	15	17	*	*5	12	13	*6	*9	15	*
794.39	*7	*	*	*	*	*	*	*	*	*	*
794.8	44	21	23	*	13	16	11	12	*	19	9

See footnotes and notes at end of table.

Table 44. Number of all-listed diagnoses for discharges from short-stay hospitals, by ICD-9-CM code, sex, age, and geographic region: United States, 1999—Con.[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table: see Methods section]

ICD-9-CM code	Sex			Age				Region			
	Total	Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	North-east	Midwest	South	West
Number of all-listed diagnoses in thousands											
795	23	*7	17	*	16	*	*	*9	*	*6	*
795.5	15	*	10	*	10	*	*	*7	*	*	*
795.7	*6	*	*	*	*5	*	*	*	*	*	*
795.79	*6	*	*	*	*	*	*	*	*	*	*
796	42	14	28	*	14	10	17	*7	12	17	*7
796.0	11	*	*8	—	*	*	*9	*	*	*	*
796.2	23	*7	16	*	11	*6	*5	*	*7	*8	*5
797	*7	*	*6	—	—	*	*7	*	*	*	*
799	319	137	182	24	47	61	187	48	84	128	59
799.0	112	44	68	14	11	26	60	17	39	36	19
799.1	16	*	11	*	*	*	10	*	*5	*	*
799.3	60	26	34	*	*	*8	50	*7	22	25	*6
799.4	41	24	17	*	*5	*7	28	11	*7	14	*9
799.9	86	38	48	*9	24	17	36	*9	10	47	21
800-999	6,038	3,056	2,982	445	1,936	1,294	2,363	1,243	1,458	2,103	1,234
800	13	*9	*	*6	*	*	*	*	*	*	*
801	36	25	12	*	20	*6	*6	*7	*8	13	*8
801.0	15	10	*	*	*7	*	*	*	*	*5	*
801.2	10	*5	*	*	*6	*	*	*	*	*	*
802	94	70	24	*	57	18	15	17	24	34	19
802.0	17	10	*8	*	*9	*	*5	*	*	*6	*
802.2	22	18	*	*	16	*	*	*	*6	*7	*
802.4	19	15	*	*	10	*5	*	*	*	*8	*
802.6	12	10	*	*	*9	*	*	*	*	*5	*
802.8	13	10	*	*	*7	*	*	*	*	*	*
803	*8	*	*	*	*	*	*	*	*	*	*
805	143	61	82	*	44	26	71	26	46	41	29
805.0	26	16	9	*	11	*6	*8	*	*8	*6	*7
805.02	10	*6	*	*	*	*	*	*	*	*	*
805.2	38	15	22	*	9	*7	21	*9	12	11	*6
805.4	60	23	36	*	15	10	35	11	20	17	12
805.6	16	*	11	*	*7	*	*6	*	*6	*	*
806	10	*6	*	*	*	*	*	*	*	*	*
807	118	70	48	*	32	34	50	25	31	41	22
807.0	101	60	40	*	27	28	43	21	28	34	17
807.00	12	*7	*	—	*	*6	*	*	*	*	*
807.01	23	15	*8	*	*	*5	12	*	*8	*9	*
807.02	14	*7	*6	*	*	*	*7	*	*5	*	*
807.03	14	*8	*5	*	*	*	*6	*	*	*	*
807.04	*8	*	*	*	*	*	*	*	*	*	*
807.05	*5	*	*	*	*	*	*	*	*	*	*
807.09	19	10	*9	*	*6	*6	*7	*	*	*6	*5
807.2	*8	*	*	—	*	*	*	*	*	*	*
807.4	*6	*	*	—	*	*	*	*	*	*	*
808	111	38	73	*	30	10	69	14	34	39	24
808.0	22	13	*9	*	10	*	*8	*	*7	*9	*
808.2	65	17	48	*	14	*	45	*9	20	21	15
808.4	11	*	*7	*	*	*	*7	*	*	*	*
808.8	10	*	*9	*	*	*	*8	*	*	*	*
810	26	16	11	*	10	*	11	*5	*8	*	*8
810.0	26	15	11	*	10	*	11	*5	*8	*	*8
810.00	16	*8	*8	*	*	*	*8	*	*	*	*6
810.02	*7	*5	*	*	*	*	*	*	*	*	*
811	13	9	*	*	*7	*	*	*	*6	*	*
811.0	12	*9	*	*	*6	*	*	*	*6	*	*
811.00	*5	*	*	*	*	*	*	*	*	*	*

See footnotes and notes at end of table.

Table 44. Number of all-listed diagnoses for discharges from short-stay hospitals, by ICD-9-CM code, sex, age, and geographic region: United States, 1999—Con.

[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table: see Methods section]

ICD-9-CM code	Sex			Age				Region			
	Total	Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	North-east	Midwest	South	West
Number of all-listed diagnoses in thousands											
812	102	38	63	17	16	16	53	22	23	34	23
812.0	53	15	38	*	*7	*9	37	12	13	17	11
812.00	16	*	12	*	*	*	12	*	*	*5	*
812.01	18	*	14	—	*	*	13	*5	*	*6	*
812.03	*6	*	*	—	*	*	*	*	*	*	*
812.09	12	*	*8	*	*	*	*9	*	*	*	*
812.2	16	*5	11	*	*	*	11	*	*	*8	*
812.20	*8	*	*6	*	*	*	*6	*	*	*	*
812.21	*8	*	*	*	*	*	*	*	*	*	*
812.4	26	13	13	14	*	*	*	*6	*	*7	*8
812.41	15	*8	*7	11	*	*	*	*	*	*	*
812.5	*5	*	*	*	*	*	*	*	*	*	*
813	104	45	59	9	34	19	41	22	26	40	16
813.0	18	*6	12	*	*	*	*8	*	*	*7	*
813.01	*9	*	*6	*	*	*	*6	*	*	*	*
813.2	*8	*	*	*	*	*	*	*	*	*	*
813.4	58	23	35	*	17	11	27	13	14	22	9
813.41	15	*	11	*	*	*	10	*	*	*6	*
813.42	29	11	18	*	10	*6	12	*5	*7	12	*
813.44	11	*6	*5	*	*	*	*	*	*	*	*
813.8	*8	*	*5	*	*	*	*	*	*	*	*
814	11	*8	*	*	*	*6	*	*	*	*	*
814.0	*8	*	*	*	*	*	*	*	*	*	*
815	14	*8	*6	*	*8	*	*	*	*	*	*
815.0	10	*6	*	*	*6	*	*	*	*	*	*
815.00	*6	*	*	*	*	*	*	*	*	*	*
816	23	13	10	*	11	*5	*6	*	*	9	*5
816.0	13	*6	*8	*	*5	*	*	*	*	*6	*
816.01	*8	*	*5	*	*	*	*	*	*	*	*
816.1	9	*7	*	*	*5	*	*	*	*	*	*
816.11	*6	*	*	*	*	*	*	*	*	*	*
820	380	103	276	*	12	28	338	81	87	149	62
820.0	95	22	74	*	*	*9	84	21	22	34	18
820.00	*8	*	*5	—	*	*	*6	*	*	*	*
820.02	11	*	*7	*	*	*	10	*	*	*	*
820.03	*9	*	*7	—	*	*	*8	*	*	*	*
820.09	66	13	53	*	*	*5	60	15	16	24	11
820.2	186	51	134	*	*7	12	167	43	42	70	30
820.20	11	*	*8	—	*	*	10	*	*	*	*
820.21	156	42	114	*	*	*8	143	35	36	58	27
820.22	19	*7	12	*	*	*	13	*	*	*8	*
820.8	96	29	67	*	*	*7	85	16	23	45	12
821	75	36	39	14	20	11	30	15	19	26	14
821.0	40	24	16	11	13	*	12	*8	12	14	*7
821.00	12	*6	*6	*	*	*	*6	*	*	*6	*
821.01	28	18	10	*8	11	*	*6	*6	9	*8	*
821.2	28	*8	20	*	*	*6	17	*6	*7	9	*6
821.20	9	*	*5	*	*	*	*5	*	*	*	*
821.23	12	*	10	*	*	*	9	*	*	*	*
822	23	10	14	*	*7	*6	10	*	*5	10	*
822.0	20	*7	13	*	*	*5	10	*	*	*8	*
823	102	59	43	*8	56	20	18	21	24	35	22
823.0	35	17	19	*	18	*7	*9	*	10	14	*6
823.00	27	12	14	*	13	*	*7	*	*6	11	*
823.02	*5	*	*	*	*	*	*	*	*	*	*
823.2	25	17	*8	*	16	*	*	*6	*5	*5	*8
823.22	18	14	*	*	12	*	*	*	*	*	*6

See footnotes and notes at end of table.

Table 44. Number of all-listed diagnoses for discharges from short-stay hospitals, by ICD-9-CM code, sex, age, and geographic region: United States, 1999—Con.[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table: see Methods section]

ICD-9-CM code	Sex			Age				Region			
	Total	Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	North-east	Midwest	South	West
Number of all-listed diagnoses in thousands											
823.3	*9	*8	*	*	*6	*	*	*	*	*	*
823.32	*6	*6	*	*	*	*	*	*	*	*	*
823.8	22	12	10	*	10	*	*5	*6	*	*8	*
823.82	15	*9	*6	*	*7	*	*	*	*	*6	*
823.9	*7	*	*	*	*	*	*	*	*	*	*
823.92	*6	*	*	*	*	*	*	*	*	*	*
824	136	58	78	*5	53	38	40	28	31	47	29
824.0	12	*6	*6	*	*7	*	*	*	*	*	*
824.2	11	*	*7	*	*	*	*	*	*	*	*
824.4	34	13	21	*	11	*9	13	*7	*9	10	*7
824.6	27	*	22	*	9	*9	*9	*	*8	9	*6
824.8	38	22	16	*	16	10	*9	10	9	13	*7
824.9	*6	*	*	*	*	*	*	*	*	*	*
825	43	24	19	*	20	11	10	*7	*9	18	9
825.0	12	*9	*	*	*7	*	*	*	*	*	*
825.2	26	11	14	*	11	*6	*8	*	*	13	*5
825.21	*7	*	*	*	*	*	*	*	*	*	*
825.25	13	*	*8	*	*	*	*6	*	*	*7	*
826	*8	*	*	*	*	*	*	*	*	*	*
826.0	*6	*	*	—	*	*	*	*	*	*	*
831	10	*7	*	*	*	*	*	*	*	*	*
831.0	10	*7	*	*	*	*	*	*	*	*	*
835	*7	*	*	*	*	*	*	*	*	*	*
835.0	*7	*	*	*	*	*	*	*	*	*	*
836	23	11	12	*	14	*	*	*	*	12	*
836.0	10	*6	*	*	*5	*	*	*	*	*	*
836.1	*6	*	*	—	*	*	*	*	*	*	*
839	10	*8	*	*	*6	*	*	*	*	*	*
839.0	*6	*	*	*	*5	*	*	*	*	*	*
840	37	17	20	*	*7	14	14	*7	*7	14	*7
840.4	27	11	17	*	*	12	9	*5	*	12	*6
840.9	*5	*	*	—	*	*	*	*	*	*	*
844	30	14	16	*	16	*6	*5	*8	*7	9	*5
844.2	16	*8	*8	*	11	*	*	*	*	*6	*
844.8	*6	*	*	*	*	*	*	*	*	*	*
845	28	12	15	*	12	*9	*6	*	*6	10	*7
845.0	26	12	14	*	11	*9	*5	*	*6	10	*6
845.00	14	*6	*8	*	*6	*	*	*	*	*	*
845.01	*5	*	*	*	*	*	*	*	*	*	*
845.09	*6	*	*	*	*	*	*	*	*	*	*
846	*7	*	*	—	*	*	*	*	*	*	*
846.0	*6	*	*	—	*	*	*	*	*	*	*
847	39	16	23	*	20	*7	11	*7	13	12	*8
847.0	20	9	11	*	12	*	*	*	*6	*7	*
847.2	14	*5	*9	*	*6	*	*5	*	*	*	*
848	*7	*	*	*	*	*	*	*	*	*	*
850	58	38	20	*7	33	*8	10	19	13	16	10
850.0	14	*8	*6	*	*7	*	*	*	*	*	*
850.1	22	15	*7	*	15	*	*	10	*	*	*
850.5	*9	*6	*	*	*	*	*	*	*	*	*
850.9	13	*8	*	*	*6	*	*	*	*	*	*
851	29	20	9	*	12	*6	10	*6	*6	10	*7
851.8	21	14	*7	*	*9	*	*7	*	*	*7	*5
851.86	*7	*	*	*	*	*	*	*	*	*	*

See footnotes and notes at end of table.

Table 44. Number of all-listed diagnoses for discharges from short-stay hospitals, by ICD-9-CM code, sex, age, and geographic region: United States, 1999—Con.

[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table: see Methods section]

ICD-9-CM code	Sex			Age				Region			
	Total	Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	North-east	Midwest	South	West
Number of all-listed diagnoses in thousands											
852	52	24	28	*	15	9	24	*7	10	18	17
852.0	12	*	*8	*	*	*	*	*	*	*	*
852.2	36	19	18	*	*9	*6	20	*	*6	12	13
852.20	13	*8	*5	*	*	*	*6	*	*	*6	*
852.21	12	*6	*6	*	*	*	*8	*	*	*	*
853	11	*6	*	*	*	*	*5	*	*	*	*
853.0	11	*6	*	*	*	*	*5	*	*	*	*
854	21	12	*8	*	13	*	*	*	*7	*6	*5
854.0	20	11	*8	*	12	*	*	*	*7	*6	*
854.02	*5	*	*	*	*	*	*	*	*	*	*
860	68	45	23	*	37	15	14	13	18	25	13
860.0	32	18	14	*	18	*6	*7	*7	9	10	*6
860.1	*6	*	*	*	*6	*	—	*	*	*	*
860.2	*8	*6	*	*	*	*	*	*	*	*	*
860.4	14	9	*	*	*5	*	*	*	*	*	*
860.5	*5	*	*	*	*	*	—	*	*	*	*
861	35	27	*8	*	18	*7	*7	*7	*8	13	*7
861.2	28	22	*6	*	13	*6	*6	*	*7	12	*
861.21	27	21	*6	*	13	*6	*6	*	*7	11	*
862	13	10	*	*	*5	*	*	*	*	*7	*
863	26	22	*	*	17	*	*	*	*6	9	*7
863.4	*5	*	*	*	*	—	*	*	*	*	*
864	21	13	*8	*	12	*	*	*	*	*8	*6
864.0	17	10	*7	*	*9	*	*	*	*	*6	*
864.05	*7	*	*	*	*	*	*	*	*	*	*
865	24	17	*7	*	13	*5	*	*	*7	*9	*
865.0	22	16	*6	*	11	*5	*	*	*7	*8	*
865.00	*6	*5	*	*	*	*	*	*	*	*	*
866	13	10	*	*	*8	*	*	*	*	*5	*
866.0	10	*8	*	*	*6	*	*	*	*	*	*
866.01	*7	*	*	*	*	*	*	*	*	*	*
867	16	13	*	*	*	*	*9	*	*	*7	*
867.0	14	12	*	*	*	*	*8	*	*	*6	*
868	17	11	*6	*	10	*	*	*	*	*5	*7
868.0	11	*7	*	*	*	*	*	*	*	*	*
868.1	*6	*	*	*	*	*	*	*	*	*	*
870	19	15	*	*	11	*	*	*	*	*9	*
870.0	*6	*	*	*	*	*	*	*	*	*	*
870.8	*8	*6	*	*	*	*	*	*	*	*	*
871	*6	*	*	*	*	*	*	*	*	*	*
872	12	*8	*	*	*7	*	*	*	*	*	*
872.0	*6	*	*	*	*	*	*	*	*	*	*
873	190	111	78	14	84	28	62	38	48	56	47
873.0	62	36	27	*	23	9	28	13	16	16	16
873.2	*6	*	*	*	*	*	*	*	*	*	*
873.4	91	54	37	*5	42	15	29	18	24	28	20
873.40	*7	*	*	*	*	*	*	*	*	*	*
873.41	*7	*	*	*	*	*	*	*	*	*	*
873.42	44	24	20	*	17	*7	18	*8	12	13	11
873.43	15	10	*5	*	*8	*	*	*	*	*	*
873.44	10	*6	*	*	*	*	*	*	*	*	*
873.49	*9	*6	*	*	*	*	*	*	*	*	*
873.5	*8	*	*	*	*	*	*	*	*	*	*
873.6	14	*9	*5	*	*7	*	*	*	*	*	*
873.63	*6	*	*	*	*	*	*	*	*	*	*

See footnotes and notes at end of table.

Table 44. Number of all-listed diagnoses for discharges from short-stay hospitals, by ICD-9-CM code, sex, age, and geographic region: United States, 1999—Con.[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table: see Methods section]

ICD-9-CM code	Sex			Age				Region			
	Total	Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	North-east	Midwest	South	West
Number of all-listed diagnoses in thousands											
874	*8	*6	*	*	*6	*	*	*	*	*	*
874.8	*6	*	*	*	*	*	*	*	*	*	*
875	*8	*8	*	*	*6	*	*	*	*	*	*
875.0	*7	*7	*	*	*5	*	*	*	*	*	*
876	*8	*	*	*	*7	*	*	*	*	*	*
876.0	*6	*	*	*	*6	*	—	*	*	*	*
879	21	15	*7	*5	12	*	*	*	*	11	*
879.2	*6	*	*	*	*	*	*	*	*	*	*
879.8	*8	*5	*	*	*	*	*	*	*	*5	*
880	10	*7	*	*	*7	*	*	*	*	*	*
880.0	*9	*5	*	*	*5	*	*	*	*	*	*
881	41	25	15	*	24	*	10	*	13	14	9
881.0	27	15	13	*	16	*	*8	*	10	*9	*5
881.00	13	*8	*5	*	*7	*	*	*	*6	*	*
881.01	*6	*	*	*	*	*	*	*	*	*	*
881.02	*9	*	*6	*	*7	*	*	*	*	*	*
881.1	*8	*6	*	*	*	*	*	*	*	*	*
882	19	13	*6	*	10	*	*	*	*5	*6	*5
882.0	9	*7	*	*	*	*	*	*	*	*	*
882.1	*6	*5	*	*	*	*	*	*	*	*	*
883	23	17	*6	*	14	*	*	*	*5	*7	*6
883.0	*8	*6	*	*	*	*	*	*	*	*	*
883.1	*6	*	*	*	*	*	*	*	*	*	*
883.2	*8	*7	*	*	*6	*	*	*	*	*	*
886	*7	*6	*	*	*	*	*	*	*	*	*
886.0	*7	*6	*	*	*	*	*	*	*	*	*
890	13	*9	*	*	10	*	*	*	*	*6	*
890.0	*8	*	*	*	*	*	*	*	*	*	*
890.1	*5	*	*	*	*	*	*	*	*	*	*
891	41	24	17	*	19	*7	11	*8	*8	16	*8
891.0	24	15	9	*	10	*	*6	*	*	9	*6
891.1	13	*7	*6	*	*6	*	*	*	*	*	*
892	14	9	*	*	*5	*	*	*	*	*7	*
892.0	*5	*	*	*	*	*	*	*	*	*	*
892.1	*6	*	*	*	*	*	*	*	*	*	*
902	*6	*	*	*	*	*	*	*	*	*	*
903	*8	*7	*	*	*	*	*	*	*	*	*
904	*7	*5	*	*	*	*	*	*	*	*	*
905	51	25	26	*	16	14	20	11	18	13	9
905.1	*6	*	*	*	*	*	*	*	*	*	*
905.2	9	*6	*	*	*	*	*	*	*	*	*
905.3	11	*	*8	*	*	*	*7	*	*	*	*
905.4	18	9	*9	*	*5	*6	*6	*	*8	*	*
906	26	14	12	11	11	*	*	*	*	19	*
906.9	13	*8	*5	*8	*	—	—	—	—	13	—
907	49	37	12	*	29	14	*	*9	15	15	10
907.0	22	15	*7	*	12	*8	*	*	*8	*	*6
907.2	25	21	*	*	17	*6	*	*	*6	10	*
908	13	*8	*5	*	*6	*	*	*	*	*	*
908.9	*8	*	*	*	*	*	*	*	*	*	*
909	*8	*5	*	*	*	*	*	*	*	*	*
910	28	19	*9	*	14	*	*6	*7	10	*6	*
910.0	26	18	*8	*	13	*	*5	*6	9	*6	*

See footnotes and notes at end of table.

Table 44. Number of all-listed diagnoses for discharges from short-stay hospitals, by ICD-9-CM code, sex, age, and geographic region: United States, 1999—Con.

[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table: see Methods section]

ICD-9-CM code	Sex			Age				Region			
	Total	Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	North-east	Midwest	South	West
Number of all-listed diagnoses in thousands											
911	14	10	*	*	*8	*	*	*	*	*	*
911.0	12	*9	*	*	*7	*	*	*	*	*	*
912	*8	*6	*	*	*	*	*	*	*	*	*
912.0	*8	*6	*	*	*	*	*	*	*	*	*
913	16	*9	*7	*	*7	*	*5	*	*5	*	*
913.0	13	*7	*6	*	*5	*	*	*	*	*	*
916	25	14	11	*6	9	*	*7	*	*8	*6	*7
916.0	19	12	*7	*	*8	*	*	*	*7	*	*5
919	19	13	*6	*	*9	*	*	*	*6	*6	*
919.0	18	13	*6	*	*8	*	*	*	*6	*	*
920	55	29	26	*	19	*8	24	13	14	18	11
921	14	*7	*7	*	*6	*	*	*	*	*	*
921.2	*5	*	*	*	*	*	*	*	*	*	*
922	71	34	37	*	24	12	32	11	24	16	20
922.1	38	19	19	*	13	*7	17	*6	12	*8	12
922.2	14	*8	*6	*	*6	*	*	*	*	*	*
922.3	14	*	9	*	*	*	*9	*	*6	*	*
922.31	9	*	*6	*	*	*	*5	*	*	*	*
923	34	13	20	*	11	*	17	*6	11	*7	*9
923.0	15	*7	*8	*	*	*	*7	*	*6	*	*
923.00	10	*6	*	*	*	*	*5	*	*	*	*
923.1	10	*	*7	*	*	*	*7	*	*	*	*
923.11	*7	*	*	*	*	*	*	*	*	*	*
923.2	*5	*	*	*	*	*	*	*	*	*	*
924	93	37	56	*	22	16	52	17	27	28	20
924.0	32	10	21	*	*	*	23	*6	9	*8	*8
924.00	*9	*	*	*	*	*	*	*	*	*	*
924.01	23	*6	17	*	*	*	20	*	*7	*6	*
924.1	22	*7	15	*	*6	*	12	*	*9	*	*
924.10	*7	*	*5	*	*	*	*	*	*	*	*
924.11	15	*5	10	*	*	*	*8	*	*7	*	*
924.8	26	13	14	*	*8	*5	12	*	*6	9	*6
933	12	*6	*6	*	*	*	*8	*	*	*6	*
933.1	11	*6	*5	*	*	*	*7	*	*	*6	*
934	10	*	*	*	*	*	*	*	*	*	*
935	14	*8	*6	*	*	*	*6	*	*	*	*
935.1	10	*6	*	*	*	*	*	*	*	*	*
941	16	12	*	*	*7	*	*	*	*5	*	*
941.2	10	*7	*	*	*	*	*	*	*	*	*
942	19	11	*7	*7	*6	*	*	*8	*	*	*
942.2	12	*8	*	*	*	*	*	*	*	*	*
943	19	11	*8	*	*8	*5	*	*5	*	*8	*
943.2	14	*7	*6	*	*6	*	*	*	*	*5	*
943.21	*6	*	*	*	*	*	*	*	*	*	*
944	17	13	*	*	*8	*5	*	*	*5	*8	*
944.2	*9	*7	*	*	*	*	*	*	*	*	*
945	24	16	*8	*8	*7	*5	*	*5	*5	10	*
945.2	13	*8	*5	*	*	*	*	*	*	*	*
945.3	9	*7	*	*	*	*	*	*	*	*	*
946	*6	*	*	*	*	*	*	*	*	*	*

See footnotes and notes at end of table.

Table 44. Number of all-listed diagnoses for discharges from short-stay hospitals, by ICD-9-CM code, sex, age, and geographic region: United States, 1999—Con.[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table: see Methods section]

ICD-9-CM code	Sex			Age				Region			
	Total	Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	North-east	Midwest	South	West
Number of all-listed diagnoses in thousands											
948	55	37	18	29	15	*7	*	12	*7	31	*5
948.0	26	16	9	12	*8	*	*	*7	*	12	*
948.00	26	16	9	12	*8	*	*	*7	*	12	*
948.1	13	10	*	*6	*	*	*	*	*	*6	*
948.10	11	*9	*	*	*	*	*	*	*	*5	*
948.2	*5	*	*	*	*	*	*	*	*	*	—
952	10	*8	*	*	*	*	*	*	*	*	*
952.0	*5	*	*	*	*	*	*	*	*	*	*
955	12	9	*	*	*8	*	*	*	*6	*	*
958	31	22	9	*	15	*	*6	*8	*8	10	*6
958.3	*8	*6	*	*	*	*	*	*	*	*	*
958.7	*6	*	*	*	*	*	*	*	*	*	*
958.8	10	*7	*	*	*	*	*	*	*	*	*
959	78	35	43	*8	30	9	30	13	22	23	19
959.0	46	23	22	*6	18	*	17	*7	16	12	11
959.01	43	22	21	*5	16	*	17	*6	15	11	11
959.1	14	*	9	*	*7	*	*	*	*	*	*
962	*8	*6	*	*	*	*	*	*	*	*	*
962.3	*7	*6	*	*	*	*	*	*	*	*	*
963	13	*6	*7	*	9	*	*	*	*	*	*
963.0	11	*5	*6	*	*9	*	*	*	*	*	*
964	*6	*	*	*	*	*	*	*	*	*	*
965	70	21	49	*6	45	13	*6	11	19	24	16
965.0	20	*8	12	*	9	*7	*	*	*	*7	*6
965.00	*7	*	*	*	*	*	*	*	*	*	*
965.09	10	*	*6	*	*	*	*	*	*	*	*
965.1	*8	*	*6	*	*6	*	*	*	*	*	*
965.4	26	*7	19	*	18	*	*	*	10	*8	*
965.6	*8	*	*7	*	*7	*	*	*	*	*	*
965.61	*8	*	*6	*	*6	*	*	*	*	*	*
966	16	*	11	*	*6	*	*	*	*	*6	*
966.1	*5	*	*	*	*	*	*	*	*	*	*
966.3	10	*	*7	*	*	*	*	*	*	*	*
967	11	*	*7	*	*7	*	*	*	*	*	*
967.0	*5	*	*	*	*	*	*	*	*	*	*
968	14	9	*	*	11	*	*	*	*	*5	*
968.0	*6	*	*	*	*5	*	*	*	*	*	*
968.5	*8	*6	*	—	*6	*	*	*	*	*	*
969	95	39	56	*	60	25	*6	17	22	34	22
969.0	35	16	20	*	23	10	*	*6	*9	12	*8
969.4	41	14	27	*	23	13	*	*7	10	15	*9
969.7	*5	*	*	*	*	*	*	*	*	*	*
971	*6	*	*	*	*	*	*	*	*	*	*
972	16	*7	*9	*	*	*	*8	*	*	*	*5
972.1	*6	*	*	*	*	*	*5	*	*	*	*
974	*6	*	*	*	*	*	*	*	*	*	*
977	17	*7	9	*	10	*	*	*	*	10	*
977.8	*7	*	*	*	*	*	*	*	*	*	*
977.9	9	*	*5	*	*5	*	*	*	*	*6	*
980	15	*8	*8	*	13	*	*	*	*	*	*
980.0	12	*7	*5	*	10	*	*	*	*	*	*
987	*8	*	*	*	*	*	*	*	*	*	*
989	13	*8	*5	*	*	*	*	*	*	*6	*
989.5	*8	*5	*	*	*	*	*	*	*	*	*

See footnotes and notes at end of table.

Table 44. Number of all-listed diagnoses for discharges from short-stay hospitals, by ICD-9-CM code, sex, age, and geographic region: United States, 1999—Con.

[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table: see Methods section]

ICD-9-CM code	Sex			Age				Region			
	Total	Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	North-east	Midwest	South	West
Number of all-listed diagnoses in thousands											
991	*8	*	*	*	*	*	*	*	*	*	*
991.6	*6	*	*	*	*	*	*	*	*	*	*
992	*5	*	*	—	*	*	*	*	*	*	*
994	11	*8	*	*	*5	*	*	*	*	*	*
995	109	44	65	13	20	29	46	17	26	51	15
995.0	*6	*	*	*	*	*	*	*	*	*	*
995.1	10	*	*5	*	*	*	*	*	*	*	*
995.2	72	28	45	*	10	21	38	12	18	32	9
995.5	11	*5	*5	*8	*	—	—	*	*	*7	*
996	670	331	338	39	133	185	313	157	136	252	125
996.0	39	21	18	*	*	*9	27	11	10	12	*6
996.01	21	9	12	*	*	*	15	*5	*5	*8	*
996.04	*9	*6	*	—	*	*	*6	*	*	*	*
996.1	26	11	14	*	*5	*8	10	*5	*6	9	*6
996.2	31	15	16	11	13	*	*	*6	11	*9	*5
996.3	10	*6	*	*	*	*	*5	*	*	*	*
996.39	*6	*	*	*	*	*	*	*	*	*	*
996.4	95	39	56	*	11	22	61	21	26	28	20
996.5	24	10	14	*	*	*7	10	*7	*	*8	*
996.59	15	*7	*8	*	*	*	*8	*	*	*	*
996.6	196	99	98	13	43	55	85	50	39	77	31
996.61	10	*5	*	—	*	*	*5	*	*	*	*
996.62	102	50	52	10	24	32	36	29	17	41	16
996.63	*8	*	*	*	*	*	*	*	*	*	*
996.64	26	13	13	—	*	*	20	*	*6	13	*
996.66	14	*7	*7	*	*	*	10	*	*	*	*
996.67	12	*6	*6	*	*	*	*	*	*	*	*
996.68	*6	*	*	*	*	*	*	*	*	*	*
996.69	11	*6	*6	*	*	*	*	*	*	*	*
996.7	208	108	100	*	32	61	110	46	31	86	45
996.72	19	10	9	*	*	*7	11	*7	*	*8	*
996.73	72	36	36	*	13	25	34	11	9	31	21
996.74	59	30	29	*	*6	15	35	18	10	23	*9
996.76	12	*9	*	*	*	*	*8	*	*	*	*
996.77	12	*5	*7	—	*	*	*8	*	*	*	*
996.78	17	*9	*8	*	*6	*	*6	*	*	*9	*
996.79	*8	*	*	*	*	*	*	*	*	*	*
996.8	41	23	18	*	18	18	*	10	*7	20	*
996.81	28	17	10	*	15	11	*	*	*	15	*
997	619	301	318	15	95	179	330	146	135	187	151
997.0	33	15	18	*	*6	*8	18	*9	*5	11	*8
997.02	17	*7	10	*	*	*	12	*	*	*	*
997.09	13	*6	*7	*	*	*	*	*	*	*5	*
997.1	133	76	56	*	*6	31	93	40	35	30	28
997.2	22	*9	13	*	*	*8	11	9	*	*5	*
997.3	118	57	61	*5	19	34	59	26	21	35	35
997.4	182	75	107	*	43	57	78	33	39	62	48
997.5	85	45	40	*	14	21	48	20	18	26	21
997.6	29	17	12	*	*	13	13	*7	*7	12	*
997.62	17	10	*7	*	*	*8	*7	*6	*	*6	*
997.69	10	*6	*	—	*	*5	*	*	*	*5	*
997.9	18	*7	11	*	*	*6	10	*	*7	*	*
997.99	14	*5	9	*	*	*	*8	*	*7	*	*
998	632	289	343	21	136	193	281	157	136	212	126
998.1	215	99	116	*8	42	51	114	52	54	68	40
998.11	110	56	54	*7	23	23	57	25	29	34	22
998.12	92	37	56	*	16	25	51	25	22	30	16
998.13	13	*6	*6	*	*	*	*6	*	*	*	*

See footnotes and notes at end of table.

Table 44. Number of all-listed diagnoses for discharges from short-stay hospitals, by ICD-9-CM code, sex, age, and geographic region: United States, 1999—Con.[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table: see Methods section]

ICD-9-CM code	Sex			Age				Region			
	Total	Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	North-east	Midwest	South	West
Number of all-listed diagnoses in thousands											
998.2	74	24	50	*	19	24	29	15	16	22	20
998.3	40	19	21	*	*6	17	16	13	*8	14	*6
998.5	176	85	91	*6	40	64	66	46	35	60	35
998.59	172	83	89	*6	39	63	65	46	33	59	35
998.6	15	*7	*7	*	*	*	*6	*	*	*	*
998.8	103	50	53	*	24	31	45	26	20	39	19
998.81	*6	*	*	*	*	*	*5	*	*	*	*
998.83	12	*7	*5	—	*	*	*	*	*	*	*
998.89	84	39	46	*	20	26	35	20	16	34	14
999	46	20	26	*	10	11	21	11	13	13	10
999.2	16	*6	11	*	*	*	*7	*	*	*	*
999.3	11	*	*6	*	*	*	*	*	*	*	*
999.8	13	*7	*6	*	*	*	*8	*	*	*	*
V01-V82	11,418	3,356	8,062	267	5,452	1,789	3,911	2,219	2,672	4,074	2,453
V01	*6	*	*	*	*	*	*	*	*	*	*
V02	222	*9	213	*	211	*	*	35	49	83	55
V02.5	209	*	206	*	204	*	*	33	47	77	52
V02.51	198	—	198	*	198	—	—	30	42	74	51
V02.59	10	*	*8	*	*6	*	*	*	*	*	*
V02.6	12	*6	*6	*	*7	*	*	*	*	*	*
V02.61	*5	*	*	—	*	*	*	*	*	*	*
V02.62	*6	*	*	—	*	*	*	*	*	*	*
V04	10	*	10	*	10	*	*	*	*	*	*
V04.3	10	—	10	—	10	—	—	*	*	*	*
V06	*6	*	*6	*	*6	—	*	*	*	*	*
V06.4	*6	—	*6	—	*6	—	—	*	—	*	*
V07	38	*	36	*	30	*	*	*8	*	12	13
V07.2	25	*	25	—	25	—	*	*6	*	*8	*8
V07.4	*7	—	*7	—	*	*	*	*	*	*	*
V08	45	29	16	*	30	13	*	22	*	16	*
V09	56	27	29	*	*8	13	33	9	12	25	9
V09.0	50	24	26	*	*7	10	30	*8	11	22	9
V10	1,001	411	589	*	66	237	694	276	260	275	191
V10.0	222	110	111	*	*5	51	165	63	54	67	38
V10.02	*8	*6	*	—	*	*	*	*	*	*	*
V10.03	*7	*	*	—	*	*	*	*	*	*	*
V10.04	13	*7	*6	*	*	*	*9	*	*	*	*5
V10.05	162	76	86	*	*	29	130	47	40	48	27
V10.06	18	9	*9	—	*	*7	10	*	*	*7	*
V10.09	*8	*	*	—	*	*	*5	*	*	*	*
V10.1	74	36	38	*	*	18	54	22	18	24	10
V10.11	74	36	38	*	*	18	54	22	18	24	10
V10.2	17	13	*	—	*	*7	9	*5	*	*7	*
V10.21	14	11	*	—	*	*6	*8	*5	*	*	*
V10.3	252	*	250	—	14	70	168	68	67	68	49
V10.4	241	144	96	*	22	41	177	63	65	62	51
V10.41	28	—	28	—	10	10	*8	*8	*6	11	*
V10.42	36	—	36	—	*	*8	25	*8	13	*6	*9
V10.43	22	—	22	—	*	*8	11	*5	*	*6	*7
V10.44	*8	—	*8	*	*	*	*6	*	*	*	*
V10.46	135	135	—	—	*	10	125	34	38	36	27
V10.47	*8	*8	—	*	*	*	*	*	*	*	*
V10.5	74	49	24	*	*	19	52	25	21	14	14
V10.51	35	25	10	*	*	*6	28	14	*9	*7	*6
V10.52	36	22	14	*	*	13	21	10	11	*7	*8

See footnotes and notes at end of table.

Table 44. Number of all-listed diagnoses for discharges from short-stay hospitals, by ICD-9-CM code, sex, age, and geographic region: United States, 1999—Con.

[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table: see Methods section]

ICD-9-CM code	Sex			Age				Region			
	Total	Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	North-east	Midwest	South	West
Number of all-listed diagnoses in thousands											
V10.7	19	*7	11	*	*	*	11	*	*6	*	*5
V10.79	14	*6	*8	*	*	*	10	*	*	*	*
V10.8	95	47	48	*	16	25	53	25	23	27	20
V10.82	27	14	13	-	*	9	15	*8	*5	*8	*6
V10.83	29	17	12	-	*	*6	21	*7	*8	*7	*8
V10.85	10	*	*6	*	*	*	*	*	*	*	*
V10.87	10	*	*8	*	*	*	*5	*	*	*	*
V10.89	*9	*5	*	*	*	*	*	*	*	*	*
V11	33	20	13	*	*7	14	13	*	12	11	*8
V11.3	15	11	*	-	*	*8	*	*	*	*6	*
V11.8	11	*7	*	*	*	*	*	*	*7	*	*
V12	699	306	392	10	67	178	443	132	194	244	128
V12.0	50	21	30	*	13	13	22	9	*8	19	14
V12.01	24	11	12	-	*	*7	14	*6	*	*6	9
V12.02	9	*	*5	-	*	*	*6	*	*	*	*
V12.09	17	*5	12	*	10	*	*	*	*	11	*
V12.2	21	*9	12	*	*	*8	10	*	*8	*8	*
V12.3	*7	*	*5	-	*	*	*6	-	*	*	*
V12.4	15	*6	*9	*	*	*	*7	*	*6	*	*
V12.49	12	*	*7	*	*	*	*5	*	*	*	*
V12.5	446	191	255	*	32	95	314	96	122	151	76
V12.50	35	16	18	*	*	9	23	*8	*6	18	*
V12.51	91	34	56	*	15	25	50	16	27	27	21
V12.52	*8	*	*	-	*	*	*	*	*	*	*
V12.59	313	136	177	*	14	59	236	70	88	103	52
V12.6	25	10	15	*	*	9	12	*	*8	9	*
V12.7	134	69	66	*	14	48	71	19	38	49	29
V12.71	71	38	33	*	9	26	36	*6	19	28	19
V12.72	18	10	*8	-	*	*6	11	*	*6	*5	*
V12.79	41	19	22	*	*	15	23	9	12	13	*7
V13	91	39	52	20	21	21	29	15	30	27	19
V13.0	36	19	18	*	*9	12	14	*6	11	12	*6
V13.01	25	15	10	*	*6	10	*8	*5	*8	*7	*
V13.09	10	*	*6	*	*	*	*5	*	*	*	*
V13.2	*8	*	*7	*	*6	*	*	*	*	*	*
V13.4	*7	*	*	-	-	*	*	*	*	*	*
V13.5	9	*	*8	-	*	*	*5	*	*	*	*
V13.6	*5	*	*	*	*	*	*	*	*	*	*
V13.7	16	10	*6	15	*	*	-	-	9	*	*
V13.8	*7	*	*6	*	*	*	*	*	*	*	*
V14	46	15	31	*	14	17	15	12	*8	13	13
V14.0	25	9	16	*	*8	10	*8	9	*	*5	*7
V14.2	*6	*	*	*	*	*	*	*	*	*	*
V14.5	*5	*	*	-	*	*	*	*	*	*	*
V15	653	369	284	21	137	220	275	100	162	194	197
V15.0	*5	*	*	*	*	*	*	*	*	*	*
V15.1	20	10	10	14	*	*	*	*5	*6	*	*7
V15.3	25	15	10	*	*6	*7	11	*	*	*7	11
V15.4	*5	*	*	*	*	*	-	*	*	*	*
V15.5	24	12	12	*	*9	*	11	*	*8	*8	*7
V15.8	569	327	242	*	113	206	247	88	141	172	169
V15.81	156	83	73	*	69	50	34	33	43	53	27
V15.82	386	231	155	-	39	145	203	53	88	110	136
V15.89	23	*8	15	*	*	*8	*9	*	*9	*7	*6
V16	29	*6	23	*	*6	13	10	*6	*7	*8	*8
V16.0	15	*	11	-	*	*6	*6	*	*	*	*
V16.3	*7	*	*7	-	*	*	*	*	*	*	*

See footnotes and notes at end of table.

Table 44. Number of all-listed diagnoses for discharges from short-stay hospitals, by ICD-9-CM code, sex, age, and geographic region: United States, 1999—Con.[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table: see Methods section]

ICD-9-CM code	Sex			Age				Region			
	Total	Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	North-east	Midwest	South	West
Number of all-listed diagnoses in thousands											
V17	85	53	32	*	17	44	23	11	19	33	22
V17.3	68	44	24	—	14	37	18	9	15	28	17
V17.4	11	*6	*	*	*	*6	*	*	*	*	*
V18	11	*	*6	*	*8	*	*	*	*	*	*
V18.0	*8	*	*	*	*7	*	*	*	*	*	*
V22	26	—	26	—	26	—	—	*5	*8	10	*
V22.1	*5	—	*5	—	*5	—	—	*	*	*	*
V22.2	20	—	20	—	20	—	—	*	*7	*6	*
V23	118	—	118	*	116	*	—	15	17	57	30
V23.2	*6	—	*6	—	*6	—	—	*	*	*	*
V23.4	12	—	12	—	12	—	—	*	*	*	*
V23.7	74	—	74	*	74	—	—	*9	12	36	18
V23.8	19	—	19	*	17	*	—	*	*	10	*
V23.89	11	—	11	—	11	—	—	*	*	*8	*
V24	*7	—	*7	—	*7	—	—	*	*	*	*
V24.0	*6	—	*6	—	*6	—	—	*	*	*	*
V25	328	*	327	*	326	*	—	34	58	172	64
V25.0	*7	—	*7	*	*7	—	—	*	*	*6	*
V25.2	316	*	316	—	315	*	—	34	58	163	62
V26	*7	*	*7	—	*5	*	*	*	*	*	*
V27	3,810	—	3,810	*7	3,797	*6	—	636	809	1,454	912
V27.0	3,583	—	3,583	*7	3,572	*	—	601	694	1,403	884
V27.1	18	—	18	—	18	—	—	*	*	*8	*
V27.2	38	—	38	—	38	*	—	*8	*7	15	*9
V27.9	169	—	169	*	167	*	—	23	104	27	15
V29	10	*6	*	10	—	—	—	*	*	*	*
V29.0	10	*6	*	10	—	—	—	*	*	*	*
V30	12	9	*	12	—	—	—	*	*	*	*
V30.1	12	9	*	12	—	—	—	*	*	*	*
V42	113	59	54	11	33	42	27	32	20	43	18
V42.0	51	25	26	*	19	25	*	12	10	20	9
V42.1	*8	*6	*	*	*	*	*	*	*	*	*
V42.2	19	*9	11	*	*	*	17	*5	*	*6	*
V42.7	14	*9	*	*	*	*6	*	*	*	*7	*
V42.8	17	*8	*9	*5	*7	*	*	*6	*	*6	*
V42.81	10	*	*5	*	*	*	*	*	*	*	*
V43	406	157	249	*	22	85	298	93	110	134	69
V43.3	123	57	66	*	10	34	79	32	32	37	22
V43.6	274	96	178	*	11	49	214	58	77	94	45
V43.64	138	50	88	—	*7	26	106	29	36	51	23
V43.65	130	44	86	—	*	23	103	28	38	41	22
V44	175	90	85	29	24	35	86	42	48	50	35
V44.0	30	20	10	*7	*5	*8	10	*8	9	*9	*
V44.1	59	32	27	17	*6	*6	30	16	12	18	13
V44.2	14	*6	*8	*	*	*	*	*	*5	*	*
V44.3	53	24	29	*	*6	12	33	12	17	14	10
V44.6	10	*	*7	*	*	*	*5	*	*	*	*
V45	1,749	999	750	34	125	468	1,122	397	468	577	307
V45.0	278	138	139	*	*7	34	235	78	73	83	44
V45.01	248	116	131	*	*5	23	218	69	64	74	41
V45.02	25	18	*7	*	*	10	14	*7	*6	*9	*
V45.1	67	35	32	*	13	22	32	14	*7	30	16
V45.2	27	10	17	12	*7	*	*	*7	*7	*6	*6
V45.3	11	*	*7	*	*	*	*5	*	*	*	*

See footnotes and notes at end of table.

Table 44. Number of all-listed diagnoses for discharges from short-stay hospitals, by ICD-9-CM code, sex, age, and geographic region: United States, 1999—Con.

[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table: see Methods section]

ICD-9-CM code	Sex			Age				Region			
	Total	Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	North-east	Midwest	South	West
Number of all-listed diagnoses in thousands											
V45.4	32	14	19	*	11	12	9	*	9	12	*8
V45.6	11	*6	*	*	*	*	*8	*	*	*	*
V45.61	9	*5	*	*	*	*	*7	*	*	*	*
V45.7	52	13	39	*	*6	17	27	10	20	9	13
V45.71	22	*	21	—	*	*7	12	*	*8	*	*7
V45.72	*8	*	*5	*	*	*	*	*	*	*	*
V45.73	23	10	13	*	*	*8	12	*5	*7	*	*6
V45.8	1,271	778	493	17	76	376	802	281	347	428	216
V45.81	687	468	219	*	14	174	497	175	187	220	104
V45.82	361	220	141	*	19	147	194	69	100	120	71
V45.89	223	91	133	15	42	56	110	36	60	88	40
V46	17	*8	*9	*	*	*	*6	*6	*	*6	*
V46.1	16	*8	*8	*	*	*	*5	*	*	*6	*
V49	84	53	31	*	14	26	42	17	16	36	15
V49.6	*9	*6	*	*	*	*	*	*	*	*	*
V49.7	75	46	28	*	11	21	41	15	14	33	13
V49.72	*7	*	*	*	*	*	*	*	*	*	*
V49.75	32	18	13	*	*	12	18	*7	*7	13	*
V49.76	27	19	*8	*	*6	*6	15	*	*	12	*7
V50	*8	*	*	*	*	*	*	*	*	*	*
V53	17	10	*7	*	*	*	12	*	*	*7	*
V53.3	13	*9	*	*	*	*	11	*	*	*6	*
V53.31	10	*6	*	*	*	*	*8	*	*	*	*
V54	30	11	19	*	*6	*5	16	*6	*	13	*6
V54.0	*7	*	*	*	*	*	*	*	*	*	*
V54.8	15	*5	9	*	*	*	*8	*	*	*7	*
V54.9	*9	*	*6	—	*	*	*5	*	*	*	*
V55	51	27	24	*6	*9	16	19	13	*8	19	10
V55.1	*8	*	*6	*	*	*	*	*	*	*	*
V55.2	*7	*	*	*	*	*	*	*	*	*	*
V55.3	29	18	12	*	*5	12	10	*7	*5	11	*6
V56	85	34	51	*	15	27	43	12	12	36	26
V56.0	81	32	49	*	14	26	41	10	12	35	24
V57	606	243	363	10	51	112	433	101	140	259	107
V57.1	169	64	104	*	9	35	121	37	15	85	31
V57.2	62	25	38	*	*5	11	45	11	*8	31	12
V57.21	62	25	38	*	*5	11	45	11	*8	31	12
V57.8	356	145	211	*	35	64	252	47	113	137	58
V57.89	355	145	210	*	35	64	251	47	113	137	58
V57.9	15	*6	*9	—	*	*	13	*6	*	*	*
V58	346	181	164	40	60	91	155	83	74	108	81
V58.0	24	13	11	*	*	10	9	*	*	15	*
V58.1	159	90	69	35	32	45	46	47	30	49	32
V58.4	24	12	13	*	*8	*6	*9	*7	*	*7	*6
V58.49	23	11	12	*	*7	*5	*8	*7	*	*6	*5
V58.6	124	60	64	*	13	28	81	24	32	32	35
V58.61	109	52	57	*	11	25	73	23	28	29	30
V58.69	13	*7	*6	*	*	*	*8	*	*	*	*
V58.8	13	*5	*8	*	*	*	*9	*	*	*	*
V58.89	12	*	*7	*	*	*	*9	*	*	*	*
V59	*6	*	*	*	*	*	*	*	*	*	*
V60	28	16	12	*	13	*8	*6	*	9	*8	*8
V60.0	16	12	*	*	10	*5	*	*	*5	*	*

See footnotes and notes at end of table.

Table 44. Number of all-listed diagnoses for discharges from short-stay hospitals, by ICD-9-CM code, sex, age, and geographic region: United States, 1999—Con.

[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table: see Methods section]

ICD-9-CM code	Sex			Age				Region			
	Total	Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	North-east	Midwest	South	West
Number of all-listed diagnoses in thousands											
V61	55	25	30	10	40	*	*	*	24	19	*7
V61.0	*6	*	*	*	*5	*	*	*	*	*	*
V61.1	13	9	*	—	10	*	*	—	*	*6	*
V61.10	12	9	*	—	10	*	*	—	*	*6	*
V61.2	18	*8	10	*8	10	*	—	*	*8	*7	*
V61.20	16	*7	*9	*8	*9	*	—	*	*7	*6	*
V61.5	*9	—	*9	—	*9	—	—	*	*	*	*
V62	35	19	15	*	22	*8	*	*6	11	12	*7
V62.8	21	10	11	*	13	*	*	*	*8	*6	*
V62.89	14	*6	*8	*	9	*	*	*	*5	*	*
V64	102	39	63	*	35	28	37	23	18	40	21
V64.1	24	9	15	*	*	*7	13	*6	*	*8	*6
V64.2	15	*9	*6	*	*	*	*7	*	*	*6	*
V64.3	18	*7	11	*	*6	*	*6	*	*	10	*
V64.4	45	14	31	*	20	14	11	10	9	17	9
V65	17	*9	*8	*	*8	*	*	*	*	*	*5
V65.3	10	*	*5	*	*	*	*	*	*	*	*
V66	44	23	22	*	*	*8	34	12	12	11	*9
V66.2	*6	*	*	*	*	*	*	*	*	*	*
V66.7	29	16	14	*	*	*	24	*7	*8	*9	*5
V67	*6	*	*	*	*	*	*	*	*	*	*
V70	15	9	*6	*	9	*	*	*5	*	*6	*
V70.7	13	*8	*	*	*7	*	*	*5	*	*	*
V71	29	13	15	*	11	*8	*5	*5	*8	*5	10
V71.0	*6	*	*	*	*	*	*	*	*	*	*
V71.09	*6	*	*	—	*	*	*	—	*	*	*
V71.4	*5	*	*	*	*	*	*	*	*	*	*
V71.7	10	*	*6	*	*	*5	*	*	*	*	*6
V72	10	*	*6	*	*	*	*	*	—	*6	*
V72.8	*6	*	*	*	*	*	*	*	—	*	*

* Figure does not meet standard of reliability or precision.

— Quantity zero.

NOTE: Estimates of 5,000-9,000 are to be used with caution; see Methods section.

Table 45. Number of all-listed procedures for discharges from short-stay hospitals, by ICD-9-CM code, sex, age, and geographic region: United States, 1999

[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table: see Methods section]

ICD-9-CM code	Sex			Age				Region			
	Total	Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	North-east	Midwest	South	West
Number of procedures in thousands											
All codes	41,315	16,370	24,945	2,077	13,791	9,981	15,467	9,797	9,025	14,072	8,421
01-05	1,048	475	572	212	337	228	270	222	284	323	219
01	134	74	60	16	37	36	44	27	28	45	34
01.0	*9	*	*5	*	*	*	*	*	*	*	*
01.02	*6	*	*	*	*	*	*	*	*	*	*
01.1	23	11	13	*	*8	*7	*6	*5	*	9	*
01.13	*7	*	*	*	*	*	*	*	*	*	*
01.18	11	*6	*	*	*	*	*	*	*	*	*
01.2	25	15	10	*	*7	*7	*7	*	*	9	*7
01.24	18	11	*7	*	*6	*	*	*	*	*7	*
01.3	35	22	13	*	*8	*8	16	*5	*7	12	10
01.31	23	13	10	*	*	*5	12	*	*	10	*7
01.39	11	*8	*	*	*	*	*	*	*	*	*
01.5	39	21	18	*	10	12	13	10	*8	12	*9
01.51	9	*	*7	*	*	*	*	*	*	*	*
01.59	28	18	11	*	*8	*9	*8	*6	*7	*8	*7
02	126	64	62	32	45	24	24	29	29	38	30
02.0	18	9	*9	*6	*7	*	*	*	*	*6	*6
02.02	*6	*	*	*	*	*	*	*	*	*	*
02.06	*8	*	*	*	*	*	*	*	*	*	*
02.1	10	*	*6	*	*	*	*	*	*	*	*
02.12	9	*	*6	*	*	*	*	*	*	*	*
02.2	24	12	12	*	9	*6	*6	*6	*	*8	*6
02.3	23	12	11	*7	*5	*	*7	*	*	*9	*6
02.34	18	*8	10	*6	*	*	*6	*	*	*6	*5
02.4	32	17	16	13	13	*	*	*	12	*8	*7
02.42	25	12	13	11	11	*	*	*	11	*6	*
02.43	*8	*	*	*	*	*	*	*	*	*	*
02.9	18	11	*7	*	*8	*	*	*8	*	*	*
02.94	11	*7	*	*	*	*	*	*5	*	*	*
03	731	309	421	159	235	152	185	154	212	217	147
03.0	146	74	72	*	23	54	67	32	34	50	30
03.09	141	70	71	*	21	53	65	31	33	48	29
03.3	313	159	154	144	71	46	51	73	74	96	70
03.31	310	158	153	144	71	45	50	72	73	95	69
03.4	13	*7	*7	*	*	*	*	*	*	*	*
03.5	22	12	10	*	*8	*	*8	*	*5	*7	*6
03.53	*7	*5	*	—	*	*	*	*	*	*	*
03.59	15	*6	*9	*	*	*	*5	*	*	*	*
03.6	*7	*	*	*	*	*	*	*	*	*	*
03.9	221	50	171	*6	126	38	52	38	94	54	35
03.90	96	22	74	*	53	15	24	18	39	25	14
03.91	70	*9	62	*	50	9	11	10	40	10	10
03.92	30	13	16	*	*6	*8	14	*6	11	*7	*5
03.93	*8	*	*	—	*	*	*	*	*	*	*
03.95	12	*	10	*	11	*	*	*	*	*	*
04	50	24	25	*5	18	13	14	10	11	20	*8
04.0	*8	*5	*	*	*	*	*	*	*	*	*
04.4	16	*8	*8	*	*6	*	*	*	*	*7	*
04.43	*8	*	*	*	*	*	*	*	*	*	*
04.49	*5	*	*	*	*	*	*	*	*	*	*
04.8	12	*	*7	*	*	*	*6	*	*	*	*
04.81	11	*	*7	*	*	*	*6	*	*	*	*
05	*8	*	*	*	*	*	*	*	*	*	*
05.3	*6	*	*	—	*	*	*	*	*	*	*
05.31	*6	*	*	—	*	*	*	*	*	*	*

See footnotes and notes at end of table.

Table 45. Number of all-listed procedures for discharges from short-stay hospitals, by ICD-9-CM code, sex, age, and geographic region: United States, 1999—Con.[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table: see Methods section]

ICD-9-CM code	Sex			Age				Region			
	Total	Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	North-east	Midwest	South	West
Number of procedures in thousands											
06-07	87	20	67	*	29	33	20	22	16	29	20
06	75	15	60	*	25	30	18	18	14	25	18
06.2	19	*	15	*	*7	*7	*	*	*	*	*
06.3	16	*	13	—	*5	*6	*	*	*	*8	*
06.39	14	*	13	—	*	*5	*	*	*	*8	*
06.4	14	*	13	*	*7	*5	*	*	*	*	*
06.8	12	*	9	—	*	*7	*	*	*	*	*
06.89	11	*	*9	—	*	*7	*	*	*	*	*
07	12	*	*7	*	*	*	*	*	*	*	*
08-16	110	59	51	10	31	28	40	16	17	59	18
08	33	19	15	*	15	9	*7	*7	*9	11	*7
08.8	20	11	*9	*	11	*	*	*	*7	*6	*
08.81	16	10	*6	*	9	*	*	*	*7	*	*
11	*7	*5	*	*	*	*	*	*	*	*	*
12	*5	*	*	*	*	*	*	*	*	*	*
13	16	*	11	*	*	*	10	*	*	10	*
13.4	*5	*	*	*	*	*	*	*	*	*	*
13.7	*6	*	*	—	*	*	*	*	*	*	*
13.71	*6	*	*	—	*	*	*	*	*	*	*
14	32	15	17	*	*7	*8	13	*	*	20	*
14.4	*5	*	*	*	*	*	*	*	*	*	*
14.7	13	*7	*5	*	*	*	*6	*	*	*8	*
14.74	*6	*	*	*	*	*	*	*	*	*	*
16	12	9	*	*	*5	*	*	*	*	*7	*
18-20	54	32	22	27	11	*8	*7	15	*7	20	12
18	21	12	*9	*7	*7	*	*	*	*	*8	*6
18.4	*8	*	*	*	*5	*	*	*	*	*	*
20	30	18	12	19	*	*	*	11	*	10	*6
20.0	18	11	*7	14	*	*	*	*7	*	*6	*
20.01	15	9	*6	11	*	*	*	*7	*	*	*
20.4	*5	*	*	*	*	*	*	*	*	*	*
21-29	284	175	109	67	101	60	56	81	51	83	68
21	66	42	24	*	24	16	23	16	13	22	15
21.0	22	12	10	*	*	*5	14	*7	*7	*5	*
21.01	10	*6	*	*	*	*	*6	*	*	*	*
21.02	*5	*	*	*	*	*	*	*	*	*	*
21.03	*5	*	*	*	*	*	*	*	*	*	*
21.2	*5	*	*	*	*	*	*	*	*	*	*
21.6	*8	*6	*	*	*	*	*	*	*	*	*
21.69	*6	*	*	—	*	*	*	*	*	*	*
21.7	*7	*	*	*	*	*	*	*	*	*	*
21.8	18	14	*	*	11	*	*	*	*	*7	*
21.88	*8	*7	*	*	*	*	*	*	*	*	*
22	42	27	16	*8	16	10	*8	15	*6	11	10
22.2	*7	*	*	*	*	*	*	*	*	*	*
22.4	*7	*6	*	*	*	*	*	*	*	*	*
22.42	*5	*	*	*	*	*	*	*	*	*	*
22.6	16	10	*6	*	*5	*	*	*6	*	*	*
22.63	10	*6	*	*	*	*	*	*	*	*	*
23	31	18	14	9	14	*	*	*7	*5	11	*7
23.0	13	*8	*5	*	*6	*	*	*	*	*5	*
23.09	12	*7	*5	*	*6	*	*	*	*	*	*
23.1	13	*8	*5	*	*7	*	*	*	*	*	*
23.19	13	*8	*	*	*7	*	*	*	*	*	*
24	*8	*	*	*	*	*	*	*	*	*	*

See footnotes and notes at end of table.

Table 45. Number of all-listed procedures for discharges from short-stay hospitals, by ICD-9-CM code, sex, age, and geographic region: United States, 1999—Con.

[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table: see Methods section]

ICD-9-CM code	Sex			Age				Region			
	Total	Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	North-east	Midwest	South	West
Number of procedures in thousands											
25	9	*6	*	*	*	*	*	*	*	*	*
26	14	*8	*6	*	*	*	*5	*	*	*	*
26.3	11	*6	*	*	*	*	*	*	*	*	*
26.31	*6	*	*	*	*	*	*	*	*	*	*
27	57	36	21	21	19	*9	*8	16	10	18	13
27.0	10	*7	*	*	*7	*	*	*	*	*	*
27.5	21	11	9	*8	*7	*	*	*	*	*7	*5
27.51	*9	*	*	*	*	*	*	*	*	*	*
27.54	*5	*	*	*5	*	—	—	*	*	*	*
27.6	16	11	*	9	*	*	*	*7	*	*	*
27.62	*7	*	*	*7	*	—	*	*	*	*	*
27.69	*6	*	*	*	*	*	*	*	*	*	*
28	36	19	17	19	13	*	*	12	*6	11	*7
28.0	*6	*	*	*	*	*	*	*	*	*	*
28.2	11	*7	*	*	*6	*	*	*	*	*	*
28.3	11	*6	*5	9	*	—	—	*	*	*	*
29	21	16	*	*	*7	*8	*	*7	*	*	*6
29.1	*5	*	*	*	*	*	*	*	*	*	*
29.4	9	*8	*	*	*	*	*	*	*	*	*
30-34	1,027	573	455	61	178	281	507	262	221	360	184
30	*9	*	*	*	*	*	*	*	*	*	*
31	157	90	66	19	34	35	69	44	29	54	30
31.1	83	47	36	*	21	19	40	23	15	30	15
31.2	13	*6	*7	*	*	*	*8	*	*	*6	*
31.29	13	*6	*7	*	*	*	*8	*	*	*6	*
31.4	50	31	19	12	*8	12	19	15	*9	14	12
31.42	41	25	16	12	*6	*8	14	13	*8	10	10
31.43	*7	*5	*	*	*	*	*	*	*	*	*
32	71	39	32	*	10	26	33	19	12	27	13
32.2	28	17	11	*	*7	12	*8	*8	*	11	*
32.29	24	15	10	*	*7	10	*7	*7	*	10	*
32.3	*6	*	*	*	*	*	*	*	*	*	*
32.4	31	15	16	*	*	10	19	*8	*6	10	*7
33	322	182	141	19	39	97	167	95	71	106	51
33.2	314	178	136	17	38	96	164	92	68	104	50
33.21	*5	*	*	*	*	*	*	*	*	*	*
33.22	54	28	26	*	*	18	28	16	9	19	9
33.23	46	26	20	*8	*6	12	20	15	10	15	*6
33.24	124	76	49	*	17	37	67	35	28	41	20
33.26	26	12	14	*	*	*8	17	*7	*5	10	*
33.27	39	23	16	*	*	13	21	10	10	13	*6
33.28	18	10	*8	*	*	*7	*8	*6	*	*5	*
34	469	257	213	21	95	118	235	103	107	170	89
34.0	196	117	80	13	57	48	77	41	44	71	39
34.01	*6	*	*	*	*	*	*	*	*	*	*
34.03	*5	*	*	*	*	*	*	*	*	*	*
34.04	152	91	62	10	45	35	62	36	36	51	30
34.09	29	15	13	*	11	*9	*8	*	*6	12	*8
34.2	47	24	23	*	*7	15	24	13	*8	16	9
34.21	14	*8	*7	*	*	*6	*5	*	*	*	*
34.22	10	*	*5	*	*	*	*6	*	*	*	*
34.24	13	*6	*7	*	*	*	*8	*	*	*	*
34.5	18	12	*6	*	*6	*	*6	*	*	*7	*
34.51	12	*8	*	*	*	*	*	*	*	*	*
34.59	*6	*	*	—	*	*	*	*	*	*	*
34.6	*8	*	*	*	*	*	*	*	*	*	*
34.7	*5	*	*	*	*	*	*	*	*	*	*

See footnotes and notes at end of table.

Table 45. Number of all-listed procedures for discharges from short-stay hospitals, by ICD-9-CM code, sex, age, and geographic region: United States, 1999—Con.[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table: see Methods section]

ICD-9-CM code	Sex			Age				Region			
	Total	Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	North-east	Midwest	South	West
Number of procedures in thousands											
34.9	184	88	96	*	18	45	119	40	45	68	31
34.91	159	76	82	*	16	36	104	34	41	57	27
34.92	25	11	14	*	*	*9	14	*5	*	11	*
35-39	6,133	3,498	2,634	179	569	2,170	3,215	1,400	1,381	2,241	1,111
35	132	62	70	25	15	30	62	46	23	36	28
35.1	16	*7	*9	*	*	*	*7	*	*	*	*
35.12	11	*6	*5	*	*	*	*5	*	*	*	*
35.2	80	40	40	*	*7	21	49	26	16	22	16
35.21	20	12	*8	*	*	*	14	*9	*	*	*
35.22	32	16	15	*	*	9	20	10	*7	*9	*6
35.23	*5	*	*	—	*	*	*	*	*	*	*
35.24	20	*9	12	*	*	*8	10	*5	*	*6	*
35.3	*7	*	*	*	*	*	*	*	*	*	*
35.33	*6	*	*	*	*	*	*	*	*	*	*
35.7	10	*	*7	*	*	*	*	*	*	*	*
35.71	*9	*	*5	*	*	*	*	*	*	*	*
35.9	*5	*	*	*	*	*	*	*	*	*	*
36	1,646	1,101	545	*	75	741	828	373	426	542	305
36.0	1,069	708	361	*	58	505	504	226	292	361	191
36.01	497	327	170	*	29	233	234	107	130	169	90
36.02	23	15	*8	—	*	11	10	*	*9	*6	*
36.04	*9	*6	*	—	*	*	*	*6	*	*	*
36.05	81	56	25	—	*	36	42	11	27	25	19
36.06	457	303	154	*	23	221	212	98	125	158	77
36.1	571	389	182	*	16	232	322	147	133	178	113
36.10	*6	*	*	—	*	*	*	*	—	*	*
36.11	61	38	23	—	*	22	37	20	10	16	15
36.12	98	65	33	—	*	34	61	27	25	30	17
36.13	108	73	35	—	*	44	62	22	28	37	22
36.14	56	43	13	*	*	22	33	*7	17	19	13
36.15	218	151	67	—	*8	93	117	61	50	66	41
36.16	16	11	*5	—	*	10	*6	*8	*	*	*
36.19	*7	*5	*	—	*	*	*	*	*	*	*
37	1,844	1,065	779	20	133	670	1,021	423	413	725	283
37.0	11	*6	*6	*	*	*	*	*	*	*	*
37.1	12	*7	*5	*	*	*	*	*	*	*6	*
37.12	11	*6	*	*	*	*	*	*	*	*6	*
37.2	1,359	813	547	12	109	576	663	301	306	540	213
37.21	36	21	15	*	*	15	14	*7	*	21	*
37.22	1,062	640	422	*	80	477	504	216	248	425	173
37.23	173	97	76	*7	11	56	99	46	38	66	24
37.25	*5	*	*	*	*	*	*	*	*	*	*
37.26	66	43	23	*	*8	21	36	26	11	20	9
37.27	11	*6	*5	*	*	*	*	*	*	*	*
37.3	32	16	16	*	*7	9	13	10	*7	10	*
37.33	*5	*	*	*	*	*	*	*	*	*	*
37.34	22	12	11	*	*	*7	10	*7	*	*7	*
37.6	43	29	14	—	*	17	23	9	11	14	*8
37.61	38	26	13	—	*	16	21	9	10	12	*7
37.7	174	83	91	*	*	25	144	44	40	65	24
37.71	31	12	19	*	*	*	28	9	*7	12	*
37.72	93	46	47	*	*	11	79	24	19	38	13
37.76	*8	*	*	*	*	*	*6	*	*	*	*
37.78	29	15	14	*	*	*7	20	*6	10	*8	*
37.8	162	77	86	*	*	17	141	41	30	68	23
37.80	*9	*	*5	—	*	*	*8	*	*	*	*
37.81	17	*6	11	*	—	*	15	*	*	*8	*

See footnotes and notes at end of table.

Table 45. Number of all-listed procedures for discharges from short-stay hospitals, by ICD-9-CM code, sex, age, and geographic region: United States, 1999—Con.

[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table: see Methods section]

ICD-9-CM code	Sex			Age				Region			
	Total	Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	North-east	Midwest	South	West
Number of procedures in thousands											
37.82	21	*7	13	*	*	*	20	*	*	10	*
37.83	93	46	47	*	*	11	80	24	19	37	14
37.87	11	*7	*	*	*	*	*8	*	*	*	*
37.9	46	32	13	*	*	16	27	12	12	13	*8
37.94	35	26	10	—	*	12	21	10	10	*9	*7
38	1,259	586	673	98	184	310	667	273	280	473	233
38.0	35	15	19	*	*	9	21	*9	*8	14	*
38.03	*6	*	*	*	*	*	*	*	*	*	*
38.08	20	*9	11	*	*	*6	13	*	*5	*9	*
38.1	156	80	76	—	*	39	115	34	42	56	24
38.12	131	67	65	—	*	32	99	26	37	48	21
38.18	19	10	9	—	*	*5	13	*6	*	*7	*
38.2	10	*	*	*	*	*	*5	*	*	*	*
38.21	*7	*	*	—	*	*	*	*	*	*	*
38.4	49	35	14	*	*	11	35	11	10	17	11
38.44	33	24	*9	—	*	*6	27	9	*7	10	*6
38.45	*7	*	*	*	*	*	*	*	*	*	*
38.6	12	*8	*	*	*	*	*7	*	*	*	*
38.7	49	18	31	—	*7	14	28	16	10	17	*6
38.8	37	16	21	10	9	*8	10	11	*6	12	*8
38.85	12	*6	*6	*9	*	*	*	*	*	*	*
38.86	9	*	*6	*	*	*	*	*	*	*6	*
38.9	905	405	499	84	157	220	443	183	197	349	176
38.91	97	50	47	17	12	20	48	21	33	31	12
38.92	*9	*5	*	*8	—	*	*	*	*	*	*
38.93	648	278	369	54	115	158	321	130	138	248	132
38.94	12	*7	*	*	*	*	*6	*	*	*	*
38.95	124	59	65	*	25	37	60	24	23	52	25
38.98	10	*	*7	*	*	*	*6	*	*	*7	*
38.99	*6	*	*	*	*	*	*	*	*	*	*
39	1,252	684	567	35	161	419	636	285	240	465	262
39.1	9	*	*6	*	*6	*	*	*	*	*	*
39.2	156	87	69	*	11	47	97	38	31	60	27
39.25	24	15	9	—	*	12	11	*7	*5	*8	*
39.27	39	21	18	*	*7	12	21	*8	*5	16	10
39.29	85	49	36	—	*	21	61	21	19	32	13
39.3	29	17	11	*	13	*8	*6	*8	*	*9	*7
39.31	19	11	*8	*	*7	*6	*	*5	*	*6	*
39.32	*8	*5	*	*	*	*	*	*	*	*	*
39.4	73	38	35	*	11	26	35	17	10	30	16
39.42	19	10	*9	—	*	*6	*8	*	*	10	*
39.49	48	25	23	*	*	18	24	12	*7	18	11
39.5	126	69	58	*	19	39	65	29	35	42	21
39.50	80	43	36	*	*8	23	47	20	21	26	12
39.51	9	*	*6	—	*	*	*	*	*	*	*
39.52	13	*7	*5	*	*	*5	*7	*	*	*	*
39.56	*7	*	*	*	*	*	*	*	*	*	*
39.57	*7	*	*	*	*	*	*	*	*	*	*
39.59	*6	*	*	*	*	*	*	*	*	*	*
39.6	344	215	129	20	25	120	180	81	67	110	86
39.61	317	197	120	18	21	109	169	77	64	95	81
39.64	17	12	*5	*	*	*6	*8	*	*	11	*
39.9	508	252	257	*	77	177	250	107	90	210	100
39.90	30	17	13	*	*	11	16	*8	*9	12	*
39.95	458	227	231	*	70	161	226	95	78	189	96
39.98	*8	*	*6	*	*	*	*	*	*	*	*
40-41	348	170	178	25	53	113	157	96	81	109	63

See footnotes and notes at end of table.

Table 45. Number of all-listed procedures for discharges from short-stay hospitals, by ICD-9-CM code, sex, age, and geographic region: United States, 1999—Con.[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table: see Methods section]

ICD-9-CM code	Sex			Age				Region			
	Total	Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	North-east	Midwest	South	West
Number of procedures in thousands											
40	203	101	102	*	29	78	93	55	46	66	37
40.1	51	20	31	*	*8	17	24	14	14	16	*8
40.11	50	19	31	*	*7	17	24	14	14	15	*7
40.2	45	16	30	*	*8	16	20	10	12	15	*7
40.23	9	*	*9	—	*	*	*	*	*	*	*
40.29	30	12	18	*	*7	*9	14	*5	9	10	*5
40.3	70	41	29	*	*6	31	33	21	12	24	13
40.4	14	10	*	*	*	*	*5	*5	*	*	*
40.41	10	*8	*	*	*	*	*	*	*	*	*
40.5	21	13	*7	—	*	*8	10	*	*6	*6	*6
40.53	*6	*	*	—	*	*	*	*	*	*	*
40.54	*5	*	*	—	*	*	*	*	*	*	*
40.59	*7	*	*	—	*	*	*	*	*	*	*
41	145	69	76	21	24	35	64	41	34	44	26
41.0	*7	*	*	*	*	*	*	*	*	*	—
41.3	108	53	55	16	12	24	56	31	25	32	19
41.31	107	53	54	16	12	24	55	31	25	32	19
41.5	25	11	14	*	*7	*9	*7	*5	*6	*9	*6
42-54	5,189	2,189	3,000	217	1,111	1,385	2,476	1,119	1,205	1,846	1,019
42	188	103	84	*9	24	61	93	49	46	61	32
42.2	114	58	56	*6	16	40	51	36	25	37	16
42.23	109	55	54	*5	16	39	48	32	25	36	16
42.3	27	16	11	*	*	11	11	*6	*6	10	*6
42.33	26	16	11	*	*	11	11	*6	*5	9	*6
42.8	*6	*	*	*	*	*	*	*	*	*	*
42.9	33	19	14	*	*	*	25	*5	13	11	*
42.92	32	19	13	*	*	*	24	*5	13	11	*
43	243	118	126	22	18	44	160	56	47	99	41
43.1	195	93	103	13	12	32	138	44	36	83	32
43.11	163	79	84	*5	10	25	123	35	30	74	24
43.19	32	14	19	*8	*	*7	15	*9	*6	10	*7
43.3	9	*7	*8	*8	—	*	*	*	*	*	*
43.4	12	*	*8	*	*	*	*8	*	*	*	*
43.41	*9	*	*6	*	*	*	*6	*	*	*	*
43.7	11	*	*6	—	*	*	*	*	*	*	*
43.9	*6	*	*	—	*	*	*	*	*	*	*
43.99	*6	*	*	—	*	*	*	*	*	*	*
44	213	102	111	11	46	59	97	41	48	79	45
44.0	*9	*5	*	—	*	*	*	*	*	*	*
44.01	*5	*	*	—	*	*	*	*	*	*	*
44.1	36	17	19	*	*7	*9	20	*5	*6	23	*
44.13	21	12	10	*	*6	*	10	*	*	14	*
44.14	12	*5	*7	—	*	*	*8	*	*	*8	*
44.2	12	*6	*6	*	*	*	*6	*	*	*	*
44.29	10	*6	*	*	*	*	*5	*	*	*	*
44.3	24	*6	17	*	*7	11	*5	*6	*6	*	*7
44.31	10	*	*8	—	*	*6	*	*	*	*	*
44.39	13	*	9	*	*	*5	*	*	*	*	*
44.4	71	40	31	—	*7	18	46	15	19	24	14
44.41	*6	*	*	—	*	*	*	*	*	*	*
44.42	11	*7	*	—	*	*	*7	*	*	*	*
44.43	53	29	24	—	*	13	35	*9	13	20	11
44.6	59	26	33	10	21	15	13	10	11	20	18
44.66	42	20	21	*7	15	12	*8	*7	*7	13	14
44.69	*8	*	*6	*	*	*	*	*	*	*	*

See footnotes and notes at end of table.

Table 45. Number of all-listed procedures for discharges from short-stay hospitals, by ICD-9-CM code, sex, age, and geographic region: United States, 1999—Con.

[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table: see Methods section]

ICD-9-CM code	Sex			Age				Region			
	Total	Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	North-east	Midwest	South	West
Number of procedures in thousands											
45	1,990	851	1,140	36	238	490	1,226	428	503	708	351
45.1	926	405	521	18	129	225	554	195	232	333	167
45.13	389	176	213	*	53	93	238	84	100	135	70
45.16	528	224	305	13	73	129	313	109	130	194	95
45.2	544	209	335	*8	59	120	358	125	132	198	88
45.23	286	111	175	*	23	61	200	63	66	105	52
45.24	66	26	40	*	*9	14	43	14	20	27	*6
45.25	187	69	118	*	27	43	112	47	46	66	28
45.3	*8	*	*	*	*	*	*5	*	*	*	*
45.33	*6	*	*	*	*	*	*	*	*	*	*
45.4	117	57	60	*	*6	23	88	23	32	43	19
45.41	*6	*	*	—	*	*	*	*	*	*	*
45.42	95	46	49	*	*	20	72	20	27	33	14
45.43	16	*7	*8	*	*	*	13	*	*	*6	*
45.6	63	26	37	*	10	17	32	14	13	22	15
45.62	59	25	34	*	9	15	30	13	12	20	14
45.7	251	115	136	*	21	82	146	50	71	86	43
45.72	13	*5	*8	*	*	*	*6	*	*	*	*
45.73	84	38	46	*	*6	22	55	14	24	30	15
45.74	9	*	*6	*	*	*	*7	*	*	*	*
45.75	37	16	21	—	*	14	20	*6	9	16	*
45.76	80	37	43	*	*5	33	41	17	24	25	14
45.79	27	15	13	*	*	*8	16	*7	*8	*8	*
45.8	*7	*	*	*	*	*	*	*	*	*	*
45.9	67	29	38	*	9	19	36	14	18	20	16
45.91	15	*	10	*	*	*	*8	*	*5	*	*
45.93	30	14	16	*	*	9	17	*7	*6	11	*6
45.94	18	*7	10	*	*	*7	*9	*	*5	*5	*
46	210	108	102	12	37	61	100	44	48	75	43
46.0	15	*8	*7	*	*	*	*7	*	*	*	*
46.01	*6	*	*	*	*	*	*	*	*	*	*
46.03	*9	*	*	*	*	*	*	*	*	*	*
46.1	51	26	25	*	*6	16	28	10	10	22	*9
46.10	26	13	13	*	*	*7	14	*5	*6	12	*
46.11	16	10	*7	*	*	*6	*9	*	*	*7	*5
46.13	*8	*	*	*	*	*	*	*	*	*	*
46.2	12	*	*7	*	*	*	*	*	*	*	*
46.20	*5	*	*	*	*	*	*	*	*	*	*
46.3	29	15	14	*	*	*7	17	*7	*7	11	*
46.32	*7	*	*	*	*	*	*	*	*	*	*
46.39	22	12	10	*	*	*5	13	*6	*5	*8	*
46.4	12	*7	*	*	*	*	*7	*	*	*	*
46.5	42	23	19	*	*9	17	15	*9	9	15	*9
46.51	*7	*	*	*	*	*	*	*	*	*	*
46.52	34	19	16	*	*6	14	13	*7	*7	13	*7
46.7	31	14	17	*	11	*8	10	*7	*7	*8	*9
46.73	15	*6	10	*	*	*	*5	*	*	*	*
46.75	*6	*	*	*	*	*	*	*	*	*	*
46.8	14	*8	*7	*	*	*	10	*	*	*6	*
46.85	*7	*	*	*	*	*	*6	*	*	*	*
47	345	174	171	72	184	57	32	63	84	104	93
47.0	290	158	132	65	158	42	24	51	70	88	81
47.01	66	32	34	14	39	*9	*	*7	19	19	21
47.09	224	126	98	52	119	33	20	44	50	69	61
47.1	54	15	39	*7	25	15	*8	12	14	16	12
47.11	*6	*	*5	*	*	*	*	*	*	*	*
47.19	47	14	33	*6	20	14	*8	11	13	15	*9

See footnotes and notes at end of table.

Table 45. Number of all-listed procedures for discharges from short-stay hospitals, by ICD-9-CM code, sex, age, and geographic region: United States, 1999—Con.[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table: see Methods section]

ICD-9-CM code	Sex			Age				Region			
	Total	Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	North-east	Midwest	South	West
Number of procedures in thousands											
48	132	63	69	*8	17	36	71	34	29	45	23
48.2	52	21	31	*	*7	11	28	13	11	18	9
48.23	22	*9	13	*	*	*6	12	*6	*5	*8	*
48.24	26	11	15	*	*	*5	14	*6	*5	9	*6
48.3	18	9	*9	*	*	*	12	*	*6	*8	*
48.36	12	*6	*7	*	*	*	9	*	*	*6	*
48.5	11	*7	*	—	*	*	*5	*	*	*	*
48.6	27	13	14	*	*	*7	18	10	*5	*6	*5
48.63	17	*8	*8	—	*	*5	10	*6	*	*	*
48.69	*8	*	*5	*	*	*	*6	*	*	*	*
48.7	*8	*	*6	*	*	*	*	*	*	*	*
48.8	11	*7	*	*	*	*	*	*	*	*	*
48.81	10	*6	*	*	*	*	*	*	*	*	*
49	46	24	22	*	15	12	17	11	10	18	*7
49.0	*7	*	*	*	*	*	*	*	*	*	*
49.01	*5	*	*	*	*	*	*	*	*	*	*
49.4	19	10	10	—	*7	*5	*8	*	*	10	*
49.46	15	*7	*8	—	*6	*	*5	*	*	*9	*
50	109	54	55	*	22	35	48	28	19	43	19
50.1	87	43	44	*	14	29	42	22	16	35	15
50.11	63	32	31	*	11	21	29	15	10	27	11
50.12	24	11	13	*	*	*8	13	*7	*6	*7	*
50.2	*5	*	*	*	*	*	*	*	*	*	*
51	676	229	447	*	185	204	283	143	139	252	142
51.1	76	28	48	*	19	21	35	18	15	28	15
51.10	60	20	40	*	15	19	26	13	12	22	13
51.11	*6	*	*	—	*	*	*	*	*	*	*
51.14	*8	*	*	—	*	*	*6	*	*	*	*
51.2	428	137	292	*	130	138	158	94	86	159	89
51.22	117	50	67	*	19	40	58	26	29	37	26
51.23	310	87	223	*	111	97	99	67	58	121	63
51.3	*9	*	*5	*	*	*	*	*	*	*	*
51.36	*6	*	*	*	*	*	*	*	*	*	*
51.4	10	*	*6	—	*	*	*7	*	*	*	*
51.41	*7	*	*	—	*	*	*	*	*	*	*
51.5	*9	*6	*	*	*	*	*	*	*	*	*
51.51	*7	*	*	*	*	*	*	*	*	*	*
51.8	127	44	83	*	28	32	66	23	28	49	27
51.85	61	20	42	*	14	16	31	10	14	24	12
51.87	25	11	14	*	*	*6	15	*	*6	9	*6
51.88	34	11	23	*	*8	*8	18	*6	*7	13	*8
51.9	10	*	*5	*	*	*	*5	*	*	*	*
51.98	*6	*	*	*	*	*	*	*	*	*	*
52	33	13	20	*	10	10	13	*6	*7	10	11
52.1	*9	*	*	*	*	*	*	*	*	*	*
52.9	9	*	*7	*	*	*	*	*	*	*	*
52.93	*5	*	*	*	*	*	*	*	*	*	*
53	250	118	132	10	41	87	111	59	58	85	48
53.0	47	41	*6	*	*8	12	24	13	11	17	*6
53.00	*6	*	*	*	*	*	*	*	*	*	*
53.03	*9	*7	*	—	*	*	*	*	*	*	*
53.04	9	9	*	—	*	*	*	*	*	*	*
53.05	14	13	*	—	*	*	*9	*	*	*	*
53.1	13	10	*	*	*	*	*	*	*	*	*
53.2	10	*	*7	*	*	*	10	*	*	*	*
53.29	*7	*	*	*	*	—	*6	*	*	*	*
53.4	51	19	32	*	10	20	19	11	12	18	9
53.41	*8	*	*5	*	*	*	*	*	*	*	*

See footnotes and notes at end of table.

Table 45. Number of all-listed procedures for discharges from short-stay hospitals, by ICD-9-CM code, sex, age, and geographic region: United States, 1999—Con.

[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table: see Methods section]

ICD-9-CM code	Sex			Age				Region			
	Total	Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	North-east	Midwest	South	West
Number of procedures in thousands											
53.49	43	16	27	*	*8	16	17	9	9	16	*8
53.5	45	15	29	*	*7	19	18	9	11	16	*8
53.51	30	11	19	*	*	13	13	*6	*8	10	*6
53.59	15	*	10	*	*	*6	*	*	*	*6	*
53.6	68	23	45	*	9	28	31	16	14	23	16
53.61	54	19	36	*	*7	23	24	13	11	19	12
53.69	14	*	10	*	*	*5	*7	*	*	*	*
53.7	11	*	*6	*	*	*	*	*	*	*	*
54	754	232	522	27	274	227	226	156	167	266	165
54.0	16	*7	10	*	*	*7	*	*	*	*7	*
54.1	61	23	38	*	27	11	19	15	16	17	13
54.11	28	12	16	*	10	*5	10	*7	*7	*8	*
54.12	11	*	*7	*	*5	*	*	*	*	*	*
54.19	23	*8	15	*	12	*	*6	*5	*7	*7	*
54.2	127	36	91	*	58	31	35	26	29	42	30
54.21	73	17	56	*	45	15	12	14	15	25	19
54.23	19	*6	13	*	*	*6	*8	*	*	*	*
54.24	15	*	10	*	*5	*	*7	*	*	*6	*
54.25	18	*6	11	*	*	*6	*7	*	*	*	*6
54.3	13	*5	*8	*	*	*5	*	*	*	*	*
54.4	47	14	33	*	16	13	16	9	10	14	14
54.5	299	56	243	*6	127	83	83	61	60	119	59
54.51	38	*7	31	*	23	9	*5	*6	*6	15	10
54.59	262	49	213	*5	105	74	78	54	55	103	49
54.6	11	*6	*5	*	*5	*	*	*	*	*	*
54.7	*9	*	*	*	*	*	*	*	*	*	*
54.9	170	81	89	*9	31	72	58	35	42	56	38
54.91	114	53	61	*	20	47	44	26	27	37	24
54.93	11	*6	*	*	*	*	*	*	*	*	*
54.95	10	*6	*	*	*	*	*	*	*	*	*
54.98	30	14	16	*	*5	15	*7	*	*8	9	*8
55-59	966	461	505	40	225	299	402	242	207	351	166
55	136	63	73	9	36	43	47	36	26	47	27
55.0	31	12	19	*	*7	11	13	10	*	11	*6
55.03	21	*8	13	*	*	*6	11	*7	*	*7	*
55.2	23	12	11	*	*9	*8	*5	*7	*	*9	*
55.23	20	11	10	*	*7	*7	*	*6	*	*7	*
55.5	46	22	23	*	*9	11	23	11	12	15	*8
55.51	43	20	22	*	*7	10	23	10	11	14	*8
55.6	15	*8	*7	*	*7	*7	*	*	*	*6	*
55.69	15	*8	*7	*	*7	*7	*	*	*	*6	*
55.8	*7	*	*	*	*	*	*	*	*	*	*
55.87	*5	*	*	*	*	*	*	*	*	*	*
55.9	*5	*	*	*	*	*	*	*	*	*	*
56	112	69	43	11	37	35	28	23	25	50	14
56.0	45	31	15	*	19	17	9	9	11	22	*
56.3	30	19	11	*	12	12	*7	*5	*	18	*
56.31	28	19	10	*	12	10	*6	*	*	17	*
56.5	*6	*	*	—	*	*	*	*	*	*	*
56.51	*5	*	*	—	*	*	*	*	*	*	*
56.7	17	*8	*9	9	*	*	*	*	*	*	*
56.74	13	*5	*7	*8	*	*	*	*	*	*	*
57	415	226	189	14	69	101	231	123	86	134	72
57.0	24	18	*6	*	*	*	16	*9	*	*8	*
57.1	48	19	29	*	*7	16	20	13	*5	20	10
57.17	16	*6	10	*	*	*7	*7	*	*	*5	*
57.18	23	*7	15	*	*	*8	11	*	*	12	*
57.3	190	102	88	*5	34	51	100	53	48	64	26

See footnotes and notes at end of table.

Table 45. Number of all-listed procedures for discharges from short-stay hospitals, by ICD-9-CM code, sex, age, and geographic region: United States, 1999—Con.[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table: see Methods section]

ICD-9-CM code	Sex			Age				Region			
	Total	Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	North-east	Midwest	South	West
Number of procedures in thousands											
57.32	174	94	80	*	32	46	91	50	45	55	24
57.33	14	*7	*8	*	*	*	9	*	*	*7	*
57.4	49	37	12	*	*	*8	40	19	10	14	*6
57.49	49	37	12	*	*	*8	40	19	10	14	*6
57.7	*8	*7	*	—	*	*	*5	*	*	*	*
57.71	*8	*7	*	—	*	*	*5	*	*	*	*
57.8	22	*	17	*	10	*6	*	*	*5	9	*
57.81	10	*	*8	*	*6	*	*	*	*	*	*
57.89	*7	*	*6	*	*	*	*	*	*	*	*
57.9	65	34	31	*	13	13	39	22	*6	14	23
57.94	56	26	30	*	13	10	33	20	*	11	21
58	47	36	11	*	*9	10	25	13	11	15	*8
58.5	11	10	*	—	*	*	*7	*	*	*	*
58.6	20	15	*5	*	*	*	13	*	*6	*7	*
59	257	68	189	*	73	110	71	47	58	107	45
59.3	*7	—	*7	—	*	*	*	*	*	*	*
59.4	14	—	14	—	*	*7	*	*	*	*	*
59.5	52	*	52	—	14	31	*8	*	12	25	10
59.7	42	*	42	*	*7	22	13	*	*8	22	*7
59.79	38	*	38	*	*7	20	11	*	*7	20	*7
59.8	128	64	64	*	44	42	41	30	32	47	19
60-64	277	277	—	32	10	66	170	60	68	94	56
60	206	206	—	*	*	51	152	43	53	68	42
60.1	9	9	—	*	—	*	*8	*	*	*	*
60.11	*9	*9	—	*	—	*	*8	*	*	*	*
60.2	134	134	—	*	*	21	112	28	37	41	28
60.29	131	131	—	*	*	20	110	26	36	41	28
60.5	50	50	—	—	*	28	22	9	10	20	11
61	11	11	—	*	*	*	*	*	*	*	*
61.0	*5	*5	—	*	*	*	*	*	*	*	*
62	12	12	—	*	*	*	*	*	*	*	*
62.3	*6	*6	—	*	*	*	*	*	*	*	*
63	9	9	—	*	*	*	*	*	*	*	*
64	39	39	—	25	*	*6	*7	*8	*8	14	*8
64.0	23	23	—	21	*	*	*	*5	*6	*8	*
64.9	12	12	—	*	*	*	*5	*	*	*5	*
64.97	*7	*7	—	—	*	*	*	*	*	*	*
65-71	2,084	—	2,084	9	1,206	622	247	320	426	905	433
65	563	—	563	*	270	229	60	88	119	240	116
65.1	*7	—	*7	*	*	*	*	*	*	*	*
65.12	*6	—	*6	*	*	*	*	*	*	*	*
65.2	45	—	45	*	37	*	*	9	*7	16	13
65.25	*7	—	*7	*	*6	—	*	*	*	*	*
65.29	34	—	34	*	28	*	*	*6	*5	13	11
65.3	21	—	21	*	12	*	*	*	*	*9	*
65.39	18	—	18	*	11	*	*	*	*	*8	*
65.4	86	—	86	*	58	20	*7	15	18	36	17
65.41	15	—	15	*	11	*	*	*	*	*8	*
65.49	71	—	71	*	47	16	*7	13	16	28	14
65.5	11	—	11	—	*	*5	*	*	*	*	*
65.51	*7	—	*7	—	*	*	*	*	*	*	*
65.6	348	—	348	—	120	187	42	51	71	157	69
65.61	310	—	310	—	105	166	39	46	67	136	61
65.62	12	—	12	—	*	*7	*	*	*	*7	*
65.63	26	—	26	—	10	13	*	*	*	14	*6
65.8	27	—	27	—	19	*6	*	*	*6	11	*7

See footnotes and notes at end of table.

Table 45. Number of all-listed procedures for discharges from short-stay hospitals, by ICD-9-CM code, sex, age, and geographic region: United States, 1999—Con.

[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table: see Methods section]

ICD-9-CM code	Sex			Age				Region			
	Total	Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	North-east	Midwest	South	West
Number of procedures in thousands											
65.81	*5	—	*5	—	*	*	*	*	*	*	*
65.89	22	—	22	—	16	*	*	*	*5	*8	*5
65.9	10	—	10	*	9	*	—	*	*	*	*
65.91	*9	—	*9	*	*8	*	—	*	*	*	*
66	399	—	399	*	390	*7	*	47	78	193	80
66.0	10	—	10	*	10	*	—	*	*	*	*
66.02	*7	—	*7	*	*7	*	—	*	*	*	*
66.2	11	—	11	—	11	*	—	*	*	*5	*
66.29	*7	—	*7	—	*7	*	—	*	*	*	*
66.3	318	—	318	—	317	*	—	33	62	160	64
66.32	188	—	188	—	188	*	—	17	32	98	41
66.39	130	—	130	—	129	*	—	16	30	61	23
66.4	10	—	10	*	*8	*	*	*	*	*5	*
66.5	*8	—	*8	—	*7	*	*	*	*	*	*
66.51	*6	—	*6	—	*5	*	*	*	*	*	*
66.6	31	—	31	*	29	*	*	*7	*6	11	*7
66.61	*7	—	*7	*	*6	*	*	*	*	*	*
66.62	18	—	18	—	18	—	—	*	*	*8	*
67	20	—	20	*	11	*	*	*	*	10	*
67.1	*6	—	*6	*	*	*	*	*	*	*	*
67.5	*5	—	*5	—	*5	—	—	*	*	*	*
68	673	—	673	—	356	250	67	110	148	288	127
68.1	18	—	18	—	*9	*	*5	*6	*	*7	*
68.12	10	—	10	—	*	*	*	*	*	*	*
68.16	*7	—	*7	—	*	*	*	*	*	*	*
68.2	38	—	38	—	30	*	*	11	*5	16	*6
68.29	35	—	35	—	29	*	*	10	*	15	*6
68.3	16	—	16	—	10	*	*	*	*	*6	*
68.4	400	—	400	—	200	165	35	62	96	177	66
68.5	192	—	192	—	101	69	21	26	42	77	47
68.51	53	—	53	—	33	19	*	*8	11	25	9
68.59	138	—	138	—	68	51	19	18	31	52	38
68.6	*5	—	*5	—	*	*	*	*	*	*	*
69	116	—	116	*	96	12	*8	25	17	44	30
69.0	67	—	67	*	52	*8	*7	14	11	23	19
69.02	34	—	34	*	34	—	—	*7	*	14	*9
69.09	31	—	31	—	16	*8	*7	*7	*6	9	*9
69.2	*7	—	*7	—	*5	*	*	*	*	*5	*
69.22	*7	—	*7	—	*5	*	*	*	*	*5	*
69.5	23	—	23	*	21	*	*	*6	*	*7	*6
69.52	17	—	17	*	17	*	—	*5	*	*6	*
69.9	14	—	14	—	13	*	*	*	*	*7	*
69.96	9	—	9	—	9	*	—	*	*	*	*
70	279	—	279	*	70	109	98	38	54	116	70
70.2	*8	—	*8	*	*	*	*	*	*	*	*
70.3	12	—	12	*	*9	*	*	*	*	*7	*
70.32	*8	—	*8	—	*6	*	*	*	*	*5	*
70.5	150	—	150	—	29	68	53	23	32	63	33
70.50	83	—	83	—	14	38	30	11	16	39	18
70.51	32	—	32	—	*7	13	12	*5	*9	10	*7
70.52	36	—	36	—	*8	16	11	*6	*8	14	*8
70.7	44	—	44	*	*9	16	19	*6	*6	14	17
70.77	30	—	30	—	*	11	14	*	*	11	11
70.79	*9	—	*9	*	*	*	*	*	*	*	*5
70.9	55	—	55	—	15	20	20	*	11	23	16
70.92	54	—	54	—	15	20	19	*	11	23	16

See footnotes and notes at end of table.

Table 45. Number of all-listed procedures for discharges from short-stay hospitals, by ICD-9-CM code, sex, age, and geographic region: United States, 1999—Con.[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table: see Methods section]

ICD-9-CM code	Sex			Age				Region			
	Total	Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	North-east	Midwest	South	West
Number of procedures in thousands											
71	34	—	34	*	14	11	*8	*7	*8	13	*7
71.3	*8	—	*8	*	*6	*	*	*	*	*	*
71.7	13	—	13	*	*	*7	*	*	*	*	*
71.71	*7	—	*7	*	*	*	*	*	*	*	*
71.79	*6	—	*6	*	*	*	*	*	*	*	*
72-75	6,174	—	6,174	11	6,155	*8	—	1,162	1,445	2,068	1,498
72	389	—	389	*	388	*	—	47	87	155	99
72.0	26	—	26	—	26	—	—	*	*	14	*6
72.1	85	—	85	*	85	—	—	9	15	53	*7
72.2	*5	—	*5	—	*5	—	—	*	*	*	*
72.4	*7	—	*7	—	*7	—	—	*	*	*	*
72.5	*7	—	*7	*	*7	—	—	*	*	*	*
72.7	254	—	254	*	253	*	—	32	64	76	82
72.71	154	—	154	*	153	—	—	22	39	48	45
72.79	100	—	100	*	100	*	—	10	25	28	37
73	2,907	—	2,907	*6	2,899	*	—	491	692	1,020	704
73.0	809	—	809	*	806	*	—	126	232	220	230
73.01	131	—	131	*	131	—	—	12	38	48	33
73.09	677	—	677	*	675	*	—	114	195	172	197
73.1	*6	—	*6	—	*6	—	—	*	*	*	*
73.4	476	—	476	*	475	*	—	78	111	147	140
73.5	802	—	802	*	800	*	—	138	165	319	180
73.51	*8	—	*8	—	*8	—	—	*	*	*	*
73.59	794	—	794	*	792	*	—	137	162	317	178
73.6	806	—	806	*	803	*	—	149	179	327	151
73.9	*5	—	*5	—	*5	*	—	*	*	*	*
74	842	—	842	*	838	*	—	143	170	335	194
74.0	10	—	10	—	10	—	—	*	*	*6	*
74.1	803	—	803	*	799	*	—	141	163	316	183
74.9	26	—	26	—	26	—	—	*	*5	12	*8
74.99	26	—	26	—	26	—	—	*	*5	12	*8
75	2,037	—	2,037	*	2,031	*	—	482	496	558	501
75.1	13	—	13	—	13	—	—	*	*	*	*5
75.3	847	—	847	*	843	*	—	265	218	173	192
75.32	216	—	216	*	216	—	—	50	70	35	61
75.34	539	—	539	*	537	*	—	198	122	110	108
75.35	63	—	63	*	63	—	—	13	21	16	13
75.37	26	—	26	*	26	—	—	*	*	11	10
75.4	37	—	37	—	37	*	—	*	11	11	11
75.5	10	—	10	—	10	—	—	*	*	*	*
75.51	*7	—	*7	—	*7	—	—	*	*	*	*
75.6	1,106	—	1,106	*	1,103	*	—	207	256	357	287
75.61	36	—	36	—	36	—	—	*	9	15	*8
75.62	123	—	123	*	122	*	—	25	32	40	27
75.69	947	—	947	*	945	*	—	178	215	302	252
75.7	20	—	20	—	20	—	—	*	*	9	*
76-84	3,219	1,569	1,650	163	903	921	1,231	700	750	1,137	631
76	87	51	36	*6	60	16	*6	19	15	37	15
76.6	29	12	17	*	25	*	*	*	*	15	*7
76.62	*6	*	*	*	*6	*	—	*	*	*	*
76.64	*7	*	*6	*	*7	*	—	*	*	*	*
76.65	*8	*	*	*	*6	*	—	*	*	*	*
76.7	35	28	*7	*	24	*6	*	*8	*9	13	*
76.72	*6	*	*	*	*	*	*	*	*	*	*
76.76	11	10	*	*	10	*	*	*	*	*	*
76.79	*6	*	*	*	*	*	*	*	*	*	*
76.9	12	*7	*6	*	*7	*	*	*	*	*	*
76.91	*6	*	*	*	*	*	*	*	*	*	*

See footnotes and notes at end of table.

Table 45. Number of all-listed procedures for discharges from short-stay hospitals, by ICD-9-CM code, sex, age, and geographic region: United States, 1999—Con.

[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table: see Methods section]

ICD-9-CM code	Sex			Age				Region			
	Total	Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	North-east	Midwest	South	West
Number of procedures in thousands											
77	293	151	142	18	96	100	79	62	67	104	61
77.2	*7	*	*	*	*	*	*	*	*	*	*
77.3	17	*7	10	*6	*6	*	*	*6	*	*	*
77.37	*5	*	*	*	*	*	*	*	*	*	*
77.4	21	13	*8	*	*	*6	*9	*5	*6	*6	*
77.49	10	*7	*	*	*	*	*	*	*	*	*
77.5	12	*	9	*	*	*	*6	*	*	*5	*
77.6	44	25	19	*	12	13	16	12	10	15	*7
77.67	*8	*	*	*	*	*	*	*	*	*	*
77.68	*5	*	*	*	*	*	*	*	*	*	*
77.69	16	10	*6	*	*	*	*9	*5	*	*	*
77.7	135	71	64	*	54	56	20	26	30	52	27
77.70	*5	*	*	*	*	*	*	*	*	*	—
77.79	121	64	57	*	50	49	18	24	29	46	23
77.8	47	24	24	*	9	15	23	*8	11	17	13
77.81	13	*6	*7	—	*	*5	*6	*	*	*5	*
77.88	*7	*	*	*	*	*	*	*	*	*	*
77.89	16	*8	*8	*	*	*	*8	*	*	*	*5
78	205	103	102	27	70	57	51	40	46	77	42
78.0	39	22	17	*	16	14	*7	*8	10	13	*9
78.03	*6	*	*	*	*	*	*	*	*	*	*
78.05	11	*	*7	*	*	*	*	*	*	*	*
78.07	9	*7	*	*	*5	*	*	*	*	*	*
78.09	*7	*	*	*	*	*	*	*	*	*	*
78.1	35	19	16	*	14	*8	*8	*7	*8	16	*
78.13	11	*	*7	—	*	*	*	*	*	*6	*
78.15	*6	*	*	*	*	*	*	*	*	*	*
78.17	13	*7	*6	*	*7	*	*	*	*	*6	*
78.4	*8	*	*	*	*	*	*	*	*	*	*
78.5	54	27	27	10	15	13	16	10	10	19	14
78.55	29	11	18	*6	*	*6	13	*5	*6	10	*8
78.57	10	*6	*	*	*6	*	*	*	*	*	*
78.59	*7	*	*	*	*	*	*	*	*	*	*
78.6	56	25	31	*	19	16	17	12	13	21	11
78.65	19	*6	13	*	*	*	*9	*	*	*8	*
78.67	12	*6	*6	*	*	*	*	*	*	*	*
78.69	18	*8	10	*	*8	*7	*	*	*	*6	*
78.9	*6	*	*	*	*	*	*	*	*	*	*
78.99	*5	*	*	*	*	*	*	*	*	*	*
79	681	311	371	54	210	126	291	155	163	236	128
79.0	73	37	36	16	21	12	24	18	19	26	9
79.02	26	12	14	*	*7	*	11	*6	*7	10	*
79.05	10	*7	*	*7	*	*	*	*	*	*	*
79.06	25	12	13	*	*9	*5	*8	*7	*5	*9	*
79.1	78	35	43	14	19	12	33	18	17	24	19
79.11	16	*7	*9	10	*	*	*	*	*	*	*6
79.12	*9	*	*6	*	*	*	*	*	*	*	*
79.15	34	13	21	*	*5	*	24	*	11	11	*8
79.16	16	9	*6	*	*9	*	*	*5	*	*6	*
79.2	11	*6	*	*	*	*	*	*	*	*	*
79.3	431	179	252	13	132	83	203	97	103	151	80
79.31	31	14	17	*	10	*7	10	*7	9	*9	*6
79.32	35	17	17	*	14	*7	10	*6	*9	12	*8
79.33	*7	*5	*	—	*	*	*	*	*	*	*
79.34	*8	*7	*	*	*	*	*	*	*	*	*
79.35	184	55	129	*	16	16	150	49	41	66	27
79.36	140	62	78	*	67	40	30	27	33	51	28
79.37	12	*8	*	—	*8	*	*	*	*	*	*

See footnotes and notes at end of table.

Table 45. Number of all-listed procedures for discharges from short-stay hospitals, by ICD-9-CM code, sex, age, and geographic region: United States, 1999—Con.[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table: see Methods section]

ICD-9-CM code	Sex		Age				Region				
	Total	Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	North-east	Midwest	South	West
Number of procedures in thousands											
79.39	13	9	*	*	*8	*	*	*	*	*	*
79.6	44	29	15	*	23	11	*	*7	12	16	*9
79.62	*6	*	*	*	*	*	*	*	*	*	*
79.66	21	13	*8	*	12	*	*	*	*	*8	*
79.7	30	16	15	*	*6	*	19	*8	*8	*7	*8
79.75	21	11	10	*	*	*	14	*	*6	*	*5
79.8	*7	*	*	*	*	*	*	*	*	*	*
80	466	246	220	10	186	173	97	90	96	181	98
80.0	*9	*	*5	*	*	*	*7	*	*	*	*
80.06	*6	*	*	*	*	*	*	*	*	*	*
80.1	26	13	13	*	10	*	*8	*	*6	9	*6
80.16	13	*6	*7	*	*5	*	*6	*	*	*	*
80.2	30	18	12	*	12	9	*8	10	*	*6	11
80.21	11	*7	*	*	*	*	*	*	*	*	*
80.26	18	10	*8	*	9	*	*	*5	*	*	*6
80.4	13	*	*8	*	*	*	*	*	*	*	*5
80.46	*7	*	*5	*	*	*	*	*	*	*	*
80.5	313	167	146	*	132	134	46	54	69	135	56
80.50	10	*	*6	—	*	*	*	*	*	*7	*
80.51	303	163	140	*	128	129	45	52	69	128	55
80.6	20	*9	11	*	11	*	*	*	*6	*5	*
80.7	13	*5	*8	*	*	*	*5	*	*	*	*
80.76	10	*	*6	*	*	*	*	*	*	*	*
80.8	33	20	13	*	9	11	11	*6	*6	11	11
80.81	*6	*	*	*	*	*	*	*	*	*	*
80.86	16	*9	*7	*	*5	*	*6	*	*	*5	*
81	1,071	476	595	17	185	314	555	233	273	349	216
81.0	258	135	124	*8	94	105	51	45	59	103	51
81.02	105	57	47	*	38	52	15	16	22	48	18
81.03	11	*6	*5	*	*	*	*5	*	*	*	*
81.04	*6	*	*	*	*	*	*	*	*	*	*
81.05	14	*6	*8	*	*5	*	*	*	*	*	*
81.06	24	16	*9	*	12	11	*	*	*	10	*
81.07	16	10	*7	—	*7	*6	*	*	*6	*7	*
81.08	67	30	37	*	23	24	19	9	16	25	16
81.09	11	*	*6	*	*	*	*	*	*	*	*
81.1	18	*8	10	*	*5	*9	*	*5	*	*6	*
81.11	*7	*	*	*	*	*	*	*	*	*	*
81.2	*7	*	*	*	*	*	*	*	*	*	*
81.4	54	28	26	*	32	*8	10	14	*9	22	10
81.40	*5	*	*	*	*	*	*	*	*	*	*
81.45	23	12	11	*	19	*	—	*6	*5	*8	*
81.47	19	11	*8	*	*9	*	*6	*	*	*8	*
81.5	597	231	365	*	20	158	417	132	168	173	123
81.51	168	73	95	*	12	54	102	41	47	44	36
81.52	106	27	79	*	*	10	95	22	27	37	20
81.53	30	14	16	—	*	*8	20	*7	*9	*8	*7
81.54	267	105	162	—	*5	80	182	56	78	79	54
81.55	25	12	12	—	*	*6	18	*6	*8	*6	*6
81.8	51	26	25	*	13	16	21	12	*8	17	13
81.80	*5	*	*	—	*	*	*	*	*	*	*
81.81	*8	*	*6	—	*	*	*6	*	*	*	*
81.83	30	15	15	—	*7	12	11	*9	*	9	*8
81.9	84	43	41	*	16	16	48	22	24	25	12
81.91	50	31	19	*	12	11	25	15	14	13	*8

See footnotes and notes at end of table.

Table 45. Number of all-listed procedures for discharges from short-stay hospitals, by ICD-9-CM code, sex, age, and geographic region: United States, 1999—Con.

[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table: see Methods section]

ICD-9-CM code	Sex			Age				Region			
	Total	Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	North-east	Midwest	South	West
Number of procedures in thousands											
81.92	24	*8	16	—	*	*	20	*5	*8	*8	*
82	27	15	11	*	16	*5	*	*6	*7	11	*
82.0	*6	*	*	*	*	*	*	*	*	*	*
82.4	9	*7	*	*	*6	*	*	*	*	*	*
82.45	*5	*	*	*	*	*	*	*	*	*	*
83	223	119	104	27	63	74	60	55	46	76	47
83.0	21	14	*7	*	*7	*8	*5	*7	*7	*	*
83.09	12	*8	*	*	*	*	*	*	*	*	*
83.1	26	12	14	*8	9	*	*	*5	*7	*8	*5
83.13	*6	*	*	*	*	*	*	*	*	*	*
83.14	15	*9	*6	*	*7	*	*	*	*	*5	*
83.2	12	*6	*6	*	*	*	*5	*	*	*	*
83.21	12	*6	*6	*	*	*	*5	*	*	*	*
83.3	18	*8	10	*	*	*7	*7	*	*	9	*
83.39	17	*8	*9	*	*	*6	*7	*	*	*9	*
83.4	24	16	*8	*	*7	11	*	*8	*	*8	*
83.45	13	11	*	*	*	*6	*	*	*	*	*
83.6	63	34	28	*	18	25	16	12	12	25	14
83.63	36	15	21	*	*	19	13	*7	*6	15	*9
83.64	13	10	*	*	*8	*	*	*	*	*7	*
83.65	12	*8	*	*	*6	*	*	*	*	*	*
83.7	10	*	*6	*	*	*	*	*	*	*	*
83.75	*5	*	*	*	*	*	*	*	*	*	*
83.8	32	18	14	*8	9	*8	*7	*9	*7	10	*6
83.82	*9	*5	*	*	*	*	*	*	*	*	*
83.85	9	*5	*	*	*	*	*	*	*	*	*
83.88	*7	*	*	*	*	*	*	*	*	*	*
83.9	15	*5	10	*	*	*	*9	*	*	*5	*
84	165	96	68	*	18	55	90	41	36	66	22
84.0	10	*7	*	*	*	*	*	*	*	*	*
84.01	*7	*6	*	*	*	*	*	*	*	*	*
84.1	137	80	58	*	12	45	80	35	32	52	19
84.11	46	29	17	*	*5	18	22	11	11	16	*8
84.12	16	11	*6	—	*	*7	*7	*	*5	*6	*
84.15	33	19	14	—	*	11	20	10	*5	13	*
84.17	37	18	19	*	*	*7	28	9	9	14	*
84.3	14	*8	*6	*	*	*5	*6	*	*	*6	*
85-86	1,351	603	749	139	408	374	430	323	283	539	206
85	211	*	208	*	50	90	71	61	37	79	34
85.0	*6	*	*5	*	*	*	*	*	*	*	*
85.1	13	—	13	—	*	*	*7	*	*	*	*
85.12	*6	—	*6	—	*	*	*	*	*	*	*
85.2	35	*	35	*	*6	15	14	12	*	14	*
85.21	23	*	22	*	*	10	*9	*7	*	11	*
85.23	11	—	11	—	*	*	*	*	*	*	*
85.3	14	*	14	*	10	*	*	*7	*	*	*
85.32	13	*	13	*	10	*	*	*6	*	*	*
85.4	87	*	87	—	11	34	42	23	20	31	13
85.41	17	*	16	—	*	*7	*9	*	*	*7	*
85.43	66	*	65	—	*9	25	32	18	16	22	10
85.5	*8	*	*8	—	*	*	*	*	*	*	*
85.7	15	*	15	—	*7	*7	*	*	*	*8	*
85.8	11	*	11	—	*	*6	*	*	*	*	*
85.9	18	*	18	*	*	11	*	*	*	*7	*
85.95	12	—	12	—	*	*7	*	*	*	*5	*

See footnotes and notes at end of table.

Table 45. Number of all-listed procedures for discharges from short-stay hospitals, by ICD-9-CM code, sex, age, and geographic region: United States, 1999—Con.[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table: see Methods section]

ICD-9-CM code	Sex			Age				Region			
	Total	Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	North-east	Midwest	South	West
Number of procedures in thousands											
86	1,140	599	541	139	358	285	358	262	246	460	173
86.0	299	147	153	33	80	91	96	70	75	102	52
86.01	11	*6	*	*	*	*	*5	*	*	*	*
86.04	111	59	52	10	40	34	28	27	27	32	25
86.05	34	16	19	*	11	12	*7	*8	*6	15	*
86.06	15	*7	*8	*	*	*	*	*	*	*7	*
86.07	103	46	57	13	15	32	43	25	26	36	17
86.09	23	12	11	*	*7	*5	*9	*6	*7	*8	*
86.1	23	10	13	*	*8	*6	*7	*8	*	*7	*
86.11	22	9	13	*	*8	*5	*7	*7	*	*7	*
86.2	380	208	171	20	101	104	155	100	90	136	53
86.22	263	149	114	11	75	78	98	70	58	95	40
86.23	9	*	*	*	*	*	*	*	*	*	*
86.27	35	15	20	—	*	*6	28	13	9	*9	*
86.28	69	39	30	*7	23	17	23	14	20	26	*9
86.3	78	30	48	16	30	16	16	14	13	41	10
86.4	14	*7	*7	*	*	*	*	*	*	*5	*
86.5	155	98	57	12	69	26	48	30	39	48	38
86.59	155	98	57	12	69	26	48	30	39	48	38
86.6	118	71	47	48	35	16	18	21	11	79	*7
86.62	*7	*	*	*	*	*	*	*	*	*	*
86.69	96	57	38	43	27	12	13	15	*8	67	*5
86.7	32	19	14	*	13	*7	*8	*6	*6	17	*
86.70	*6	*	*	*	*	*	*	*	*	*5	*
86.74	17	10	*6	*	*8	*	*	*	*	*7	*
86.8	37	*8	29	*	16	13	*	*7	*	21	*
86.83	16	*	15	*	*7	*8	*	*	*	*8	*
86.84	10	*	*7	*	*6	*	*	*	*	*8	*
86.89	*6	*	*	*	*	*	*	*	*	*	*
87-99	12,964	6,269	6,695	878	2,463	3,384	6,239	3,758	2,584	3,907	2,715
87	1,274	592	682	57	224	307	686	290	322	359	303
87.0	429	201	228	22	72	92	243	169	60	94	106
87.03	421	197	224	20	70	90	240	168	59	91	102
87.2	48	23	25	*	13	15	20	*	15	20	9
87.21	19	*9	11	*	*	*7	*7	*	*5	*8	*
87.22	*7	*	*	*	*	*	*	*	*	*	*
87.23	*5	*	*	*	*	*	*	*	*	*	*
87.24	12	*	*8	—	*	*	*7	*	*	*7	*
87.4	460	227	233	20	55	102	282	44	181	133	102
87.41	104	53	51	*	14	27	61	41	20	23	20
87.44	281	144	137	16	31	60	175	*	135	70	74
87.49	72	29	43	*	10	15	45	*	26	39	*7
87.5	133	42	90	*	39	43	50	23	24	49	38
87.53	121	38	83	*	35	40	45	20	21	46	34
87.6	73	29	44	*7	10	14	43	18	11	18	26
87.61	19	*7	11	*	*	*	13	*5	*	*6	*
87.62	26	10	16	*	*	*	13	*7	*	*6	*8
87.63	11	*	*8	*	*	*	*5	*	*	*	*
87.64	11	*5	*5	*	*	*	*7	*	*	*	*6
87.69	*5	*	*	*	*	*	*	*	*	*	*
87.7	123	67	56	*6	34	39	44	31	30	42	20
87.73	27	16	11	*	*9	11	*7	*9	*	*8	*5
87.74	72	38	34	*	20	22	28	17	20	27	*7
87.76	*5	*	*	*	*	*	*	*	*	*	*
87.77	*5	*	*	*	*	*	*	*	*	*	*
87.79	*8	*	*	*	*	*	*	*	*	*	*

See footnotes and notes at end of table.

Table 45. Number of all-listed procedures for discharges from short-stay hospitals, by ICD-9-CM code, sex, age, and geographic region: United States, 1999—Con.

[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table: see Methods section]

ICD-9-CM code	Sex			Age				Region			
	Total	Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	North-east	Midwest	South	West
Number of procedures in thousands											
88	3,765	1,928	1,837	91	530	1,267	1,877	1,048	780	1,234	703
88.0	254	112	142	9	68	67	109	108	33	48	65
88.01	248	111	138	9	68	67	104	107	33	46	63
88.1	53	25	28	*	13	17	22	*	24	19	9
88.19	51	24	27	*	12	17	21	*	24	18	*8
88.2	37	21	16	*	*6	*9	22	*	19	*8	10
88.26	16	*	12	*	*	*	13	*	9	*	*
88.27	*6	*	*	*	*	*	*	*	*	*	*
88.28	*6	*	*	*	*	*	*	*	*	*	*
88.3	106	50	55	*6	23	26	51	45	11	33	16
88.38	97	47	50	*	23	23	47	44	10	28	15
88.4	394	194	200	*6	42	119	227	79	111	153	51
88.41	106	47	59	*	15	31	59	17	27	44	17
88.42	114	62	51	*	*7	38	65	22	34	44	14
88.43	18	*6	12	*	*5	*	*7	*	*	*5	*
88.44	*6	*	*	*	*	*	*	*	*	*	—
88.45	21	*8	13	*	*	*7	11	*	*7	*8	*
88.47	15	*6	*9	*	*	*	9	*	*	*6	*
88.48	91	52	39	*	*5	26	60	24	27	34	*7
88.49	18	*8	11	*	*	*7	*8	*	*	*6	*6
88.5	1,639	950	689	9	145	706	779	351	374	672	243
88.51	10	*6	*	*	*	*	*6	*	*	*	*
88.52	*6	*	*	*	*	*	*	*	*	*	*
88.53	645	364	281	*	58	283	302	142	139	278	87
88.54	19	11	*8	*	*	*6	9	*5	*6	*	*
88.55	20	12	*8	*	*	*7	11	*5	*	9	*
88.56	835	496	339	*	71	358	403	176	203	317	140
88.57	100	57	42	*	*9	47	44	19	19	52	10
88.6	25	12	13	*	*8	*6	11	*8	*6	*8	*
88.66	10	*	*6	—	*	*	*	*	*	*	*
88.67	*8	*5	*	*	*	*	*	*	*	*	*
88.7	1,022	451	571	41	180	250	551	375	154	237	256
88.71	96	46	50	*	*5	23	65	30	13	21	33
88.72	547	260	287	17	55	143	333	211	76	144	116
88.73	*8	*	*	*	*	*	*	*	*	*	*
88.74	20	*7	13	*	*6	*	*8	*6	*6	*5	*
88.75	58	26	32	*7	12	11	27	21	10	11	16
88.76	135	55	81	9	41	33	52	50	15	21	49
88.77	95	42	53	*	14	29	51	34	24	15	21
88.78	32	—	32	*	32	*	—	*9	*5	*9	*9
88.79	31	12	19	*	13	*5	11	12	*	10	*6
88.9	233	110	123	17	45	67	105	80	48	56	50
88.91	133	65	67	11	27	34	60	50	29	26	27
88.93	44	22	22	*	*9	13	20	14	10	9	11
88.94	14	*6	*9	*	*	*5	*6	*8	*	*	*
88.97	32	12	19	*	*6	11	13	*7	*	13	*8
89	1,089	518	571	51	140	295	602	306	237	275	271
89.0	36	16	20	*	*7	11	18	*5	*	*	28
89.06	15	*7	*8	—	*	*6	*8	—	*	*	13
89.07	*6	*	*	—	*	*	*	—	—	—	*6
89.08	*6	*	*	*	*	*	*	*	*	—	—
89.09	*8	*	*	—	*	*	*	—	*	*	*8
89.1	110	50	60	27	23	19	42	50	*6	35	19
89.14	81	37	45	14	12	15	39	34	*	27	16
89.17	*6	*	*	*	*	*	*	*	*	*	*
89.19	18	*8	10	*8	*8	*	*	*9	*	*6	*
89.2	22	10	12	*	*	*	14	*7	*6	*6	*
89.22	12	*7	*5	—	*	*	10	*	*	*	*

See footnotes and notes at end of table.

Table 45. Number of all-listed procedures for discharges from short-stay hospitals, by ICD-9-CM code, sex, age, and geographic region: United States, 1999—Con.[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table: see Methods section]

ICD-9-CM code	Sex		Age				Region				
	Total	Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	North-east	Midwest	South	West
Number of procedures in thousands											
89.3	56	23	33	11	10	13	22	13	12	12	19
89.37	24	11	14	*	*6	*8	*9	*7	*	*	12
89.38	11	*	*6	*	*	*	*	*	*	*	*
89.39	18	*6	12	*	*	*	10	*	*	*8	*
89.4	150	73	77	*	17	58	74	56	20	38	36
89.41	42	24	17	*	*	21	16	19	*	*	16
89.44	106	48	58	*	12	37	57	36	17	33	20
89.5	537	258	278	*8	51	143	334	123	152	131	130
89.50	19	*8	11	*	*	*	13	10	*	*7	*
89.51	*5	*	*	*	*	*	*	*	*	*	*
89.52	340	165	174	*	31	90	215	22	139	92	86
89.54	162	76	86	*	17	45	97	85	9	29	38
89.59	10	*6	*	*	*	*	*6	*	*	*	*
89.6	178	88	90	*	27	48	99	52	38	52	36
89.61	10	*	*6	*	*	*	*7	*	*	*	*
89.62	17	*7	10	*	*	*	*7	*6	*	*6	*
89.64	60	36	24	*	*5	18	37	14	17	18	10
89.65	80	34	46	*	15	19	44	27	12	24	17
89.68	*8	*	*	*	*	*	*	*	*	*	*
90	65	23	42	*5	15	18	27	*	*8	54	*
90.5	48	18	30	*	12	15	16	*	*	44	*
90.52	*5	*	*	*	*	*	*	*	*	*	*
90.59	40	13	27	*	11	14	13	*	—	40	*
90.8	10	*	*6	—	*	*	*8	*	*6	*	*
90.81	10	*	*6	—	*	*	*7	*	*6	*	*
91	21	*7	15	*	*8	*6	*6	*	*	18	*
91.3	13	*	10	*	*	*	*	*	*	12	*
91.39	10	*	*8	*	*	*	*	—	—	10	—
92	322	147	175	10	44	104	164	121	49	80	71
92.0	109	49	60	*	14	37	57	43	16	25	25
92.02	16	*	11	*	*	*	*9	*5	*	*5	*
92.03	*7	*	*	*	*	*	*	*	*	*	*
92.04	14	*5	*9	*	*	*	*8	*5	*	*	*
92.05	70	34	36	—	*6	27	36	30	9	15	16
92.1	116	51	65	*	17	31	63	44	16	23	32
92.11	*6	*	*	*	*	*	*	*	*	*	*
92.14	38	18	20	*	*	*8	24	17	*	10	9
92.15	58	24	34	*	10	17	31	22	*8	10	19
92.19	*7	*	*	*	*	*	*	*	*	*	*
92.2	90	43	47	*	12	32	42	31	16	30	13
92.24	20	10	10	*	*	*6	11	11	*	*	*
92.27	18	*8	9	*	*	*6	*8	*7	*	*6	*
92.29	46	22	24	*	*6	16	22	11	11	17	*7
92.3	*7	*	*	*	*	*	*	*	*	*	*
93	1,521	683	838	201	185	306	828	434	233	441	414
93.0	46	21	25	*	*7	9	29	21	*6	*5	14
93.01	40	17	23	*	*	*8	26	18	*	*	13
93.08	*5	*	*	*	*	*	*	*	*	*	*
93.1	32	12	20	*	*	*8	21	24	*	*	*
93.11	*8	*	*	*	*	*	*5	*7	*	*	*
93.12	12	*	*7	*	*	*	*8	11	*	*	—
93.2	45	19	26	*	*6	13	26	25	*9	*	*7
93.22	33	13	19	*	*	*8	21	20	*7	*	*
93.3	336	139	197	*7	24	65	241	78	57	126	76
93.32	*5	*	*	*	*	*	*	*	*	*	*
93.38	12	*	*8	*	*	*	*9	*	*	*	*7
93.39	315	130	185	*7	20	60	229	73	55	120	67
93.4	16	*7	9	*	*	*	*8	*	*5	*	*
93.46	*8	*	*6	*	*	*	*6	*	*	*	*

See footnotes and notes at end of table.

Table 45. Number of all-listed procedures for discharges from short-stay hospitals, by ICD-9-CM code, sex, age, and geographic region: United States, 1999—Con.

[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table: see Methods section]

ICD-9-CM code	Sex			Age				Region			
	Total	Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	North-east	Midwest	South	West
Number of procedures in thousands											
93.5	88	45	43	14	23	19	32	22	18	26	22
93.53	19	9	10	*6	*6	*	*6	*6	*	*	*
93.54	20	*9	11	*	*5	*	10	*	*	*	*6
93.57	15	11	*	*	*	*	*	*	*	*6	*
93.59	24	10	14	*	*	*8	10	*5	*	*8	*7
93.7	36	18	18	*	*	10	23	*6	10	14	*7
93.75	35	18	17	*	*	9	23	*5	10	13	*6
93.8	369	155	215	11	48	74	235	94	62	151	62
93.81	27	12	15	*	*6	*8	11	*6	*5	12	*
93.83	232	98	133	*5	27	49	151	72	46	85	28
93.89	109	43	66	*	14	17	74	16	11	53	30
93.9	550	267	283	163	68	107	213	159	65	105	221
93.90	56	28	28	*6	*7	16	28	18	16	11	11
93.94	219	110	108	93	27	34	65	86	16	33	83
93.96	220	104	116	55	27	44	94	44	26	37	114
93.99	46	21	24	*9	*6	11	20	9	*	23	11
94	718	438	280	32	449	186	51	363	166	146	43
94.0	10	*7	*	*	*6	*	*	*	—	*8	*
94.08	*6	*	*	*	*	*	*	*	—	*	*
94.1	*7	*	*	*	*	*	*	*	—	*	*
94.2	154	80	74	*7	90	41	17	109	22	18	*6
94.23	27	16	11	*	15	*8	*	25	—	*	*
94.25	95	50	45	*	60	23	*8	74	9	*6	*
94.27	28	12	16	—	11	10	*7	*6	11	10	*
94.3	108	56	52	9	67	25	*7	61	26	20	*
94.38	39	19	19	*	26	10	*	39	—	*	—
94.39	63	32	31	*6	39	14	*	16	26	20	*
94.4	126	67	59	13	80	24	*9	66	30	29	*
94.42	14	*7	*7	*	*8	*	*	*6	*5	*	—
94.44	99	51	48	*9	62	20	*8	48	25	24	*
94.46	*6	*5	*	—	*	*	*	*6	—	*	—
94.6	311	223	88	*	202	93	16	122	89	67	33
94.61	13	10	*	—	10	*	*	*7	*	*	*
94.62	134	106	28	—	69	53	11	47	43	28	16
94.63	32	21	11	—	19	10	*	*5	16	*5	*6
94.64	*7	*	*	*	*6	*	*	*	*	*	*
94.65	51	31	20	—	39	12	*	27	*8	12	*
94.66	10	*5	*	—	*8	*	*	*	*	*6	*
94.67	*9	*6	*	*	*7	*	—	*	*6	*	—
94.68	44	31	13	—	34	10	—	27	*	10	*
94.69	12	9	*	—	10	*	*	*	*	*	*
95	*7	*	*	*5	*	*	*	*	*	*	*
95.4	*5	*	*	*	*	*	*	*	*	*	*
96	1,375	660	715	126	284	270	694	359	291	444	281
96.0	529	270	259	45	82	108	294	130	118	172	109
96.04	445	227	219	40	69	92	244	109	95	151	91
96.05	*8	*6	*	*	*	*	*	*	*	*	*
96.07	70	36	34	*	11	13	42	18	20	17	15
96.2	*8	*	*	*	*	*	*	*	*	*	*
96.3	28	12	16	*	10	*6	*8	9	*6	*7	*6
96.33	17	*6	11	*	*9	*	*	*6	*	*	*
96.38	*7	*	*	*	*	*	*	*	*	*	*
96.4	86	*	81	*	81	*	*	9	18	29	29
96.49	82	*	80	*	80	*	*	*9	17	27	28
96.5	46	26	20	*	12	15	15	10	10	18	*9
96.56	25	13	13	*	*	*8	12	*6	*5	*8	*6
96.59	19	12	*6	*	*8	*5	*	*	*	10	*

See footnotes and notes at end of table.

Table 45. Number of all-listed procedures for discharges from short-stay hospitals, by ICD-9-CM code, sex, age, and geographic region: United States, 1999—Con.[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table: see Methods section]

ICD-9-CM code	Sex			Age				Region			
	Total	Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	North-east	Midwest	South	West
Number of procedures in thousands											
96.6	108	50	58	*9	11	14	75	54	18	24	12
96.7	567	293	274	61	86	124	296	142	120	190	115
96.70	21	14	*7	*	*	*	13	*6	*	*8	*
96.71	352	180	172	34	56	81	180	88	81	111	71
96.72	194	99	95	25	27	39	103	48	36	70	39
97	97	46	51	12	19	18	49	25	18	41	13
97.0	33	14	19	*	*	*	25	*9	*6	15	*
97.02	23	9	14	*	*	*	17	*7	*5	*9	*
97.2	*9	*	*5	*	*	*	*	*	*	*	*
97.23	*7	*	*	*	*	*	*	*	*	*	*
97.4	13	*8	*5	*	*	*	*	*	*	*5	*
97.49	*9	*5	*	*	*	*	*	*	*	*	*
97.5	*8	*	*	*	*	*	*	*	*	*	*
97.6	12	*5	*7	*	*	*	*6	*	*	*5	*
97.62	*9	*	*6	—	*	*	*	*	*	*	*
97.8	13	*8	*5	*	*	*	*	*	*	*6	*
97.89	*7	*	*	*	*	*	*	*	*	*	*
98	32	17	15	*6	*9	*7	10	*9	*	12	*7
98.0	*9	*	*	*	*	*	*	*	*	*	*
98.02	*7	*	*	*	*	*	*	*	*	*	*
98.1	*6	*	*	*	*	*	*	*	*	*	*
98.5	13	*6	*6	*	*	*	*5	*	*	*6	*
98.51	11	*6	*5	*	*	*	*	*	*	*5	*
99	2,678	1,207	1,471	281	555	598	1,244	799	475	803	602
99.0	1,023	440	583	48	131	223	621	223	206	365	228
99.00	12	*6	*6	*	*	*	*7	*	*	*	*
99.02	50	22	29	*	*	17	30	9	12	11	18
99.03	37	16	21	*	*	*6	24	*	*9	19	*6
99.04	807	335	472	32	104	167	504	185	156	293	172
99.05	50	27	23	9	11	14	17	12	*9	14	16
99.07	58	28	31	*	*6	14	35	10	14	21	14
99.1	386	157	229	35	117	86	148	148	58	95	84
99.10	66	41	25	*	*6	25	34	20	10	17	18
99.11	68	—	68	*	68	*	—	14	11	25	18
99.14	*6	*	*	*	*	*	*	*	*	*	*
99.15	99	47	52	23	13	19	44	36	23	22	19
99.17	40	16	23	*	10	13	15	21	*	*5	*9
99.18	47	19	28	*5	10	10	22	26	*	16	*
99.19	60	31	29	*	10	17	32	27	9	*8	15
99.2	995	470	524	145	265	238	347	364	163	251	217
99.20	45	32	14	—	*	22	20	*9	12	16	*9
99.21	347	154	193	57	111	68	111	147	40	81	79
99.23	85	35	50	*8	16	23	39	31	18	23	13
99.24	*7	*	*6	*	*6	*	*	*	*	*	*
99.25	211	116	95	42	39	62	68	67	41	63	39
99.29	293	129	164	37	88	61	107	108	49	62	75
99.3	*6	*	*	—	*	*	*	*	*	*	*
99.4	16	*	16	—	16	*	*	*	*	*6	*5
99.47	10	—	10	—	10	—	—	*	*	*	*
99.48	*6	—	*6	—	*6	—	—	*	—	*	*
99.5	*6	*	*	*	*	*	*	*	*	*	*
99.6	161	90	71	*	10	41	106	40	37	45	39
99.60	52	26	25	*	*	12	34	10	12	12	17
99.61	36	20	16	—	*	10	25	10	*7	12	*8
99.62	64	39	24	*	*	17	42	17	16	19	12
99.69	*8	*	*	*	*	*	*5	*	*	*	*

See footnotes and notes at end of table.

Table 45. Number of all-listed procedures for discharges from short-stay hospitals, by ICD-9-CM code, sex, age, and geographic region: United States, 1999—Con.

[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table: see Methods section]

ICD-9-CM code	Sex			Age				Region			
	Total	Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	North-east	Midwest	South	West
Number of procedures in thousands											
99.7	12	*	*8	*	*	*	*	*	*	*	*
99.71	*7	*	*	*	*	*	*	*	*	*	*
99.8	51	28	22	44	*	*	*6	*8	*	18	21
99.83	41	22	19	41	*	*	—	*7	*	10	20
99.84	*6	*	*	*	*	*	*5	*	*	*5	—
99.9	24	12	11	*	*6	*	11	*	*	15	*
99.99	23	12	11	*	*6	*	11	*	*	15	*

* Figure does not meet standard of reliability or precision.

— Quantity zero.

NOTE: Estimates of 5,000-9,000 are to be used with caution; see Methods section.

Appendix

Technical Notes

Abstract Form

Figure I shows the medical abstract form used for the 1999 NHDS. The hospital number, medical record number, date of birth, and patient ZIP Code are confidential information and are not released to the public.

Population Estimates

Population estimates used to compute rates are shown in table I. These estimates are for the U.S. civilian population, including institutionalized persons, as of July 1, 1999. They were provided by the U.S. Bureau of the Census and are consistent with population estimates in Population Division Release US-99-SIS-7 (*U.S. Population Estimates by Age, Sex, Race, and Hispanic Origin: 1999*). These estimates have been adjusted for the net underenumeration using the 1990 National Population Adjustment Matrix. This is a change from population data provided in reports of 1990–96 NHDS data, which were not adjusted for the underenumeration.

Reliability of Estimates

The statistics presented in this report are based on sample surveys, and therefore may differ from the figures that would be obtained if a complete census had been taken. The standard error is primarily a measure of sampling variability that occurs by chance because only a sample rather than an entire universe is surveyed. Standard errors for the estimates from NHDS were calculated with SUDAAN software, which takes into account the complex sample designs. A description of the software and the approach it uses has been published (19).

The chances are about 32 in 100 that an estimate from the sample would differ from a complete census by more than the standard error. The chances are 5 in 100 that the difference would be more than twice the standard error, and

Table I. Civilian population by selected characteristics: United States, 1999

	Both sexes	Male	Female
	Population in thousands		
All ages	275,538	134,644	140,894
Northeast	52,212	25,285	26,928
Midwest	63,568	31,022	32,546
South	97,622	47,313	50,310
West	62,135	31,025	31,111
Under 15 years	60,259	30,829	29,430
Under 1 year	3,967	2,027	1,940
1 to 4 years	15,651	8,000	7,651
5 to 14 years	40,641	20,802	19,839
Northeast	10,720	5,486	5,234
Midwest	13,750	7,038	6,711
South	21,438	10,956	10,481
West	14,351	7,348	7,003
15 to 44 years	122,029	60,964	61,065
15 to 24 years	38,452	19,546	18,907
25 to 34 years	38,540	19,070	19,470
35 to 44 years	45,037	22,347	22,689
Northeast	22,829	11,344	11,486
Midwest	28,002	13,995	14,007
South	42,995	21,214	21,781
West	28,202	14,411	13,791
45 to 64 years	59,011	28,613	30,398
45 to 54 years	35,812	17,527	18,285
55 to 64 years	23,198	11,086	12,112
Northeast	11,460	5,509	5,950
Midwest	13,703	6,666	7,036
South	21,114	10,151	10,963
West	12,735	6,287	6,448
65 years and over	34,239	14,238	20,001
65 to 74 years	18,044	8,148	9,896
75 to 84 years	12,047	4,853	7,194
85 years and over	4,149	1,237	2,911
Northeast	7,203	2,945	4,257
Midwest	8,113	3,322	4,791
South	12,076	4,992	7,084
West	6,847	2,979	3,869

NOTE: Population estimates have been adjusted for net underenumeration using the 1990 National Population Adjustment Matrix. Estimates are consistent with those in US-99-SIS-7 (*U.S. Population Estimates by Age, Sex, Race and Hispanic Origin: 1999*).

about 1 in 100 that the difference would be more than 2.5 times as large as the standard error. The relative standard error (RSE) expresses the standard error as a percent of the estimate and can be multiplied by the estimate to obtain the standard error.

To derive approximate standard errors for the estimates in this report, first calculate the approximate RSE of the estimate using the parameters shown in table II. Then multiply the approximate RSE by the estimate. The approximate RSE of an estimate X may be estimated from the formula:

$$\text{RSE}(X) = \sqrt{a + b/X}$$

where a and b are defined in table II.

When multiplied by 100, $\text{RSE}(X)$ is expressed as a percent of X .

The approximate RSE for a percent, $100p$, in which the numerator is a subclass of the denominator, may be calculated directly using the formula:

$$\text{RSE}(p) = \sqrt{b(1-p) / (pX)}$$

where $100p$ is the percent of interest, X is the denominator of the percent, and b is the parameter b in the formula for approximating the $\text{RSE}(X)$. The values for b are given in table II. When multiplied by 100, $\text{RSE}(p)$ is expressed as a percent of p . This approximation is valid if the RSE of the denominator is less than 5 percent or the RSE's of the

Table II. Estimated parameters for approximate relative standard error equations for National Hospital Discharge Survey statistics by selected characteristics, 1999

Characteristic	First-listed diagnoses		Days of care		All-listed diagnoses		All-listed procedures	
	<i>a</i>	<i>b</i>	<i>a</i>	<i>b</i>	<i>a</i>	<i>b</i>	<i>a</i>	<i>b</i>
Total	0.001560	352.57572	0.002604	1000.94965	0.003105	412.75822	0.003187	357.68835
Sex								
Male	0.001633	359.91941	0.002981	1317.06683	0.004093	363.50935	0.003464	340.74739
Female	0.001578	334.61786	0.002666	1087.27514	0.003769	326.47722	0.003684	301.79493
Age group								
Under 15 years	0.016494	223.07202	0.025039	652.94193	0.018165	238.34924	0.025237	253.97288
15-44 years	0.001763	325.12768	0.003308	1033.44779	0.001589	343.08995	0.002402	305.64621
45-64 years	0.002044	377.51124	0.003097	1341.59977	0.002337	337.00450	0.004366	273.42803
65 years and over	0.002189	338.47209	0.002729	2033.88218	0.002026	346.43798	0.003618	341.62920
Geographic region								
Northeast	0.005121	193.54615	0.010383	269.39682	0.007518	197.33709	0.007950	230.43315
Midwest	0.009636	233.35485	0.013644	402.06132	0.011474	211.10520	0.012990	172.54412
South	0.003298	353.03917	0.005486	920.65327	0.003989	367.90190	0.008102	286.55150
West	0.004267	367.67568	0.008367	937.57802	0.004568	435.54443	0.004604	393.92609
Principal expected source of payment								
Medicare	0.002462	359.77067	0.002871	2258.16638	0.002217	378.71984	0.004041	346.83166
Medicaid	0.005774	284.19933	0.009001	992.65007	0.004929	286.51731	0.006838	249.47826
Worker's compensation and other government payments	0.009213	295.76741	0.018237	1143.72306	0.010336	366.61738	0.010730	305.51886
HMO/PPO	0.004928	253.10321	0.008001	652.40977	0.004700	302.50753	0.008371	244.03377
Blue Cross/Blue Shield and other private insurance	0.004259	294.39929	0.007039	601.48207	0.007244	273.28557	0.007154	273.35592
Self-pay	0.004763	256.70174	0.008699	982.59488	0.005202	255.34030	0.007336	228.72794
No charge and other	0.037061	130.96480	0.044592	525.47202	0.031448	148.70879	0.043054	130.39311

NOTES: The relative standard error (RSE) can be determined from the equation $RSE(x) = \sqrt{a + b/x}$. HMO/PPO is defined as Health Maintenance Organization and Preferred Provider Organization.

numerator and denominator are both less than 10 percent (20,21).

The approximate RSE of a population-based rate is equivalent to the approximate RSE of the numerator (obtained by the method described above using figures from table II), if the denominator is an estimate produced by the U.S. Bureau of the Census for the total population or one or more of the age-sex groups of the total population.

The approximate RSE of a ratio (X/Y) in which the numerator (X) and the denominator (Y) are both estimated from the same survey, but the numerator is not a subclass of the denominator, is calculated using the formula:

$$RSE(X/Y) = \sqrt{[RSE(X)]^2 + [RSE(Y)]^2}$$

When multiplied by 100, the RSE(X/Y) is expressed as a percent of the ratio X/Y. This approximation is valid if the RSE of the denominator is less than 5 percent or the RSE's of both the

numerator and denominator are both less than 10 percent (20,21).

Definitions of Selected Terms

Age—Patient's age at the birthday prior to admission to the hospital.

Average length of stay—The mean length of stay for discharges. It is calculated by dividing the number of days of care by the number of discharges.

Bed size of hospital—The number of beds, cribs, and pediatric bassinets regularly maintained (set up and staffed for use) for patients, not including bassinets for newborn infants. In this report, the classification of hospitals by bed size is based on the number of beds at or near midyear as reported by the hospital.

Days of care—The total number of days a patient spent in the hospital. A stay of less than 1 day (patient admission and discharge on the same day) is counted as one day. For patients

admitted and discharged on different days, the number of days of care is computed by counting all the days from (and including) the day of admission to (but not including) the day of discharge.

Death rate—Calculated by dividing the number of deaths for a category by the total number of discharges in that category and multiplying by 100. Discharges whose status is unknown (0.6 percent) were assumed to be discharged alive.

Diagnosis—A disease or injury (or other factor that influences a patient's health status or causes contact with health services) that is listed on the medical record of the patient. The diagnosis established after study to be chiefly responsible for the admission of the patient to the hospital is the principal diagnosis. The first-listed diagnosis is the one specified as the principal diagnosis on the face sheet or discharge summary of the medical record. However, if the principal diagnosis is not specified, the first one

listed on the face sheet or discharge summary of the medical record is used. The number of first-listed diagnoses is the same as the number of discharges. Data for all-listed diagnoses include first-listed diagnoses and other diagnoses appearing on the medical record. A minimum of one and a maximum of seven diagnoses are coded per discharge in the NHDS. The HIV data (table 22) are for discharges with an HIV diagnosis listed in any position among the coded diagnoses.

Discharge—A completed inpatient hospitalization. A hospitalization may be completed by death or by releasing the patient to the customary place of residence, a nursing home, another hospital, or other locations.

Hospital—Hospitals with an average length of stay for all inpatients of less than 30 days (short stay) or those whose specialty is general (medical or surgical) or children's general. Federal hospitals, hospital units of institutions, and hospitals with less than six beds staffed for patient use are excluded.

Newborn infant—Inpatients admitted to a hospital by birth. Sick newborn infants are those that had at least one of the ICD-9-CM (1) diagnostic codes 001-999.

Ownership of hospital—The type of ownership is determined by the organization that controls and operates the hospital. Hospitals are grouped as follows:

Proprietary—Hospitals operated by individuals, partnerships, or corporations for profit.

Government—Hospitals operated by State or local governments.

Nonprofit—Hospitals operated by a nonprofit organization or church.

Principal expected source of payment—The primary expected method of payment for the hospitalization. Types of payment are as follows:

Medicare—The Centers for Medicare and Medicaid Services' (formerly the Health Care Financing Administration) health insurance program for the aged and disabled.

Medicaid—A jointly funded Federal-State health insurance

program providing medical care to those unable to afford it.

Worker's compensation—A State or municipal disability insurance or industrial accident insurance. Does not include company health insurance plans or direct payment by an employer.

Other government—Payment that cannot be classified in one of the other three government categories (Medicare, Medicaid, or worker's compensation). This includes payments under Title V Program, payments made under the State funded Maternal and Child Health Program, State funded Crippled Children's Program, payments made under TRICARE/CHAMPUS, casualty insurance paid by the State, vocational rehabilitation, Federal or State research grant (medical research), or legal hold (prisoner in medical detention).

HMO/PPO—Any Health Maintenance Organization or Preferred Provider Organization sponsored by consumers, communities, physicians, or hospitals.

Blue Cross/Blue Shield and other private—A Blue Cross insurance plan, Blue Cross Association, or Blue Shield plans.

Other private—A private insurance plan not specified as an HMO/PPO. This includes Blue Cross/Blue Shield, medical coverage provided by life insurance companies, casualty insurance companies, health insurance companies, and independent plans such as employer/union sponsored plans and/or self-funded plans (partial or total).

Self-pay—The majority of the total costs is expected by the patient, spouse, family, or next of kin.

No charge—Patients admitted with the understanding that payment would not be expected because the medical services are free. This includes hospital-sponsored special research or "teaching" patients.

Other—Includes any nonprofit source of payment, such as church welfare, United Way (United Appeal), or Shriner's Crippled Children Services.

Procedure—Surgical procedures (such as tonsillectomy), diagnostic procedures (such as cystoscopy), and therapeutic treatments (such as injection or infusion of cancer chemotherapeutic substance) reported on a patient's medical record. Data are for all-listed procedures, which include all occurrences of the procedure coded, regardless of their order in the medical record. Up to four procedures are coded per discharge in the NHDS. The procedures classified as nonsurgical in this report are listed in table III by ICD-9-CM code numbers. All other procedures were considered surgical. This classification was begun in 1979 for the presentation of NHDS data. The distinction between surgical and nonsurgical procedures has become less meaningful in the last two decades with the development of minimally invasive and noninvasive procedures. The procedures classified as nonsurgical may have as much operative or anesthetic risk and require highly trained personnel or special equipment similar to those classified as surgical.

Region—Hospitals are classified by location in one of the four geographic regions of the United States that correspond to those used by the U.S. Bureau of the Census.

<i>Region</i>	<i>States included</i>
Northeast	Maine, New Hampshire, Vermont, Massachusetts, Rhode Island, Connecticut, New York, New Jersey, and Pennsylvania
Midwest	Michigan, Ohio, Illinois, Indiana, Wisconsin, Minnesota, Iowa, Missouri, North Dakota, South Dakota, Nebraska, and Kansas
South	Delaware, Maryland, District of Columbia, Virginia, West Virginia, North Carolina, South Carolina, Georgia,

Table III. Code numbers for procedures considered nonsurgical in the National Hospital Discharge Survey by ICD-9-CM category, 1999

[Procedure categories and code numbers are based on the *International Classification of Diseases, 9th Revision, Clinical Modification (ICD-9-CM)*]

Procedure category and ICD-9-CM code	Nonsurgical codes
Operations on the nervous system 01-05	01.18-01.19,03.31,03.39,04.19,05.19
Operations on the endocrine system 06-07	06.19,07.19
Operations on the eye 08-16	08.19,09.19,09.41-09.49,10.29,11.29,12.29,14.19,15.09,16.21,16.29
Operations on the ear 18-20	18.01,18.11,18.19,20.31,20.39
Operations on the nose, mouth, and pharynx 21-29	21.00-21.02,21.21,21.29,22.19,24.19,25.09,26.19,27.29,28.19,29.11,29.19
Operations on the respiratory system 30-34	31.41-31.42,31.48-31.49,33.21-33.23,33.29,34.21-34.22,34.28-34.29
Operations on the cardiovascular system 35-39	37.26-37.27,37.29,38.29,39.95
Operations on the hemic and lymphatic system 40-41	40.19,41.38-41.39
Operations on the digestive system 42-54	42.22-42.23,42.29,44.11-44.13,44.19,45.11-45.13,45.19,45.21-45.24,45.28-45.29,48.21-48.23,48.29,49.21,49.29,50.19,51.10-51.11,51.19,52.19,54.21,54.29
Operations on the urinary system 55-59	55.21-55.22,55.29,56.31,56.35,56.39,57.31-57.32,57.39,57.94-57.95,58.21-58.22,58.29,59.29
Operations on the male genital organs 60-64	60.18-60.19,61.19,62.19,63.09,64.19,64.94
Operations on the female genital organs 65-71	65.19,66.19,67.19,68.11,68.19,69.92,70.21-70.22,70.29,71.19
Obstetrical procedures 72-75	73.4-73.59,73.91-73.92,75.31-75.32,75.34-75.35,75.94
Operations on the musculoskeletal system 76-84	76.19,78.80-78.89,80.20-80.29,81.98,83.29,84.41-84.43,84.45-84.47
Operations on the integumentary system 85-86	85.19,86.19,86.92
Miscellaneous diagnostic and therapeutic procedures 87-99	87.01-99.99

Florida, Kentucky, Tennessee, Alabama, Mississippi, Arkansas, Louisiana, Oklahoma, and Texas

West Montana, Idaho, Wyoming, Colorado, New Mexico, Arizona, Utah, Nevada, Washington, Oregon, California, Hawaii, and Alaska

Changes in ICD-9-CM Codes

Although the ICD-9-CM has been used for coding NHDS data since 1979, the classification system undergoes annual updating. Assignment of new

diagnostic and procedure codes, fourth- and fifth-digit expansion of codes, as well as code deletions, are contained in addenda developed by the ICD-9-CM Coordination and Maintenance Committee and approved by the Director of NCHS and the Administrator of the Centers for Medicare and Medicaid Services (formerly the Health Care Financing Administration). Addenda to the ICD-9-CM become effective October 1 of each year and have been released annually since 1986.

The data in this report were coded using the addenda for October 1, 1986-98. When hospitals reported data using code changes introduced in the

October 1, 1999, addendum, the new codes were converted to their previous code assignments so that the annual estimates would be based on consistent coding.

Tables IV and V show the new ICD-9-CM codes from the 1986-98 addenda, the date they were introduced, and the code to which the diagnosis or procedure had been assigned before development of the new code. Specific titles and more detailed information about the coding system can be found in appropriate volumes of the ICD-9-CM (1).

Table IV. Changes in diagnosis codes

Current code(s) assignment	Effective October 1	Previous code(s) assignment
005.81	1995	005.8
005.89	1995	005.8
007.4	1997	007.8
008.00-008.09	1992	008.0
008.43-008.47	1992	008.49
008.61-008.69	1992	008.6
031.2	1997	031.8
038.10	1997	038.1
038.11	1997	038.1
038.19	1997	038.1
041.00-041.09	1992	041.0
041.04 (code title restated)	1997	041.04
041.10-041.19	1992	041.1
041.81-041.85	1992	041.8
041.86	1995	041.84
041.89	1992	041.8
042	1994	042.0-042.2, 042.9, 043.0-043.3, 043.9, 044.0, 044.9 (Codes deleted)
042.0-042.9	1986	279.19
043.0-043.9	1986	279.19
044.0-044.9	1986	279.19
070.20-070.21	1991	070.2
070.22	1994	070.20
070.23	1994	070.21
070.30-070.31	1991	070.3
070.32	1994	070.30
070.33	1994	070.31
070.41-070.43	1991	070.4
070.44	1994	070.41
070.49	1991	070.4
070.51-070.53	1991	070.5
070.54	1994	070.51
070.59	1991	070.5
077.98-077.99	1993	077.9
078.10-078.11,078.19	1993	078.1
078.88	1993	078.89
079.4	1993	079.8
079.50-079.53,079.59	1993	079.8
079.6	1996	079.89
079.81	1995	079.89
079.88-079.89	1993	079.8
079.98-079.99	1993	079.9
088.81,088.89	1989	088.8
088.82	1993	088.89
099.40-099.49	1992	099.4
099.50-099.59	1992	078.89
112.84-112.85	1992	112.89
114.4-114.5	1993	114.3
176.0-176.9	1991	173.0-173.9
203.00	1991	203.0
203.01	1991	V10.79
203.10	1991	203.1
203.11	1991	V10.79
203.80	1991	203.8
203.81	1991	V10.79
204.00	1991	204.0
204.01	1991	V10.61
204.10	1991	204.1
204.11	1991	V10.61
204.20	1991	204.2
204.21	1991	V10.61
204.80	1991	204.8
204.81	1991	V10.61
204.90	1991	204.9
204.91	1991	V10.61
205.00	1991	205.0
205.01	1991	V10.62
205.10	1991	205.1

Table IV. Changes in diagnosis codes—Con.

Current code(s) assignment	Effective October 1	Previous code(s) assignment
205.11	1991	V10.62
205.20	1991	205.2
205.21	1991	V10.62
205.30	1991	205.3
205.31	1991	V10.62
205.80	1991	205.8
205.81	1991	V10.62
205.90	1991	205.9
205.91	1991	V10.62
206.00	1991	206.0
206.01	1991	V10.63
206.10	1991	206.1
206.11	1991	V10.63
206.20	1991	206.2
206.21	1991	V10.63
206.80	1991	206.8
206.81	1991	V10.63
206.90	1991	206.9
206.91	1991	V10.63
207.00	1991	207.0
207.01	1991	V10.69
207.10	1991	207.1
207.11	1991	V10.69
207.20	1991	207.2
207.21	1991	V10.69
207.80	1991	207.8
207.81	1991	V10.69
208.00	1991	208.0
208.01	1991	V10.60
208.10	1991	208.1
208.11	1991	V10.60
208.20	1991	208.2
208.21	1991	V10.60
208.80	1991	208.8
208.81	1991	V10.60
208.90	1991	208.9
208.91	1991	V10.60
237.70-237.72	1990	237.7
250.02	1993	250.90
250.03	1993	250.91
250.12	1993	250.10
250.13	1993	250.11
250.22	1993	250.20
250.23	1993	250.21
250.32	1993	250.30
250.33	1993	250.31
250.42	1993	250.40
250.43	1993	250.41
250.52	1993	250.50
250.53	1993	250.51
250.62	1993	250.60
250.63	1993	250.61
250.72	1993	250.70
250.73	1993	250.71
250.82	1993	250.80
250.83	1993	250.81
250.92	1993	250.90
250.93	1993	250.91
275.40-275.42	1997	275.4
275.49	1997	275.4
278.00-278.01	1995	278.0
283.10-283.11,283.19	1993	283.1
291.81	1996	291.8
291.89	1996	291.8
293.84	1996	293.89

Table IV. Changes in diagnosis codes—Con.

Current code(s) assignment	Effective October 1	Previous code(s) assignment
300.82	1996	300.81
305.1	1994	305.10, 305.11, 305.12, 305.13 (Codes deleted)
312.81-312.82,312.89	1994	312.8
315.32	1996	315.39
320.81-320.89	1992	320.8
333.92-333.93	1994	333.99
337.20-337.22,337.29	1993	337.9
337.3	1998	337.9
342.00-342.02	1994	342.0
342.10-342.12	1994	342.1
342.80-342.82	1994	342.9
342.90-342.92	1994	342.9
344.00-344.04,344.09	1994	344.0
344.30-344.32	1994	344.3
344.40-344.42	1994	344.4
344.81,344.89	1993	344.8
345.00-345.01	1989	345.0
345.10-345.11	1989	345.1
345.40-345.41	1989	345.4
345.50-345.51	1989	345.5
345.60-345.61	1989	345.6
345.70-345.71	1989	345.7
345.80-345.81	1989	345.8
345.90-345.91	1989	345.9
346.00-346.01	1992	346.0
346.10-346.11	1992	346.1
346.20-346.21	1992	346.2
346.80-346.81	1992	346.8
346.90-346.91	1992	346.9
355.71	1993	354.4
355.79	1993	355.7
371.82	1992	371.89
374.87	1990	374.89
403.00-403.01	1989	403.0
403.10-403.11	1989	403.1
403.90-403.91	1989	403.9
404.00-404.03	1989	404.0
404.10-404.13	1989	404.1
404.90-404.93	1989	404.9
410.00-410.02	1989	410.0
410.10-410.12	1989	410.1
410.20-410.22	1989	410.2
410.30-410.32	1989	410.3
410.40-410.42	1989	410.4
410.50-410.52	1989	410.5
410.60-410.62	1989	410.6
410.70-410.72	1989	410.7
410.80-410.82	1989	410.8
410.90-410.92	1989	410.9
411.81	1989	410.9
411.89	1989	411.8
414.00-414.01	1994	414.0
414.02-414.03	1994	996.03
414.04-414.05	1996	414.00
415.11	1995	997.3 & 415.1
415.19	1995	415.1
429.71	1989	410.0-410.9
429.79	1989	410.0-410.9
433.00-433.01	1993	433.0
433.10-433.11	1993	433.1
433.20-433.21	1993	433.2
433.30-433.31	1993	433.3
433.80-433.81	1993	433.8
433.90-433.91	1993	433.9
434.00-434.01	1993	434.0
434.10-434.11	1993	434.1

Table IV. Changes in diagnosis codes—Con.

Current code(s) assignment	Effective October 1	Previous code(s) assignment
434.90-434.91	1993	434.9
435.3	1995	435.0 & 435.1
437.7	1992	780.9
438.0	1997	294.9 & 438
438.10	1997	784.5 & 438
438.11	1997	784.3 & 438
438.12	1997	784.4 & 438
438.19	1997	784.5 & 438
438.20	1997	342.90 & 438
438.21	1997	342.91 & 438
438.22	1997	342.92 & 438
438.30	1997	344.40 & 438
438.31	1997	344.41 & 438
438.32	1997	344.42 & 438
438.40	1997	344.30 & 438
438.41	1997	344.31 & 438
438.42	1997	344.32 & 438
438.50-438.52	1997	344.89 & 438
438.53	1998	438.50
438.81	1997	784.69 & 438
438.82	1997	787.2 & 438
438.89	1997	438
438.9	1997	438
440.20-440.22	1992	440.2
440.23	1993	440.20 & 707.1 or 707.8 or 707.9
440.24	1993	440.20 & 785.4
440.29	1993	440.2
440.30-440.32	1994	996.1
441.00-441.03	1994	441.0
441.6	1993	441.1 & 441.3
441.7	1993	441.2 & 441.4
446.20-446.21,446.29	1990	446.2
451.82-451.84	1993	451.89
458.2	1995	997.9 & 458.9
458.8	1997	458.9
466.11	1996	466.1
466.19	1996	466.1
474.0 (code title restated)	1997	474.0
474.00-474.02	1997	474.0
482.30-482.39	1992	482.3
482.4	1998	482.4
482.41	1998	482.4
482.49	1998	482.4
482.81-482.89	1992	482.8
482.84	1997	482.83
483.0	1992	483
483.1	1996	078.88 & 484.8
483.8	1992	483
491.20-491.21	1991	491.2
493.20	1989	493.90
493.21	1989	493.91
512.1	1994	997.3
518.6	1997	518.89
518.81	1987	799.1
518.82-518.89	1987	518.8
518.83	1998	518.81
518.84	1998	518.81
519.00-519.02, 519.09	1998	519.0
524.00-524.09	1992	524.0
524.10-524.19	1992	524.1
524.60-524.69	1991	524.6
524.70-524.79	1992	524.8
530.10-530.11, 530.19	1993	530.1
530.81	1993	530.1
530.82-530.84, 530.89	1993	530.8
535.00-535.01	1991	535.0
535.10-535.11	1991	535.1

Table IV. Changes in diagnosis codes—Con.

Current code(s) assignment	Effective October 1	Previous code(s) assignment
535.20-535.21	1991	535.2
535.30-535.31	1991	535.3
535.40-535.41	1991	535.4
535.50-535.51	1991	535.5
535.60-535.61	1991	535.6
536.3	1994	536.8
536.40-536.42, 536.49	1998	997.4
537.82	1990	537.89
537.83	1991	537.82
556.0-556.6	1994	556
556.8-556.9	1994	556
562.02	1991	562.00
562.03	1991	562.01
562.12	1991	562.10
562.13	1991	562.11
564.81	1998	564.8
564.89	1998	564.8
569.60-569.61	1995	569.6
569.62	1998	569.69
569.69	1995	569.6
569.84	1990	557.1
569.85	1991	569.84
574.60	1996	574.00 & 574.30
574.61	1996	574.01 & 574.31
574.70	1996	574.10 & 574.40
574.71	1996	574.11 & 574.41
574.80	1996	574.00 & 574.10 574.30 & 574.40
574.81	1996	574.01 & 574.11 574.31 & 574.41
574.90	1996	574.20 & 574.50
574.91	1996	574.21 & 574.51
575.10-575.11	1996	575.1
575.12	1996	575.0 & 575.1
593.70-593.73	1994	593.7
596.51-596.53	1992	596.5
596.54	1992	344.61
596.55-596.59	1992	596.5
599.81-599.89	1992	599.8
645.0	1991	645
651.30-651.31, 651.33	1989	651.00-651.01, 651.03
651.40-651.41, 651.43	1989	651.10-651.11, 651.13
651.50-651.51, 651.53	1989	651.20-651.21, 651.23
651.60-651.61, 651.63	1989	651.80-651.81, 651.83
654.20-654.21, 654.23	1990	654.2, 654.9
654.90-654.94	1990	654.2, 654.9
655.70 & 655.71	1997	655.8
655.73	1997	655.8
657.0	1991	657
659.60, 659.61, 659.63	1992	659.80-659.81, 659.83
659.70, 659.71, 659.73	1998	656.30, 656.31, 656.33
665.10, 665.11	1992	665.10, 665.11, 665.12, 665.14 ¹
670.0	1991	670
672.0	1991	672
677	1994	No previous code assignment
686.00 & 686.01	1997	686.0
686.09	1997	686.0
690.10	1995	690
690.11	1995	691.8 & 704.8
690.12	1995	691.8
690.18	1995	690
690.8	1995	690
692.72-692.74	1992	692.79
692.82-692.83	1992	692.89
702.0-702.8	1991	702
702.11,702.19	1994	702.1

¹The title for the subcategory 665.1 has been changed, making the fifth-digit subclassification 665.12 and 665.14 invalid.

Table IV. Changes in diagnosis codes-Con.

Current code(s) assignment	Effective October 1	Previous code(s) assignment
704.02	1993	704.09
709.00-709.01, 709.09	1994	709.0
710.5	1992	288.3, 729.1
728.86	1995	729.4
733.10-733.16, 733.19	1993	733.1
738.10-738.19	1992	738.1
747.60-747.64, 747.69	1993	747.6
747.82	1993	747.89
752.51-752.52	1996	752.5
752.61-752.63	1996	752.6
752.64-752.65	1996	752.8
752.69	1996	752.8
753.10-753.17, 753.19	1990	753.1
753.20-753.23	1996	753.2
753.29	1996	753.2
756.70-756.71	1997	756.7
756.79	1997	756.7
758.81	1996	758.8
758.89	1996	758.9
759.81-759.82	1989	759.8
759.83	1994	759.89
759.89	1989	759.8
760.75	1991	760.79
760.76	1994	760.79
763.81-763.83, 763.89	1998	763.8
764.00-764.09	1988	764.0
764.10-764.19	1988	764.1
764.20-764.29	1988	764.2
764.90-764.99	1988	764.9
765.00-765.09	1988	765.0
765.10-765.19	1988	765.1
780.01-780.02	1992	780.0
780.03	1993	780.01
780.09	1992	780.0
780.31	1997	780.3
780.39	1997	780.3
780.57	1992	780.51, 780.53
780.71	1998	780.7
780.79	1998	780.7
781.8	1994	781.9
786.03	1998	786.09
786.04	1998	786.09
786.05	1998	786.09
786.06	1998	786.09
786.07	1998	786.09
787.01-787.03	1994	787.0
787.91	1995	558.9
787.99	1995	787.9
788.20-788.21, 788.29	1993	788.2
788.30-788.39	1992	788.3
788.41-788.43	1993	788.4
788.61-788.62, 788.69	1993	788.6
789.00-789.07, 789.09	1994	789.0
789.30-789.37, 789.39	1994	789.3
789.40-789.47, 789.49	1994	789.4
789.60-789.67, 789.69	1994	789.6
790.91	1993	790.9
790.92	1993	286.9
790.93, 790.99	1993	790.9
790.94	1997	790.99
795.71	1994	795.8 (Code deleted)
795.79	1994	795.7
795.8	1986	795.7
796.5	1997	796.9
864.05	1992	864.09
864.15	1992	864.19
909.5	1994	909.9

Table IV. Changes in diagnosis codes—Con.

Current code(s) assignment	Effective October 1	Previous code(s) assignment
922.31-922.33	1996	922.3
925.1-925.2	1993	925
959.0 (code title restated)	1997	959.0
959.01	1997	854.00
959.09	1997	959.0
965.61	1998	965.6
965.69	1998	965.6
989.81-989.84	1995	989.8
989.89	1995	989.8
995.50-995.55	1996	995.5
995.59	1996	995.5
995.60-995.69	1993	995.0
995.80	1996	995.81
995.81 (Code title restated)	1996	995.81
995.82-995.85	1996	995.81
995.86	1998	995.89
996.04	1994	996.09
996.51-996.59	1987	996.5
996.55	1998	996.52
996.56	1998	996.59
996.60-996.69	1989	996.6
996.68	1998	996.69
996.70-996.79	1989	996.7
996.80-996.84	1987	996.8
996.85	1990	996.8
996.86	1987	996.8
996.89	1987	996.8
997.00-997.01	1995	997.0
997.02	1995	997.9 & 430-434, 436
997.09	1995	997.0
997.91	1995	997.9
997.99	1995	997.9
998.11-998.12	1996	998.1
998.13	1996	998.89
998.51	1996	998.5
998.59	1996	998.5
998.81-998.82, 998.89	1994	998.8
998.83	1996	998.89
V02.51	1998	V02.5
V02.52	1998	V02.5
V02.59	1998	V02.5
V02.60-V02.62	1997	V02.6
V02.69	1997	V02.6
V03.81-V03.82, V03.89	1994	V03.8
V05.3-V05.4	1993	V05.8
V06.5-V06.6	1994	V06.8
V07.31, V07.39	1994	V07.3
V07.4	1992	V07.8
V08	1994	044.9, 795.8 (Codes deleted)
V09.0-V09.91	1993	No previous code assignments
V10.48	1998	V10.49
V12.00-V12.03, V12.09	1994	V12.0
V12.40-V12.41	1997	V12.4
V12.49	1997	V12.4
V12.50-V12.52	1995	V12.5
V12.59	1995	V12.5
V12.70-V12.72, V12.79	1994	V12.7
V13.00-V13.01, V13.09	1994	V13.0
V13.61	1998	V13.6
V13.69	1998	V13.6
V15.41-V15.42	1996	V15.4
V15.49	1996	V15.4
V15.82	1994	305.13 (Codes deleted)
V15.84-V15.86	1995	V15.89
V16.40-V16.43	1997	V16.4
V16.49	1997	V16.4

Table IV. Changes in diagnosis codes—Con.

Current code(s) assignment	Effective October 1	Previous code(s) assignment
V16.51	1998	V16.5
V16.59	1998	V16.5
V18.61	1998	V18.6
V18.69	1998	V18.6
V23.7	1989	V23.8
V23.81	1998	V23.8
V23.82	1998	V23.8
V23.83	1998	V23.8
V23.84	1998	V23.8
V23.89	1998	V23.8
V25.43	1992	V25.49
V25.5	1992	V25.8
V26.51	1998	No previous code assignment
V26.52	1998	No previous code assignment
V28.6	1997	V28.8
V29.0-V29.1, V29.8	1992	V71.8
V29.2	1994	V29.8
V29.3	1998	V29.8
V29.8	1992	V71.8
V29.9	1992	V71.9
V30.00-V30.01	1989	V30.0
V31.00-V31.01	1989	V31.0
V32.00-V32.01	1989	V32.0
V33.00-V33.01	1989	V33.0
V34.00-V34.01	1989	V34.0
V35.00-V35.01	1989	V35.0
V36.00-V36.01	1989	V36.0
V37.00-V37.01	1989	V37.0
V39.00-V39.01	1989	V39.0
V42.81-V42.83	1997	V42.8
V42.89	1997	V42.8
V43.60-V43.66, V43.69	1994	V43.6
V43.81-V43.82	1995	V43.8
V43.83	1998	V43.89
V43.89	1995	V43.8
V44.50	1998	V44.5
V44.51	1998	V44.5
V44.52	1998	V44.5
V44.59	1998	V44.5
V45.00	1994	V45.89
V45.01	1994	V45.0
V45.02, V45.09	1994	V45.89
V45.51	1994	V45.5
V45.52, V45.59	1994	V45.89
V45.61	1997	V45.6
V45.63	1997	V45.6
V45.71	1997	611.8
V45.72	1997	569.89
V45.73	1997	593.89
V45.82	1994	V45.89
V45.83	1995	V45.89
V49.60-V49.67	1994	V49.5
V49.70-V49.77	1994	V49.5
V50.41-V50.42, V50.49	1994	V50.8
V53.01-V53.02	1997	V53.0
V53.09	1997	V53.0
V53.31	1994	V53.3
V53.32, V53.39	1994	V53.9
V56.1	1995	V58.89
V56.1 (code title restated)	1998	V56.1
V56.2	1998	V56.1
V57.21-V57.22	1994	V57.2
V58.41, V58.49	1994	V58.4
V58.61	1995	V67.51
V58.62	1998	V58.69
V58.69	1995	V67.51
V58.81, V58.89	1994	V58.8

Table IV. Changes in diagnosis codes—Con.

Current code(s) assignment	Effective October 1	Previous code(s) assignment
V58.82	1995	V58.89
V59.01-V59.02	1995	V59.0
V59.09	1995	V59.0
V59.6	1995	V59.8
V61.10-V61.12	1996	V61.1
V61.22	1996	V61.21
V62.83	1996	V65.49
V64.4	1997	No previous code assignment
V65.40-V65.45, V65.49	1994	V65.4
V66.7	1996	No previous code assignment
V69.0-V69.3	1994	No previous code assignments
V69.8-V69.9	1994	No previous code assignments
V72.81-V72.85	1993	V72.8
V73.88-V73.89	1993	V73.8
V73.98-V73.99	1993	V73.9
V76.10-V76.12	1997	V76.1
V76.19	1997	V76.1
V76.44	1998	V76.49
V76.45	1998	V76.49
E854.8	1995	E858.8
E869.4	1994	E869.8
E880.1	1995	E884.9
E884.3-E884.4	1995	E884.2
E884.5-E884.6	1995	E884.9
E906.5	1995	E906.3
E908.0-E908.4	1995	E908
E908.8-E908.9	1995	E908
E909.0-E909.4	1995	E909
E909.8-E909.9	1995	E909
E920.5	1995	E920.4
E922.4	1997	E917.9
E924.2	1995	E924.0
E955.6	1997	E955.9
E967.2	1996	E967.0
E967.3	1996	No previous code assignment
E967.4-E967.8	1996	E967.1
E968.5	1995	E968.8
E968.6	1997	E968.8
E985.6	1997	E985.4

Table V. Changes in procedure codes

Current code(s) assignment	Effective October 1	Previous code(s) assignment
02.96	1992	89.19
03.90	1987	03.99 (Insertion of Catheter)
05.25	1995	39.7
11.75	1989	11.79
11.76	1989	11.62
20.96-20.98	1986	20.95
22.12	1988	22.11
26.12	1988	26.11
29.31	1991	83.02
29.32	1991	29.3
29.33	1991	29.3
29.39	1991	29.3
31.45	1988	31.43-31.44
31.95	1989	31.75
32.01	1989	32.0
32.09	1989	32.0
32.22	1995	32.29, 32.9
32.28	1989	32.29
33.27	1987	33.22 + 33.27
33.28	1987	33.27
33.29	1987	33.28-33.29
33.50	1995	33.5
33.51	1995	33.5
33.52	1995	33.5
33.6	1990	33.5 + 37.5
34.05	1994	34.99
35.84	1988	35.82
35.96	1986	35.03
36.00-36.03	1986	36.0
36.04	1986	39.97
36.05	1987	36.01
36.05	1986	36.01, ¹ 36.02
36.06	1995	36.01, 36.02, 36.03, 36.05
36.09	1986	36.0
36.09	1991	36.00 (Code deleted)
36.17	1996	36.19
36.31	1998	36.3
36.32	1998	36.3
36.39	1998	36.3
37.26-37.27	1988	37.29
37.34	1988	37.33
37.35	1997	37.33
37.65	1995	37.62
37.66	1995	37.62
37.67	1998	37.4
37.70 (Leads only)	1987	37.70 (Leads/Device)
37.71-37.72 (Leads only)	1987	37.74 (Leads/Device)
37.73 (Leads only)	1987	37.73 (Leads/Device)
37.74 (Leads only)	1987	37.76 (Leads/Device)
37.75 (Leads only)	1987	37.89 (Leads/Device)
37.76 (Leads only)	1987	37.81 (Leads/Device)
37.77 (Leads only)	1987	37.83-37.84 (Leads/Device)
37.78	1987	37.71-37.72
37.79	1987	86.09
37.80-37.87	1992	89.49 (Code deleted, this procedure is included in the code for pacemaker insertion/replacement)
37.80 (Device only)	1987	37.73-37.77 (Leads/Device)
37.81 (Device only)	1987	37.73-37.77 (Leads/Device)
37.82 (Device only)	1987	37.73-37.77 (Leads/Device)
37.83 (Device only)	1987	37.73-37.77 (Leads/Device)
37.85-37.87	1987	37.85
37.89	1987	37.86 + 37.89
37.94-37.98	1986	37.99
38.22	1986	38.29
38.44 (Abdominal Aorta Only)	1986	38.44 (Entire Aorta)
38.45 (Thoracic Aorta Added)	1986	38.44-38.45
38.95	1989	38.93
39.28	1991	39.29

See footnotes at end of table.

Table V. Changes in procedure codes—Con.

Current code(s) assignment	Effective October 1	Previous code(s) assignment
39.50	1995	39.59
39.65	1988	39.61
39.66	1990	39.65
39.90	1996	39.50
41.00-41.03	1988	41.0
41.04	1994	99.79
41.05	1997	No previous code assignment
41.06	1997	No previous code assignment
42.25	1988	42.24
42.33	1989	42.32, 42.39
42.33	1990	42.91
43.11	1989	43.1
43.19	1989	43.1, 43.2
43.41	1989	43.41, 43.49
44.21	1986	44.2
44.22	1986	44.99
44.29	1986	44.2
44.43	1989	43.49, 45.32
44.44	1989	38.86
44.49	1989	43.0
44.93-44.94	1986	44.99
45.16	1988	45.14 (45.15 before 1987)
45.30	1989	45.31, 45.32
45.42	1988	45.41
45.43	1989	45.49
45.75 (Hartmann Resection Added)	1988	48.66 (Code deleted)
45.95	1987	45.93
46.13	1992	46.12 (Code deleted)
46.32	1989	46.39
46.85	1989	46.99
47.01	1996	47.0
47.09	1996	47.0
47.11	1996	47.1
47.19	1996	47.1
48.36	1995	45.42
49.31	1989	49.3
49.39	1989	49.3
51.10	1989	51.97
51.11	1989	51.11, 51.97
51.14	1989	51.12
51.15	1989	51.97
51.21	1996	51.22, 51.23
51.22	1991	51.21 (Code deleted), 51.22
51.23	1991	51.22
51.24	1996	51.22, 51.23
51.64	1989	51.69
51.84-51.88	1989	51.97
51.97	1986	52.91, 51.99, or 51.82
51.98	1986	51.99
52.13	1989	51.97, 52.91
52.14	1989	52.11
52.21	1989	52.2
52.22	1989	52.2
52.84	1996	99.29
52.85	1996	99.29
52.86	1996	99.29
52.93	1989	52.93 + 52.91
52.94	1989	52.09
52.97	1989	52.91
52.98	1989	52.91
52.99	1989	52.93, 52.94, 52.99
54.24	1987	54.23
54.25	1993	54.98
54.51	1996	54.5
54.59	1996	54.5
55.03-55.04	1986	55.02
56.33-56.34	1987	56.33
56.35	1987	45.12

See footnotes at end of table.

Table V. Changes in procedure codes—Con.

Current code(s) assignment	Effective October 1	Previous code(s) assignment
57.17-57.18	1989	57.21
57.22	1989	57.22, 57.82
58.31	1990	58.3
58.39	1990	58.3
58.93	1986	57.99
59.03	1996	59.02
59.12	1996	59.11
59.72	1995	59.79
59.96	1986	59.95
60.21	1995	60.2
60.29	1995	60.2
60.95	1991	60.99
64.97	1986	64.95
65.01	1996	65.0
65.09	1996	65.0
65.13	1996	65.12
65.14	1996	65.19
65.23	1996	65.21
65.24	1996	65.22
65.25	1996	65.29
65.31	1996	65.3
65.39	1996	65.3
65.41	1996	65.4
65.49	1996	65.4
65.53	1996	65.51
65.54	1996	65.52
65.63	1996	65.61
65.64	1996	65.62
65.74	1996	65.71
65.75	1996	65.72
65.76	1996	65.73
65.81	1996	65.8
65.89	1996	65.8
66.01	1992	66.0
66.02	1992	66.73
68.15	1987	68.14
68.16	1987	68.13
68.23	1996	68.29
68.51	1996	68.5
68.59	1996	68.5
68.9	1992	68.4
74.3	1992	69.11 (Code deleted)
75.37	1998	99.29
77.56	1989	77.89, 78.49, 81.18
77.57	1989	77.89, 80.48, 81.18, 83.85
77.58	1989	77.59, 81.18
78.10	1991	78.40
78.11	1991	78.41
78.12	1991	78.42
78.13	1991	78.43
78.14	1991	78.44
78.15	1991	78.45
78.16	1991	78.46
78.17	1991	78.47
78.18	1991	78.48
78.19	1991	78.49
78.20	1991	78.10, 78.20, 78.30
78.21 (Invalid code)	1991	78.11, 78.31
78.22	1991	78.12, 78.22, 78.32
78.23	1991	78.13, 78.23, 78.33
78.24	1991	78.14, 78.34
78.25	1991	78.15, 78.25, 78.35
78.27	1991	78.17, 78.27, 78.37
78.28	1991	78.18, 78.38
78.29	1991	78.11, 78.16, 78.19, 78.29, 78.39
78.39	1991	78.31
78.90 ²	1987	78.40
78.91 ²	1987	78.41

See footnotes at end of table.

Table V. Changes in procedure codes—Con.

Current code(s) assignment	Effective October 1	Previous code(s) assignment
78.92 ²	1987	78.42
78.93 ²	1987	78.43
78.94 ²	1987	78.44
78.95 ²	1987	78.45
78.96 ²	1987	78.46
78.97 ²	1987	78.47
78.98 ²	1987	78.48
78.99 ²	1987	78.49
80.50-80.59	1986	80.5
81.03	1989	81.02
81.04-81.05	1989	81.03, 81.04, 81.05
81.06-81.07	1989	81.06, 81.07
81.08	1989	81.06, 81.07, 81.08
81.09	1989	81.08
81.40	1989	81.69
81.51	1989	81.51, 81.59
81.52	1989	81.61, 81.62, 81.63, 81.64
81.53	1989	81.51, 81.59, 81.61, 81.62, 81.63, 81.64
81.54-81.55	1989	81.41 (Code deleted)
81.56	1989	81.48
81.57	1989	81.31, 81.39
81.59	1989	81.39
81.72	1989	81.79
81.73-81.74	1989	81.86 (Code deleted)
81.75	1989	81.87 (Code deleted)
81.79	1989	81.79, 81.87
81.80	1989	81.81
81.97	1992	81.59
85.95	1987	85.99
85.96	1987	85.99
86.06	1987	86.09
86.07	1990	86.09
86.27	1986	86.22-86.23
86.28	1988	86.22
86.67	1998	86.65
86.93	1987	86.89
88.90	1986	88.39
88.91	1986	89.15
88.92	1986	89.39
88.93	1986	89.15
88.94	1986	89.39
88.95	1986	89.29
88.97	1989	88.99
88.98	1989	88.90
88.99	1986	89.39
89.10	1989	89.15
89.17-89.18	1988	89.15
89.19	1989	89.15
89.50	1991	89.54
92.3	1995	01.59, 04.07, 07.63, 07.68
92.30	1998	92.3
92.31	1998	92.3
92.32	1998	92.3
92.33	1998	92.3
92.39	1998	92.3
93.90	1988	93.92
94.61-94.69	1989	94.25
96.29	1998	96.39
96.6	1986	96.35
96.70	1991	93.92 (Code deleted)
96.71	1991	93.92 (Code deleted)
96.72	1991	93.92 (Code deleted)
97.05	1989	51.97
98.51-98.52	1989	59.96 (Code deleted)
98.59	1989	59.96 (Code deleted)
99.00	1995	99.02
99.10	1998	99.29
99.15	1986	99.29

See footnotes at end of table.

Table V. Changes in procedure codes—Con.

Current code(s) assignment	Effective October 1	Previous code(s) assignment
99.20	1998	99.29
99.28	1994	99.25
99.71-99.74, 99.79 ³	1988	99.07
99.85	1987	93.35
99.86	1987	93.39
99.88	1988	99.83

¹Before October 1986 contents of current code 36.05 would have been assigned to 36.0.

²Codes 78.90-78.99 were retitled as "Insertion of bone growth stimulator" in October 1987; the previous contents of codes 78.90-78.99 were reassigned to codes 78.40-78.49.

³Codes 99.71-99.79 were deleted in October 1987; their contents were not transferred elsewhere. In the October 1988 revision, codes 99.71-99.79 were reclassified as "Therapeutic apheresis." Codes 99.75-99.78 have not yet been reassigned.

Vital and Health Statistics series descriptions

- SERIES 1. **Programs and Collection Procedures**—These reports describe the data collection programs of the National Center for Health Statistics. They include descriptions of the methods used to collect and process the data, definitions, and other material necessary for understanding the data.
- SERIES 2. **Data Evaluation and Methods Research**—These reports are studies of new statistical methods and include analytical techniques, objective evaluations of reliability of collected data, and contributions to statistical theory. These studies also include experimental tests of new survey methods and comparisons of U.S. methodology with those of other countries.
- SERIES 3. **Analytical and Epidemiological Studies**—These reports present analytical or interpretive studies based on vital and health statistics. These reports carry the analyses further than the expository types of reports in the other series.
- SERIES 4. **Documents and Committee Reports**—These are final reports of major committees concerned with vital and health statistics and documents such as recommended model vital registration laws and revised birth and death certificates.
- SERIES 5. **International Vital and Health Statistics Reports**—These reports are analytical or descriptive reports that compare U.S. vital and health statistics with those of other countries or present other international data of relevance to the health statistics system of the United States.
- SERIES 6. **Cognition and Survey Measurement**—These reports are from the National Laboratory for Collaborative Research in Cognition and Survey Measurement. They use methods of cognitive science to design, evaluate, and test survey instruments.
- SERIES 10. **Data From the National Health Interview Survey**—These reports contain statistics on illness; unintentional injuries; disability; use of hospital, medical, and other health services; and a wide range of special current health topics covering many aspects of health behaviors, health status, and health care utilization. They are based on data collected in a continuing national household interview survey.
- SERIES 11. **Data From the National Health Examination Survey, the National Health and Nutrition Examination Surveys, and the Hispanic Health and Nutrition Examination Survey**—Data from direct examination, testing, and measurement on representative samples of the civilian noninstitutionalized population provide the basis for (1) medically defined total prevalence of specific diseases or conditions in the United States and the distributions of the population with respect to physical, physiological, and psychological characteristics, and (2) analyses of trends and relationships among various measurements and between survey periods.
- SERIES 12. **Data From the Institutionalized Population Surveys**—Discontinued in 1975. Reports from these surveys are included in Series 13.
- SERIES 13. **Data From the National Health Care Survey**—These reports contain statistics on health resources and the public's use of health care resources including ambulatory, hospital, and long-term care services based on data collected directly from health care providers and provider records.
- SERIES 14. **Data on Health Resources: Manpower and Facilities**—Discontinued in 1990. Reports on the numbers, geographic distribution, and characteristics of health resources are now included in Series 13.
- SERIES 15. **Data From Special Surveys**—These reports contain statistics on health and health-related topics collected in special surveys that are not part of the continuing data systems of the National Center for Health Statistics.
- SERIES 16. **Compilations of Advance Data From Vital and Health Statistics**—Advance Data Reports provide early release of information from the National Center for Health Statistics' health and demographic surveys. They are compiled in the order in which they are published. Some of these releases may be followed by detailed reports in Series 10–13.
- SERIES 20. **Data on Mortality**—These reports contain statistics on mortality that are not included in regular, annual, or monthly reports. Special analyses by cause of death, age, other demographic variables, and geographic and trend analyses are included.
- SERIES 21. **Data on Natality, Marriage, and Divorce**—These reports contain statistics on natality, marriage, and divorce that are not included in regular, annual, or monthly reports. Special analyses by health and demographic variables and geographic and trend analyses are included.
- SERIES 22. **Data From the National Mortality and Natality Surveys**—Discontinued in 1975. Reports from these sample surveys, based on vital records, are now published in Series 20 or 21.
- SERIES 23. **Data From the National Survey of Family Growth**—These reports contain statistics on factors that affect birth rates, including contraception, infertility, cohabitation, marriage, divorce, and remarriage; adoption; use of medical care for family planning and infertility; and related maternal and infant health topics. These statistics are based on national surveys of women of childbearing age.
- SERIES 24. **Compilations of Data on Natality, Mortality, Marriage, Divorce, and Induced Terminations of Pregnancy**—These include advance reports of births, deaths, marriages, and divorces based on final data from the National Vital Statistics System that were published as supplements to the *Monthly Vital Statistics Report* (MVSR). These reports provide highlights and summaries of detailed data subsequently published in *Vital Statistics of the United States*. Other supplements to the MVSR published here provide selected findings based on final data from the National Vital Statistics System and may be followed by detailed reports in Series 20 or 21.

For answers to questions about this report or for a list of reports published in these series, contact:

Data Dissemination Branch
National Center for Health Statistics
Centers for Disease Control and Prevention
6525 Belcrest Road, Room 1064
Hyattsville, MD 20782-2003
(301) 458-4636
E-mail: nchsquery@cdc.gov
Internet: www.cdc.gov/nchs/