



Trends in the Prevalence of Selected Risk Behaviors and Obesity for Hispanic Students National YRBS: 1991–2009

The national Youth Risk Behavior Survey (YRBS) monitors priority health risk behaviors that contribute to the leading causes of death, disability, and social problems among youth and adults in the United States. The national YRBS is conducted every two years during the spring semester and provides data representative of 9th through 12th grade students in public and private schools throughout the United States.

1991	1993	1995	1997	1999	2001	2003	2005	2007	2009	Changes from 1991–2009 ¹	Change from 2007–2009 ²
Rarely or never wore a seat belt (when riding in a car driven by someone else)											
31.3 (24.8–38.6) ³	19.5 (16.1–23.5)	18.4 (13.7–24.2)	20.3 (14.6–27.4)	14.4 (10.7–19.2)	14.5 (11.9–17.5)	20.2 (13.8–28.5)	10.6 (8.6–13.0)	12.9 (9.7–17.0)	8.8 (7.5–10.3)	Decreased, 1991–2009	Decreased
Rode with a driver who had been drinking alcohol one or more times (in a car or other vehicle during the 30 days before the survey)											
46.9 (40.9–53.0)	42.3 (39.8–45.0)	49.5 (41.9–57.1)	42.8 (38.8–46.8)	39.5 (35.4–43.8)	38.3 (34.0–42.9)	36.4 (32.2–40.8)	36.1 (32.8–39.6)	35.5 (32.3–38.8)	34.2 (31.8–36.7)	Decreased, 1991–2009	No change
Carried a weapon on at least 1 day (for example, a gun, knife, or club during the 30 days before the survey)											
25.8 (21.5–30.6)	24.4 (21.9–27.1)	24.7 (20.7–29.2)	23.3 (20.6–26.3)	18.7 (16.1–21.5)	16.5 (15.0–18.1)	16.5 (14.0–19.3)	19.0 (16.9–21.3)	18.5 (16.2–21.1)	17.2 (15.4–19.1)	Decreased, 1991–2001 No change, 2001–2009	No change
Did not go to school because they felt unsafe at school or on their way to or from school on at least 1 day (during the 30 days before the survey)											
NA ⁴	10.1 (8.5–11.9)	8.5 (6.0–11.7)	7.2 (5.6–9.1)	11.2 (8.2–15.1)	10.2 (8.9–11.7)	9.4 (8.0–11.1)	10.2 (8.7–12.1)	9.6 (7.8–11.8)	8.1 (6.9–9.4)	No change, 1993–2009	No change
Attempted suicide one or more times (during the 12 months before the survey)											
7.9 (6.2–9.9)	13.6 (12.1–15.1)	13.4 (10.4–17.3)	10.7 (8.6–13.3)	12.8 (10.9–15.1)	12.1 (10.5–13.9)	10.6 (9.3–12.1)	11.3 (9.8–12.9)	10.2 (9.0–11.6)	8.1 (7.1–9.3)	Increased, 1991–1993 Decreased, 1993–2009	Decreased
Smoked cigarettes on a least 1 day (during the 30 days before the survey)											
25.3 (22.5–28.2)	28.7 (25.8–31.8)	34.0 (28.7–39.6)	34.0 (31.3–36.9)	32.7 (29.0–36.6)	26.6 (22.4–31.2)	18.4 (16.1–20.9)	22.0 (18.7–25.8)	16.7 (13.5–20.4)	18.0 (16.0–20.2)	Increased, 1991–1995 Decreased, 1995–2009	No change
Had at least one drink of alcohol on at least 1 day (during the 30 days before the survey)											
54.3 (48.3–60.1)	50.8 (45.3–56.4)	54.7 (49.3–59.9)	53.9 (50.0–57.8)	52.8 (47.9–57.6)	49.2 (46.2–52.3)	45.6 (42.8–48.4)	46.8 (44.0–49.7)	47.6 (44.0–51.3)	42.9 (40.1–45.8)	Decreased, 1991–2009	Decreased

Where can I get more information? Visit www.cdc.gov/healthyyouth or call 800–CDC–INFO (800–232–4636).



U.S. Department of Health and Human Services
Centers for Disease Control and Prevention
National Center for Chronic Disease Prevention and Health Promotion
Division of Adolescent and School Health



1991	1993	1995	1997	1999	2001	2003	2005	2007	2009	Changes from 1991–2009 ¹	Change from 2007–2009 ²
Used marijuana one or more times (during the 30 days before the survey)											
14.4 (11.2–18.3)	19.4 (16.8–22.3)	27.8 (22.3–34.1)	28.6 (24.7–33.0)	28.2 (23.8–33.0)	24.6 (23.0–26.3)	23.8 (21.5–26.2)	23.0 (20.7–25.6)	18.5 (15.8–21.5)	21.6 (19.6–23.8)	Increased, 1991–1997 Decreased, 1997–2009	No change
Ever used methamphetamines one or more times (also called “speed”, “crystal”, “crank”, or “ice”, during their life)											
NA	NA	NA	NA	11.3 (7.2–17.2)	9.1 (7.3–11.2)	8.3 (6.5–10.5)	8.8 (6.8–11.3)	5.7 (4.1–7.9)	5.7 (4.5–7.1)	Decreased, 1999–2009	No change
Ever had sexual intercourse											
53.1 (49.4–56.7)	56.0 (51.8–60.2)	57.6 (48.6–66.1)	52.2 (48.4–55.8)	54.1 (49.0–59.0)	48.4 (43.8–53.0)	51.4 (48.1–54.8)	51.0 (46.5–55.4)	52.0 (48.3–55.6)	49.1 (46.0–52.2)	No change, 1991–2009	No change
Had sexual intercourse with four or more persons (during their life)											
16.8 (14.3–19.7)	18.6 (15.7–22.0)	17.6 (14.1–21.7)	15.5 (13.2–18.1)	16.6 (13.2–20.7)	14.9 (13.2–16.7)	15.7 (13.5–18.1)	15.9 (13.6–18.5)	17.3 (15.2–19.5)	14.2 (12.7–15.9)	No change, 1991–2009	Decreased
Used a condom during last sexual intercourse (among students who were currently sexually active)											
37.4 (31.3–44.0)	46.1 (41.6–50.6)	44.4 (33.4–56.0)	48.3 (42.6–54.0)	55.2 (48.1–62.0)	53.5 (48.2–58.7)	57.4 (51.9–62.8)	57.7 (53.4–61.8)	61.4 (56.7–65.9)	54.9 (51.7–58.0)	Increased, 1991–2007 Decreased, 2007–2009	Decreased
Used birth control pills or Depo-Provera before last sexual intercourse to prevent pregnancy (among students who were currently sexually active)											
NA	NA	NA	NA	11.0 (7.3–16.3)	12.2 (8.6–17.2)	14.5 (10.8–19.2)	11.3 (8.7–14.4)	10.1 (7.9–12.7)	14.0 (11.4–17.1)	No change, 1999–2009	Increased
Attended physical education daily in an average week (when they were in school)											
46.6 (40.7–52.5)	39.7 (35.6–44.0)	33.1 (25.7–41.5)	38.4 (34.5–42.5)	40.4 (31.2–50.2)	38.7 (32.5–45.3)	36.7 (28.9–45.3)	38.3 (27.8–50.1)	36.0 (28.4–44.3)	40.5 (32.8–48.7)	No change, 1991–2009	No change
Obese (students who were $\geq 95^{\text{th}}$ percentile for body mass index, by age and sex, based on reference data)											
NA	NA	NA	NA	13.6 (11.4–16.1)	15.1 (12.4–18.3)	16.4 (14.0–19.2)	16.8 (14.5–19.4)	16.6 (14.7–18.7)	15.1 (13.5–16.8)	No change, 1999–2009	No change

¹ Based on trend analyses using a logistic regression model controlling for sex and grade.

² Based on t-test analyses, $p < 0.05$.

³ 95% confidence interval.

⁴ Not available.

Where can I get more information? Visit www.cdc.gov/healthyyouth or call 800–CDC–INFO (800–232–4636).



U.S. Department of Health and Human Services
Centers for Disease Control and Prevention
National Center for Chronic Disease Prevention and Health Promotion
Division of Adolescent and School Health

