

Worktable 293. Age-adjusted death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by race and sex: United States, 1999-2005

[Age-adjusted death rates per 100,000 U.S. standard population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992) and year	All races 1/			White 2/			Black 2/		
	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female
All causes									
2005.....	798.8	951.1	677.6	785.3	933.2	666.5	1,016.5	1,252.9	845.7
2004.....	800.8	955.7	679.2	786.3	936.9	666.9	1,027.3	1,269.4	855.3
2003.....	832.7	994.3	706.2	817.0	973.9	693.1	1,065.9	1,319.1	885.6
2002.....	845.3	1,013.7	715.2	829.0	992.9	701.3	1,083.3	1,341.4	901.8
2001.....	854.5	1,029.1	721.8	836.5	1,006.1	706.7	1,101.2	1,375.0	912.5
2000.....	869.0	1,053.8	731.4	849.8	1,029.4	715.3	1,121.4	1,403.5	927.6
1999.....	875.6	1,067.0	734.0	854.6	1,040.0	716.6	1,135.7	1,432.6	933.6
Salmonella infections (A01-A02)									
2005.....	0.0	*	*	0.0	*	*	*	*	*
2004.....	0.0	0.0	*	0.0	*	*	*	*	*
2003.....	0.0	0.0	0.0	0.0	*	*	*	*	*
2002.....	0.0	*	*	*	*	*	*	*	*
2001.....	0.0	0.0	0.0	0.0	*	*	*	*	*
2000.....	0.0	*	*	0.0	*	*	*	*	*
1999.....	0.0	0.0	*	0.0	*	*	*	*	*
Shigellosis and amebiasis (A03,A06)									
2005.....	*	*	*	*	*	*	*	*	*
2004.....	*	*	*	*	*	*	*	*	*
2003.....	*	*	*	*	*	*	*	*	*
2002.....	*	*	*	*	*	*	*	*	*
2001.....	*	*	*	*	*	*	*	*	*
2000.....	*	*	*	*	*	*	*	*	*
1999.....	*	*	*	*	*	*	*	*	*
Certain other intestinal infections (A04,A07-A09)									
2005.....	1.8	1.9	1.8	1.9	2.0	1.9	1.2	1.3	1.2
2004.....	1.5	1.4	1.5	1.5	1.5	1.6	0.9	0.9	0.9
2003.....	1.0	1.0	1.1	1.1	1.0	1.1	0.7	0.7	0.8
2002.....	0.8	0.8	0.8	0.9	0.9	0.9	0.6	0.7	0.6
2001.....	0.5	0.5	0.5	0.6	0.5	0.6	0.4	0.5	0.4
2000.....	0.5	0.5	0.5	0.5	0.5	0.5	0.3	0.4	0.3
1999.....	0.4	0.4	0.4	0.4	0.4	0.4	0.3	0.3	0.3

Worktable 293. Age-adjusted death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by race and sex: United States, 1999-2005

[Age-adjusted death rates per 100,000 U.S. standard population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992) and year

	All races 1/			White 2/			Black 2/		
	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female
Scarlet fever and erysipelas (A38,A46)									
2005.....	*	*	*	*	*	*	*	*	*
2004.....	*	*	*	*	*	*	*	*	*
2003.....	*	*	*	*	*	*	*	*	*
2002.....	*	*	*	*	*	*	*	*	*
2001.....	*	*	*	*	*	*	*	*	*
2000.....	*	*	*	*	*	*	*	*	*
1999.....	*	*	*	*	*	*	*	*	*

Meningococcal infection (A39)

2005.....	0.0	0.0	0.0	0.0	0.0	0.0	*	*	*
2004.....	0.0	0.0	0.0	0.0	0.0	0.0	0.1	*	*
2003.....	0.0	0.0	0.0	0.0	0.0	0.0	0.1	*	*
2002.....	0.0	0.1	0.0	0.0	0.1	0.0	0.1	*	*
2001.....	0.1	0.1	0.1	0.1	0.0	0.1	0.1	*	*
2000.....	0.0	0.1	0.1	0.0	0.1	0.1	0.1	0.1	*
1999.....	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	*

Septicemia (A40-A41)

2005.....	11.2	12.3	10.4	10.2	11.2	9.4	22.6	25.8	20.5
2004.....	11.2	12.3	10.3	10.2	11.3	9.4	22.5	25.7	20.4
2003.....	11.6	12.7	10.9	10.5	11.5	9.9	23.8	27.3	21.6
2002.....	11.7	12.9	10.9	10.7	11.7	9.9	24.2	27.8	22.1
2001.....	11.4	12.7	10.5	10.3	11.5	9.5	23.6	27.7	21.2
2000.....	11.3	12.4	10.5	10.2	11.2	9.4	24.6	28.0	22.5
1999.....	11.3	12.5	10.4	10.0	11.1	9.3	24.6	28.9	21.9

Syphilis (A50-A53)

2005.....	0.0	0.0	*	*	*	*	0.1	0.2	*
2004.....	0.0	0.0	0.0	*	*	*	0.1	*	*
2003.....	0.0	0.0	*	*	*	*	*	*	*
2002.....	0.0	0.0	0.0	*	*	*	0.1	*	*
2001.....	0.0	*	*	0.0	*	*	*	*	*
2000.....	0.0	0.0	*	0.0	*	*	0.1	*	*
1999.....	0.0	*	0.0	0.0	*	*	*	*	*

Worktable 293. Age-adjusted death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by race and sex: United States, 1999-2005

[Age-adjusted death rates per 100,000 U.S. standard population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992) and year

	All races 1/			White 2/			Black 2/		
	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female
Acute poliomyelitis (A80)									
2005.....	*	*	*	*	*	*	*	*	*
2004.....	*	*	*	*	*	*	*	*	*
2003.....	*	*	*	*	*	*	*	*	*
2002.....	*	*	*	*	*	*	*	*	*
2001.....	*	*	*	*	*	*	*	*	*
2000.....	*	*	*	*	*	*	*	*	*
1999.....	*	*	*	*	*	*	*	*	*
Arthropod-borne viral encephalitis (A83-A84,A85.2)									
2005.....	*	*	*	*	*	*	*	*	*
2004.....	*	*	*	*	*	*	*	*	*
2003.....	*	*	*	*	*	*	*	*	*
2002.....	*	*	*	*	*	*	*	*	*
2001.....	*	*	*	*	*	*	*	*	*
2000.....	*	*	*	*	*	*	*	*	*
1999.....	*	*	*	*	*	*	*	*	*
Measles (B05)									
2005.....	*	*	*	*	*	*	*	*	*
2004.....	*	*	*	*	*	*	*	*	*
2003.....	*	*	*	*	*	*	*	*	*
2002.....	*	*	*	*	*	*	*	*	*
2001.....	*	*	*	*	*	*	*	*	*
2000.....	*	*	*	*	*	*	*	*	*
1999.....	*	*	*	*	*	*	*	*	*
Viral hepatitis (B15-B19)									
2005.....	1.8	2.4	1.2	1.7	2.3	1.1	2.6	3.4	1.9
2004.....	1.8	2.4	1.2	1.7	2.3	1.1	2.6	3.8	1.7
2003.....	1.8	2.5	1.2	1.7	2.4	1.1	2.7	3.8	1.8
2002.....	2.0	2.7	1.3	1.9	2.6	1.2	2.8	3.8	1.9
2001.....	2.0	2.6	1.3	1.8	2.5	1.2	2.7	3.4	2.1
2000.....	1.9	2.6	1.3	1.8	2.4	1.2	2.7	3.8	1.8
1999.....	1.8	2.4	1.2	1.6	2.2	1.1	2.6	3.4	1.9

Worktable 293. Age-adjusted death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by race and sex: United States, 1999-2005

[Age-adjusted death rates per 100,000 U.S. standard population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992) and year

	All races 1/			White 2/			Black 2/		
	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female
Human immunodeficiency virus (HIV) disease (B20-B24)									
2005.....	4.2	6.2	2.3	2.2	3.6	0.8	19.4	28.2	12.0
2004.....	4.5	6.6	2.4	2.3	3.8	0.9	20.4	29.2	13.0
2003.....	4.7	7.1	2.4	2.5	4.2	0.9	21.3	31.3	12.8
2002.....	4.9	7.4	2.5	2.6	4.3	0.9	22.5	33.3	13.4
2001.....	5.0	7.5	2.5	2.6	4.4	0.9	22.8	33.8	13.4
2000.....	5.2	7.9	2.5	2.8	4.6	1.0	23.3	35.1	13.2
1999.....	5.3	8.2	2.5	2.9	4.9	1.0	23.6	36.1	13.1

Malaria (B50-B54)

2005.....	*	*	*	*	*	*	*	*	*
2004.....	*	*	*	*	*	*	*	*	*
2003.....	*	*	*	*	*	*	*	*	*
2002.....	*	*	*	*	*	*	*	*	*
2001.....	*	*	*	*	*	*	*	*	*
2000.....	*	*	*	*	*	*	*	*	*
1999.....	*	*	*	*	*	*	*	*	*

Other and unspecified infectious and parasitic diseases and their sequelae (A00,A05,A20-A36,A42-A44,A48-A49,A54-A79,A81-A82,A85.0-A85.1,A85.8,A86-B04,B06-B09,B25-B49,B55-B99)

2005.....	2.5	3.1	2.0	2.5	3.0	1.9	3.1	3.9	2.5
2004.....	2.5	3.1	2.0	2.4	3.0	2.0	3.2	4.1	2.6
2003.....	2.5	3.1	2.1	2.4	3.0	2.0	3.2	4.0	2.6
2002.....	2.3	2.9	1.9	2.2	2.7	1.8	3.2	4.1	2.5
2001.....	2.0	2.4	1.7	1.9	2.3	1.6	2.8	3.4	2.4
2000.....	2.0	2.4	1.7	1.9	2.3	1.6	2.9	3.7	2.4
1999.....	2.6	3.1	2.3	2.4	2.9	2.1	4.6	5.5	4.0

Malignant neoplasms (C00-C97)

2005.....	183.8	225.1	155.6	182.6	222.3	155.2	222.7	293.7	179.6
2004.....	185.8	227.7	157.4	184.4	224.4	157.0	227.2	301.2	182.5
2003.....	190.1	233.3	160.9	188.5	230.1	160.2	233.3	308.8	187.7
2002.....	193.5	238.9	163.1	191.7	235.2	162.4	238.8	319.6	190.3
2001.....	196.0	243.7	164.7	193.9	239.2	163.9	243.1	330.9	191.3
2000.....	199.6	248.9	167.6	197.2	243.9	166.9	248.5	340.3	193.8
1999.....	200.8	251.9	167.6	197.9	246.5	166.4	252.5	346.1	196.9

Worktable 293. Age-adjusted death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by race and sex: United States, 1999-2005

[Age-adjusted death rates per 100,000 U.S. standard population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992) and year	All races 1/			White 2/			Black 2/		
	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female
Malignant neoplasms of lip, oral cavity and pharynx (C00-C14)									
2005.....	2.5	3.9	1.4	2.4	3.6	1.4	3.5	6.5	1.4
2004.....	2.6	4.0	1.5	2.5	3.7	1.5	3.8	6.9	1.6
2003.....	2.6	4.0	1.5	2.5	3.8	1.4	3.8	6.9	1.6
2002.....	2.7	4.1	1.5	2.5	3.8	1.5	3.6	6.3	1.7
2001.....	2.7	4.1	1.6	2.6	3.9	1.5	3.9	6.7	1.8
2000.....	2.7	4.1	1.6	2.5	3.7	1.6	4.1	7.3	1.8
1999.....	2.7	4.1	1.6	2.5	3.8	1.5	4.3	7.5	2.1
Malignant neoplasm of esophagus (C15)									
2005.....	4.4	7.9	1.7	4.5	8.0	1.6	5.0	8.8	2.3
2004.....	4.3	7.7	1.7	4.3	7.7	1.6	5.4	9.0	2.8
2003.....	4.4	7.7	1.7	4.3	7.7	1.7	5.7	9.7	2.9
2002.....	4.4	7.8	1.7	4.3	7.7	1.6	5.9	10.1	2.9
2001.....	4.4	7.8	1.8	4.3	7.6	1.7	6.4	11.3	3.0
2000.....	4.4	7.7	1.8	4.3	7.6	1.7	6.5	11.0	3.3
1999.....	4.4	7.7	1.8	4.2	7.4	1.6	6.5	11.5	3.1
Malignant neoplasm of stomach (C16)									
2005.....	3.8	5.2	2.7	3.3	4.5	2.4	7.2	10.6	4.9
2004.....	4.0	5.5	2.8	3.5	4.9	2.5	7.5	11.1	5.2
2003.....	4.1	5.7	3.0	3.6	5.0	2.6	7.8	11.3	5.4
2002.....	4.2	5.9	3.0	3.7	5.2	2.5	8.4	11.9	6.1
2001.....	4.4	6.1	3.1	3.8	5.4	2.7	8.3	11.9	5.8
2000.....	4.6	6.4	3.3	4.0	5.6	2.8	9.0	12.7	6.5
1999.....	4.6	6.6	3.3	4.1	5.8	2.8	9.2	13.5	6.4
Malignant neoplasms of colon, rectum and anus (C18-C21)									
2005.....	17.5	20.9	14.8	16.9	20.3	14.3	24.8	30.2	21.2
2004.....	18.0	21.5	15.3	17.6	21.0	14.9	24.7	30.0	21.2
2003.....	19.1	22.9	16.2	18.6	22.4	15.7	26.4	31.9	22.8
2002.....	19.7	23.7	16.7	19.2	23.1	16.2	26.8	33.1	22.8
2001.....	20.1	24.2	17.2	19.6	23.7	16.6	27.6	33.1	24.1
2000.....	20.8	25.1	17.7	20.3	24.5	17.2	28.2	34.9	24.0
1999.....	20.9	25.3	17.8	20.4	24.8	17.3	28.4	34.2	24.7

Worktable 293. Age-adjusted death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by race and sex: United States, 1999-2005

[Age-adjusted death rates per 100,000 U.S. standard population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992) and year

	All races 1/			White 2/			Black 2/		
	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female
Malignant neoplasms of liver and intrahepatic bile ducts (C22)									
2005.....	5.2	7.7	3.2	4.8	7.1	3.0	7.0	11.3	3.8
2004.....	5.1	7.4	3.2	4.7	6.7	3.0	6.9	10.7	4.1
2003.....	5.0	7.4	3.1	4.6	6.7	2.8	6.6	10.2	4.0
2002.....	4.9	7.1	3.0	4.5	6.5	2.8	6.3	9.9	3.6
2001.....	4.7	6.9	3.0	4.3	6.3	2.7	6.2	9.1	4.1
2000.....	4.7	6.8	3.0	4.2	6.2	2.7	6.1	9.5	3.6
1999.....	4.5	6.5	3.0	4.2	5.9	2.7	5.9	8.7	3.8
Malignant neoplasm of pancreas (C25)									
2005.....	10.8	12.3	9.5	10.6	12.2	9.3	13.6	15.3	12.3
2004.....	10.6	12.3	9.3	10.4	12.2	9.1	13.6	15.4	12.1
2003.....	10.5	12.0	9.3	10.3	11.9	9.0	13.7	15.3	12.4
2002.....	10.5	12.1	9.2	10.3	12.0	9.0	13.6	15.3	12.3
2001.....	10.5	12.1	9.3	10.4	12.0	9.1	13.6	15.0	12.5
2000.....	10.6	12.2	9.3	10.4	12.0	9.0	14.1	15.8	12.7
1999.....	10.6	12.3	9.3	10.4	12.0	9.1	14.1	16.4	12.3
Malignant neoplasm of larynx (C32)									
2005.....	1.2	2.2	0.5	1.2	2.1	0.5	2.3	4.6	0.8
2004.....	1.2	2.2	0.4	1.1	2.1	0.4	2.2	4.6	0.7
2003.....	1.3	2.4	0.5	1.2	2.2	0.5	2.5	5.1	0.7
2002.....	1.3	2.3	0.5	1.2	2.1	0.5	2.5	4.8	0.8
2001.....	1.3	2.5	0.5	1.2	2.3	0.5	2.5	5.0	0.9
2000.....	1.4	2.5	0.5	1.3	2.3	0.5	2.8	5.5	0.9
1999.....	1.4	2.5	0.5	1.3	2.3	0.5	2.6	5.1	0.8
Malignant neoplasms of trachea, bronchus and lung (C33-C34)									
2005.....	52.6	69.0	40.5	53.1	68.7	41.5	58.4	86.4	40.0
2004.....	53.2	70.1	40.9	53.6	69.4	41.9	59.8	90.0	39.9
2003.....	54.1	71.7	41.3	54.5	71.1	42.3	60.8	92.4	40.2
2002.....	54.9	73.2	41.6	55.3	72.5	42.6	61.9	95.0	40.1
2001.....	55.3	75.2	41.0	55.6	74.2	42.1	62.5	99.1	38.7
2000.....	56.1	76.7	41.3	56.2	75.7	42.3	64.0	101.1	39.8
1999.....	55.5	76.9	40.2	55.4	75.6	41.0	64.8	103.0	39.9

Worktable 293. Age-adjusted death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by race and sex: United States, 1999-2005

[Age-adjusted death rates per 100,000 U.S. standard population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992) and year	All races 1/			White 2/			Black 2/		
	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female
Malignant melanoma of skin (C43)									
2005.....	2.7	4.0	1.8	3.1	4.5	2.1	0.4	0.5	0.4
2004.....	2.7	3.9	1.7	3.0	4.5	1.9	0.4	0.5	0.4
2003.....	2.7	3.9	1.8	3.0	4.4	2.0	0.4	0.4	0.4
2002.....	2.6	3.8	1.7	2.9	4.2	2.0	0.4	0.5	0.3
2001.....	2.7	3.9	1.8	3.0	4.3	2.0	0.4	0.3	0.4
2000.....	2.7	3.8	1.8	3.0	4.3	2.0	0.6	0.6	0.5
1999.....	2.6	3.8	1.7	3.0	4.2	2.0	0.4	0.5	0.3
Malignant neoplasm of breast (C50)									
2005.....	13.5	0.3	24.1	13.1	0.3	23.4	19.3	0.5	32.8
2004.....	13.8	0.3	24.4	13.4	0.3	23.9	19.0	0.5	32.2
2003.....	14.2	0.3	25.3	13.8	0.3	24.7	20.1	0.5	34.0
2002.....	14.5	0.3	25.6	14.1	0.3	25.0	20.2	0.6	34.0
2001.....	14.8	0.3	26.0	14.4	0.3	25.5	20.4	0.6	34.4
2000.....	15.3	0.4	26.8	14.9	0.4	26.3	20.5	0.6	34.5
1999.....	15.2	0.3	26.6	14.8	0.3	26.0	20.8	0.7	35.1
Malignant neoplasm of cervix uteri (C53)									
2005.....	1.3	...	2.4	1.1	...	2.2	2.5	...	4.4
2004.....	1.3	...	2.4	1.1	...	2.2	2.6	...	4.5
2003.....	1.3	...	2.5	1.2	...	2.2	2.7	...	4.7
2002.....	1.3	...	2.6	1.2	...	2.3	2.9	...	4.9
2001.....	1.4	...	2.7	1.3	...	2.5	2.8	...	4.8
2000.....	1.5	...	2.8	1.3	...	2.5	3.2	...	5.4
1999.....	1.5	...	2.8	1.4	...	2.6	3.1	...	5.4
Malignant neoplasms of corpus uteri and uterus, part unspecified (C54-C55)									
2005.....	2.3	...	4.1	2.2	...	3.8	4.2	...	7.1
2004.....	2.3	...	4.1	2.2	...	3.9	4.1	...	6.9
2003.....	2.3	...	4.1	2.2	...	3.9	4.2	...	7.0
2002.....	2.4	...	4.2	2.2	...	3.9	4.4	...	7.3
2001.....	2.4	...	4.2	2.3	...	4.0	4.2	...	7.0
2000.....	2.4	...	4.1	2.2	...	3.9	4.4	...	7.2
1999.....	2.4	...	4.1	2.2	...	3.9	4.2	...	6.8

Worktable 293. Age-adjusted death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by race and sex: United States, 1999-2005

[Age-adjusted death rates per 100,000 U.S. standard population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992) and year

	All races 1/			White 2/			Black 2/		
	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female
Malignant neoplasm of ovary (C56)									
2005.....	4.8	...	8.6	5.0	...	9.0	4.3	...	7.2
2004.....	4.9	...	8.8	5.1	...	9.1	4.5	...	7.6
2003.....	5.0	...	8.8	5.1	...	9.2	4.4	...	7.4
2002.....	5.1	...	9.0	5.3	...	9.4	4.4	...	7.4
2001.....	5.1	...	9.0	5.2	...	9.3	4.5	...	7.5
2000.....	5.1	...	8.9	5.3	...	9.3	4.3	...	7.2
1999.....	5.0	...	8.8	5.1	...	9.1	4.4	...	7.3
Malignant neoplasm of prostate (C61)									
2005.....	9.5	24.5	...	8.8	22.6	...	19.1	53.3	...
2004.....	9.8	25.4	...	9.0	23.4	...	19.7	55.5	...
2003.....	10.1	26.5	...	9.3	24.4	...	20.4	57.4	...
2002.....	10.6	27.9	...	9.8	25.7	...	22.0	62.0	...
2001.....	10.9	29.1	...	10.0	26.6	...	23.0	66.1	...
2000.....	11.3	30.4	...	10.3	27.8	...	23.9	68.1	...
1999.....	11.6	31.3	...	10.7	28.7	...	24.2	69.0	...
Malignant neoplasms of kidney and renal pelvis (C64-C65)									
2005.....	4.1	5.9	2.7	4.2	6.0	2.8	4.0	6.0	2.6
2004.....	4.1	5.9	2.7	4.2	6.0	2.8	4.0	5.8	2.7
2003.....	4.2	6.0	2.7	4.3	6.2	2.8	4.1	6.0	2.7
2002.....	4.2	6.1	2.8	4.3	6.3	2.8	4.0	5.8	2.8
2001.....	4.3	6.2	2.8	4.3	6.3	2.8	4.2	6.4	2.8
2000.....	4.2	6.2	2.8	4.3	6.3	2.8	4.1	6.3	2.8
1999.....	4.1	5.9	2.7	4.1	5.9	2.7	4.1	6.1	2.8
Malignant neoplasm of bladder (C67)									
2005.....	4.3	7.5	2.2	4.5	7.9	2.2	3.7	5.7	2.6
2004.....	4.4	7.6	2.2	4.5	8.0	2.3	3.6	4.9	2.7
2003.....	4.3	7.3	2.2	4.4	7.7	2.2	3.7	5.4	2.7
2002.....	4.4	7.5	2.4	4.5	7.8	2.4	3.9	5.5	3.0
2001.....	4.3	7.5	2.3	4.5	7.9	2.2	3.7	5.1	2.9
2000.....	4.3	7.6	2.3	4.5	7.9	2.3	3.7	5.6	2.7
1999.....	4.4	7.6	2.3	4.5	7.9	2.3	4.0	5.8	3.0

Worktable 293. Age-adjusted death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by race and sex: United States, 1999-2005

[Age-adjusted death rates per 100,000 U.S. standard population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992) and year	All races 1/			White 2/			Black 2/		
	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female
Malignant neoplasms of meninges, brain and other parts of central nervous system (C70-C72)									
2005.....	4.3	5.3	3.5	4.7	5.7	3.8	2.5	3.2	2.0
2004.....	4.3	5.2	3.5	4.7	5.6	3.8	2.5	3.0	2.0
2003.....	4.4	5.3	3.6	4.7	5.7	3.9	2.7	3.4	2.1
2002.....	4.4	5.5	3.6	4.8	5.8	3.9	2.6	3.2	2.2
2001.....	4.5	5.5	3.6	4.8	5.9	3.9	2.6	3.1	2.2
2000.....	4.5	5.6	3.7	4.9	6.0	3.9	2.7	3.3	2.3
1999.....	4.6	5.6	3.8	5.0	6.0	4.1	2.7	3.3	2.3
Malignant neoplasms of lymphoid, hematopoietic and related tissue (C81-C96)									
2005.....	18.2	23.5	14.3	18.5	24.0	14.5	17.6	22.5	14.3
2004.....	18.4	23.6	14.6	18.6	24.0	14.7	18.9	23.6	15.8
2003.....	19.1	24.5	15.1	19.4	25.1	15.2	18.8	22.9	16.1
2002.....	19.5	25.1	15.5	19.9	25.6	15.7	19.4	24.4	16.1
2001.....	20.0	25.5	16.0	20.2	25.9	16.2	20.0	25.3	16.5
2000.....	20.4	26.0	16.5	20.7	26.4	16.7	19.9	25.0	16.6
1999.....	20.6	26.1	16.7	20.8	26.5	16.8	20.3	25.6	16.8
Hodgkin's disease (C81)									
2005.....	0.4	0.5	0.4	0.4	0.5	0.4	0.4	0.5	0.3
2004.....	0.4	0.5	0.3	0.4	0.6	0.3	0.4	0.5	0.4
2003.....	0.5	0.5	0.4	0.5	0.6	0.4	0.4	0.5	0.3
2002.....	0.5	0.6	0.4	0.5	0.6	0.4	0.4	0.6	0.3
2001.....	0.5	0.6	0.4	0.5	0.6	0.4	0.4	0.5	0.4
2000.....	0.5	0.6	0.4	0.5	0.6	0.4	0.3	0.4	0.3
1999.....	0.5	0.6	0.4	0.5	0.7	0.4	0.4	0.5	0.3
Non-Hodgkin's lymphoma (C82-C85)									
2005.....	6.9	8.7	5.5	7.2	9.1	5.8	4.7	6.0	3.8
2004.....	7.0	8.8	5.7	7.3	9.2	6.0	5.0	6.2	4.1
2003.....	7.3	9.3	5.9	7.7	9.7	6.1	4.9	5.8	4.2
2002.....	7.6	9.6	6.2	7.9	10.0	6.5	5.4	6.8	4.4
2001.....	7.9	9.9	6.4	8.2	10.2	6.7	5.6	7.2	4.5
2000.....	8.2	10.2	6.7	8.6	10.7	7.1	5.2	6.6	4.2
1999.....	8.3	10.3	6.9	8.7	10.7	7.2	5.6	7.4	4.4

Worktable 293. Age-adjusted death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by race and sex: United States, 1999-2005

[Age-adjusted death rates per 100,000 U.S. standard population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992) and year

	All races 1/			White 2/			Black 2/		
	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female
Leukemia (C91-C95)									
2005.....	7.1	9.6	5.4	7.4	10.0	5.6	5.9	7.8	4.7
2004.....	7.2	9.6	5.5	7.4	9.9	5.6	6.7	8.7	5.3
2003.....	7.4	9.9	5.6	7.6	10.2	5.7	6.4	8.3	5.2
2002.....	7.5	10.0	5.7	7.7	10.3	5.8	6.6	8.6	5.3
2001.....	7.6	10.1	5.9	7.8	10.4	6.0	6.7	8.7	5.4
2000.....	7.7	10.3	5.9	7.9	10.5	6.1	6.7	9.0	5.2
1999.....	7.6	10.2	6.0	7.8	10.4	6.1	6.7	8.7	5.4

Multiple myeloma and immunoproliferative neoplasms (C88,C90)

2005.....	3.7	4.7	3.0	3.5	4.4	2.8	6.5	8.1	5.5
2004.....	3.7	4.6	3.1	3.4	4.3	2.8	6.9	8.3	6.0
2003.....	3.8	4.8	3.2	3.6	4.6	2.9	7.2	8.4	6.3
2002.....	4.0	5.0	3.3	3.7	4.8	3.0	6.9	8.3	6.1
2001.....	4.0	4.9	3.3	3.7	4.6	3.0	7.3	8.8	6.4
2000.....	4.0	4.9	3.4	3.7	4.6	3.1	7.7	8.9	6.8
1999.....	4.0	4.9	3.4	3.7	4.7	3.1	7.5	8.9	6.7

Other and unspecified malignant neoplasms of lymphoid, hematopoietic and related tissue (C96)

2005.....	0.0	0.0	0.0	0.0	0.0	*	*	*	*
2004.....	0.0	0.0	0.0	0.0	0.0	0.0	*	*	*
2003.....	0.0	0.0	0.0	0.0	0.0	0.0	*	*	*
2002.....	0.0	0.0	0.0	0.0	0.0	0.0	*	*	*
2001.....	0.0	0.0	0.0	0.0	0.0	0.0	*	*	*
2000.....	0.0	0.0	0.0	0.0	0.0	0.0	*	*	*
1999.....	0.0	0.0	0.0	0.0	0.0	0.0	*	*	*

All other and unspecified malignant neoplasms (C17,C23-C24,C26-C31,C37-C41,C44-C49,C51-C52,C57-C60,C62-C63,C66,C68-C69,C73-C80,C97)

2005.....	20.6	24.8	17.5	20.7	24.9	17.5	23.2	28.4	19.6
2004.....	20.8	25.0	17.6	20.7	25.0	17.6	23.9	29.6	19.9
2003.....	21.4	25.7	18.2	21.4	25.7	18.2	24.3	30.0	20.5
2002.....	21.8	26.5	18.5	21.8	26.4	18.4	25.7	31.4	21.9
2001.....	22.2	26.7	18.9	22.0	26.5	18.8	26.2	32.8	21.9
2000.....	22.8	27.5	19.4	22.6	27.3	19.3	26.4	33.0	22.1
1999.....	24.2	29.3	20.6	24.0	29.1	20.4	28.5	35.2	23.9

Worktable 293. Age-adjusted death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by race and sex: United States, 1999-2005

[Age-adjusted death rates per 100,000 U.S. standard population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992) and year	All races 1/			White 2/			Black 2/		
	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female
In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)									
2005.....	4.5	5.5	3.8	4.6	5.7	3.8	4.1	4.6	3.9
2004.....	4.5	5.7	3.8	4.6	5.8	3.9	4.4	5.2	3.9
2003.....	4.6	5.7	3.9	4.7	5.9	4.0	4.3	5.1	3.7
2002.....	4.6	5.7	3.9	4.7	5.8	4.0	4.4	5.1	4.1
2001.....	4.9	6.0	4.1	4.9	6.2	4.1	4.6	5.6	4.1
2000.....	4.9	6.0	4.2	5.0	6.1	4.3	4.6	5.4	4.1
1999.....	4.9	6.1	4.1	4.9	6.2	4.1	4.8	5.4	4.4
Anemias (D50-D64)									
2005.....	1.5	1.5	1.5	1.3	1.2	1.3	3.3	3.4	3.2
2004.....	1.5	1.5	1.5	1.3	1.3	1.3	3.3	3.5	3.1
2003.....	1.6	1.6	1.5	1.3	1.3	1.3	3.5	3.9	3.2
2002.....	1.6	1.6	1.6	1.3	1.3	1.4	3.7	3.9	3.5
2001.....	1.6	1.6	1.6	1.4	1.4	1.4	3.5	3.6	3.4
2000.....	1.6	1.6	1.6	1.4	1.4	1.4	3.4	3.4	3.5
1999.....	1.6	1.7	1.6	1.4	1.4	1.4	3.6	3.7	3.5
Diabetes mellitus (E10-E14)									
2005.....	24.6	28.4	21.6	22.5	26.5	19.3	46.9	50.8	43.8
2004.....	24.5	28.2	21.7	22.3	26.2	19.2	48.0	51.3	45.3
2003.....	25.3	28.9	22.5	23.0	27.0	19.9	49.2	50.7	47.5
2002.....	25.4	28.6	23.0	23.1	26.8	20.3	49.5	49.4	48.6
2001.....	25.3	28.1	23.1	23.0	26.2	20.5	49.2	49.9	48.1
2000.....	25.0	27.8	23.0	22.8	26.0	20.4	49.5	48.7	49.4
1999.....	25.0	27.8	23.0	22.6	25.8	20.2	49.7	49.4	49.5
Nutritional deficiencies (E40-E64)									
2005.....	1.0	1.0	1.1	1.0	0.9	1.0	1.7	1.6	1.7
2004.....	1.0	0.9	1.0	0.9	0.9	1.0	1.7	1.9	1.5
2003.....	1.1	1.1	1.2	1.1	1.0	1.1	1.9	2.1	1.7
2002.....	1.3	1.2	1.3	1.2	1.2	1.3	2.1	2.1	2.1
2001.....	1.3	1.2	1.4	1.2	1.1	1.3	2.1	2.4	1.8
2000.....	1.5	1.4	1.6	1.5	1.3	1.5	2.4	2.6	2.2
1999.....	1.6	1.5	1.6	1.5	1.4	1.5	2.7	3.1	2.4

Worktable 293. Age-adjusted death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by race and sex: United States, 1999-2005

[Age-adjusted death rates per 100,000 U.S. standard population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992) and year

	All races 1/			White 2/			Black 2/		
	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female
Malnutrition (E40-E46)									
2005.....	1.0	0.9	1.0	0.9	0.8	1.0	1.6	1.5	1.6
2004.....	0.9	0.9	1.0	0.9	0.8	0.9	1.6	1.8	1.5
2003.....	1.1	1.0	1.1	1.0	0.9	1.1	1.8	2.1	1.7
2002.....	1.2	1.1	1.2	1.1	1.1	1.2	2.0	2.0	2.0
2001.....	1.2	1.1	1.3	1.2	1.0	1.2	1.9	2.2	1.7
2000.....	1.4	1.3	1.5	1.4	1.2	1.4	2.2	2.5	2.1
1999.....	1.5	1.4	1.5	1.4	1.3	1.4	2.5	2.9	2.2
Other nutritional deficiencies (E50-E64)									
2005.....	0.1	0.0	0.1	0.1	0.0	0.1	*	*	*
2004.....	0.1	0.0	0.0	0.1	0.0	0.1	*	*	*
2003.....	0.1	0.1	0.0	0.1	0.1	0.1	*	*	*
2002.....	0.1	0.1	0.1	0.1	0.1	0.1	0.1	*	*
2001.....	0.1	0.1	0.1	0.1	0.1	0.1	0.1	*	*
2000.....	0.1	0.1	0.1	0.1	0.1	0.1	0.1	*	0.1
1999.....	0.1	0.1	0.1	0.1	0.1	0.1	0.2	*	0.2
Meningitis (G00,G03)									
2005.....	0.2	0.3	0.2	0.2	0.2	0.2	0.4	0.5	0.4
2004.....	0.3	0.3	0.2	0.2	0.2	0.2	0.5	0.5	0.4
2003.....	0.3	0.3	0.2	0.2	0.3	0.2	0.6	0.5	0.6
2002.....	0.2	0.3	0.2	0.2	0.3	0.2	0.5	0.7	0.4
2001.....	0.3	0.3	0.2	0.2	0.3	0.2	0.5	0.8	0.4
2000.....	0.3	0.3	0.2	0.2	0.3	0.2	0.6	0.8	0.6
1999.....	0.3	0.3	0.3	0.3	0.3	0.2	0.7	0.9	0.6
Parkinson's disease (G20-G21)									
2005.....	6.4	9.8	4.4	6.8	10.4	4.6	3.0	4.9	1.9
2004.....	6.1	9.3	4.1	6.4	9.8	4.4	2.7	4.3	1.9
2003.....	6.2	9.3	4.2	6.5	9.8	4.5	2.7	4.4	1.8
2002.....	5.9	9.0	4.1	6.3	9.5	4.3	2.5	4.0	1.7
2001.....	5.9	9.1	4.0	6.2	9.7	4.3	2.5	3.9	1.8
2000.....	5.7	8.8	3.9	6.0	9.3	4.2	2.5	4.2	1.6
1999.....	5.4	8.4	3.7	5.7	8.9	3.9	2.4	3.7	1.7

Worktable 293. Age-adjusted death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by race and sex: United States, 1999-2005

[Age-adjusted death rates per 100,000 U.S. standard population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992) and year	All races 1/			White 2/			Black 2/		
	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female
Alzheimer's disease (G30)									
2005.....	22.9	18.5	25.1	23.7	19.0	26.1	19.4	16.4	20.7
2004.....	21.8	17.7	23.8	22.5	18.3	24.7	18.5	15.2	19.9
2003.....	21.4	17.5	23.3	22.2	18.2	24.3	16.6	14.7	17.3
2002.....	20.2	16.6	22.0	21.1	17.3	22.9	15.2	13.2	15.9
2001.....	19.1	16.1	20.5	19.8	16.7	21.3	14.6	13.1	15.1
2000.....	18.1	15.2	19.3	18.8	15.8	20.1	13.0	11.4	13.7
1999.....	16.5	14.4	17.4	17.2	15.0	18.1	11.3	10.2	11.8
Major cardiovascular diseases (I00-I78)									
2005.....	277.3	329.5	235.6	271.0	323.2	229.0	367.9	435.3	317.4
2004.....	286.5	340.0	243.8	279.8	333.6	236.7	381.9	451.1	331.0
2003.....	306.1	363.4	260.7	298.9	356.4	253.0	406.1	479.1	352.4
2002.....	317.4	377.2	270.5	310.4	370.7	262.7	416.1	487.4	363.8
2001.....	326.5	388.1	278.8	319.2	381.1	270.8	427.5	505.3	372.4
2000.....	339.7	407.3	288.4	332.5	400.7	280.3	438.7	517.6	383.3
1999.....	349.3	419.0	296.5	341.9	411.8	288.4	447.6	532.2	388.6
Diseases of heart (I00-I09,I11,I13,I20-I51)									
2005.....	211.1	260.9	172.3	207.8	258.0	168.2	271.3	329.8	228.3
2004.....	217.0	267.9	177.3	213.3	264.6	172.9	280.6	342.1	236.5
2003.....	232.3	286.6	190.3	228.2	282.9	185.4	300.2	364.3	253.8
2002.....	240.8	297.4	197.2	236.7	294.1	192.1	308.4	371.0	263.2
2001.....	247.8	305.4	203.9	243.5	301.8	198.7	316.9	384.5	269.8
2000.....	257.6	320.0	210.9	253.4	316.7	205.6	324.8	392.5	277.6
1999.....	266.5	331.0	218.1	261.9	327.1	212.8	334.3	407.2	283.7
Acute rheumatic fever and chronic rheumatic heart diseases (I00-I09)									
2005.....	1.1	0.8	1.3	1.1	0.8	1.3	0.8	0.7	0.8
2004.....	1.1	0.8	1.3	1.1	0.8	1.3	0.8	0.6	1.0
2003.....	1.2	1.0	1.4	1.3	1.0	1.4	1.1	0.9	1.2
2002.....	1.2	0.9	1.5	1.3	0.9	1.5	0.9	0.8	1.0
2001.....	1.2	0.9	1.5	1.2	0.9	1.5	1.1	0.7	1.3
2000.....	1.3	1.0	1.5	1.3	1.0	1.6	1.0	0.7	1.2
1999.....	1.3	0.9	1.6	1.4	0.9	1.6	1.0	0.8	1.2

Worktable 293. Age-adjusted death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by race and sex: United States, 1999-2005

[Age-adjusted death rates per 100,000 U.S. standard population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992) and year	All races 1/			White 2/			Black 2/		
	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female
Hypertensive heart disease (I11)									
2005.....	9.4	10.1	8.5	7.9	8.3	7.2	24.1	28.4	20.4
2004.....	9.5	10.1	8.5	8.0	8.4	7.2	24.2	28.4	20.6
2003.....	9.6	10.1	8.8	8.0	8.3	7.3	25.0	28.7	21.9
2002.....	9.1	9.5	8.3	7.6	7.8	7.0	24.4	28.0	21.2
2001.....	8.7	8.9	8.1	7.3	7.3	6.8	23.2	26.3	20.5
2000.....	8.6	8.7	8.1	7.1	7.1	6.7	23.7	26.3	21.2
1999.....	8.3	8.3	7.9	6.9	6.7	6.6	22.9	25.7	20.5
Hypertensive heart and renal disease (I13)									
2005.....	1.0	1.1	0.9	0.8	0.9	0.8	3.2	3.8	2.8
2004.....	1.0	1.1	0.9	0.8	0.8	0.7	3.2	3.6	2.9
2003.....	1.1	1.1	1.0	0.8	0.9	0.7	3.4	3.7	3.1
2002.....	1.0	1.0	0.9	0.7	0.8	0.7	3.5	3.9	3.2
2001.....	1.0	1.1	1.0	0.8	0.8	0.7	3.3	3.6	3.0
2000.....	1.0	1.1	0.9	0.8	0.9	0.7	3.3	3.8	3.1
1999.....	1.2	1.4	1.1	0.9	1.0	0.9	4.2	4.5	3.9
Ischemic heart diseases (I20-I25)									
2005.....	144.4	187.4	111.7	143.8	187.7	110.0	171.3	213.9	140.9
2004.....	150.2	194.1	116.7	149.2	194.2	114.7	179.8	223.9	148.7
2003.....	162.9	209.9	127.2	161.7	209.6	125.0	195.0	243.3	160.9
2002.....	170.8	220.4	133.6	169.8	220.5	131.2	203.0	250.6	169.7
2001.....	177.8	228.5	139.9	176.5	228.4	137.4	211.6	262.0	176.7
2000.....	186.8	241.4	146.5	185.6	241.6	143.9	218.3	269.5	183.2
1999.....	194.6	251.2	152.9	193.4	251.2	150.5	224.9	278.7	188.0
Acute myocardial infarction (I21-I22)									
2005.....	49.1	63.6	37.8	49.1	64.1	37.2	57.6	69.7	48.8
2004.....	52.2	67.8	40.2	52.2	68.3	39.5	62.0	74.8	52.7
2003.....	57.9	75.0	44.8	57.8	75.3	44.0	68.6	84.3	57.6
2002.....	62.1	80.6	48.1	61.9	81.0	47.1	73.9	89.5	63.2
2001.....	65.4	84.9	50.9	65.1	85.4	49.9	76.8	93.1	66.0
2000.....	69.9	91.3	54.1	69.6	91.8	53.1	81.0	98.9	69.2
1999.....	73.2	96.1	56.3	73.1	96.8	55.4	83.1	102.1	70.5

Worktable 293. Age-adjusted death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by race and sex: United States, 1999-2005

[Age-adjusted death rates per 100,000 U.S. standard population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992) and year

	All races 1/			White 2/			Black 2/		
	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female
Other acute ischemic heart diseases (I24)									
2005.....	1.2	1.4	0.9	1.1	1.3	0.9	2.0	2.6	1.5
2004.....	0.8	1.0	0.6	0.8	1.0	0.6	1.4	2.1	1.0
2003.....	1.1	1.3	0.9	1.0	1.2	0.8	1.6	2.4	1.1
2002.....	1.2	1.5	0.9	1.1	1.4	0.9	1.7	2.3	1.3
2001.....	1.2	1.5	0.9	1.1	1.4	0.9	1.8	2.3	1.3
2000.....	1.2	1.5	0.9	1.2	1.5	0.9	1.8	2.5	1.3
1999.....	1.2	1.5	0.9	1.1	1.4	0.9	1.8	2.4	1.3
Other forms of chronic ischemic heart disease (I20,I25)									
2005.....	94.2	122.4	73.0	93.6	122.3	72.0	111.8	141.6	90.5
2004.....	97.1	125.3	75.8	96.3	125.0	74.6	116.4	146.9	95.0
2003.....	103.8	133.5	81.5	102.9	133.0	80.1	124.8	156.7	102.2
2002.....	107.6	138.3	84.5	106.8	138.0	83.2	127.5	158.9	105.2
2001.....	111.2	142.1	88.1	110.2	141.6	86.7	133.0	166.7	109.3
2000.....	115.7	148.6	91.5	114.8	148.3	90.0	135.4	168.1	112.6
1999.....	120.2	153.6	95.6	119.2	153.0	94.2	140.0	174.2	116.2
Atherosclerotic cardiovascular disease, so described (I25.0)									
2005.....	20.3	26.6	15.0	19.1	25.0	14.0	34.5	46.9	25.3
2004.....	21.1	27.5	15.7	19.9	25.9	14.7	36.1	48.9	27.0
2003.....	22.7	29.6	17.1	21.4	27.8	16.1	38.2	52.7	28.0
2002.....	23.5	30.4	17.8	22.1	28.5	16.7	40.3	54.8	30.0
2001.....	24.1	30.8	18.5	22.5	28.8	17.3	41.8	56.0	31.7
2000.....	25.2	32.3	19.4	23.6	30.2	18.2	43.5	58.7	32.9
1999.....	26.4	33.5	20.6	24.7	31.2	19.3	45.8	61.5	34.9
All other forms of chronic ischemic heart disease (I20,I25.1-I25.9)									
2005.....	73.9	95.8	58.0	74.5	97.3	57.9	77.3	94.7	65.2
2004.....	76.0	97.8	60.1	76.5	99.1	59.9	80.3	98.0	68.0
2003.....	81.1	104.0	64.4	81.5	105.2	64.0	86.6	104.0	74.2
2002.....	84.1	108.0	66.7	84.7	109.5	66.4	87.2	104.1	75.2
2001.....	87.1	111.3	69.6	87.6	112.8	69.4	91.1	110.6	77.6
2000.....	90.5	116.3	72.0	91.2	118.1	71.8	91.9	109.5	79.7
1999.....	93.8	120.1	75.0	94.5	121.8	74.9	94.1	112.7	81.3

Worktable 293. Age-adjusted death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by race and sex: United States, 1999-2005

[Age-adjusted death rates per 100,000 U.S. standard population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992) and year	All races 1/			White 2/			Black 2/		
	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female
Other heart diseases (I26-I51)									
2005.....	55.2	61.5	49.9	54.1	60.2	49.0	71.9	83.0	63.4
2004.....	55.2	61.7	49.9	54.1	60.3	49.0	72.5	85.6	63.2
2003.....	57.6	64.5	52.0	56.4	63.1	50.9	75.7	87.7	66.7
2002.....	58.5	65.5	52.8	57.3	64.1	51.7	76.6	87.8	68.1
2001.....	59.1	66.1	53.4	57.7	64.4	52.3	77.8	91.8	68.2
2000.....	60.0	67.9	53.8	58.6	66.3	52.6	78.4	92.2	69.0
1999.....	60.9	69.2	54.5	59.3	67.2	53.1	81.2	97.5	70.1
Acute and subacute endocarditis (I33)									
2005.....	0.4	0.5	0.3	0.4	0.5	0.3	0.7	0.9	0.6
2004.....	0.4	0.5	0.3	0.4	0.5	0.3	0.7	0.8	0.6
2003.....	0.4	0.5	0.3	0.4	0.5	0.3	0.8	0.9	0.6
2002.....	0.4	0.5	0.3	0.4	0.5	0.3	0.7	0.8	0.6
2001.....	0.4	0.5	0.3	0.4	0.5	0.3	0.8	1.1	0.6
2000.....	0.4	0.5	0.3	0.4	0.5	0.3	0.7	0.9	0.7
1999.....	0.4	0.5	0.3	0.4	0.5	0.3	0.8	0.9	0.6
Diseases of pericardium and acute myocarditis (I30-I31,I40)									
2005.....	0.3	0.3	0.3	0.3	0.3	0.2	0.5	0.5	0.5
2004.....	0.3	0.3	0.2	0.2	0.3	0.2	0.5	0.6	0.4
2003.....	0.3	0.3	0.3	0.3	0.3	0.2	0.5	0.5	0.5
2002.....	0.3	0.3	0.3	0.3	0.3	0.3	0.5	0.5	0.4
2001.....	0.3	0.3	0.3	0.3	0.3	0.3	0.5	0.5	0.4
2000.....	0.3	0.3	0.2	0.3	0.3	0.2	0.4	0.5	0.4
1999.....	0.3	0.4	0.3	0.3	0.3	0.3	0.5	0.5	0.5
Heart failure (I50)									
2005.....	18.9	19.9	18.0	18.9	20.0	18.1	21.7	23.0	20.4
2004.....	18.9	19.9	18.0	19.0	20.0	18.1	20.9	23.4	19.3
2003.....	19.4	20.6	18.4	19.5	20.7	18.4	21.8	23.6	20.5
2002.....	19.5	20.4	18.6	19.5	20.5	18.6	21.8	23.3	20.5
2001.....	20.2	21.2	19.2	20.2	21.2	19.3	22.1	24.4	20.6
2000.....	20.3	21.5	19.2	20.3	21.6	19.3	22.0	24.2	20.7
1999.....	20.3	21.4	19.3	20.3	21.4	19.3	22.5	25.3	20.9

Worktable 293. Age-adjusted death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by race and sex: United States, 1999-2005

[Age-adjusted death rates per 100,000 U.S. standard population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992) and year

	All races 1/			White 2/			Black 2/		
	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female
All other forms of heart disease (I26-I28,I34-I38,I42-I49,I51)									
2005.....	35.6	40.7	31.3	34.5	39.5	30.4	49.0	58.6	41.9
2004.....	35.7	41.0	31.4	34.5	39.5	30.4	50.4	60.9	43.0
2003.....	37.5	43.1	33.0	36.2	41.6	31.9	52.6	62.8	45.1
2002.....	38.4	44.2	33.7	37.1	42.8	32.6	53.6	63.1	46.6
2001.....	38.2	44.1	33.6	36.8	42.4	32.4	54.4	65.8	46.5
2000.....	39.0	45.5	34.0	37.7	43.9	32.8	55.2	66.6	47.2
1999.....	39.9	46.9	34.6	38.4	45.0	33.3	57.4	70.8	48.2
Essential (primary) hypertension and hypertensive renal disease (I10,I12)									
2005.....	8.0	7.8	8.0	7.0	6.6	7.1	18.4	19.9	17.1
2004.....	7.7	7.5	7.6	6.6	6.5	6.6	18.2	19.0	17.3
2003.....	7.4	7.1	7.5	6.4	6.1	6.5	17.6	18.6	16.6
2002.....	7.0	6.8	7.0	6.0	5.8	6.0	16.9	17.7	16.1
2001.....	6.8	6.7	6.8	5.8	5.6	5.9	16.6	17.9	15.4
2000.....	6.5	6.5	6.5	5.6	5.5	5.5	16.3	17.1	15.6
1999.....	6.2	6.2	6.1	5.3	5.2	5.2	16.0	17.3	15.0
Cerebrovascular diseases (I60-I69)									
2005.....	46.6	46.9	45.6	44.7	44.7	44.0	65.2	70.5	60.7
2004.....	50.0	50.4	48.9	48.0	48.1	47.2	69.9	74.9	65.5
2003.....	53.5	54.1	52.3	51.4	51.7	50.5	74.3	79.5	69.8
2002.....	56.2	56.5	55.2	54.2	54.2	53.4	76.3	81.7	71.8
2001.....	57.9	59.0	56.4	55.8	56.5	54.5	78.8	85.4	73.7
2000.....	60.9	62.4	59.1	58.8	59.8	57.3	81.9	89.6	76.2
1999.....	61.6	63.2	59.8	59.6	60.8	58.0	81.8	89.6	76.2
Atherosclerosis (I70)									
2005.....	3.8	3.9	3.7	3.9	4.0	3.7	3.4	3.4	3.3
2004.....	3.9	4.1	3.7	4.0	4.2	3.8	3.5	3.5	3.3
2003.....	4.4	4.5	4.3	4.5	4.5	4.4	4.2	4.7	3.8
2002.....	4.7	4.9	4.6	4.8	5.0	4.7	4.4	4.9	4.0
2001.....	5.0	5.1	4.8	5.1	5.2	5.0	4.3	4.6	4.1
2000.....	5.2	5.5	5.0	5.3	5.6	5.1	4.8	5.4	4.4
1999.....	5.5	5.7	5.3	5.6	5.8	5.4	5.0	5.8	4.5

Worktable 293. Age-adjusted death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by race and sex: United States, 1999-2005

[Age-adjusted death rates per 100,000 U.S. standard population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992) and year	All races 1/			White 2/			Black 2/		
	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female
Other diseases of circulatory system (I71-I78)									
2005.....	7.8	10.0	6.1	7.6	10.0	5.9	9.6	11.7	8.0
2004.....	7.9	10.2	6.3	7.9	10.2	6.1	9.7	11.6	8.3
2003.....	8.4	11.2	6.4	8.4	11.3	6.3	9.8	11.8	8.3
2002.....	8.7	11.5	6.6	8.6	11.6	6.5	10.0	12.0	8.6
2001.....	9.0	12.0	6.9	8.9	12.0	6.7	10.8	12.8	9.4
2000.....	9.4	12.9	7.1	9.3	13.0	6.8	10.9	13.0	9.5
1999.....	9.5	12.8	7.1	9.4	13.0	7.0	10.5	12.4	9.2
Aortic aneurysm and dissection (I71)									
2005.....	4.6	6.5	3.1	4.6	6.6	3.1	4.6	6.0	3.6
2004.....	4.6	6.7	3.1	4.7	6.8	3.2	4.5	6.1	3.3
2003.....	5.1	7.5	3.3	5.1	7.7	3.3	4.6	6.1	3.5
2002.....	5.2	7.7	3.3	5.2	7.9	3.4	4.6	6.1	3.5
2001.....	5.4	8.0	3.6	5.5	8.2	3.6	5.0	6.4	3.9
2000.....	5.7	8.7	3.7	5.8	9.0	3.6	5.2	6.8	4.1
1999.....	5.8	8.8	3.7	5.9	9.1	3.7	4.9	6.3	3.9
Other diseases of arteries, arterioles and capillaries (I72-I78)									
2005.....	3.2	3.5	2.9	3.1	3.4	2.8	5.0	5.6	4.5
2004.....	3.3	3.5	3.1	3.2	3.4	3.0	5.2	5.6	5.0
2003.....	3.3	3.7	3.1	3.2	3.6	3.0	5.2	5.7	4.9
2002.....	3.5	3.8	3.3	3.4	3.7	3.1	5.4	6.0	5.0
2001.....	3.6	3.9	3.3	3.4	3.8	3.2	5.9	6.4	5.5
2000.....	3.7	4.2	3.4	3.6	4.0	3.2	5.7	6.2	5.3
1999.....	3.7	4.0	3.4	3.5	3.9	3.3	5.6	6.0	5.3
Other disorders of circulatory system (I80-I99)									
2005.....	1.6	1.6	1.5	1.5	1.5	1.4	2.7	3.1	2.3
2004.....	1.5	1.6	1.5	1.5	1.5	1.4	2.7	2.9	2.6
2003.....	1.6	1.6	1.6	1.5	1.4	1.5	2.9	3.2	2.8
2002.....	1.6	1.6	1.6	1.5	1.5	1.6	2.9	3.2	2.6
2001.....	1.6	1.7	1.6	1.5	1.6	1.5	2.7	3.1	2.5
2000.....	1.7	1.7	1.6	1.6	1.6	1.5	2.9	3.0	2.8
1999.....	1.5	1.5	1.4	1.4	1.4	1.3	2.4	2.6	2.2

[Age-adjusted death rates per 100,000 U.S. standard population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Both sexes	All races 1/		Both sexes	White 2/		Both sexes	Black 2/	
	Male	Female		Male	Female		Male	Female

2005.....	20.3	23.9	17.9	20.2	23.6	18.0	21.7	26.9	18.4
2004.....	19.8	23.7	17.3	19.6	23.3	17.2	22.3	29.1	18.2
2003.....	22.0	26.1	19.4	21.9	25.7	19.5	23.3	30.9	18.7
2002.....	22.6	27.0	19.9	22.6	26.7	19.9	24.0	30.8	20.0
2001.....	22.0	26.6	19.2	21.7	26.0	19.1	24.1	32.3	19.4
2000.....	23.7	28.9	20.7	23.5	28.4	20.7	25.6	33.9	20.8
1999.....	23.5	28.5	20.6	23.3	28.1	20.6	25.4	33.4	20.8

2005	0.6	0.6	0.6	0.6	0.6	0.6	0.2	0.3	0.2
2004	0.4	0.4	0.3	0.4	0.4	0.4	0.2	0.2	0.1
2003	0.6	0.6	0.6	0.6	0.6	0.6	0.3	0.3	0.3
2002	0.2	0.3	0.2	0.2	0.3	0.2	0.1	*	*
2001	0.1	0.1	0.1	0.1	0.1	0.1	0.1	*	*
2000	0.6	0.7	0.6	0.6	0.7	0.6	0.3	0.4	0.3
1999	0.6	0.6	0.6	0.6	0.6	0.6	0.3	0.3	0.3

2005.....	19.7	23.3	17.3	19.6	23.0	17.3	21.5	26.6	18.2
2004.....	19.4	23.3	16.9	19.2	22.9	16.8	22.1	28.9	18.0
2003.....	21.4	25.5	18.8	21.3	25.0	18.9	23.0	30.6	18.4
2002.....	22.4	26.7	19.6	22.3	26.4	19.7	23.9	30.7	19.9
2001.....	21.9	26.5	19.1	21.6	25.9	19.0	24.1	32.2	19.4
2000.....	23.1	28.2	20.1	22.9	27.7	20.0	25.3	33.5	20.5
1999.....	22.9	27.9	20.0	22.7	27.5	19.9	25.1	33.2	20.5

[illegible]

Worktable 293. Age-adjusted death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by race and sex: United States, 1999-2005

[Age-adjusted death rates per 100,000 U.S. standard population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992) and year	All races 1/			White 2/			Black 2/		
	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female
Acute bronchitis and bronchiolitis (J20-J21)									
2005.....	0.1	0.1	0.1	0.1	0.1	0.1	0.1	*	*
2004.....	0.1	0.1	0.1	0.1	0.1	0.1	0.1	*	*
2003.....	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	*
2002.....	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	*
2001.....	0.1	0.1	0.1	0.1	0.1	0.1	0.1	*	*
2000.....	0.1	0.1	0.1	0.1	0.1	0.1	0.1	*	0.1
1999.....	0.1	0.1	0.1	0.1	0.1	0.1	0.2	0.2	0.2
Unspecified acute lower respiratory infection (J22)									
2005.....	0.0	0.0	0.0	0.0	0.0	0.0	*	*	*
2004.....	0.0	0.0	0.0	0.0	0.0	0.0	*	*	*
2003.....	0.0	0.0	0.0	0.0	0.0	0.0	*	*	*
2002.....	0.0	0.0	0.0	0.0	0.0	0.0	*	*	*
2001.....	0.0	0.0	0.0	0.0	0.0	0.0	*	*	*
2000.....	0.0	0.0	0.0	0.1	0.0	0.0	*	*	*
1999.....	0.1	0.0	0.1	0.1	0.0	0.1	*	*	*
Chronic lower respiratory diseases (J40-J47)									
2005.....	43.2	51.2	38.1	45.4	52.8	40.7	30.6	44.1	22.8
2004.....	41.1	49.5	36.0	43.2	51.1	38.4	28.2	40.9	20.9
2003.....	43.3	52.3	37.8	45.4	53.8	40.3	30.1	44.4	22.0
2002.....	43.5	53.5	37.4	45.4	54.9	39.7	31.2	46.3	22.6
2001.....	43.7	54.0	37.6	45.6	55.5	39.8	30.9	46.3	22.4
2000.....	44.2	55.8	37.4	46.0	57.2	39.5	31.6	47.5	22.7
1999.....	45.4	58.7	37.7	47.1	60.0	39.7	33.5	51.5	23.4
Bronchitis, chronic and unspecified (J40-J42)									
2005.....	0.3	0.3	0.3	0.3	0.3	0.3	0.2	0.3	0.1
2004.....	0.2	0.3	0.2	0.3	0.3	0.2	0.2	0.3	0.2
2003.....	0.3	0.3	0.3	0.3	0.3	0.3	0.3	0.3	0.3
2002.....	0.3	0.4	0.3	0.3	0.4	0.3	0.2	0.3	0.2
2001.....	0.3	0.4	0.3	0.3	0.4	0.3	0.2	0.3	0.2
2000.....	0.4	0.5	0.4	0.4	0.5	0.4	0.3	0.3	0.3
1999.....	0.4	0.5	0.4	0.4	0.5	0.4	0.4	0.5	0.3

Worktable 293. Age-adjusted death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by race and sex: United States, 1999-2005

[Age-adjusted death rates per 100,000 U.S. standard population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992) and year

	All races 1/			White 2/			Black 2/		
	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female
Emphysema (J43)									
2005.....	4.6	5.8	3.8	4.9	6.0	4.1	3.1	4.9	1.9
2004.....	4.6	5.8	3.8	4.9	6.1	4.1	2.7	4.6	1.6
2003.....	5.1	6.3	4.3	5.4	6.6	4.6	3.2	5.3	2.0
2002.....	5.4	7.0	4.4	5.7	7.2	4.8	3.4	5.8	2.0
2001.....	5.8	7.4	4.7	6.1	7.8	5.1	3.3	5.3	2.2
2000.....	6.0	7.8	4.9	6.4	8.1	5.3	3.8	6.2	2.4
1999.....	6.5	8.6	5.2	6.9	8.9	5.6	4.0	6.7	2.3
Asthma (J45-J46)									
2005.....	1.3	1.0	1.5	1.0	0.7	1.3	3.0	2.7	3.3
2004.....	1.3	1.0	1.4	1.0	0.8	1.2	3.0	3.0	3.1
2003.....	1.4	1.1	1.6	1.1	0.9	1.3	3.2	2.9	3.3
2002.....	1.5	1.2	1.7	1.2	0.9	1.4	3.4	3.3	3.4
2001.....	1.5	1.2	1.7	1.2	0.9	1.5	3.6	3.2	3.8
2000.....	1.6	1.3	1.8	1.3	1.0	1.5	3.9	3.5	4.1
1999.....	1.7	1.3	2.0	1.4	1.0	1.7	3.9	3.5	4.1
Other chronic lower respiratory diseases (J44,J47)									
2005.....	37.0	44.1	32.5	39.2	45.8	35.0	24.3	36.3	17.5
2004.....	35.0	42.3	30.5	37.0	43.9	32.8	22.2	33.0	16.0
2003.....	36.5	44.5	31.7	38.6	46.0	34.0	23.4	36.0	16.3
2002.....	36.2	44.9	31.1	38.2	46.4	33.2	24.1	36.9	16.9
2001.....	36.1	45.0	30.8	37.9	46.4	32.9	23.7	37.5	16.2
2000.....	36.1	46.1	30.3	37.8	47.5	32.3	23.7	37.5	15.9
1999.....	36.8	48.3	30.2	38.5	49.6	32.0	25.3	40.7	16.7
Pneumoconioses and chemical effects (J60-J66,J68)									
2005.....	0.3	0.8	0.0	0.4	0.9	0.0	0.2	0.5	*
2004.....	0.3	0.9	0.0	0.4	0.9	0.0	0.2	0.6	*
2003.....	0.4	1.0	0.0	0.4	1.0	0.0	0.2	0.5	*
2002.....	0.4	1.0	0.0	0.4	1.0	0.0	0.2	0.6	*
2001.....	0.4	1.0	0.0	0.4	1.1	0.0	0.2	0.7	*
2000.....	0.4	1.1	0.0	0.4	1.1	0.0	0.2	0.7	*
1999.....	0.4	1.0	0.0	0.4	1.1	0.0	0.3	0.8	*

Worktable 293. Age-adjusted death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by race and sex: United States, 1999-2005

[Age-adjusted death rates per 100,000 U.S. standard population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992) and year	All races 1/			White 2/			Black 2/		
	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female
Pneumonitis due to solids and liquids (J69)									
2005.....	5.6	7.6	4.4	5.6	7.6	4.4	6.4	9.0	4.9
2004.....	5.6	7.7	4.3	5.6	7.7	4.3	6.0	8.6	4.6
2003.....	5.9	8.0	4.6	5.8	8.0	4.6	6.7	9.1	5.3
2002.....	6.1	8.4	4.7	6.1	8.5	4.7	6.6	9.2	5.1
2001.....	6.1	8.5	4.8	6.1	8.5	4.8	7.0	9.4	5.7
2000.....	6.1	8.6	4.6	6.0	8.6	4.6	6.8	9.4	5.2
1999.....	5.6	8.1	4.3	5.6	8.0	4.3	6.8	9.9	5.1
Other diseases of respiratory system (J00-J06, J30-J39, J67, J70-J98)									
2005.....	8.9	10.6	7.7	9.0	10.7	7.8	8.7	10.7	7.5
2004.....	9.2	11.1	7.8	9.3	11.3	7.9	8.6	10.4	7.4
2003.....	8.7	10.5	7.5	8.8	10.6	7.6	8.1	10.0	6.9
2002.....	8.7	10.4	7.6	8.8	10.5	7.6	8.4	9.9	7.4
2001.....	9.3	11.0	8.1	9.3	11.1	8.1	9.4	10.7	8.6
2000.....	9.2	11.3	7.9	9.3	11.3	8.0	9.2	11.2	8.0
1999.....	9.2	11.2	7.8	9.2	11.2	7.8	9.4	11.2	8.4
Peptic ulcer (K25-K28)									
2005.....	1.1	1.4	0.9	1.1	1.3	0.9	1.3	1.8	0.9
2004.....	1.2	1.4	1.1	1.2	1.4	1.1	1.3	1.7	1.1
2003.....	1.3	1.6	1.1	1.3	1.5	1.2	1.4	2.1	1.0
2002.....	1.4	1.7	1.2	1.4	1.7	1.2	1.4	2.0	1.1
2001.....	1.6	1.8	1.4	1.6	1.8	1.4	1.6	2.3	1.1
2000.....	1.6	2.0	1.4	1.6	2.0	1.4	1.7	2.3	1.3
1999.....	1.7	2.0	1.4	1.7	2.0	1.4	1.9	2.4	1.4
Diseases of appendix (K35-K38)									
2005.....	0.1	0.2	0.1	0.1	0.2	0.1	0.2	0.2	0.1
2004.....	0.1	0.2	0.1	0.1	0.2	0.1	0.2	0.3	0.1
2003.....	0.1	0.2	0.1	0.1	0.2	0.1	0.2	0.3	0.2
2002.....	0.2	0.2	0.1	0.1	0.2	0.1	0.2	0.4	0.1
2001.....	0.1	0.2	0.1	0.1	0.2	0.1	0.2	0.4	0.2
2000.....	0.1	0.2	0.1	0.1	0.2	0.1	0.3	0.4	0.2
1999.....	0.1	0.2	0.1	0.1	0.2	0.1	0.2	0.4	0.1

Worktable 293. Age-adjusted death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by race and sex: United States, 1999-2005

[Age-adjusted death rates per 100,000 U.S. standard population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992) and year	All races 1/			White 2/			Black 2/		
	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female
Hernia (K40-K46)									
2005.....	0.5	0.5	0.5	0.5	0.5	0.5	0.5	0.6	0.5
2004.....	0.5	0.5	0.5	0.5	0.5	0.5	0.5	0.6	0.5
2003.....	0.6	0.6	0.5	0.6	0.6	0.5	0.6	0.7	0.6
2002.....	0.5	0.5	0.6	0.6	0.5	0.6	0.5	0.6	0.5
2001.....	0.5	0.5	0.5	0.5	0.5	0.5	0.6	0.8	0.5
2000.....	0.5	0.5	0.5	0.6	0.6	0.6	0.6	0.6	0.6
1999.....	0.5	0.6	0.5	0.5	0.6	0.5	0.6	0.6	0.6
Chronic liver disease and cirrhosis (K70,K73-K74)									
2005.....	9.0	12.4	5.8	9.2	12.7	6.0	7.7	11.5	4.8
2004.....	9.0	12.5	5.8	9.2	12.8	6.0	7.9	11.7	4.9
2003.....	9.3	13.0	6.0	9.5	13.3	6.1	8.4	12.4	5.2
2002.....	9.4	12.9	6.3	9.6	13.2	6.3	8.5	12.0	5.7
2001.....	9.5	13.2	6.2	9.6	13.3	6.3	9.3	13.8	5.6
2000.....	9.5	13.4	6.2	9.6	13.5	6.1	9.4	13.5	6.3
1999.....	9.6	13.5	6.1	9.6	13.6	6.0	10.1	14.9	6.3
Alcoholic liver disease (K70)									
2005.....	4.2	6.4	2.2	4.3	6.6	2.3	3.5	5.4	1.9
2004.....	4.2	6.3	2.2	4.3	6.5	2.2	3.5	5.6	1.9
2003.....	4.2	6.4	2.1	4.2	6.6	2.1	3.8	6.0	2.0
2002.....	4.2	6.3	2.2	4.2	6.5	2.2	3.8	5.8	2.3
2001.....	4.3	6.6	2.2	4.3	6.6	2.2	4.4	6.9	2.4
2000.....	4.3	6.8	2.2	4.3	6.9	2.0	4.6	7.0	2.8
1999.....	4.3	6.9	2.1	4.3	6.8	2.0	5.1	8.1	2.8
Other chronic liver disease and cirrhosis (K73-K74)									
2005.....	4.7	6.0	3.6	4.9	6.2	3.7	4.2	6.1	2.8
2004.....	4.8	6.1	3.7	4.9	6.2	3.8	4.3	6.2	2.9
2003.....	5.2	6.5	3.9	5.3	6.7	4.0	4.6	6.4	3.2
2002.....	5.2	6.5	4.1	5.4	6.7	4.2	4.6	6.2	3.4
2001.....	5.2	6.6	4.0	5.3	6.7	4.1	4.9	6.9	3.2
2000.....	5.2	6.6	4.0	5.3	6.7	4.0	4.8	6.5	3.5
1999.....	5.2	6.7	4.0	5.3	6.7	4.0	5.0	6.8	3.5

Worktable 293. Age-adjusted death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by race and sex: United States, 1999-2005

[Age-adjusted death rates per 100,000 U.S. standard population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992) and year	All races 1/			White 2/			Black 2/		
	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female
Cholelithiasis and other disorders of gallbladder (K80-K82)									
2005.....	1.0	1.1	0.9	1.0	1.1	0.9	1.2	1.3	1.2
2004.....	1.0	1.2	0.9	1.0	1.2	0.9	1.2	1.2	1.2
2003.....	1.0	1.2	0.9	1.0	1.1	0.9	1.1	1.3	1.0
2002.....	1.0	1.2	0.9	1.0	1.2	0.9	1.1	1.1	1.1
2001.....	1.0	1.2	0.9	1.0	1.2	0.9	1.3	1.4	1.2
2000.....	1.0	1.2	0.9	1.0	1.2	0.9	1.2	1.1	1.2
1999.....	1.0	1.2	0.9	1.0	1.2	0.9	1.1	1.1	1.0
Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)									
2005.....	14.3	17.5	12.3	12.9	16.2	10.8	29.7	34.1	26.9
2004.....	14.2	17.3	12.3	12.8	15.9	10.8	29.6	34.2	26.7
2003.....	14.4	17.8	12.4	13.0	16.3	10.9	30.4	36.1	26.8
2002.....	14.2	17.6	12.1	12.7	16.1	10.7	29.7	35.3	26.3
2001.....	14.0	17.4	12.0	12.5	16.0	10.5	29.5	33.8	27.1
2000.....	13.5	16.9	11.5	12.1	15.6	10.0	28.7	33.1	26.0
1999.....	13.0	16.5	11.1	11.5	15.0	9.6	28.4	33.7	25.4
Acute and rapidly progressive nephritic and nephrotic syndrome (N00-N01,N04)									
2005.....	0.0	0.0	0.0	0.0	0.1	0.0	*	*	*
2004.....	0.0	0.0	0.0	0.0	0.0	0.0	0.1	*	*
2003.....	0.1	0.0	0.0	0.0	0.0	0.0	*	*	*
2002.....	0.0	0.1	0.0	0.0	0.1	0.0	0.1	*	*
2001.....	0.0	0.1	0.0	0.0	0.1	0.0	0.1	*	*
2000.....	0.1	0.1	0.0	0.0	0.1	0.0	0.1	*	*
1999.....	0.1	0.1	0.1	0.0	0.1	0.1	0.1	*	*
Chronic glomerulonephritis, nephritis and nephropathy not specified as acute or chronic, and renal sclerosis unspecified (N02-N03,N05-N07,N26)									
2005.....	0.3	0.4	0.2	0.3	0.3	0.2	0.6	0.7	0.5
2004.....	0.2	0.2	0.2	0.2	0.2	0.1	0.3	0.4	0.3
2003.....	0.2	0.2	0.1	0.2	0.2	0.1	0.3	0.4	0.2
2002.....	0.2	0.2	0.2	0.2	0.2	0.1	0.3	0.4	0.3
2001.....	0.2	0.2	0.2	0.2	0.2	0.1	0.2	0.3	0.2
2000.....	0.2	0.3	0.2	0.2	0.2	0.1	0.4	0.4	0.3
1999.....	0.2	0.3	0.2	0.2	0.3	0.2	0.4	0.5	0.3

Worktable 293. Age-adjusted death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by race and sex: United States, 1999-2005

[Age-adjusted death rates per 100,000 U.S. standard population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992) and year

	All races 1/			White 2/			Black 2/		
	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female
Renal failure (N17-N19)									
2005.....	14.0	17.1	12.0	12.6	15.8	10.5	29.0	33.3	26.3
2004.....	13.9	17.0	12.0	12.5	15.6	10.6	29.1	33.8	26.3
2003.....	14.2	17.5	12.1	12.7	16.1	10.7	30.0	35.6	26.5
2002.....	13.9	17.3	11.9	12.5	15.8	10.5	29.3	34.8	25.9
2001.....	13.7	17.1	11.8	12.3	15.8	10.2	29.2	33.4	26.7
2000.....	13.2	16.6	11.3	11.8	15.2	9.8	28.2	32.6	25.5
1999.....	12.8	16.1	10.8	11.3	14.6	9.4	27.9	33.0	24.9

Other disorders of kidney (N25,N27)

2005.....	0.0	*	*	0.0	*	*	*	*	*
2004.....	0.0	*	0.0	0.0	*	*	*	*	*
2003.....	0.0	*	*	0.0	*	*	*	*	*
2002.....	0.0	*	0.0	0.0	*	0.0	*	*	*
2001.....	0.0	*	0.0	0.0	*	*	*	*	*
2000.....	0.0	0.0	0.0	0.0	*	*	*	*	*
1999.....	0.0	*	0.0	0.0	*	*	*	*	*

Infections of kidney (N10-N12,N13.6,N15.1)

2005.....	0.2	0.2	0.3	0.2	0.2	0.3	0.3	0.2	0.4
2004.....	0.2	0.2	0.3	0.2	0.2	0.3	0.3	0.2	0.3
2003.....	0.3	0.2	0.3	0.3	0.2	0.3	0.3	0.3	0.4
2002.....	0.3	0.2	0.3	0.3	0.2	0.3	0.3	0.2	0.3
2001.....	0.3	0.2	0.3	0.3	0.2	0.3	0.3	0.3	0.3
2000.....	0.3	0.2	0.3	0.3	0.2	0.3	0.4	0.3	0.5
1999.....	0.3	0.2	0.3	0.3	0.3	0.4	0.3	0.3	0.4

Hyperplasia of prostate (N40)

2005.....	0.2	0.5	...	0.2	0.5	...	0.1	0.4	...
2004.....	0.2	0.5	...	0.2	0.5	...	0.1	0.4	...
2003.....	0.2	0.5	...	0.2	0.5	...	0.1	0.4	...
2002.....	0.1	0.4	...	0.1	0.4	...	0.2	0.5	...
2001.....	0.1	0.5	...	0.2	0.5	...	0.1	0.4	...
2000.....	0.2	0.5	...	0.2	0.5	...	0.2	0.5	...
1999.....	0.2	0.5	...	0.2	0.5	...	0.2	0.5	...

Worktable 293. Age-adjusted death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by race and sex: United States, 1999-2005

[Age-adjusted death rates per 100,000 U.S. standard population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992) and year

	All races 1/			White 2/			Black 2/		
	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female
Inflammatory diseases of female pelvic organs (N70-N76)									
2005.....	0.0	...	0.1	0.0	...	0.1	*	...	*
2004.....	0.0	...	0.1	0.0	...	0.0	0.1	...	0.1
2003.....	0.0	...	0.1	0.0	...	0.1	0.1	...	0.1
2002.....	0.0	...	0.1	0.0	...	0.0	0.1	...	0.1
2001.....	0.0	...	0.1	0.0	...	0.1	0.1	...	0.1
2000.....	0.0	...	0.1	0.0	...	0.1	0.1	...	0.1
1999.....	0.0	...	0.1	0.0	...	0.0	0.1	...	0.2
Pregnancy, childbirth and the puerperium (000-099)									
2005.....	0.3	...	0.5	0.2	...	0.4	0.7	...	1.3
2004.....	0.2	...	0.5	0.2	...	0.4	0.6	...	1.2
2003.....	0.2	...	0.4	0.1	...	0.3	0.5	...	0.9
2002.....	0.1	...	0.3	0.1	...	0.2	0.4	...	0.8
2001.....	0.1	...	0.3	0.1	...	0.2	0.4	...	0.8
2000.....	0.1	...	0.3	0.1	...	0.2	0.4	...	0.7
1999.....	0.2	...	0.3	0.1	...	0.2	0.4	...	0.8
Pregnancy with abortive outcome (000-007)									
2005.....	0.0	...	0.0	*	...	*	*	...	*
2004.....	0.0	...	0.0	*	...	*	*	...	*
2003.....	0.0	...	0.0	*	...	*	*	...	*
2002.....	0.0	...	0.0	*	...	*	*	...	*
2001.....	0.0	...	0.0	0.0	...	0.0	*	...	*
2000.....	0.0	...	0.0	*	...	*	*	...	*
1999.....	0.0	...	0.0	*	...	*	*	...	*
Other complications of pregnancy, childbirth and the puerperium (010-099)									
2005.....	0.2	...	0.5	0.2	...	0.4	0.6	...	1.2
2004.....	0.2	...	0.5	0.2	...	0.4	0.6	...	1.1
2003.....	0.2	...	0.3	0.1	...	0.3	0.4	...	0.8
2002.....	0.1	...	0.2	0.1	...	0.2	0.3	...	0.7
2001.....	0.1	...	0.3	0.1	...	0.2	0.4	...	0.7
2000.....	0.1	...	0.3	0.1	...	0.2	0.3	...	0.6
1999.....	0.1	...	0.3	0.1	...	0.2	0.4	...	0.7

Worktable 293. Age-adjusted death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by race and sex: United States, 1999-2005

[Age-adjusted death rates per 100,000 U.S. standard population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992) and year

	All races 1/			White 2/			Black 2/		
	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female
Certain conditions originating in the perinatal period (P00-P96)									
2005.....	4.9	5.4	4.4	3.8	4.2	3.4	10.5	11.5	9.5
2004.....	4.8	5.4	4.2	3.8	4.3	3.3	10.3	11.3	9.2
2003.....	4.9	5.5	4.3	4.0	4.4	3.5	10.2	11.4	9.0
2002.....	4.9	5.4	4.3	3.9	4.3	3.4	10.1	10.9	9.2
2001.....	4.7	5.3	4.2	3.7	4.1	3.3	10.3	11.3	9.1
2000.....	5.1	5.6	4.5	4.0	4.4	3.5	11.3	12.6	9.9
1999.....	5.2	5.7	4.6	4.0	4.4	3.6	11.8	13.1	10.4

Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)

2005.....	3.5	3.7	3.3	3.5	3.6	3.3	3.9	4.1	3.7
2004.....	3.6	3.7	3.4	3.5	3.7	3.4	4.0	4.4	3.7
2003.....	3.6	3.8	3.4	3.6	3.7	3.4	4.0	4.4	3.6
2002.....	3.7	3.8	3.5	3.7	3.8	3.5	4.1	4.4	3.9
2001.....	3.6	3.8	3.4	3.6	3.8	3.4	4.1	4.3	3.9
2000.....	3.8	4.1	3.6	3.7	4.0	3.5	4.4	4.8	4.1
1999.....	3.8	3.9	3.6	3.7	3.9	3.6	4.3	4.7	4.1

Symptoms, signs and abnormal clinical and laboratory findings, not elsewhere classified (R00-R99)

2005.....	10.4	11.1	9.4	10.0	10.6	9.1	14.6	17.1	12.5
2004.....	10.0	10.7	9.0	9.6	10.2	8.7	14.0	16.0	12.1
2003.....	10.6	11.4	9.5	10.2	11.0	9.2	14.8	17.4	12.5
2002.....	10.3	11.4	9.0	9.9	10.8	8.7	14.5	18.0	11.8
2001.....	11.3	12.7	9.8	11.0	12.2	9.5	15.1	17.7	12.8
2000.....	11.4	13.1	9.7	11.0	12.6	9.3	16.5	19.7	13.6
1999.....	9.7	11.0	8.4	9.1	10.1	7.9	15.4	19.1	12.4

All other diseases (Residual) (D65-E07, E15-E34, E65-F99, G04-G12, G23-G25, G31-H93, K00-K22, K29-K31, K50-K66, K71-K72, K75-K76, K83-M99, N13.0-N13.5, N13.7-N13.9, N14, N15.0, N15.8-N15.9, N20-N23, N28-N39, N41-N64, N80-N98)

2005.....	70.5	70.5	68.6	69.7	69.5	67.9	87.7	91.5	83.5
2004.....	67.1	67.3	65.2	66.2	66.1	64.5	84.9	91.8	79.3
2003.....	68.3	68.2	66.6	67.4	66.9	65.8	85.6	91.6	80.3
2002.....	67.1	67.3	65.2	66.1	65.9	64.3	84.7	91.5	79.1
2001.....	65.9	66.1	64.0	64.7	64.5	63.0	84.6	92.0	78.5
2000.....	64.6	65.4	62.4	63.2	63.6	61.2	83.8	92.3	77.2
1999.....	63.0	64.0	60.6	61.6	62.2	59.5	81.0	90.0	74.5

Worktable 293. Age-adjusted death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by race and sex: United States, 1999-2005

[Age-adjusted death rates per 100,000 U.S. standard population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992) and year

	All races 1/			White 2/			Black 2/		
	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female
Accidents (unintentional injuries) (V01-X59,Y85-Y86)									
2005.....	39.1	54.2	25.0	40.1	55.2	25.8	38.7	58.1	22.8
2004.....	37.7	52.1	24.5	38.8	53.2	25.3	36.3	54.4	21.7
2003.....	37.3	51.8	24.1	38.2	52.8	24.7	36.1	53.9	21.6
2002.....	36.9	51.5	23.5	37.5	52.0	24.0	36.9	56.2	21.3
2001.....	35.7	50.2	22.5	36.0	50.3	22.8	37.6	56.9	22.0
2000.....	34.9	49.3	22.0	35.1	49.2	22.2	37.7	58.1	21.3
1999.....	35.3	49.8	22.3	35.2	49.3	22.4	40.1	61.3	23.1
Transport accidents (V01-V99,Y85)									
2005.....	16.2	23.4	9.3	16.7	23.9	9.7	15.4	24.1	8.0
2004.....	16.2	23.1	9.7	16.7	23.6	10.0	15.7	24.4	8.4
2003.....	16.4	23.4	9.8	16.8	23.9	10.0	15.8	24.5	8.6
2002.....	16.7	23.9	9.9	17.1	24.2	10.2	16.0	25.0	8.5
2001.....	16.6	23.7	9.8	16.8	24.0	10.0	16.5	25.5	8.9
2000.....	16.6	23.7	10.0	16.8	23.8	10.2	16.7	26.1	8.7
1999.....	16.6	23.6	10.2	16.8	23.6	10.4	17.1	26.5	9.5
Motor vehicle accidents (V02-V04,V09.0,V09.2,V12-V14,V19.0-V19.2,V19.4-V19.6,V20-V79,V80.3-V80.5,V81.0-V81.1,V82.0-V82.1,V83-V86,V87.0-V87.8,V88.0-V88.8,V89.0,V89.2)									
2005.....	15.2	21.7	8.9	15.6	22.2	9.2	14.5	22.5	7.6
2004.....	15.2	21.4	9.3	15.6	21.8	9.5	14.8	22.7	8.1
2003.....	15.3	21.6	9.3	15.7	21.9	9.5	14.9	22.7	8.3
2002.....	15.7	22.1	9.6	16.0	22.4	9.8	15.0	23.2	8.2
2001.....	15.3	21.8	9.3	15.6	22.0	9.5	15.4	23.5	8.5
2000.....	15.4	21.7	9.5	15.6	21.8	9.8	15.7	24.4	8.4
1999.....	15.2	21.3	9.6	15.3	21.3	9.7	15.9	24.3	9.0
Other land transport accidents (V01,V05-V06,V09.1,V09.3-V09.9,V10-V11,V15-V18,V19.3,V19.8-V19.9,V80.0-V80.2,V80.6-V80.9,V81.2-V81.9,V82.2-V82.9,V87.9,V88.9,V89.1,V89.3,V89.9)									
2005.....	0.4	0.7	0.2	0.4	0.6	0.2	0.5	1.0	0.2
2004.....	0.4	0.6	0.2	0.4	0.6	0.2	0.5	0.9	0.2
2003.....	0.4	0.7	0.2	0.5	0.7	0.2	0.5	1.0	0.2
2002.....	0.4	0.6	0.1	0.4	0.6	0.1	0.5	0.9	0.1
2001.....	0.5	0.8	0.2	0.5	0.7	0.2	0.6	1.0	0.3
2000.....	0.5	0.8	0.3	0.5	0.8	0.2	0.6	1.0	0.3
1999.....	0.8	1.1	0.4	0.8	1.1	0.4	0.7	1.2	0.3

Worktable 293. Age-adjusted death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by race and sex: United States, 1999-2005

[Age-adjusted death rates per 100,000 U.S. standard population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992) and year

	All races 1/			White 2/			Black 2/		
	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female
Water, air and space, and other and unspecified transport accidents and their sequelae (V90-V99,Y85)									
2005.....	0.6	1.0	0.2	0.7	1.1	0.2	0.4	0.7	0.1
2004.....	0.6	1.1	0.3	0.7	1.1	0.3	0.4	0.8	0.1
2003.....	0.7	1.1	0.3	0.7	1.2	0.3	0.4	0.8	0.1
2002.....	0.6	1.1	0.2	0.7	1.2	0.2	0.5	0.9	0.1
2001.....	0.7	1.2	0.3	0.8	1.3	0.3	0.5	1.0	0.1
2000.....	0.7	1.2	0.2	0.7	1.3	0.2	0.4	0.8	*
1999.....	0.7	1.1	0.3	0.7	1.2	0.3	0.5	1.0	0.2
Nontransport accidents (W00-X59,Y86)									
2005.....	22.9	30.9	15.7	23.4	31.4	16.1	23.3	34.0	14.8
2004.....	21.5	29.0	14.9	22.1	29.6	15.3	20.6	30.0	13.3
2003.....	20.9	28.4	14.3	21.4	28.9	14.6	20.2	29.5	13.0
2002.....	20.2	27.6	13.5	20.5	27.8	13.8	20.9	31.2	12.9
2001.....	19.1	26.4	12.7	19.2	26.4	12.8	21.1	31.4	12.8
2000.....	18.4	25.6	12.0	18.3	25.3	12.0	21.0	32.0	12.5
1999.....	18.7	26.3	12.1	18.4	25.7	12.0	22.9	34.8	13.6
Falls (W00-W19)									
2005.....	6.4	8.3	4.9	6.7	8.6	5.2	3.7	5.2	2.6
2004.....	6.3	8.2	4.8	6.6	8.6	5.1	3.6	5.5	2.3
2003.....	5.9	7.6	4.5	6.1	7.9	4.8	3.4	4.7	2.4
2002.....	5.6	7.4	4.3	5.9	7.6	4.6	3.1	4.9	2.0
2001.....	5.3	7.2	3.9	5.6	7.5	4.1	3.3	5.1	2.0
2000.....	4.8	6.6	3.6	5.0	6.8	3.7	2.9	4.6	1.8
1999.....	4.8	6.6	3.5	5.0	6.8	3.7	3.4	5.3	2.1
Accidental discharge of firearms (W32-W34)									
2005.....	0.3	0.5	0.1	0.3	0.4	0.1	0.4	0.7	*
2004.....	0.2	0.4	0.0	0.2	0.4	0.1	0.3	0.6	*
2003.....	0.2	0.5	0.1	0.2	0.5	0.1	0.3	0.6	*
2002.....	0.3	0.4	0.1	0.2	0.4	0.1	0.4	0.7	*
2001.....	0.3	0.5	0.1	0.3	0.5	0.1	0.3	0.5	*
2000.....	0.3	0.5	0.1	0.3	0.5	0.1	0.3	0.6	0.1
1999.....	0.3	0.5	0.1	0.3	0.5	0.1	0.4	0.7	0.1

Worktable 293. Age-adjusted death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by race and sex: United States, 1999-2005

[Age-adjusted death rates per 100,000 U.S. standard population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992) and year	All races 1/			White 2/			Black 2/		
	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female
Accidental drowning and submersion (W65-W74)									
2005.....	1.2	1.9	0.5	1.2	1.8	0.5	1.4	2.5	0.4
2004.....	1.1	1.8	0.5	1.1	1.7	0.5	1.3	2.2	0.5
2003.....	1.1	1.8	0.5	1.1	1.7	0.5	1.2	2.2	0.4
2002.....	1.2	2.0	0.5	1.1	1.8	0.4	1.6	2.7	0.5
2001.....	1.1	1.8	0.5	1.1	1.7	0.5	1.5	2.6	0.5
2000.....	1.2	2.0	0.5	1.2	1.8	0.5	1.5	2.6	0.6
1999.....	1.3	2.0	0.5	1.2	1.9	0.5	1.7	3.0	0.5
Accidental exposure to smoke, fire and flames (X00-X09)									
2005.....	1.1	1.4	0.8	0.9	1.2	0.7	2.2	3.1	1.6
2004.....	1.1	1.4	0.8	0.9	1.2	0.7	2.4	3.3	1.8
2003.....	1.2	1.4	0.9	1.1	1.3	0.8	2.4	3.2	1.9
2002.....	1.1	1.4	0.8	0.9	1.3	0.7	2.4	3.3	1.7
2001.....	1.2	1.5	0.9	1.0	1.3	0.7	2.7	3.8	1.9
2000.....	1.2	1.6	0.9	1.0	1.3	0.7	3.0	4.2	2.2
1999.....	1.2	1.6	0.9	1.0	1.4	0.7	3.0	4.0	2.3
Accidental poisoning and exposure to noxious substances (X40-X49)									
2005.....	7.9	10.7	5.1	8.4	11.3	5.5	8.2	11.9	4.9
2004.....	7.1	9.5	4.7	7.6	10.1	5.0	6.9	9.9	4.5
2003.....	6.7	9.1	4.3	7.1	9.6	4.5	6.8	9.9	4.1
2002.....	6.1	8.4	3.7	6.3	8.7	3.9	6.8	10.1	3.9
2001.....	4.9	7.0	2.9	5.0	7.1	2.9	6.3	9.5	3.7
2000.....	4.5	6.6	2.5	4.5	6.6	2.5	6.0	9.3	3.3
1999.....	4.4	6.4	2.3	4.3	6.3	2.3	6.2	9.8	3.2
Other and unspecified nontransport accidents and their sequelae (W20-W31,W35-W64,W75-W99,X10-X39,X50-X59,Y86)									
2005.....	6.1	8.2	4.2	6.0	8.1	4.1	7.5	10.6	5.2
2004.....	5.7	7.6	4.0	5.7	7.7	4.0	6.0	8.4	4.2
2003.....	5.8	7.9	4.1	5.8	7.9	4.1	6.2	8.9	4.2
2002.....	5.9	7.9	4.2	5.9	7.9	4.2	6.7	9.4	4.7
2001.....	6.3	8.4	4.5	6.2	8.3	4.5	7.0	9.8	4.9
2000.....	6.3	8.5	4.4	6.3	8.4	4.4	7.1	10.8	4.5
1999.....	6.7	9.1	4.8	6.7	8.9	4.8	8.2	12.2	5.4

Worktable 293. Age-adjusted death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by race and sex: United States, 1999-2005

[Age-adjusted death rates per 100,000 U.S. standard population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992) and year

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992) and year	All races 1/			White 2/			Black 2/		
	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female
Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)									
2005.....	10.9	18.0	4.4	12.0	19.6	4.9	5.2	9.2	1.9
2004.....	10.9	18.0	4.5	12.0	19.6	5.0	5.3	9.6	1.8
2003.....	10.8	18.0	4.2	11.8	19.6	4.6	5.2	9.2	1.9
2002.....	10.9	18.4	4.2	12.0	20.0	4.7	5.3	9.8	1.6
2001 3/.....	10.7	18.2	4.0	11.7	19.6	4.5	5.5	9.8	1.8
2000.....	10.4	17.7	4.0	11.3	19.1	4.3	5.5	10.0	1.8
1999.....	10.5	17.8	4.0	11.3	19.0	4.4	5.6	10.3	1.6

Intentional self-harm (suicide) by discharge of firearms (X72-X74)

2005.....	5.7	10.5	1.4	6.3	11.5	1.6	2.7	5.3	0.5
2004.....	5.6	10.3	1.5	6.3	11.4	1.7	2.7	5.4	0.5
2003.....	5.8	10.7	1.4	6.4	11.8	1.6	2.7	5.3	0.6
2002.....	5.9	11.1	1.4	6.5	12.1	1.6	3.0	5.9	0.5
2001.....	5.9	11.0	1.4	6.5	12.0	1.6	3.1	6.0	0.6
2000.....	5.9	11.0	1.5	6.5	11.9	1.6	3.1	6.0	0.7
1999.....	6.0	11.1	1.5	6.5	12.0	1.7	3.3	6.5	0.6

Intentional self-harm (suicide) by other and unspecified means and their sequelae (*U03,X60-X71,X75-X84,Y87.0)

2005.....	5.2	7.5	3.0	5.7	8.1	3.3	2.5	3.9	1.4
2004.....	5.3	7.6	3.1	5.7	8.2	3.3	2.6	4.2	1.3
2003.....	5.0	7.3	2.8	5.4	7.8	3.1	2.5	3.9	1.3
2002.....	5.0	7.4	2.8	5.5	7.9	3.1	2.4	3.9	1.1
2001 3/.....	4.8	7.1	2.6	5.2	7.6	2.9	2.4	3.8	1.1
2000.....	4.5	6.7	2.5	4.9	7.1	2.7	2.4	4.0	1.1
1999.....	4.5	6.6	2.5	4.8	7.1	2.7	2.3	3.8	1.0

Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)

2005.....	6.1	9.6	2.5	3.7	5.3	1.9	21.1	37.3	6.1
2004.....	5.9	9.2	2.5	3.6	5.3	1.9	20.1	35.1	6.3
2003.....	6.0	9.4	2.6	3.7	5.3	2.0	21.0	36.7	6.4
2002.....	6.1	9.4	2.8	3.7	5.3	2.0	21.0	36.4	6.9
2001 3/.....	7.1	10.8	3.3	4.9	7.1	2.6	21.2	36.2	7.4
2000.....	5.9	9.0	2.8	3.6	5.2	2.1	20.5	35.4	7.1
1999.....	6.0	9.1	2.9	3.8	5.4	2.1	20.1	34.2	7.3

Worktable 293. Age-adjusted death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by race and sex: United States, 1999-2005

[Age-adjusted death rates per 100,000 U.S. standard population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992) and year

	All races 1/			White 2/			Black 2/		
	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female
Assault (homicide) by discharge of firearms (*U01.4,X93-X95)									
2005.....	4.2	7.0	1.2	2.2	3.5	0.9	16.0	29.8	3.0
2004.....	4.0	6.6	1.1	2.2	3.4	0.8	15.0	28.0	2.9
2003.....	4.1	6.8	1.2	2.2	3.4	0.9	15.7	29.0	3.2
2002.....	4.1	6.8	1.3	2.2	3.4	1.0	15.6	28.7	3.5
2001.....	3.9	6.6	1.3	2.2	3.5	1.0	14.8	27.4	3.0
2000.....	3.8	6.3	1.2	2.1	3.2	0.9	14.6	26.9	3.1
1999.....	3.8	6.3	1.3	2.1	3.3	1.0	14.3	26.1	3.5

Assault (homicide) by other and unspecified means and their sequelae (*U01.0-*U01.3,*U01.5-*U01.9,*U02,X85-X92,X96-Y09,Y87.1)

2005.....	1.9	2.6	1.3	1.5	1.9	1.0	5.1	7.5	3.1
2004.....	2.0	2.5	1.4	1.5	1.9	1.1	5.1	7.1	3.4
2003.....	2.0	2.6	1.4	1.5	1.9	1.1	5.3	7.7	3.2
2002.....	2.0	2.6	1.4	1.5	1.9	1.1	5.4	7.7	3.4
2001 3/.....	3.1	4.3	2.0	2.6	3.6	1.6	6.4	8.8	4.3
2000.....	2.1	2.7	1.5	1.5	2.0	1.1	6.0	8.5	3.9
1999.....	2.2	2.8	1.6	1.6	2.1	1.2	5.8	8.1	3.8

Legal intervention (Y35,Y89.0)

2005.....	0.1	0.3	*	0.1	0.2	*	0.3	0.6	*
2004.....	0.1	0.3	*	0.1	0.2	*	0.2	0.5	*
2003.....	0.1	0.3	*	0.1	0.2	*	0.3	0.6	*
2002.....	0.1	0.3	*	0.1	0.2	*	0.3	0.6	*
2001.....	0.1	0.3	*	0.1	0.2	*	0.2	0.5	*
2000.....	0.1	0.2	*	0.1	0.2	*	0.3	0.6	*
1999.....	0.1	0.3	*	0.1	0.2	*	0.3	0.7	*

Events of undetermined intent (Y10-Y34,Y87.2,Y89.9)

2005.....	1.6	1.9	1.2	1.6	1.9	1.3	1.8	2.5	1.2
2004.....	1.7	2.2	1.2	1.7	2.2	1.3	1.9	2.7	1.1
2003.....	1.7	2.3	1.2	1.8	2.3	1.3	1.9	2.9	1.0
2002.....	1.7	2.2	1.2	1.7	2.2	1.2	2.1	3.1	1.2
2001.....	1.5	2.0	0.9	1.5	2.0	1.0	1.8	2.9	0.9
2000.....	1.3	1.9	0.8	1.3	1.8	0.8	1.9	3.0	1.0
1999.....	1.4	2.0	0.9	1.4	1.8	0.9	2.0	3.2	0.9

Worktable 293. Age-adjusted death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by race and sex: United States, 1999-2005

[Age-adjusted death rates per 100,000 U.S. standard population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992) and year

	All races 1/			White 2/			Black 2/		
	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female
Discharge of firearms, undetermined intent (Y22-Y24)									
2005.....	0.1	0.1	0.0	0.1	0.1	0.0	0.1	0.1	*
2004.....	0.1	0.1	0.0	0.1	0.1	0.0	0.1	0.1	*
2003.....	0.1	0.1	0.0	0.1	0.1	0.0	0.1	0.2	*
2002.....	0.1	0.1	0.0	0.1	0.1	0.0	0.1	0.2	*
2001.....	0.1	0.1	0.0	0.1	0.1	0.0	0.1	0.2	*
2000.....	0.1	0.1	0.0	0.1	0.1	0.0	0.1	0.3	*
1999.....	0.1	0.2	0.0	0.1	0.2	0.0	0.2	0.3	*
Other and unspecified events of undetermined intent and their sequelae (Y10-Y21,Y25-Y34,Y87.2,Y89.9)									
2005.....	1.5	1.8	1.2	1.5	1.8	1.3	1.7	2.4	1.1
2004.....	1.6	2.0	1.2	1.7	2.1	1.2	1.8	2.6	1.1
2003.....	1.7	2.2	1.2	1.7	2.2	1.3	1.8	2.7	1.0
2002.....	1.6	2.0	1.1	1.6	2.0	1.2	2.0	2.9	1.2
2001.....	1.4	1.9	0.9	1.4	1.8	1.0	1.7	2.7	0.9
2000.....	1.3	1.8	0.8	1.2	1.7	0.8	1.8	2.8	0.9
1999.....	1.3	1.8	0.8	1.2	1.7	0.9	1.8	2.9	0.9
Operations of war and their sequelae (Y36,Y89.1)									
2005.....	0.0	0.0	*	0.0	0.0	*	*	*	*
2004.....	0.0	0.0	*	0.0	0.0	*	*	*	*
2003.....	*	*	*	*	*	*	*	*	*
2002.....	0.0	0.0	*	*	*	*	*	*	*
2001.....	*	*	*	*	*	*	*	*	*
2000.....	*	*	*	*	*	*	*	*	*
1999.....	0.0	0.0	*	0.0	0.0	*	*	*	*
Complications of medical and surgical care (Y40-Y84,Y88)									
2005.....	0.9	0.9	0.8	0.8	0.9	0.8	1.5	1.5	1.5
2004.....	1.0	1.1	0.9	0.9	1.1	0.8	1.5	1.5	1.6
2003.....	1.0	1.1	0.9	0.9	1.0	0.8	1.5	1.5	1.6
2002.....	1.0	1.0	1.0	0.9	1.0	0.9	1.6	1.6	1.6
2001.....	1.1	1.1	1.0	1.0	1.1	0.9	2.0	2.3	1.7
2000.....	1.1	1.2	1.0	1.0	1.2	0.9	1.9	1.9	1.9
1999.....	1.0	1.1	0.9	1.0	1.1	0.9	1.8	2.0	1.7

Worktable 293. Age-adjusted death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by race and sex: United States, 1999-2005

[Age-adjusted death rates per 100,000 U.S. standard population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992) and year

	All races 1/			White 2/			Black 2/		
	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female
Alcohol-induced causes (E24.4, F10, G31.2, G62.1, G72.1, I42.6, K29.2, K70, K86.0, R78.0, X45, X65, Y15) 4/									
2005.....	7.0	11.0	3.4	7.2	11.1	3.4	6.8	11.4	3.3
2004.....	7.0	11.0	3.3	7.1	11.0	3.3	7.2	12.3	3.1
2003.....	7.0	11.0	3.3	7.0	11.0	3.3	7.4	12.8	3.3
2002.....	6.9	11.0	3.3	6.9	10.9	3.2	7.8	13.1	3.6
2001.....	7.0	11.2	3.3	6.9	10.9	3.3	8.9	15.1	3.9
2000.....	7.0	11.4	3.2	6.9	11.1	3.0	9.1	15.3	4.3
1999.....	7.1	11.5	3.2	6.8	11.0	3.0	9.8	16.7	4.5

Drug-induced causes (D52.1, D59.0, D59.2, D61.1, D64.2, E06.4, E16.0, E23.1, E24.2, E27.3, E66.1, F11.0-F11.5, F11.7-F11.9, F12.0-F12.5, F12.7-F12.9, F13.0-F13.5, F13.7-F13.9, F14.0-F14.5, F14.7-F14.9, F15.0-F15.5, F15.7-F15.9, F16.0-F16.5, F16.7-F16.9, F17.0, F17.3-F17.5, F17.7-F17.9, F18.0-F18.5, F18.7-F18.9, F19.0-F19.5, F19.7-F19.9, G21.1, G24.0, G25.1, G25.4, G25.6, G44.4, G62.0, G72.0, I95.2, J70.2-J70.4, L10.5, L27.0-L27.1, M10.2, M32.0, M80.4, M81.4, M83.5, M87.1, R78.1-R78.5, X40-X44, X60-X64, X85, Y10-Y14) 4/

2005.....	11.3	14.4	8.1	11.9	15.1	8.7	11.2	15.8	7.2
2004.....	10.4	13.3	7.6	11.1	13.9	8.1	10.1	14.3	6.6
2003.....	9.9	12.8	7.0	10.4	13.4	7.4	9.9	14.1	6.4
2002.....	9.0	11.7	6.3	9.4	12.1	6.7	9.9	14.2	6.1
2001.....	7.6	10.1	5.1	7.8	10.2	5.3	9.2	13.6	5.4
2000.....	7.0	9.5	4.6	7.1	9.4	4.7	9.0	13.5	5.2
1999.....	6.8	9.4	4.4	6.8	9.2	4.4	9.3	14.3	5.1

Injury by firearms (*U01.4, W32-W34, X72-X74, X93-X95, Y22-Y24, Y35.0) 4/

2005.....	10.2	18.3	2.7	8.9	15.7	2.6	19.4	36.4	3.6
2004.....	10.0	17.7	2.7	8.8	15.4	2.7	18.4	34.5	3.6
2003.....	10.3	18.4	2.7	9.0	16.0	2.6	19.0	35.6	3.8
2002.....	10.4	18.6	2.8	9.2	16.2	2.7	19.3	36.0	4.1
2001.....	10.3	18.5	2.8	9.2	16.3	2.7	18.4	34.5	3.8
2000.....	10.2	18.1	2.8	9.0	15.9	2.7	18.4	34.2	3.9
1999.....	10.3	18.4	2.9	9.1	16.2	2.7	18.4	34.1	4.3

* Figure does not meet standards of reliability or precision.

0.0 Quantity more than zero but less than 0.05.

... Category not applicable.

1/ Includes races other than white and black.

2/ Multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming in 2005, by California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and by California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005."

3/ Figures include September 11, 2001 related deaths for which death certificates were filed as of 10/24/02; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001."

4/ Included in selected categories.