

Pediatric Nutrition Surveillance 2009 Report

This report summarizes selected data on child health and nutrition indicators received from states, U.S. territories, and Indian Tribal Organizations that contributed to the Centers for Disease Control and Prevention's (CDC) Pediatric Nutrition Surveillance System.

This report was developed by CDC's Division of Nutrition, Physical Activity, and Obesity in the National Center for Chronic Disease Prevention and Health Promotion.

Suggested Citation

Polhamus B, Dalenius K, Mackintosh H, Smith B, Grummer-Strawn L. *Pediatric Nutrition Surveillance 2009 Report*. Atlanta: U.S. Department of Health and Human Services, Centers for Disease Control and Prevention; 2011.

Acknowledgments

We gratefully acknowledge and thank all contributors to the Pediatric Nutrition Surveillance System (PedNSS). The efforts of state, territorial, and Indian Tribal Organization surveillance coordinators; informatics staff; and local clinic staff to collect data and use nutrition surveillance systems make the national PedNSS possible.

This report is available online at <http://www.cdc.gov/pednss>.

For more information, contact

Division of Nutrition, Physical Activity, and Obesity
National Center for Chronic Disease Prevention and Health Promotion
Centers for Disease Control and Prevention

4770 Buford Highway NE, Mail Stop K-25
Atlanta, GA 30341-3717
800-CDC-INFO (232-4636)
TTY: 888-232-6348

Pediatric Nutrition Surveillance

The Pediatric Nutrition Surveillance System (PedNSS) is a public health surveillance system that monitors the nutritional status of low-income children in federally funded maternal and child health programs. Data on birthweight, anemia, breastfeeding, short stature, underweight, overweight, and obesity are collected for children who attend public health clinics for routine care, nutrition education, and supplemental food.

The goal of the PedNSS is to collect, analyze, and disseminate surveillance data to guide public health policy and action. PedNSS information is used to set public health priorities and to plan, implement, and evaluate nutrition programs.

Data are collected at the clinic level, aggregated at the state level, and then submitted to the Centers for Disease Control and Prevention (CDC) for analysis. A national nutrition surveillance report is produced, and an additional surveillance report is produced for each contributor. A *contributor* is defined as a state, U.S. territory, or Indian Tribal Organization (ITO).

In 2009, a total of 55 contributors, including 46 states, the District of Columbia, Puerto Rico, the U.S. Virgin Islands, and 6 ITOs, participated in the PedNSS (Figure 1). These contributors submitted data for nearly 9 million children from birth to age 4 years, which is nearly 4 million more children than in 2000. This gain is due to increases both in the number of contributors to the PedNSS and in the number of children reported by each contributor.

Figure 1. This is a map of the United States indicating contributors to PedNSS in 2009. Contributors include Alabama, Arizona, Arkansas, California, Colorado, Connecticut, Florida, Georgia, Hawaii, Idaho, Illinois, Indiana, Iowa, Kansas, Kentucky, Louisiana, Maine, Maryland, Massachusetts, Michigan, Minnesota, Mississippi, Missouri, Montana, Nebraska, Nevada, New Hampshire, New Jersey, New Mexico, New York, North Carolina, North Dakota, Ohio, Oregon, Pennsylvania, Rhode Island, South Carolina, South Dakota, Tennessee, Texas, Utah, Vermont, Virginia, Washington, West Virginia and Wisconsin. Contributors not shown on the map include the District of Columbia, Puerto Rico, the U.S. Virgin Islands, and the following Indian Tribal Organizations: Cheyenne River Sioux (SD), Inter Tribal Council of Arizona, Navajo Nation (AZ), Rosebud Sioux (SD), Standing Rock Sioux (ND), and Three Affiliated Tribes (ND).

Fluctuations in the number of contributors or the demographic characteristics of the contributors' populations can affect trends. The number of PedNSS contributors differs slightly from year to year because some contributors did not provide data every year during the 10-year period from 2000–2009 (see Table 1 at the end of this report).

Data for the 2009 PedNSS were collected from children enrolled in federally funded programs that serve low-income children. These programs include the Special Supplemental Nutrition Program for Women, Infants, and Children (WIC) (85.8%) and other programs (14.1%), such as the Early and Periodic Screening, Diagnosis, and Treatment program and the Maternal and Child Health Bureau Title V program. This report summarizes 2009 data and highlights data trends from 2000–2009.

Demographic Characteristics

Of the children in the 2009 PedNSS, 40.6% were Hispanic, 31.8% were non-Hispanic white, 18.7% were non-Hispanic black, 2.6% were Asian or Pacific Islander, 0.9% were American Indian or Alaska Native, and 5.5% were of multiple or unspecified races and ethnicities. From 2000 through 2009, the proportion of Hispanic children in the PedNSS increased from 28.3% to 40.6%. During the same period, the proportion of non-Hispanic white and non-Hispanic black children declined.

This demographic shift should be considered when interpreting PedNSS trends. Most children in the 2009 PedNSS were younger than age 2 years (57.1%); of those, 34.5% were younger than age 1 year and 22.6% were aged 12–23 months. The remaining children, 42.9%, were aged 2–4 years. The age distribution of children in the PedNSS has been stable since 2000.

Pediatric Health Indicators

Low Birthweight

Low birthweight (<2,500 grams) is an important determinant of neonatal and postneonatal mortality.¹ Low-birthweight infants who survive are at increased risk for health problems that include neurodevelopmental disabilities and respiratory disorders.² Of the infants in the 2009 PedNSS, 8.9% were low birthweight, compared with 8.2% of all U.S. infants.³

In the 2009 PedNSS, the prevalence of low birthweight was higher for black infants (13.2%) than for white (8.5%), Asian or Pacific Islander (8.3%), American Indian or Alaska Native (8.0%), or Hispanic (7.0%) infants. *Healthy People 2010* objective 16-10a proposes reducing low birthweight to no more than 5.0% of all live births.⁴ The overall prevalence of low birthweight remained stable from 2000 (8.9%) through 2009 (8.9%) (Figure 2).

Figure 2: This is a line graph showing trends in the prevalence of low birthweight by race and ethnicity. Low birthweight is defined as birth weight < 2500 grams, among infants born during the reporting period. Year 2010 target: reduce low birthweight to 5% of live births. Data for this table came from 2009 National PedNSS Table 18D and is available at http://www.cdc.gov/pednss/pednss_tables/tables_numeric.htm.

	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009
White, Non- Hispanic	8.3	8.4	8.4	8.6	8.8	9.1	8.9	8.7	8.6	8.5
Black, Non- Hispanic	12.6	12.7	12.6	12.9	13.1	13.6	13.5	13.4	13.1	13.2
Hispanic	6.7	6.7	7.1	7.3	7.7	7.4	7.2	7.3	7.4	7.0
American Indian/Alaska Native	6.7	7.3	6.8	7.4	7.9	7.9	8.1	7.9	7.8	8.0
Asian/Pacific Islander	8.0	8.3	8.0	8.3	8.4	8.5	8.3	8.	8.3	8.3
Total	8.9	9.0	9.0	9.0	9.3	9.4	9.2	9.1	9.0	8.9

High Birthweight

High birthweight (>4,000 grams) puts infants at increased risk for death and for birth injuries such as shoulder dystocia.⁵ In the 2009 PedNSS, 6.4% of infants had high birthweights, compared with 8.3% in 2000. The prevalence was lower than in 2007 (7.7%), the most recent U.S. rate available.⁶

In 2009, the prevalence of high birthweight was higher for American Indian or Alaska Native (8.9%) infants than for white (7.4%), Hispanic (6.7%), Asian or Pacific Islander (4.5%), or black (3.9%) infants. The largest absolute decreases in high birthweight during 2000–2009 occurred among American Indian or Alaska Native (2.8%) and Hispanic (2.3%) infants.

Breastfeeding

The nutritional, immunologic, and economic advantages of breastfeeding are well recognized.⁷ In the 2009 PedNSS, 61.7% of infants were ever breastfed, 27.0% were breastfed for at least 6 months, and 18.5% were breastfed for at least 12 months.*

The 2009 PedNSS population failed to meet *Healthy People 2010* objectives 16-19 a–c, which are to increase the proportion of infants ever breastfed to 75.0%, the proportion of infants breastfed at 6 months to 50.0%, and the proportion of infants breastfed at 1 year to 25.0%.⁴

However, among PedNSS contributors, Colorado, Hawaii, Idaho, the Navajo Nation, Oregon, Utah, the U.S. Virgin Islands, and Washington met the *Healthy People 2010* objective for ever breastfeeding, and several other contributors came close (see Table 2 at the end of this report).

Nationally representative provisional data from the 2007 National Immunization Survey (NIS) indicate that 75.0% of all U.S. infants were ever breastfed, 43.0% were breastfed for 6 months, and 22.4% were breastfed for 12 months.⁸

From 2000 through 2009, the absolute increase in the prevalence of breastfeeding initiation for infants in the PedNSS was 13.7%. The breastfeeding initiation rate was 48.0% in 2000. Improved breastfeeding rates were evident among all racial and ethnic groups (Figure 3). In 2009, Hispanic infants had the highest prevalence of breastfeeding initiation (74.4%), and black infants had the lowest prevalence (48.8%). Data from the NIS indicate that the proportion of infants ever breastfed in the United States increased from 70.9% in 2000 to 75.0% in 2007.⁸

Figure 3. This is a line graph showing trends in percentage of infants ever breastfed by race and ethnicity among infants born during the reporting period. Data for this table came from 2009 National PedNSS Table 19D.

	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009
White, non-Hispanic	48.5	50.6	51.0	52.6	53.8	55.4	56.6	55.9	57.7	58.5
Black, non-Hispanic	34.5	36.1	40.0	41.6	42.6	46.4	47.3	46.9	49.7	48.8
Hispanic	65.5	68.4	67.5	68.0	68.7	71.1	72.9	72.4	73.8	74.4
American Indian/Alaska Native	53.2	52.2	58.6	55.7	62.4	61.5	65.8	63.1	64.3	63.8
Asian	54.5	55.3	51.0	53.6	58.8	59.4	59.7	60.4	63.5	65.4
Total	48.0	50.1	52.5	55.1	56.1	58.5	60.1	59.8	61.7	61.7

The proportion of infants who were breastfed for at least 6 months was 19.7% in 2000 and 27.0% in 2009, an absolute increase of 7.3%. Increases in infants being breastfed for at least 6 months were evident among all racial and ethnic groups. Hispanic infants had the highest prevalence of breastfeeding for at least 6 months (38.4%), and black infants had the lowest prevalence (19.7%). Data from the NIS indicate that the proportion of infants who were breastfed for at least 6 months in the United States increased from 34.2% in 2000 to 43.0% in 2007.⁸

In 2004, the PedNSS began monitoring *exclusive breastfeeding*, defined as an infant receiving only breast milk. In 2009, a total of 23 PedNSS contributors[†] (42%) reported this supplementary data, which showed that 9.9% of infants were exclusively breastfed for at least 3 months.

Data from the NIS indicate that 33.0% of infants in the United States were exclusively breastfed for at least 3 months in 2007. Exclusive breastfeeding has a strong protective effect against lower respiratory tract infections, middle ear infections, eczema, and childhood obesity.⁹

Anemia

Anemia (low hemoglobin or low hematocrit)[†] is an indicator of iron deficiency, which is associated with developmental delays and behavioral problems in children.^{10,11} In the 2009 PedNSS, the prevalence of anemia was 14.9%. The highest prevalence was among children aged 6–23 months (17.8%) and those aged 12–17 months (18.2%). The lowest prevalence was among children aged 3–4 years (11.2%). The overall prevalence of anemia among children in the PedNSS increased slightly from 13.7% in 2000 to 14.9% in 2009. During this 10-year period, the overall prevalence of anemia declined to 13.0% in 2002 but increased in subsequent years.

The prevalence of anemia varied among racial and ethnic groups in the PedNSS. In 2009, the highest prevalence was among black children (23.1%) and the lowest prevalence was among white children (11.4%). From 2000 through 2009, the largest increase in anemia was among black children (3.1%). The prevalence of anemia declined slightly among Asian or Pacific Islander children (Figure 4).

Figure 4: This is a line graph showing trends in prevalence of anemia among children aged < 5 years by race and ethnicity. Anemia is defined as Hemoglobin or Hematocrit < 5th percentile, CDC MMWR vol. 47 (No. RR-3), 1998. Data for this table came from 2009 National PedNSS Table 18D and is available at http://www.cdc.gov/pednss/pednss_tables/tables_numeric.htm.

	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009
White, non-Hispanic	10.6	10.4	10.1	10.5	10.8	10.8	11.1	11.9	11.4	11.4
Black, non-Hispanic	20.0	19.6	19.1	19.1	19.4	20.0	21.0	22.8	22.6	23.1
Hispanic	13.3	13.0	12.5	13.3	13.1	13.2	13.5	14.0	14.0	14.0
American Indian/Alaska Native	11.0	9.9	10.0	10.9	11.5	10.7	11.5	12.4	12.2	13.1
Asian/Pacific Isld	15.1	12.8	12.4	12.6	12.7	12.4	12.5	13.0	13.6	13.7
Total	13.7	13.4	13.0	13.4	13.5	13.6	14.0	15.0	14.8	14.9

[†] Defined as hemoglobin (Hb) concentration or hematocrit (Hct) level <5th percentile. Children aged 6–23 months are considered anemic if their Hb concentration is <11.0 g/dL or their Hct level is <32.9%. Children aged 2–4 years are considered anemic if their Hb concentration is <11.1 g/dL or their Hct level is <33.0%. Values are adjusted for altitude. Hb concentration and Hct level are not reported for children younger than age 6 months.¹²

Short Stature

Short stature[§] (low length or height for a child's age) may reflect the long-term health and nutritional status of a child or a population.¹³ Although short stature can be associated with short parental stature or low birthweight,¹³ it also can result from growth retardation because of chronic malnutrition, recurrent illness, or both. In the 2009 PedNSS, 6.0% of children from birth to age 4 years were of short stature, compared with 3.7% of all U.S. children of the same age.¹⁴

Compared with the general population, the prevalence of short stature was higher in the PedNSS population, which may reflect the nutritional risk of children participating in the WIC program. The prevalence of short stature in the 2009 PedNSS was above both the expected level (5.0%) and the *Healthy People 2010* objective 19-4 of 5.0% among low-income children from birth to age 4 years.⁴ Sixteen contributors achieved the *Healthy People 2010* objective of 5.0% in 2009 (see Table 2 at the end of this report).

The prevalence of short stature among children in the PedNSS remained stable from 2000 (6.0%) through 2009 (6.0%). Some variation was evident among racial and ethnic groups. Short stature increased among white and Hispanic children and decreased among Asian or Pacific Islander, black, and American Indian or Alaska Native children (Figure 5). In 2009, the highest prevalence of short stature was among black infants younger than age 1 year (10.9%), which may reflect the high prevalence of low birthweight among this group (data not shown).

Figure 5: This is a line graph showing trends in prevalence of short stature among children aged < 5 years by race and ethnicity. Short stature is defined as \leq 5th percentile length or height-for-age, CDC Growth Charts, 2000. Year 2010 target: reduce short stature among low-income children aged < 5 years to 5%. Data for this table came from 2009 National PedNSS Table 18D and is available at http://www.cdc.gov/pednss/pednss_tables/tables_numeric.htm.

	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009
White, non-Hispanic	6.0	6.3	6.3	6.5	6.6	6.6	6.5	6.4	6.2	6.2
Black, non-Hispanic	7.2	7.5	7.0	7.1	6.9	6.9	7.1	7.0	6.6	6.6
Hispanic	5.2	5.5	6.0	6.0	6.0	5.9	5.6	5.9	5.5	5.6
American Indian/Alaska Native	5.1	5.2	5.1	5.1	5.0	5.1	5.0	5.0	4.7	4.7
Asian/Pacific Islander	6.7	7.0	6.4	6.0	6.7	6.6	6.4	6.3	6.3	6.3
Total	6.0	6.3	6.3	6.4	6.4	6.4	6.4	6.2	6.0	6.0

[§] Based on sex-specific percentiles from the 2000 CDC growth chart for the United States. For children younger than age 2 years, short stature is defined as <5th percentile of length-for-age. For children aged 2–4 years, short stature is defined as <5th percentile of height-for-age.

Underweight

Data from contributors on underweight[†] (low weight-for-length or body mass index^{**} [BMI] for age) in children from birth to age 4 years indicate that acute malnutrition was not a public health problem in the PedNSS population. In 2009, the prevalence of underweight (4.3%) was less than the expected level (5.0%). The prevalence of underweight for all U.S. children in this age group was 3.4%.¹⁴

The highest prevalence of underweight in the PedNSS was among Asian or Pacific Islander (5.9%) and black (5.6%) children. Black infants younger than age 1 year had an underweight rate of 7.6%, which may reflect the high rate of low birthweight in this group. The overall prevalence of underweight among children in the PedNSS decreased from 5.4% in 2000 to 4.3% in 2009.

Overweight and Obesity

Overweight^{††} and obesity^{‡‡} in young children have increased in recent decades, and the associated health consequences call for preventive efforts.¹⁵ The Expert Committee on the Prevention, Assessment, and Treatment of Child and Adolescent Overweight and Obesity recommends the use of two cutoff points to screen for overweight and obesity in children aged 2 years or older.¹⁶ Children with a BMI-for-age at or above the 95th percentile are considered obese, and those with a BMI-for-age in the 85th–95th percentile are considered overweight.¹⁶

In the 2009 PedNSS, the prevalence of obesity among children from birth to age 4 years was 14.7%, compared with 12.4% for U.S. children aged 2–5 years in 2003 through 2006.¹⁷ In the PedNSS, the highest prevalence of obesity was among American Indian or Alaska Native (20.7%) and Hispanic (17.9%) children. The lowest prevalence was among white (12.3%), black (11.9%), and Asian or Pacific Islander (11.9%) children (Figure 6).

Figure 6: This is a vertical stacked bar graph that shows prevalence of obesity and overweight among children aged 2 to 5 years by race and ethnicity for 2007. Obesity is defined as \geq 95th percentile BMI-for-age; overweight is defined as \geq 85th to \leq 95th percentile BMI-for-age, CDC Growth Charts, 2000. Data for this table came from 2009 National PedNSS Table 8D and is available at http://www.cdc.gov/pednss/pednss_tables/tables_numeric.htm.

	White	Black	Hispanic	American Indian	Asian	Total
Obese	12.3	11.9	17.9	20.7	11.9	14.7
Overweight	15.9	14.5	17.6	20.2	14.0	16.4

[†] Based on sex-specific percentiles from the 2000 CDC growth chart for the United States. For children younger than age 2 years, underweight is defined as <5th percentile of weight-for-length. For children aged 2 years or older, underweight is defined as <5th percentile of BMI-for-age.

^{**} BMI is calculated as follows: Weight (kg) \div Stature (cm) \times 10,000 or Weight (lb) \div Stature (in) \times 703.

^{††} Based on sex-specific percentiles from the 2000 CDC growth chart for the United States. For children aged 2 years or older, overweight is defined as the 85th–95th percentile of BMI-for-age.

^{‡‡} Based on sex-specific percentiles from the 2000 CDC growth chart for the United States. For children aged 2 years or older, obesity is defined as \geq 95th percentile of BMI-for-age.

The prevalence of obesity among children from birth to age 4 years increased from 13.2% in 2000 to 14.7% in 2009 (Figure 7). During this 10-year period, obesity increased among all U.S. racial and ethnic groups except Asians or Pacific Islanders.

However, overall obesity rates have remained stable since 2003 (14.7%), and this trend was observed among all racial and ethnic groups except American Indians or Alaska Natives. This group experienced a 3.0% increase in the prevalence of obesity from 2003 through 2009.

Figure 7: This is a line graph showing trends in prevalence of obesity* among children aged 2 to 5 years by race and ethnicity. Obesity is defined as \geq 95th percentile BMI-for-age, CDC Growth Charts, 2000. Data for this table came from 2009 National PedNSS Table 18D and is available at http://www.cdc.gov/pednss/pednss_tables/tables_numeric.htm.

	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009
White, non- Hispanic	11.3	11.5	11.8	12.2	12.6	12.4	12.3	12.4	12.5	12.3
Black, non- Hispanic	11.1	11.0	11.9	12.1	12.2	12.0	11.9	11.9	11.9	11.9
Hispanic	17.2	17.0	19.0	18.4	18.2	18.0	18.0	18.2	18.2	17.9
American Indian/Alaska Native	17.4	17.1	17.5	17.7	19.0	19.0	19.4	19.5	20.2	20.7
Asian/Pacific Islander	13.8	13.4	13.4	13.0	12.5	12.3	12.4	11.9	12.2	11.9
Total	13.2	13.1	14.3	14.7	14.8	14.7	14.8	14.9	14.8	14.7

Among all contributors, only Colorado, Hawaii, Utah, and the U.S. Virgin Islands had a prevalence of obesity less than 12.0%, and 15 contributors had a prevalence of obesity of 16.0% or higher (Figure 8).

Figure 8: This is a map of the United States with states showing the prevalence of obesity among children aged 2 to 5 years by contributor. Obesity is defined as \geq 95th percentile BMI-for-age, CDC Growth Charts, 2000. Four states, Colorado, Hawaii, Idaho, Utah and the U.S. Virgin Islands had a prevalence of obesity less than 12%. States with a prevalence of obesity from 12% up to 13.9% include the District of Columbia, Florida, Kansas, Louisiana, Michigan, Minnesota, , Mississippi, Missouri, Montana, Nevada, New Mexico, Ohio, Pennsylvania, South Carolina, Vermont, West Virginia, and Wisconsin. Contributors with a prevalence of 14.0% to 15.9% include Alabama, Arizona, Arkansas, Georgia, Illinois, Indiana, Maine, Maryland, Nebraska, New Hampshire, New York, North Carolina, North Dakota, Oregon, Tennessee, Washington. Contributors with a prevalence of overweight from 16% or higher include California, Connecticut, Massachusetts, New Jersey, Rhode Island, South Dakota, Texas Virginia, and the following Indian Tribal Organizations and U.S. territories: Cheyenne River Sioux Tribe (SD), Inter-Tribal Council of Arizona, Navajo Nation, Rosebud Sioux (SD), Standing Rock Sioux (ND), and Three Affiliated Tribes (ND). States with no data include Alaska, Wyoming, Oklahoma, and Delaware. Data for this map came from 2009 National PedNSS Table 6D and is available at http://www.cdc.gov/pednss/pednss_tables/tables_numeric.htm.

Although the map shows no clear geographic pattern of obesity prevalence, six ITOs that participated in the PedNSS were among those contributors with the highest prevalence of obesity. No contributor had a prevalence of obesity at or less than the expected level of 5.0% (see Table 2 at the end of this report).

Pediatric Behavioral Indicators

Television Viewing

To prevent obesity and a variety of other problems during childhood, the American Academy of Pediatrics (AAP) recommends that parents limit the amount of time that children spend watching television or other media to no more than 1–2 hours per day for children aged 2 years or older. AAP also discourages exposure to any television for infants and children younger than age 2 years.¹⁸

In 2004, the PedNSS began monitoring the proportion of children from birth to age 4 years who view 2 hours or less of television (including videotapes and DVDs) per day. Mothers of children in this age range report data on their children's television viewing.

In 2009, a total of 24 PedNSS contributors^{§§} (44%) reported supplementary television viewing data, which indicated that 77.3% of children aged 2–4 years viewed 2 hours or less of television per day. The proportion of children meeting the AAP's recommendation varied by race and ethnicity.

Rates were lowest among black (67.9%) and Hispanic (72.7%) children and highest among white (84.4%) children.

Smoking in the Household

Exposure to secondhand tobacco smoke is a risk factor associated with low birthweight, sudden infant death, poor growth, and decreased lung function in young children.¹⁹ In 2004, the PedNSS began monitoring smoking in the household, defined as whether anyone in the child's household currently smokes inside the home. Mothers of children report these data.

In 2009, a total of 32 PedNSS contributors^{¶¶} (58%) reported supplementary data showing that 10.2% of children were exposed to household smoke. Exposure was highest for white (19.3%) and black (10.7%) children and lowest among Asian or Pacific Islander (4.7%) children. Exposure among children younger than age 1 year was lower (9.2%) than among children from birth to age 4 years (11.1%).

^{§§} Contributors of supplementary television viewing data include Indiana, Iowa, Kansas, Kentucky, Maryland, Michigan, Missouri, Nebraska, New Hampshire, New Jersey, New York, North Dakota, Rhode Island, South Carolina, Utah, West Virginia, Wisconsin, the District of Columbia, the U.S. Virgin Islands, and the following ITOs: the Inter Tribal Council of Arizona, the Cheyenne River Sioux (SD), the Rosebud Sioux (SD), the Standing Rock Sioux (ND), and the Three Affiliated Tribes (ND).

^{¶¶} Contributors of supplementary data on exposure to household smoking include Arizona, Arkansas, California, Connecticut, Illinois, Indiana, Iowa, Kansas, Kentucky, Maine, Maryland, Michigan, Missouri, Nebraska, New Hampshire, New Jersey, New York, North Dakota, Rhode Island, South Carolina, Utah, Virginia, West Virginia, Wisconsin, the District of Columbia, the U.S. Virgin Islands, and the following ITOs: the Inter Tribal Council of Arizona, the Cheyenne River Sioux Tribe (SD), the Navajo Nation (AZ), the Rosebud Sioux Tribe (SD), the Standing Rock Sioux Tribe (ND), and the Three Affiliated Tribes (ND).

Pediatric Health Progress Review

Advances in nutrition and health indicators were observed in the PedNSS population from 2000 through 2009 (Figure 9). The prevalence of high birthweight decreased, with the greatest improvement occurring among American Indian or Alaska Native and Hispanic children. Substantial improvements occurred in the prevalence of infants ever breastfed. The largest improvement in the prevalence of ever being breastfed occurred among black children. Overall, short stature remained stable during 2000–2009.

The 2009 PedNSS also indicated areas of concern. Although the low birthweight rate was stable during 2000–2009, it remained high. The *Healthy People 2010* objective to reduce the prevalence of low birthweight to 5.0%⁴ continues to be unmet. The prevalence of anemia stayed about the same, but it remains high among all racial and ethnic groups.

Obesity is a major public health problem that has steadily increased in the United States. The proportion of children from birth to age 4 years who were obese increased by 1.5 percentage points in 2009 compared with the proportion in 2000. This change is a relative increase of about 11.4%.

Although Hispanic and American Indian or Alaska Native children had the highest prevalence of obesity, increases occurred among all racial and ethnic groups except Asians or Pacific Islanders. Although overweight and obesity rates increased among children from birth to age 4 years during 2000–2009, the prevalence has remained stable since 2003.

While advances have been made in breastfeeding initiation, few contributors achieved the *Healthy People 2010* objective of 75% of infants ever being breastfed.⁴ The prevalence of breastfeeding remained lowest for black infants.

Figure 9: This is a bar graph showing percent change in several pediatric health indicators from 2000 to 2009. The prevalence of low birthweight has not changed. The prevalence of high birthweight has seen a relative decrease of 23%. Ever breastfed rates have improved by 29%; breastfed 6 months rates have improved by 37%. Anemia rates have increased by 9%, short stature prevalence rates have not changed, and obesity prevalence rates have seen a relative increase of 11%.

Recommendations

The PedNSS data indicate that national and state public health programs are needed to support the following actions:

- Prevent low birthweight by promoting preconception nutrition care and outreach activities to identify pregnancy in its early stages.
- Foster early entry into comprehensive prenatal care, including the WIC program.
- Promote and support breastfeeding initiatives in public health programs, medical care systems, work sites, and communities.

- Identify successful programs and policies to support exclusive breastfeeding, especially among populations with low prevalence.
- Promote adequate dietary iron intake and screening of children at risk for iron deficiency.
- Implement interventions in specific target areas to prevent obesity and chronic diseases that have been recommended by CDC's Division of Nutrition, Physical Activity and Obesity, including
 - Increasing breastfeeding initiation, duration, and exclusivity.
 - Increasing physical activity.
 - Increasing the consumption of fruits and vegetables.
 - Decreasing the consumption of sugar-sweetened beverages.
 - Reducing the consumption of high-energy-dense foods (foods high in calories per gram weight).
 - Decreasing television viewing.

References

1. Mathews TJ, MacDorman MF. Infant mortality statistics from the 2005 period linked birth/infant death set. *National Vital Statistics Reports* 2008;57(2):1–32.
2. Institute of Medicine. *Preventing Low Birthweight*. Washington, DC: National Academy Press; 1985.
3. Hamilton BE, Martin JA, Ventura SJ. Births: preliminary data for 2008. *National Vital Statistics Reports* 2010;58(16):1–18. Available at http://www.cdc.gov/nchs/data/nvsr/nvsr58/nvsr58_16.pdf.
4. U.S. Department of Health and Human Services. *Healthy People 2010*. Volume II. 2nd edition. Washington, DC: U.S. Government Printing Office; 2000. Available at <http://www.healthypeople.gov/publications>.
5. Jolly MC, Sebire NJ, Harris JP, Regan L, Robinson S. Risk factors for macrosomia and its clinical consequences: a study of 350,311 pregnancies. *European Journal of Obstetrics & Gynecology and Reproductive Biology* 2003;11: 9–14.
6. Martin JA, Hamilton BE, Sutton PD, Ventura SJ, Mathews TJ, Kirmeyer S, et al. Births: final data for 2007. *National Vital Statistics Reports* 2010;58(24):1–125. Available at http://www.cdc.gov/nchs/data/nvsr/nvsr58/nvsr58_24.pdf.
7. Gartner LM, Morton J, Lawrence RA, Naylor AJ, O'Hare D, Schanler RJ. Breastfeeding and the use of human milk. *Pediatrics* 2005;115(2):496–506.
8. Centers for Disease Control and Prevention. *Breastfeeding Among U.S. Children Born 1999–2007, CDC National Immunization Survey*. Available at http://www.cdc.gov/breastfeeding/data/NIS_data/index.htm.
9. Ip S, Chung M, Raman G, Chew P, Magula N, DeVine D, et al. *Breastfeeding and Maternal and Infant Health Outcomes in Developed Countries*. Rockville, MD: Agency for Healthcare Research and Quality; 2007. Evidence Report/Technology Assessment No. 153. Available at <http://www.ahrq.gov/downloads/pub/evidence/pdf/brfout/brfout.pdf>.
10. Pollitt E. Iron deficiency and cognitive function. *Annual Review of Nutrition* 1993;13:521–537.

11. Idjradinata P, Pollitt E. Reversal of developmental delays in iron-deficient anaemic infants treated with iron. *Lancet* 1993;341(8836):1–4.
12. Centers for Disease Control and Prevention. Recommendations to prevent and control iron deficiency in the United States. *MMWR Recommendations and Reports* 1998;47(RR-3): 1–30.
13. World Health Organization Expert Committee on Physical Status. *Physical Status: The Use and Interpretation of Anthropometry*. Geneva: World Health Organization; 1996.
14. Mei Z, Ogden CL, Flegal KM, Grummer-Strawn LM. Comparison of the prevalence of shortness, underweight, and overweight among U.S. children aged 0 to 59 months by using the Centers for Disease Control and Prevention 2000 and the WHO 2006 growth charts. *Journal of Pediatrics* 2008;153:622–628.
15. Krebs NF, Jacobson MS; American Academy of Pediatrics Committee on Nutrition. Prevention of pediatric overweight and obesity. *Pediatrics* 2003;112(2):424–430.
16. Barlow SE; Expert Committee. Expert committee recommendations regarding the prevention, assessment, and treatment of child and adolescent overweight and obesity: summary report. *Pediatrics* 2007;120(suppl 4):S164-S192. Available at http://pediatrics.aappublications.org/cgi/content/full/120/Supplement_4/S164.
17. Ogden CL, Carroll MD, Flegal KM. High body mass index for age among U.S. children and adolescents, 2003–2006. *Journal of the American Medical Association* 2008;299(20):2401–2405. Available at <http://jama.ama-assn.org/content/299/20/2401.full>.
18. American Academy of Pediatrics, Committee on Public Education: Children, adolescents, and television. *Pediatrics* 2001;107(2):423–426.
19. Centers for Disease Control and Prevention. Vital Signs: nonsmokers' exposure to secondhand smoke—United States, 1999–2008. *Morbidity and Mortality Weekly Report* 2010;59(35):1141–1146.

Table 1. Contributors to the Pediatric Nutrition Surveillance System, 2000–2009

Alabama: 2003 – 2009
 American Samoa: 2000–2001
 Arizona: 2000 – 2009
 Arkansas: 2000 – 2009
 California: 2000 – 2009
 Cheyenne River Sioux Tribe (SD): 2000 – 2009
 Chickasaw Nation (OK): 2000 – 2005
 Colorado: 2000 – 2009
 Connecticut: 2001 – 2009
 District of Columbia: 2000 – 2009
 Florida: 2000 – 2009
 Georgia: 2000 – 2009
 Hawaii: 2000 – 2005, 2007 – 2009
 Idaho: 2000 – 2009
 Illinois: 2000 – 2009
 Indiana: 2000 – 2009
 Inter Tribal Council of Arizona: 2000 – 2009
 Iowa: 2000 – 2009
 Kansas: 2000 – 2009
 Kentucky: 2000 – 2009
 Louisiana: 2000 – 2004, 2007
 Maine: 2000 – 2005
 Maryland: 2000 – 2002, 2004 – 2009
 Massachusetts: 2006 – 2009

Michigan: 2000 – 2009
 Minnesota: 2000 – 2009
 Mississippi: 2000, 2007 – 2009
 Missouri: 2000 – 2009
 Montana: 2003 – 2009
 Navajo Nation: 2000 – 2009
 Nebraska: 2000 – 2009
 Nevada: 2000 – 2009
 New Hampshire: 2000 – 2009
 New Jersey: 2000 – 2009
 New Mexico: 2000 – 2002, 2004 – 2009
 New York: 2000, 2002 – 2009
 North Carolina: 2000 – 2009
 North Dakota: 2000 – 2005, 2007 – 2009
 Ohio Child Health: 2006 – 2009
 Ohio WIC: 2000 – 2009
 Oregon: 2000 – 2003, 2006 – 2009
 Pennsylvania: 2000 – 2009
 Puerto Rico: 2002 – 2009
 Rhode Island: 2006 – 2009
 Rosebud Sioux Tribe (SD): 2000 – 2009
 South Carolina: 2000 – 2009
 South Dakota: 2000 – 2009
 Standing Rock Sioux Tribe (ND): 2000 – 2006, 2008-2009
 Tennessee: 2000 – 2009
 Texas: 2003 – 2009
 Three Affiliated Tribes (ND): 2007 – 2009
 Utah: 1999 – 2005
 Vermont: 2000 – 2009
 Virgin Islands: 2008 – 2009
 Virginia: 2003 – 2009
 Washington: 1999 – 2002, 2006 – 2009
 West Virginia: 2000 – 2009
 Wisconsin: 2000 – 2009
 Wyoming: 2001-2005

	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009
Number of Contributors	46	45	47	48	49	50	49	47	51	52
Total Records Submitted (x1000)	5,019	4,943	5,519	6,359	6,930	7,118	7,559	7,996	8,165	8,939

State-specific prevalence of selected nutrition indicators for children aged <5 years, 2009 Pediatric Nutrition Surveillance System

Contributors	LBW*	HBW†	Ever Breastfed*	Breastfed 6 Months**	Anemia‡	Short Stature§	Obesity¶
Alabama	11.8	4.3	37.2	NA	19.0	6.2	14.4
Arizona	8.0	6.6	65.5	27.8	13.4	6.2	14.3
Arkansas	10.0	5.2	45.3	15.5	18.6	7.1	14.2
California	6.2	7.3	NA	NA	14.0	5.1	17.0
Cheyenne River Sioux	7.5	5.0	27.7	14.4	9.7	1.1	19.0
Colorado	9.3	4.5	76.4	29.6	13.1	7.5	9.0
Connecticut	8.9	7.0	64.9	26.9	8.7	3.4	16.0
District of Columbia	11.0	5.1	52.5	28.6	27.9	7.6	13.6
Florida	9.5	5.7	70.6	28.0	17.4	4.5	13.7
Georgia	10.1	5.2	50.3	20.6	14.2	6.1	14.2
Hawaii	9.3	6.5	84.4	41.3	11.9	6.4	9.3
Idaho	7.4	7.2	82.9	29.3	11.3	6.5	11.9
Illinois	9.1	7.0	63.1	24.0	11.2	6.7	14.6
Indiana	9.1	6.2	61.9	23.5	14.7	6.2	14.3
Inter Tribal Council of	6.5	8.7	66.1	24.0	14.1	5.3	24.2
Iowa	8.2	8.1	NA	NA	8.0	5.5	15.0
Kansas	8.5	6.4	69.8	20.9	10.1	6.3	13.2
Kentucky	9.9	6.3	37.8	12.2	12.9	6.0	15.8
Louisiana	11.9	3.5	29.0	12.8	20.7	8.6	12.4
Maine	NA	NA	NA	27.6	13.2	3.6	14.7
Maryland	10.1	5.6	60.1	29.2	22.4	5.8	15.8
Massachusetts	8.7	7.9	74.0	28.3	11.3	4.5	16.8
Michigan	9.4	6.8	53.5	18.5	15.0	5.6	13.7
Minnesota	7.4	8.7	73.6	32.3	10.9	4.9	13.1
Mississippi	NA	NA	NA	NA	16.2	9.4	13.9
Missouri	9.3	6.4	56.7	14.7	17.0	6.6	13.9
Montana	8.0	7.6	73.7	32.9	8.5	5.7	12.5
Navajo Nation (AZ)	7.0	7.2	75.4	33.9	NA	4.8	17.3
Nebraska	8.4	6.6	69.9	23.7	14.6	6.2	14.2
Nevada	8.2	6.0	61.5	25.6	9.8	6.5	13.9
New Hampshire	8.3	10.0	68.8	26.0	13.8	6.2	14.4
New Jersey	8.8	6.3	59.2	32.9	18.8	6.1	18.4
New Mexico	9.4	4.6	NA	NA	12.9	7.2	12.0
New York	8.9	6.6	74.4	38.8	11.8	4.3	14.4
North Carolina	9.9	6.5	60.5	21.9	13.4	5.5	15.2
North Dakota	8.0	8.4	58.7	20.3	9.3	4.5	14.1
Ohio	10.4	5.9	47.4	16.1	13.5	6.5	12.3
Oregon	6.8	9.7	91.2	43.2	14.1	5.0	15.0
Pennsylvania	9.9	6.2	45.3	20.2	20.6	6.4	12.0
Puerto Rico	NA	NA	55.3	NA	5.6	10.3	18.1
Rhode Island	9.2	7.7	60.3	21.4	17.1	6.0	16.2
Rosebud Sioux (SD)	4.5	8.7	57.4	NA	24.0	1.1	18.7
South Carolina	10.3	5.1	43.4	13.8	18.1	6.8	13.3
South Dakota	6.9	8.3	59.4	19.2	8.8	5.9	16.4
Standing Rock Sioux	6.7	10.0	42.6	NA	16.7	2.0	26.1
Tennessee	10.4	5.3	39.7	15.7	7.8	6.1	14.0
Texas	8.9	5.6	72.1	42.3	21.2	6.7	16.0
Three Affiliated Tribes	NA	NA	NA	NA	19.5	1.6	31.5
Utah	8.6	5.9	75.2	NA	11.2	7.4	8.8
Vermont	8.0	10.3	73.4	30.2	6.6	4.6	13.2
U.S. Virgin Islands	9.2	4.9	89.6	61.4	11.5	4.7	11.9
Virginia	10.0	6.0	58.4	22.0	15.8	6.1	18.0
Washington	7.1	9.2	85.7	39.5	10.4	5.0	14.4
West Virginia	9.8	5.8	44.8	13.5	4.7	5.1	13.4
Wisconsin	8.6	7.7	66.9	27.5	11.1	5.6	13.7
National PedNSS	8.9	6.4	61.7	27.0	14.9	6.0	14.7

Abbreviations: LBW = low birthweight; HBW = high birthweight; NA = not available.

* Defined as birthweight <2,500 grams.

† Defined as birthweight >4,000 grams.

*Infants born during the reporting period are included in analysis.

**Infants who turned 6 months of age during the reporting period by/on their date of visit are included in analysis.

‡ Adjusted for altitude. For children aged 6 months to 2 years: Hemoglobin (Hb) is <11.0 g/d, or hematocrit (Hct) is <32.9%. For children aged 2–5 years: Hb is <11.1 g/dL, or Hct is <33.0%.

§ Based on the 2000 CDC Growth Chart. Defined as <5th percentile length-for-age for children aged <2 years or height-for-age for children aged 2 years or older.