

Maternity Practices in Infant Nutrition and Care in the Pacific Islands —2011 mPINC Survey



This report provides data from the 2011 mPINC survey for the Pacific Islands of American Samoa, Guam, and Saipan. It describes specific opportunities to improve mother-baby care at hospitals and birth centers on these Pacific Islands in order to more successfully meet national quality of care standards for perinatal care.

More information is at www.cdc.gov/mpinc

Breastfeeding is a National Priority

Breastfeeding is associated with decreased risk for infant morbidity and mortality as well as maternal morbidity,¹ and provides optimal infant nutrition. *Healthy People 2020*² establishes breastfeeding initiation, continuation, and exclusivity as national priorities.

Changes in Maternity Care Practices Improve Breastfeeding Rates

Maternity practices in hospitals and birth centers can influence breastfeeding behaviors during a period critical to successful establishment of lactation.³ Abundant literature, including a Cochrane review, document that institutional changes in maternity care practices to make them more supportive of breastfeeding increase initiation and continuation of breastfeeding.⁴

Breastfeeding Support in Pacific Islands Facilities

Strengths

	Availability of Prenatal Breastfeeding Instruction All facilities (100%) on these Pacific Islands include breastfeeding education as a routine element of their prenatal classes.	Prenatal education about breastfeeding is important because it provides mothers with a better understanding of the benefits and requirements of breastfeeding, resulting in improved breastfeeding rates.
	Documentation of Mothers' Feeding Decisions Staff at all (100%) facilities on these Pacific Islands consistently ask about and record mothers' infant feeding decisions.	Standard documentation of infant feeding decisions is important to adequately support maternal choice.

Needed Improvements

	Appropriate Use of Breastfeeding Supplements No facilities (0%) on these Pacific Islands adhere to standard clinical practice guidelines against routine supplementation with formula, glucose water, or water.	The American Academy of Pediatrics (AAP) and the American College of Obstetricians and Gynecologists (ACOG) Guidelines for Perinatal Care recommend against routine supplementation because supplementation with formula and/or water makes infants more likely to receive formula at home and stop breastfeeding prematurely.
	Inclusion of Model Breastfeeding Policy Elements Only 40% of facilities on these Pacific Islands have comprehensive breastfeeding policies including all model breastfeeding policy components recommended by the Academy of Breastfeeding Medicine (ABM).	The ABM model breastfeeding policy elements are the result of extensive research on best practices to improve breastfeeding outcomes. Facility policies determine the nature of care that is available to patients. Facilities with comprehensive policies consistently have the highest rates of exclusive breastfeeding, regardless of patient population characteristics such as ethnicity, income, and payer status.
	Provision of Hospital Discharge Planning Support Only 20% of facilities on these Pacific Islands provide hospital discharge care including a phone call to the patient's home, opportunity for follow-up visit, and referral to community breastfeeding support.	The American Academy of Pediatrics (AAP) clinical practice guidelines recommend examination of the newborn by a qualified health care professional within 48 hours of hospital discharge in order to assess breastfeeding. Ensuring post discharge ambulatory support improves breastfeeding outcomes.
	Initiation of Mother and Infant Skin-to-Skin Care Only 60% of facilities on these Pacific Islands initiate skin-to-skin care for at least 30 minutes upon delivery of the newborn.	Upon delivery, the newborn should be placed skin-to-skin with the mother and allowed uninterrupted time to initiate and establish breastfeeding in order to improve infant health outcomes and reduce the risk of impairment of the neonatal immune system from unnecessary non-breast milk feeds.

Every two years, CDC administers the national Maternity Practices in Infant Nutrition and Care (mPINC) survey to all hospitals and birth centers in the U.S. that provide maternity care.

Data from this survey can be used to establish evidence-based, breastfeeding-supportive maternity practices as standards of care in hospitals and birth centers across the US. Improved care will help meet *Healthy People 2020* breastfeeding objectives and will help improve maternal and child health nationwide.



Pacific Islands Summary —2011 mPINC Survey

Survey Method At each facility, the person who is the most knowledgeable about the facility's maternity practices related to healthy newborn feeding and care completes the CDC mPINC survey.

Response Rate 100% of the 5 eligible facilities on these Pacific Islands responded to the 2011 mPINC Survey. Each participating facility received its facility-specific mPINC benchmarking report in October 2012.

Pacific Islands's Composite Quality Practice Score* **71**

(out of 100)

Pacific Islands's Composite Rank† **21**

(out of 53)

mPINC Dimension of Care	PI Quality Practice Subscore*	Ideal Response to mPINC Survey Question	Percent of PI Facilities with Ideal Response	PI Item Rank†
Labor and Delivery Care	61	Initial skin-to-skin contact is ≥30 min w/in 1 hour (vaginal births)	60	21
		Initial skin-to-skin contact is ≥30 min w/in 2 hours (cesarean births)	25	48
		Initial breastfeeding opportunity is w/in 1 hour (vaginal births)	40	50
		Initial breastfeeding opportunity is w/in 2 hours (cesarean births)	25	50
		Routine procedures are performed skin-to-skin	60	4
Feeding of Breastfed Infants	68	Initial feeding is breast milk (vaginal births)	60	49
		Initial feeding is breast milk (cesarean births)	25	53
		Supplemental feedings to breastfeeding infants are rare	0	52
		Water and glucose water are not used	80	32
Breastfeeding Assistance	87	Infant feeding decision is documented in the patient chart	100	---
		Staff provide breastfeeding advice & instructions to patients	80	44
		Staff teach breastfeeding cues to patients	60	51
		Staff teach patients not to limit suckling time	20	52
		Staff directly observe & assess breastfeeding	80	40
		Staff use a standard feeding assessment tool	80	9
		Staff rarely provide pacifiers to breastfeeding infants	100	---
Contact Between Mother and Infant	81	Mother-infant pairs are not separated for postpartum transition	40	46
		Mother-infant pairs room-in at night	100	---
		Mother-infant pairs are not separated during the hospital stay	80	3
		Infant procedures, assessment, and care are in the patient room	0	36
		Non-rooming-in infants are brought to mothers at night for feeding	100	---
Facility Discharge Care	86	Staff provide appropriate discharge planning (referrals & other multi-modal support)	20	41
		Discharge packs containing infant formula samples and marketing products are not given to breastfeeding patients	100	---
Staff Training	37	New staff receive appropriate breastfeeding education	0	51
		Current staff receive appropriate breastfeeding education	0	51
		Staff received breastfeeding education in the past year	20	52
		Assessment of staff competency in breastfeeding management & support is at least annual	60	18
Structural & Organizational Aspects of Care Delivery	74	Breastfeeding policy includes all 10 model policy elements	40	5
		Breastfeeding policy is effectively communicated	100	---
		Facility documents infant feeding rates in patient population	75	17
		Facility provides breastfeeding support to employees	60	38
		Facility does not receive infant formula free of charge	20	17
		Breastfeeding is included in prenatal patient education	100	---
Facility has a designated staff member responsible for coordination of lactation care	60	41		

* Quality Practice scores range from 0 to 100 for each question, dimension of care, facility, and state. The highest, best possible score for each is 100. Each facility and state's "Composite Quality Practice Score" is made up of subscores for practices in each of 7 dimensions of care.

† Ranks range from 1 to 52, with 1 being the highest rank. In case of a tie, both are given the same rank.

- State ranks are not shown for survey questions with 90% or more facilities reporting ideal responses.

References

- Ip S, Chung M, Raman G, et al. Breastfeeding and maternal and infant health outcomes in developed countries. Rockville, MD: US Dept of Health and Human Services, Agency for Healthcare Research and Quality; 2007.
- US Dept of Health and Human Services. Healthy People 2020 Summary of Objectives: Maternal, Infant, and Child Health. Available at <http://www.healthypeople.gov/2020/topics/objectives/2020/pdfs/MaternalChildHealth.pdf>
- DiGirolamo AM, Grummer-Strawn LM, Fein S. Effect of maternity care practices on breastfeeding. Pediatrics 2008;122, Supp 2:S43-9.
- Fairbank L, O'Meara S, Renfrew MJ, Woolridge M, Snowden AJ, Lister-Sharp D. A systematic review to evaluate the effectiveness of interventions to promote the initiation of breastfeeding. Health Technology Assessment 2000;4:1-171.

Improvement is Needed in Maternity Care Practices and Policies on these Pacific Islands.

Many opportunities exist to protect, promote, and support breastfeeding mothers and infants on these Pacific Islands.

Potential opportunities:

- Examine Pacific Islands regulations for maternity facilities and evaluate their evidence base.
- Sponsor a Pacific Islands-wide summit of key decision-making staff at maternity facilities to highlight the importance of evidence-based practices for breastfeeding.
- Encourage and support hospital staff across these Pacific Islands to be trained in providing care that supports mothers to breastfeed.
- Establish links among maternity facilities and community breastfeeding support networks on these Pacific Islands.
- Implement evidence-based practices in medical care settings across these Pacific Islands that support mothers' efforts to breastfeed.
- Integrate maternity care into related hospital-wide Quality Improvement efforts across these Pacific Islands.
- Promote utilization of the Joint Commission's Perinatal Care Core Measure Set including exclusive breast milk feeding at hospital discharge in Pacific Islands hospital data collection systems.

Questions about the mPINC survey?

Information about the mPINC survey, benchmark reports, scoring methods, and complete references are available at: www.cdc.gov/mpinc

For more information:

Division of Nutrition, Physical Activity, and Obesity
Centers for Disease Control and Prevention
Atlanta, GA USA

February 2013