



Source: [Interactive Atlas of Heart Disease](#)

Facts on Women and Heart Disease

- Heart disease is the leading cause of death for women in the United States, killing 289,758 women in 2013— that's about 1 in every 4 female deaths.¹
- Although heart disease is sometimes thought of as a "man's disease," around the same number of women and men die each year of heart disease in the United States. Despite increases in awareness over the past decade, only 54% of women recognize that heart disease is their number 1 killer.²
- Heart disease is the leading cause of death for African American and white women in the United States. Among Hispanic women, heart disease and cancer cause roughly the same number of deaths each year. For American Indian or Alaska Native and Asian or Pacific Islander women, heart disease is second only to cancer.³
- About 5.8% of all white women, 7.6% of black women, and 5.6% of Mexican American women have coronary heart disease.⁴
- Almost two-thirds (64%) of women who die suddenly of coronary heart disease have no previous symptoms.⁴ Even if you have no symptoms, you may still be at risk for heart disease.

Symptoms

While some women have no symptoms, others experience angina (dull, heavy to sharp chest pain or discomfort), pain in the neck/jaw/throat or pain in the upper abdomen or back. These may occur during rest, begin during physical activity, or be triggered by mental stress.⁶

Women are more likely to describe chest pain that is sharp, burning and more frequently have pain in the neck, jaw, throat, abdomen or back.⁶

Sometimes heart disease may be silent and not diagnosed until a woman experiences signs or symptoms of a heart attack, heart failure, an arrhythmia,⁶ or stroke.

These symptoms may include

- Heart Attack: Chest pain or discomfort, upper back pain, indigestion, heartburn, nausea/vomiting, extreme fatigue, upper body discomfort, and shortness of breath.
- Arrhythmia: Fluttering feelings in the chest (palpitations).⁶
- Heart Failure: Shortness of breath, fatigue, swelling of the feet/ankles/legs/abdomen.
- Stroke: Sudden weakness, paralysis (inability to move) or numbness of the face/arms/legs, especially on one side of the body. Other symptoms may include: confusion, trouble speaking or understanding speech, difficulty seeing in one or both eyes, shortness of breath, dizziness, loss of balance or coordination, loss of consciousness, or sudden and severe headache.⁷

Risk Factors

High blood pressure, high LDL cholesterol, and smoking are key risk factors for heart disease. About half of Americans (49%) have at least one of these three risk factors.⁵

Several other medical conditions and lifestyle choices can also put people at a higher risk for heart disease, including:

- Diabetes
- Overweight and obesity
- Poor diet
- Physical inactivity
- Excessive alcohol use

Screening

To reduce your chances of getting heart disease it's important to⁸

- Know your blood pressure. Having uncontrolled blood pressure can result in heart disease. High blood pressure has no symptoms so it's important to have your blood pressure checked regularly.
- Talk to your healthcare provider about whether you should be tested for diabetes. Having uncontrolled diabetes raises your chances of heart disease.
- Quit smoking.
- Discuss checking your cholesterol and triglycerides with your healthcare provider.
- Make healthy food choices. Being overweight and obese raises your risk of heart disease.
- Limit alcohol intake to one drink a day.
- Lower your stress level and find healthy ways to cope with stress.

CDC's Public Health Efforts

- [State Public Health Actions to Prevent and Control Chronic Diseases](#)
- [Million Hearts®](#)
- [WISEWOMAN](#)

For More Information

For more information on women and heart disease, visit the following Web sites—

- [Centers for Disease Control and Prevention](#)
- [U.S. Department of Health and Human Services, Office on Women's Health](#)
- [American Heart Association](#)
- [National Heart, Lung, and Blood Institute](#)

References

1. Xu, JQ, Murphy, SL., Kochanek, KD, Bastian, BA. Deaths: Final data for 2013. National Vital Statistics Report. 2016;64(2).
2. Mosca L, Mochari-Greenberger H, Dolor RJ, Newby LK, Robb KJ. Twelve-year follow-up of American women's awareness of cardiovascular disease risk and barriers to heart health. Circulation: Cardiovascular Quality Outcomes. 2010;3:120-7.
3. Heron M. [Deaths: Leading causes for 2008 \[PDF-2.7M\]](#). National vital statistics reports. 2012;60(6).
4. Roger VL, Go AS, Lloyd-Jones DM, Benjamin EJ, Berry JD, Borden WB, et al. [Heart disease and stroke statistics—2012 update: a report from the American Heart Association](#). Circulation. 2012;125(1):e2–220.
5. CDC. [Million Hearts: strategies to reduce the prevalence of leading cardiovascular disease risk factors](#). United States, 2011. MMWR 2011;60(36):1248–51.
6. National Heart, Lung and Blood Institute. What Are the Signs and Symptoms of Heart Disease? [cited 2013 July 19, 2013]; Available from: www.nhlbi.nih.gov/health/health-topics/hdw/signs.html.
7. National Heart Lung and Blood Institute. What are the Signs and Symptoms of a Stroke? [cited 2013 July 19, 2013]; Available from: www.nhlbi.nih.gov/health/health-topics/topics/stroke/signs.html.
8. U.S. Department of Health and Human Services, Office on Women's Health. Heart Disease: Frequently Asked Questions. 2009. [cited 2013 July 19, 2013]; Available from: <http://www.womenshealth.gov/publications/our-publications/fact-sheet/heart-disease.pdf> [PDF-1.7M].

*Links to non-Federal organizations are provided solely as a service to our users. Links do not constitute an endorsement of any organization by CDC or the Federal Government, and none should be inferred. The CDC is not responsible for the content of the individual organization Web pages found at this link.