

---

# Canonical Insertion-Deletion Markers for Rapid DNA Typing of *Francisella tularensis*

Pär Larsson,\*† Kerstin Svensson,\*† Linda Karlsson,\* Dimitri Guala,\* Malin Granberg,\*  
Mats Forsman,\* and Anders Johansson\*†

To develop effective and accurate typing of strains of *Francisella tularensis*, a potent human pathogen and a putative bioterrorist agent, we combined analysis of insertion-deletion (indel) markers with multiple-locus variable-number tandem repeat analysis (MLVA). From 5 representative *F. tularensis* genome sequences, 38 indel markers with canonical properties, i.e., capable of sorting strains into major genetic groups, were selected. To avoid markers with a propensity for homoplasy, we used only those indels with 2 allelic variants and devoid of substantial sequence repeats. MLVA included sequences with much diversity in copy number of tandem repeats. The combined procedure allowed subspecies division, delineation of clades A.I and A.II of subspecies *tularensis*, differentiation of Japanese strains from other strains of subspecies *holarctica*, and high-resolution strain typing. The procedure uses limited amounts of killed bacterial preparations and, because only 1 single analytic method is needed, is time- and cost-effective.

*Francisella tularensis* is a highly infectious, facultative intracellular pathogen and the causative agent of the zoonotic disease tularemia. Based on virulence tests and biochemical assays, *F. tularensis* is divided into 4 subspecies, a division that has recently been corroborated by genetic typing (1,2). Each subspecies shows a discrete natural geographic distribution and also varying degrees of virulence (3). Human disease caused by *F. tularensis* subsp. *tularensis* may be fulminate or even lethal, whereas disease caused by other subspecies is less severe, although often incapacitating and protracted (4). In addition, recent molecular and epidemiologic analyses of natural isolates of *F. tularensis* subsp. *tularensis* suggest a population split of the subspecies into 2 major groups of isolates, which differ in virulence and geographic distribution (5–7).

\*Swedish Defence Research Agency, Umeå, Sweden; and †Umeå University, Umeå, Sweden

Robust and rapid typing schemes for *F. tularensis* are needed, not only because of their use in clinical and public health work but also because of a rising concern associated with risks for bioterrorism (4,8). Because of its virulence, *F. tularensis* is included among the top 6 “category A” potential bioterrorism agents believed to have the greatest potential for adverse public health effect with mass casualties. If deliberate release of the organism is suspected, the need to understand the pathogenic potency of an isolate and also its putative origin will be urgent.

In standard medical practice, subspecies determination of *F. tularensis* typically involves biochemical fermentations. Such analyses are labor-intensive, hampered by the fastidious growth characteristics of the organism on artificial media, and associated with a substantial risk for laboratory-acquired infections (2,9).

Several DNA-based methods have been found useful for typing of *F. tularensis* at the subspecies level (1,10–13). Among these, pulsed-field gel electrophoresis (PFGE) is more widely adopted and was recently proposed for diagnostic and epidemiologic work on *F. tularensis* by PulseNet laboratories throughout the United States (7). PFGE typing is, however, far from ideal for the purpose. It involves making concentration-adjusted suspensions of live bacteria, which has the potential for creating infectious aerosols, is time-consuming, produces complex banding pattern data, and has a restrictive discriminatory capacity when applied to *F. tularensis* (7,14–17).

High-resolution typing of *F. tularensis* is currently attainable only by the use of multilocus variable-number tandem repeat analysis (MLVA). The method capitalizes on differences among strains in copy numbers of sequence repeats at multiple genomic loci. MLVA has been successfully applied in epidemiologic studies on tularemia (5,6,18,19). Killed bacterial preparations can be used in

the assay and, in contrast to PFGE, MLVA produces discrete-character numeric data, which are well suited for easy transfer among laboratories. For discrimination of strains of *F. tularensis*, MLVA is the obvious choice.

A limitation inherent in MLVA is the risk for erroneous estimates of relationships among strains at larger genetic distances. The high rates at which MLVA markers mutate (20,21), and possible functional constraints on these sequences, may cause homoplasy effects, i.e., share of mutational changes for reasons other than common ancestry (22,23), implicating a risk for spurious strain affiliation. In work on *Bacillus anthracis*, the issue was addressed by analysis of single-nucleotide polymorphisms (SNPs), which exhibited canonical properties for resolving major genetic lineages (24). In a hierarchical typing approach, which conformed with concepts of traditional bacterial taxonomy, a 2-step procedure was suggested, including assay of canonical SNPs for resolution of major genetic clades and MLVA for high-resolution typing (24). A limitation of the procedure is that it involves 2 assays, thus increasing time and cost.

When aiming to construct an improved typing strategy for *F. tularensis*, we focused on insertion-deletion (indel) markers. By definition, indels are caused by insertion or deletion of  $\geq 1$  base pairs of a DNA molecule. Among indels, the evolutionary rates diverge widely. When used as a complement to MLVA, more slowly evolving indels, i.e., loci displaying a relatively low degree of variability, would be preferable. A practical reason to use canonical indel markers was that fragment analysis can be used for simultaneous assay of both indel and MLVA markers, thereby minimizing time and cost.

We identified indel markers with canonical properties in *F. tularensis* and used them to resolve major genetic lineages of the species. We also developed a strategy that combines indel analysis with MLVA for rapid and accurate discrimination of isolates of the species.

## Material and Methods

### Genome Sequences, Strains, and DNA Preparations

We used genome sequences for the 5 strains, U112 (aka FSC040, ATCC 15482), FSC147 (GIEM 543), SCHU S4 (FSC237), OSU18, and LVS (FSC155) (online Appendix Table 1, available from [www.cdc.gov/EID/content/13/11/1725-appT1.htm](http://www.cdc.gov/EID/content/13/11/1725-appT1.htm)), for in silico work, and in total, 23 isolates (online Appendix Table 2, available from [www.cdc.gov/EID/content/13/11/1725-appT2.htm](http://www.cdc.gov/EID/content/13/11/1725-appT2.htm), and online Appendix Table 3, available from [www.cdc.gov/EID/content/13/11/1725-appT3.htm](http://www.cdc.gov/EID/content/13/11/1725-appT3.htm)) were selected for the experimental work. These were chosen to represent each of the 4 currently recognized *F. tularensis* subspecies and were selected from the *Francisella* Strain Collection (FSC)

maintained at the Swedish Defence Research Agency, Umeå, Sweden. Bacteria were grown on modified Thayer-Martin agar (25), suspended in phosphate-buffered saline, and immediately heat killed. DNA was prepared by using silica and guanidine isothiocyanate buffer (26). Extended information on strains and, when appropriate, GenBank accession numbers, are available in online Appendix Tables 1–3.

### Identification and Selection of Indel Markers

Multiple alignment of genomic sequences for *F. tularensis* strains U112, FSC147, SCHU S4, OSU18, and LVS was performed by using Mauve 2.0  $\beta$  multiple alignment software (27) and the progressive alignment option. The output file produced by Mauve was parsed by using a custom Perl script to retrieve multiple aligned sequences for indel loci that fulfilled the following criteria: 1) the loci should exist in all compared strains, 2) only 2 allelic variants should exist, 3) at least 25 bp of sequences lacking other indels should flank identified loci, 4) indels should be 5- to 200-bp long, and 5) direct repeated sequences of substantial length should not be present at indel loci because such sequences may increase the risk for homoplastic mutation.

### Primer Design and PCRs

Oligonucleotide primers for PCR amplification were designed by using the Primer3 tool (28) and a Perl script to supply aligned sequences and required coordinate information. To reduce experimental cost, the forward primer of each primer pair was synthesized with an additional 17-bp M13 tail added to the 5' end of the primer (Table). This enabled the use of fluorescently labeled M13 PCR primers to simultaneously amplify marker loci and label the PCR amplicons. The M13 primers were labeled terminally with D2-PA, D3-PA, or D4-PA dyes at the 5' end (Proligo Primers and Probes, Hamburg, Germany).

PCR amplification was performed in 96-well microtiter plates. Each reaction mixture contained 0.15 mmol/L dNTP, 0.6 U DyNAzymeII polymerase (F-501L, Finnzymes, Espoo, Finland), 1  $\mu$ L PCR buffer for DyNAzyme DNA polymerase (Finnzymes), 2  $\mu$ L of template DNA (20 ng/ $\mu$ L), 0.3 pmol/L forward primer, 0.8 pmol/L reverse primer, and 0.8 pmol/L labeled M13 primer. Filtered sterile water was added to a final volume of 25  $\mu$ L. The PCR reactions were performed in a MyCycler thermal cycler (BioRad, Hercules, CA, USA) with the following program: 95°C for 2 min; 15 cycles of 95°C for 30 s, 56°C for 30 s, 72°C for 45 s; 20 cycles of 95°C for 30 s, 51°C for 30 s, and 72°C for 45s; and then a 7-min final extension step at 72°C. MLVA was performed as previously described, except modified to use fluorescence-labeled forward primers (6). The physical distribution of 38 selected indel markers identified in this

Table. Insertion-deletion loci, genomic locations, and primers

Ftind locus*	Positions†	Pattern	Forward primer sequence (5'→3')‡	Reverse primer sequence (5'→3')
1	1152573–1152844	12222	TCTCGTGACAGAGCTTTACAA	GGGAGAATTGATTATGGCTTAC
2	895732–896067	12222	AGCAGCGTATCGAAGAGATAG	TAAATCTAGTTGGCTGAGTAATAAAGTC
3	769704–770059	12222	CAAACCTAATTGCTCCAGAAC	GCAGCATATCTTTGGTCATCTAT
4	520340–520556	12222	TTTGAAAAGCTAGAAAAAGATGC	ACCAAGAATATTTAAAGCCAAATC
5	1628363–1628558	12222	AACTAAGTTGTTTTAGTGGGTTCC	CAATTTTATACCCAGTTAATATTTGA
6	562346–562675	12222	CAACAATCTCACCATTACCTAAAA	GCTAGGCAAGCCATTATATTTATC
7	688418–688771	12222	CCAAAAATATACCAAAATATCCTATCA	ATTATGCAATATCACAAGTTCCA
8	198167–198521	12222	GTGACCTAATCAAGAGCAACTAA	ATCTGCATACTTGAGTAAATGCTT
9	1830520–1830768	11211	CTCAAGAAATTAAGGGATGAGTT	ATTTGCTCAGTACCTGCTAATGTA
10	1113820–1114081	11211	CATTCTAGTRATAGTCCTGCT	ATTAAGCTTCAACACTATCATCATCT
11	1238526–1238784	11211	TACTTTTAATGCTTCAGCGACA	AATCACCATAAACCAGACAAC
12	725006–725258	11211	GCCTATGCTGGTAAAGTTGG	TCACCAATAGCTTCCATAACAC
13	1490938–1491179	12211	AACTCCTGGTTTCCCACAC	GCTACAAAACCTCACTATGTTCCAGAC
14	625186–625399	12211	GACTGAACAACAACCTGGATTATCAC	TGTAGTCCATTAGGGCAGTAATCTT
15	573074–573303	12111	GGTTTTGTTGCTAAATCTGC	ACGCTGATCATCAATCATTC
16	1628145–1628393	12111	TCCTTTAAAGAAACGGCATA	TCTGTACGGAACCCACTAAA
17	239966–240157	12111	CATGAAAACCTGGTTATAGCTGA	GCGCAAGATCAGCTTAGTT
18	439229–439434	12111	AGAGTTAACCCATTCAACAAGA	GGCAAGGTTTCTGGATAGAC
19	408363–408515	12111	TTTGATAGCTCAAATGCAAGA	AGTAGCTTGCCTCTTTTCT
20	602863–603177	11122	AAATCATTTAACAATTGGTATCTTT	TAGCTCTGAGTTAGAAAACTCG
21	271531–271863	11122	TCTTCTGTATAAGATGCGCTAAA	GGTTAAGTTAGGGCAATGTAAGAT
22	5648–5976	11122	TGACAAAGAAGACTAAGCACAAAT	GGTTTGATAAATGCAAACTATATGAT
23	1062332–1062553	11122	TCAACCGGCTTTATGAGAGTA	TATTACGAGACCGAAAAATACGATA
24	1641399–1641720	11122	AATTCAAAAAGCGATAAGTAACCT	GCCAGCAACATACTCTTTTGT
25	267938–268267	11122	AAATTAAGCAAGGACAGGTTTAT	TCCATAGTTATTTCAACTTGGTTT
26	1828819–1829145	11122	AGCTGCTAAATCTAAACTCTTTGC	GCTCCCTCAACTAGATCTATCATC
27	960872–961191	11122	AATCGCATACATTCTGCTGTA	GCTTTTCCAAATGAGGATATTA
28	1136267–1136582	11122	AAAAGTAGCTGCAGAAGTATACCC	TTCTCAAAATGTAACATGCTTCT
29	1190422–1190738	11122	CTTGAGCTTACGCCCTTTTAT	ATGTCCGCAATATTGTCCTAAC
30	871284–871614	11112	CTGCATTTTCAACATTACTCAGAT	ATTCATAAAGATCATCCATTCTCTC
31	518787–519092	11112	AGCTGTAGTGATATAAAGAAAAGTTACAT	CTATTTTCGTAGCGAGTAAGAATTT
32	1709427–1709741	11112	TTATGCAAATAACTATCCAAGTGT	TTACCATTAGCTTCAAAGTCTGT
33	511958–512251	11112	TACAAGCGTACCATCTAAGTCA	CATATTGGGATGTCAAGCA
34	99015–99303	11112	TTGATATAACCAACATAAACACTGC	TGAGTATAGAAATACAAAGCTACGC
35	772225–772590	11121	TGTGTAGTAACCCAGGAACCTTTAT	AATTTGATGCCATATGAGAGAAT
36	282847–283070	11121	TTTGGTATGAGTATTCTGGTCCTA	GTATTTTGGTTTAGCTTACGGATT
37	1486225–1486603	11121	AATATTTGCAACCAATGATGATAC	CAGTATCTTTGATGTTAGGGACAA
38	95621–95874	11121	GCTACGACAGGTCTATCTTTCTC	CAACTTATGATTGGTGATGATGT

\*Ftind, *F. tularensis* insertion-deletion marker.†Location of the DNA amplified by PCR in the chromosome of *Francisella tularensis* strain SCHU S4.

‡Sequences given for forward primers represent the target-specific parts of the primers used. For inexpensive fluorescent labeling, each forward primer was synthesized with a 17-bp extension at the 5'-end, corresponding to an M13 sequence (5'-GTAAACGACGGCCAGT-3').

study and 25 MLVA markers throughout the genome of strain SCHU S4 (29) is illustrated in Figure 1.

### PCR Amplicon Separation

PCR reaction mixtures, 2 µL from each, were pooled and diluted 15-fold. One µL of diluted sample was added to 40 µL of sample loading solution, containing DNA Size Standard-600 (Beckman Coulter Inc., Fullerton CA, USA), and sealed with a drop of mineral oil. Finally, PCR amplicons were separated and detected by using a CEQ 8800 Genetic Analysis System (Beckman Coulter Inc.). Binning of indel fragment size-calls was straightforward because of highly precise size determinations (online Appendix

Table 2). Maximum size divergence between size-call and genome sequence data was 3 bp among 38 selected indel markers for strains U112, FSC147, SCHU S4, or LVS.

### Statistical Analysis

Simpson's index of diversity (1 – D) (30) was determined for each investigated marker as a measure of both richness and evenness, calculated as

$$1 - \left[ \frac{1}{N(N-1)} \right] \sum_{j=1}^s n_j(n_j - 1)$$

where  $N$  is the number of strains,  $s$  is the number of recorded states for a marker, and  $n_j$  is the number of strains




Figure 1. Locations of 38 insertion-deletion and 25 multilocus variable-number tandem repeat analysis (MLVA) markers on the physical genome map of *Francisella tularensis* subsp. *tularensis* strain SCHU S4. Positions are given with reference to the predicted origin of replication set at position 0. Indel and MLVA marker locations are depicted by wedges on the outside and inside of the circle, respectively. Two asterisks indicate the duplicate occurrence of the MLVA loci Ft-M14 at 2 different locations because it is part of a large sized genome duplication (1,25).

belonging to the  $j$ th marker state. Both distance-based clustering, by using Hamming distance (31) and the neighbor-joining method, and maximum parsimony (MP) were performed with PAUP\* version 4c10 (32). MP analyses were performed by using 50 replicates without branch swapping and 10,000 bootstrap pseudoreplicates. Nodes supported by <50% bootstrap pseudoreplicates were collapsed in depictions of the obtained consensus topologies. Indel size and distribution of repeat size frequency were analyzed by using the R statistical package (33).

## Results

### Identification and Selection of Indel Loci

In the genomic sequences of each of 5 *F. tularensis* strains (online Appendix Table 1), a total of 280 indel loci were identified, all exhibiting only 2 allelic variants and a size range of 5–200 bp. Small-sized indels predominated; 70% were shorter than 20 bp (Figure 2, panel A). To enable the selection of loci free from such repeat nucleotide sequences, which may have a propensity to initiate deletion or insertion mutations, indels were analyzed with regard to the size of associated repeats. Two repeat size peaks were identified, 1 at  $10 \text{ bp} \pm 1 \text{ bp}$  and another  $\leq 3 \text{ bp}$  (Figure 2, panel B). In 62 loci, no repeats were found. After exclusion




Figure 2. Properties of 280 insertion-deletion (indel) loci identified by analysis of 5 *Francisella tularensis* genome sequences. The diagrams show distributions of indel sizes (A), repeat sizes detected at these loci (B), and 10 allelic diversity patterns (C); the number 1 or 2 represents each of the 2 allelic variants. A string of numbers includes, in order, strain U112 (subsp. *novicida*), FSC147 (subsp. *mediasiatica*), SCHU S4 (subsp. *tularensis*), OSU18 (subsp. *holarctica*), and LVS (subsp. *holarctica*). Empty and filled bars correspond to the presence or absence of repeats >3 bp long, respectively.



of loci associated with repeats >3 bp in length, 158 loci were retained for typing purposes.

To facilitate selection of indel loci represented in various strains, we analyzed the diversity of the 280 allelic variants among the 5 *F. tularensis* genomes included. Among the genomes, only 10 discrete allelic diversity patterns were found, depicted in Figure 2, panel C, as allelic variant 1 or 2 in each of the genomes in order of strains U112, FSC147, SCHU S4, OSU18, and LVS (e.g., 1,2,1,1,1 denotes that a deletion was present in the genome sequence of strain FSC147, but not in any of the others). After loci associated with repeats >3 bp in length were excluded, 7 allelic patterns were retained and used as a basis for selecting indel loci for the assay (Figure 2, panel C).

By these measures, a subset of 38 loci was selected (Table; online Appendix Table 2). These loci showed maximum diversity, represented each allelic pattern among the 5 genomes, and also exhibited a physical separation on the SCHU S4 chromosome (Figure 1).

#### Analysis by the Combined Procedure of 24 Strains of *F. tularensis*

Twenty-four strains, representing all 4 subspecies of *F. tularensis* and clades A.I and A.II of *F. tularensis* subsp. *tularensis*, underwent indel analysis and MLVA (online Appendix Tables 2, 3). Of these, 23 yielded indel PCR amplicons in the range of 145–399 bp, representing an allele of each of 38 loci analyzed. In the remaining strain, isolate FSC454, PCR amplification failed for 7 indel loci tested. FSC454 is an atypical *Francisella* isolate of uncertain taxonomic status recently isolated in Spain (R. Escudero, pers. comm.). FSC454 was excluded from further analyses.

Another atypical strain, ATCC 6223, yielded aberrant amplification results. This strain has lost virulence for mammals, a key characteristic of *F. tularensis*. It exhibits unusual colony morphologic features and a slow growth rate. When subjected to PCR amplification, the genome of strain ATCC 6223 yielded 2 DNA amplicons for an indel locus denoted Ftind-32. Ftind-32 and ATCC 6223 were retained for further analysis, and both alleles were considered.

A graphic representation of the observed amplification patterns at indel and MLVA loci is shown in Figure 3. A difference in mutational stability was apparent between indel and MLVA loci. Indel loci showed a binary pattern that grouped *F. tularensis* in agreement with traditional taxonomy based on phenotype. In accordance with previous genetic typing by MLVA, PFGE, or sequencing of 7 housekeeping genes, the indel analysis distinguished 2 major subpopulations of type A strains (denoted A.I and A.II) and also showed Japan-derived *F. tularensis* strains to be distinct from strains of *F. tularensis* subsp. *holarctica* isolated in other parts of the Northern Hemisphere. Furthermore, indel analysis identified additional subpopulations




Figure 3. Heat map of marker states for 38 insertion-deletion (indel) and 25 multilocus variable-number tandem repeat analysis (MLVA) loci examined. Each *Francisella tularensis* strain is represented by a single row of colored boxes and each DNA loci by a single column. Relative genetic similarity is represented by the similarity of the colors on the gradient scale ranging from blue to yellow. For the binary indel markers, the state of each marker in the genome of strain *F. tularensis* subsp. *novicida* U112 represents the index and is depicted in yellow. Blue indicates the amplification of an allelic variant distinct from that of the index genome. For strain ATCC 6223, both alleles were amplified at loci Ftind-32, and the corresponding box is thus divided into a yellow and a blue part. For MLVA loci, blue represents the largest allele size for each multistate marker; yellow represents the smallest.

among *F. tularensis* subsp. *holarctica* strains. Geographic origins of these subpopulations suggest dispersal over large distances. Two strains from the United States, OSU18 (represented by genome sequence data only) and FSC035, were identical at all indel loci and constitute a distinct genetic entity. Strains FSC012 from the United States and FSC519 from Sweden formed another entity. Finally, 6 strains originating in Sweden or Russia represented a third subpopulation. Compared with indel analysis, MLVA showed much more extensive polymorphisms, which was helpful for characterizing individual strains. Simpson's index of diversity ranged between 0.17 and 0.97 for the MLVA loci and between 0.09 and 0.52 for the indel loci, which reflects the fact that only 2 allele states were present for the indel loci while the MLVA loci were more diverse, with up to 16 alleles (for MLVA marker Ft-M3).

#### Phylogenetic Inferences Based on MLVA and Indel Data

Genetic relationships among *F. tularensis* strains were inferred by MP analysis of the MLVA data, indel data, or both indel and MLVA data (Figure 4). The use of MLVA data alone resulted in weak support for delineation of deeper branching patterns, few nodes having >50% support in bootstrap analysis (Figure 4, panel A). For such purposes, indel data alone were more valuable (Figure 4, panel B). The use of combined indel and MLVA data resulted in well-supported deep nodes and discrimination of the strains included in this study (Figure 4, panel C).




Figure 4. Cladograms depicting relationships among *Francisella tularensis* strains obtained by maximum parsimony and bootstrap analysis that used indel, multilocus variable-number tandem repeat analysis (MLVA), or combined data. Nodes supported by <50% of bootstrap pseudoreplicates were collapsed. A) Cladogram obtained solely from the use of MLVA data. B) Cladogram from the use of indel data. C) Cladogram from the combined use of indel and MLVA data. The dual bootstrap support values presented represent the use of each of 2 alleles, found at locus Ftind-32 of strain ATCC 6223.

In strain ATCC 6223, dual bootstrap support values (Figure 4, panels B, C) represent values obtained by using each of the 2 alleles amplified for locus Ftind-32. The same topology was obtained regardless of which allele was included, and the allele used had minor effect on bootstrap support values. Results were highly similar when using inference by neighbor joining (data not shown).

## Discussion

By combining canonical indels with MLVA, robust subspecies and major clade typing of *F. tularensis* was successfully combined with high-resolution typing among strains. By the use of killed bacterial preparations, the 2 marker sets were rapidly assayed by fragment analysis.

The present canonical indel/MLVA typing concept adapts well to the principles of diagnostic work inherent in public health laboratories. The concept generates portable straight numeric data and, similar to the tests of biochemical reactions, 2 alternative states are determined at multiple indels (Figure 3). The MLVA output consists of multistate discrete numbers and has proven superior to PFGE for reliable resolving of discrete strains of the species (6,7).

Typing of *F. tularensis* provides useful public health information. This is especially relevant to North America, where subpopulations varying in virulence occur naturally in the same geographic region. According to a recent report, major genetic subpopulations within the type A tularemia population (A.I and A.II) seem connected with different mortality rates in humans (7). Potential clinical correlates to type B subpopulations remain to be studied. Ongoing work shows that >90 European isolates all fall within the subpopulations described here (unpub. data).

A most conspicuous need for rapid and reliable characterization of isolates of *F. tularensis* relates to bioterrorism. Whenever tularemia appears in an area believed to be free from the agent, characterization of isolates will become urgent. Such characterization abilities may also prove useful in understanding how *F. tularensis* may spread under peaceful circumstances. Reminders of the agent's potential for infection include the unexplained introduction of the disease on Martha's Vineyard in 1937 and more recently in northern Spain in 1997–1998, along with the highly publicized 2004 laboratory infections with respiratory type A tularemia at Boston University (5,17,34).

In public health laboratories, indel/MLVA typing may replace more risky and time-consuming biochemical characterization, which is based on growth of *F. tularensis*. After initial culture of the agent, noninfectious DNA is rapidly analyzed by PCR and fragment analysis for determination of indel and MLVA data.

A major achievement of the present study was the identification of canonical indels for combined use with MLVA. From studies of *Bacillus* spp., only SNPs have

been predicted to exhibit mutation rates sufficiently slow to be useful for unambiguous assignment of bacteria at deeper taxonomic levels (24). SNPs with canonical properties are not yet recognized in *F. tularensis*, and their combined use with MLVA has thus not yet been evaluated. An SNP-based approach does conform with well-developed evolutionary models to support data analysis (35), models that do not exist for indel mutations. A drawback is, however, that the involvement of 2 different analytic methods in a combined MLVA/SNP-based analysis makes it more complicated. By use of fragment analysis for both steps, the indel/MLVA approach is more effective. This study indicates that canonical indels can be integrated into evolutionary analyses for measuring large genetic distances while MLVA provides a detailed examination at short distances.

When selecting indels for the presented typing procedure, we took precautions to avoid DNA-marker discovery bias and homoplastic markers, problems that had been carefully addressed in work on other bacterial pathogens (36,37). To minimize discovery bias, we used *F. tularensis* genomes classified as being distantly related by independent methods. Genomes selected represented all 4 subspecies of *F. tularensis* that also form major genetic clades, according to MLVA, PFGE, microarray, and various arbitrarily primed-PCR analyses (1). To avoid homoplasy, including gene conversion, we excluded indels associated with repeat sequences. Our genome sequence data and the overall tree structure obtained from analysis of indel data lent support to a paucity of homoplasy effects. Except for locus Ftind-32 in the type strain ATCC 6223, which exhibited 2 PCR amplicons, only 4 of 280 identified loci showed incongruent evolutionary allele patterns. These 4 loci were all found among those repeat-containing loci that were excluded according to our selection criteria.

The reason behind a deviant result of strain ATCC 6223 at 1 locus is unknown but may be related to laboratory-induced mutations. ATCC 6223 was originally isolated in 1920 from a human lymph node in Utah, became avirulent by laboratory passage in the early years, but still retained properties that made it useful for antigen production. Recent microarray studies showed that it lacks portions of the genetic repertoire shared by all other *F. tularensis* strains (10).

MLVA discriminates among individual isolates within subspecies but may cause false estimates of relationships at deeper phylogenetic levels. Although in a previous study that used the present 25-marker MLVA scheme, discrimination of *F. tularensis* subspecies and major genetic clades was achieved, bootstrap support at these deeper levels was weak (6). Also in the present study, deep structural relationships among strains inferred by MP analysis of MLVA data were found to be weakly resolved. Conversely, strong support was shown for deep-level nodes obtained by us-

ing indel data. A combined analysis with both MLVA and indel data retained the deep-level support and yielded the most resolved topology. Furthermore, despite the inability of the indel or MLVA data to provide support for a separate clade of Japanese strains, such separation was supported by the combined analysis. This demonstrates that topologic constraints imposed by canonical indel data reduced the number of alternative positions of a combined tree and consequently increased the support for a clade.

When the present approach is used for routine purposes, the number of DNA markers might well be reduced yet retain a high level of discrimination and robustness. However, such a reduction needs to be evaluated to ensure proper marker selection. The inclusion by international collaboration of large numbers of geographically distributed strains will be facilitated by the unambiguous nature of data collected and the use of low quantities of killed bacteria. For ordinary clinical purposes, only a few indel markers may be required to rapidly receive relevant information, i.e., whether an isolate belongs to a subspecies or major genetic clade. A reference laboratory may wish to add more markers for tracing outbreaks and for forensic applications. Tailored combinations of these markers can be easily integrated into multiplex assays with 4–8 markers per PCR amplification and subsequent multicolor fragment analysis to decrease analytical time and cost.

In essence, we used 5 genome sequences representative of the species *F. tularensis* to identify 158 canonical indel DNA-markers, of which 38 were selected to provide robust information specific to each major genetic clade. By combining analysis of these indel markers with MLVA, discrimination of individual strains was achieved. The usefulness of indels with canonical properties may not be restricted to *F. tularensis*. The current availability of multiple genome sequences should allow testing this typing strategy for other clinically relevant pathogens.

## Acknowledgments

We thank Thomas Bretin, Christine Munk, and Paul Keim for timely access to the preliminary genome sequence of strain FSC147. We thank Arne Tärnvik for helpful comments on the manuscript. We are indebted also to numerous colleagues for kindly providing strains to the FSC in Umeå, Sweden.

This work was supported by funding from the Swedish Ministry of Defence, project no. A4854, the Swedish Society for Medical Research, and the County Council of Västerbotten.

Dr Larsson is a doctoral research fellow at the Department of Clinical Microbiology, Umeå University, and at the Swedish Defence Research Agency, Umeå, Sweden. His main research interests are comparative genome analyses of pathogenic bacteria to extract information useful for developing diagnostic assays and identification of virulence factors.



## References

- Johansson A, Forsman M, Sjöstedt A. The development of tools for diagnosis of tularemia and typing of *Francisella tularensis*. APMIS. 2004;112:898–907.
- Sjöstedt A. Genus I. *Francisella* Dorofe'ev 1947, 176<sup>AL</sup>. In: Brenner DJ, Krieg NR, Staley JT, Garrity GM, editors. *Bergey's manual of systematic bacteriology*, 2nd ed. New York: Springer; 2005. p. 200–10.
- Olsufjev NG, Meshcheryakova IS. Subspecific taxonomy of *Francisella-tularensis*. Int J Syst Bacteriol. 1983;33:872–4.
- Dennis DT, Inglesby TV, Henderson DA, Bartlett JG, Ascher MS, Eitzen E, et al. Tularemia as a biological weapon: medical and public health management. JAMA. 2001;285:2763–73.
- Farlow J, Wagner DM, Dukerich M, Stanley M, Chu M, Kubota K, et al. *Francisella tularensis* in the United States. Emerg Infect Dis. 2005;11:1835–41.
- Johansson A, Farlow J, Larsson P, Dukerich M, Chambers E, Bysström M, et al. Worldwide genetic relationships among *Francisella tularensis* isolates determined by multiple-locus variable-number tandem repeat analysis. J Bacteriol. 2004;186:5808–18.
- Staples JE, Kubota KA, Chalcraft LG, Mead PS, Petersen JM. Epidemiologic and molecular analysis of human tularemia, United States, 1964–2004. Emerg Infect Dis. 2006;12:1113–8.
- Rotz LD, Khan AS, Lillibridge SR, Ostroff SM, Hughes JM. Public health assessment of potential biological terrorism agents. Emerg Infect Dis. 2002;8:225–30.
- Burke DS. Immunization against tularemia: analysis of the effectiveness of live *Francisella tularensis* vaccine in prevention of laboratory-acquired tularemia. J Infect Dis. 1977;135:55–60.
- Broekhuijsen M, Larsson P, Johansson A, Byström M, Eriksson U, Larsson E, et al. Genome-wide DNA microarray analysis of *Francisella tularensis* strains demonstrates extensive genetic conservation within the species but identifies regions that are unique to the highly virulent *F. tularensis* subsp. *tularensis*. J Clin Microbiol. 2003;41:2924–31.
- Kugeler KJ, Pappert R, Zhou Y, Petersen JM. Real-time PCR for *Francisella tularensis* types A and B. Emerg Infect Dis. 2006;12:1799–801.
- Samrakandi MM, Zhang C, Zhang M, Nietfeldt J, Kim J, Iwen PC, et al. Genome diversity among regional populations of *Francisella tularensis* subspecies *tularensis* and *Francisella tularensis* subspecies *holarctica* isolated from the US. FEMS Microbiol Lett. 2004;237:9–17.
- Tomaso H, Scholz HC, Neubauer H, Al Dahouk S, Seibold E, Landt O, et al. Real-time PCR using hybridization probes for the rapid and specific identification of *Francisella tularensis* subspecies *tularensis*. Mol Cell Probes. 2007;21:12–6.
- Gerner-Smidt P, Hise K, Kincaid J, Hunter S, Rolando S, Hyytiä-Trees E, et al. PulseNet USA: a five-year update. Foodborne Pathog Dis. 2006;3:9–19.
- van Belkum A, van Leeuwen W, Kaufmann ME, Cookson B, Forey F, Etienne J, et al. Assessment of resolution and intercenter reproducibility of results of genotyping *Staphylococcus aureus* by pulsed-field gel electrophoresis of *Sma*I macrorestriction fragments: a multicenter study. J Clin Microbiol. 1998;36:1653–9.
- Garcia Del Blanco N, Dobson ME, Vela AI, De La Puente VA, Gutierrez CB, Hadfield TL, et al. Genotyping of *Francisella tularensis* strains by pulsed-field gel electrophoresis, amplified fragment length polymorphism fingerprinting, and 16S rRNA gene sequencing. J Clin Microbiol. 2002;40:2964–72.
- Barry MA. Report of pneumonic tularemia in three Boston University researchers, November 2004–March 2005. Boston: Communicable Disease Control, Boston Public Health Commission; 2005 [cited 10 Sep 2007]. Available from [http://www.bphc.org/reports/pdfs/report\\_202](http://www.bphc.org/reports/pdfs/report_202)
- Farlow J, Smith KL, Wong J, Abrams M, Lytle M, Keim P. *Francisella tularensis* strain typing using multiple-locus, variable-number tandem repeat analysis. J Clin Microbiol. 2001;39:3186–92.
- Johansson A, Göransson I, Larsson P, Sjöstedt A. Extensive allelic variation among *Francisella tularensis* strains in a short-sequence tandem repeat region. J Clin Microbiol. 2001;39:3140–6.
- Vogler AJ, Keys C, Nemoto Y, Colman RE, Jay Z, Keim P. Effect of repeat copy number on variable-number tandem repeat mutations in *Escherichia coli* O157:H7. J Bacteriol. 2006;188:4253–63.
- Vogler AJ, Keys CE, Allender C, Bailey I, Girard J, Pearson T, et al. Mutations, mutation rates, and evolution at the hypervariable VNTR loci of *Yersinia pestis*. Mutat Res. 2007;616:145–58.
- Bayliss CD, Field D, Moxon ER. The simple sequence contingency loci of *Haemophilus influenzae* and *Neisseria meningitidis*. J Clin Invest. 2001;107:657–62.
- Field D, Magnasco MO, Moxon ER, Metzgar D, Tanaka MM, Wills C, et al. Contingency loci, mutator alleles, and their interactions. Synergistic strategies for microbial evolution and adaptation in pathogenesis. Ann N Y Acad Sci. 1999;870:378–82.
- Keim P, Van Ert MN, Pearson T, Vogler AJ, Huynh LY, Wagner DM. Anthrax molecular epidemiology and forensics: using the appropriate marker for different evolutionary scales. Infect Genet Evol. 2004;4:205–13.
- Sandström G, Tärnvik A, Wolf-Watz H, Löfgren S. Antigen from *Francisella tularensis*: nonidentity between determinants participating in cell-mediated and humoral reactions. Infect Immun. 1984;45:101–6.
- Sjöstedt A, Eriksson U, Berglund L, Tärnvik A. Detection of *Francisella tularensis* in ulcers of patients with tularemia by PCR. J Clin Microbiol. 1997;35:1045–8.
- Darling AC, Mau B, Blattner FR, Perna NT. Mauve: multiple alignment of conserved genomic sequence with rearrangements. Genome Res. 2004;14:1394–403.
- Rozen S, Skaletsky H. Primer3 on the WWW for general users and for biologist programmers. Methods Mol Biol. 2000;132:365–86.
- Larsson P, Oyston PC, Chain P, Chu MC, Duffield M, Fuxelius HH, et al. The complete genome sequence of *Francisella tularensis*, the causative agent of tularemia. Nat Genet. 2005;37:153–9.
- Hunter PR, Gaston MA. Numerical index of the discriminatory ability of typing systems: an application of Simpson's index of diversity. J Clin Microbiol. 1988;26:2465–6.
- Hamming R. Error-detecting and error-correcting codes. Bell System Technical Journal. 1950;29:147–60.
- Swofford D. PAUP\*: phylogenetic analysis using parsimony (\*and other methods), version 4. Sunderland (MA): Sinauer Associates; 2003.
- R Development Core Team. R: A language and environment for statistical computing, Vienna, Austria, 2007 [cited 2007 Sep 10]. Available from <http://www.r-project.org>
- Petersen JM, Schrieffer ME. Tularemia: emergence/re-emergence. Vet Res. 2005;36:455–67.
- Graur D, Li W-H. Fundamentals of molecular evolution. 2nd ed. Sunderland (MA): Sinauer Associates; 2000.
- Alland D, Whittam TS, Murray MB, Cave MD, Hazbon MH, Dix K, et al. Modeling bacterial evolution with comparative-genome-based marker systems: application to *Mycobacterium tuberculosis* evolution and pathogenesis. J Bacteriol. 2003;185:3392–9.
- Pearson T, Busch JD, Ravel J, Read TD, Rhoton SD, U'Ren JM, et al. Phylogenetic discovery bias in *Bacillus anthracis* using single-nucleotide polymorphisms from whole-genome sequencing. Proc Natl Acad Sci U S A. 2004;101:13536–41.

Address for correspondence: Anders Johansson, Department of Infectious Diseases, Umeå University Hospital, SE-901 85 Umeå, Sweden; email: [anders.johansson@infdis.umu.se](mailto:anders.johansson@infdis.umu.se)



Appendix Table 1. *Francisella tularensis* genomic sequences used for indel identification

<i>F. tularensis</i> subspecies	Strain	FSC* no.	ATCC† no.	Source of genomic sequence	Status	GenBankaccession no.	Reference
<i>novicida</i>	U112	040‡	15482	University of Washington, Seattle, WA, USA	Finished	CP000439	Unpub.
<i>tularensis</i>	SCHU S4	237	–	Swedish/British/US consortium§	Finished	AJ749949	(1)
<i>mediasiatica</i>	GIEM 543	147	–	Los Alamos National Laboratory, Los Alamos, NM, USA	Unfinished	Pending	Unpub.
<i>holarctica</i>	OSU18	–	–	Baylor College of Medicine, Waco, TX, USA	Finished	CP000437	(2)
<i>holarctica</i>	LVS	458‡	29684	Lawrence Livermore National Laboratory, CA, USA	Finished	AM233362	Unpub.

\**Francisella* Strain Collection.

†American Type Culture Collection.

‡Strains sequenced were not obtained from the *Francisella* Strain Collection.

§Member institutes of the sequencing consortium were: Swedish Defence Research Agency, Sweden; Uppsala and Umeå Universities, Sweden, Defence Science and Technology Laboratories, Salisbury, UK; Walter Reed Army Institute of Research, Bethesda, Maryland, USA.

References

1. Sandström G, Tärnvik A, Wolf-Watz H, Löfgren S. [Antigen from \*Francisella tularensis\*: nonidentity between determinants participating in cell-mediated and humoral reactions.](#) Infect Immun. 1984;45:101–6.Medline

2. Petrosino JF, Xiang Q, Karpathy SE, Jiang H, Yerrapragada S, Liu Y, et al. [Chromosome rearrangement and diversification of \*Francisella tularensis\* revealed by the type B \(OSU18\) genome sequence.](#) J Bacteriol. 2006;188:6977–85.

Appendix Table 2. *Francisella tularensis* strains and PCR-amplicon sizes at 38 indel loci\*

Species (genetic clade), source	FSC no.	Other designation	Amplified DNA size (bp) at indel locus*									
			Ftind1	Ftind2	Ftind3	Ftind4	Ftind5	Ftind6	Ftind7	Ftind8	Ftind9	Ftind10
<i>F. tularensis</i> subsp. <i>novicida</i>												
Water, 1950, Utah	040	U112, ATCC 15482	316	288	387	228	202	394	382	380	388	286
Human, 2004, UK-Germany-Brazil	595	F58	292	355	388	230	205	394	382	380	389	285
<i>F. tularensis</i> subsp. <i>tularensis</i> (A.II)												
Rabbit, 1953, Nevada	054	Nevada 14	293	354	376	236	216	350	372	375	389	285
Human lymph node, 1920, Utah	230	ATCC6223	292	354	376	237	216	351	372	375	388	285
Foal, 1959, Montana	604	8859	293	355	377	237	216	350	372	375	388	284
<i>F. tularensis</i> subsp. <i>tularensis</i> (A.I)												
Human ulcer, 1941, Ohio	237	SCHU S4	292	354	376	236	215	351	373	374	270	281
Tick, 1935, British Columbia, Canada	041	Vavenby	293	354	377	236	216	350	372	374	269	281
Human pleural fluid, 1940, Ohio	046	Fox Downs	292	354	376	237	217	350	372	375	270	280
<i>F. tularensis</i> subsp. <i>mediasiatica</i>												
Miday gerbil, 1965, Kazakhstan	147	GIEM 543	292	354	376	237	216	346	372	374	388	285
Tick, 1982, Central Asia	148	240	292	354	376	236	216	346	372	374	388	286
Hare, 1965, Central Asia	149	120	291	354	376	237	216	346	372	374	389	285
<i>F. tularensis</i> subsp. <i>holarctica</i> (Japanese)												
Human lymph node, 1926, Japan	017	S-2	292	354	374	237	216	350	372	374	388	285
Human, 1958, Japan	021	Tsuchiya	293	354	374	236	216	350	372	374	388	286
Human, 1950, Japan	022	Ebina	292	354	374	236	216	351	372	374	388	285
<i>F. tularensis</i> subsp. <i>holarctica</i> (non-Japanese)												
Beaver, 1976, Hamilton, Montana	035	B423A	293	354	377	236	215	351	373	374	388	286
Tick, 1941, Montana	012	425F4G	293	354	375	237	216	351	372	374	388	286
Human, 2004, Örebro, Sweden	519		293	355	377	236	216	350	372	373	388	285
Live vaccine strain, Russia	458	LVS, ATCC 29684	292	354	376	236	215	351	373	374	388	285

Human, 1995, Ljusdal, Sweden	171		292	354	376	237	216	350	372	374	388	285
Human, 2003, Örebro, Sweden	398		292	354	376	237	216	350	372	373	388	285
Human, 2003, Örebro, Sweden	412		292	354	376	236	216	351	372	373	388	285
Human, 2003, Örebro, Sweden	429		293	355	377	236	216	350	372	373	388	285
Tick, 1949, Moscow area, Russia	257	GIEM 503/840	293	355	377	237	216	350	372	373	388	285

Amplified DNA size (bp) at indel locus\*

Species (genetic clade), source	FSC no.	Other designation	Ftind11	Ftind12	Ftind13	Ftind14	Ftind15	Ftind16	Ftind17	Ftind18	Ftind19	Ftind20
<i>F. tularensis</i> subsp. <i>novicida</i>												
Water, 1950, Utah	040	U112, ATCC 15482	317	288	274	268	249	268	212	225	171	337
Human, 2004, UK-Germany-Brazil	595	F58	317	287	279	268	249	268	212	225	171	336
<i>Francisella tularensis</i> subsp. <i>tularensis</i> (A.II)												
Rabbit, 1953, Nevada	054	Nevada 14	317	273	264	256	249	268	212	225	171	336
Human lymph node, 1920, Utah	230	ATCC6223	317	273	263	256	249	268	212	225	171	336
Foal, 1959, Montana	604	8859	317	272	264	256	249	268	212	225	171	337
<i>F. tularensis</i> subsp. <i>tularensis</i> (A.I)												
Human ulcer, 1941, Ohio	237	SCHU S4	279	272	262	256	249	268	212	225	171	337
Tick, 1935, British Columbia, Canada	041	Vavenby	279	272	263	256	249	268	212	225	171	337
Human pleural fluid, 1940, Ohio	046	Fox Downs	279	272	262	256	249	268	212	225	171	337
<i>F. tularensis</i> subsp. <i>mediasiatica</i>												
Miday gerbil, 1965, Kazakhstan	147	GIEM 543	317	289	262	256	234	243	192	203	145	337
Tick, 1982, Central Asia	148	240	317	268	263	256	234	243	192	203	145	336
Hare, 1965, Central Asia	149	120	317	289	262	256	234	243	192	203	145	336
<i>F. tularensis</i> subsp. <i>holarctica</i> (Japanese)												
Human lymphnode, 1926, Japan	017	S-2	317	288	274	267	250	268	212	225	171	323
Human, 1958, Japan	021	Tsuchiya	318	288	275	267	250	268	212	225	171	323
Human, 1950, Japan	022	Ebina	317	288	274	267	250	268	212	225	171	323



<i>F. tularensis</i> subsp. <i>holarctica</i> (non-Japanese)												
Beaver, 1976, Hamilton, Montana	035	B423A	317	288	274	267	249	268	212	225	171	324
Tick, 1941, Montana	012	425F4G	318	288	274	267	249	268	212	225	171	323
Human, 2004, Örebro, Sweden	519		317	288	273	267	249	268	212	225	171	323
Live vaccine strain, Russia	458	LVS, ATCC 29684	317	289	273	268	249	268	212	225	171	324
Human, 1995, Ljusdal, Sweden	171		317	288	273	267	249	268	212	225	171	324
Human, 2003, Örebro, Sweden	398		317	287	273	267	249	268	212	225	171	324
Human, 2003, Örebro, Sweden	412		317	288	274	267	249	268	212	225	171	323
Human, 2003, Örebro, Sweden	429		317	288	273	267	249	268	212	225	171	323
Tick, 1949, Moscow area, Russia	257	GIEM 503/840	317	288	273	267	249	268	212	225	171	324
Amplified DNA size (bp) at indel locus*												
Species (genetic clade), source	FSC no.	Other designation	Ftind21	Ftind22	Ftind23	Ftind24	Ftind25	Ftind26	Ftind27	Ftind28	Ftind29	Ftind30
<i>F. tularensis</i> subsp. <i>novicida</i>												
Water, 1950, Utah	040	U112, ATCC 15482	352	345	241	341	348	347	339	337	338	352
Human, 2004, UK-Germany-Brazil	595	F58	352	347	241	341	349	346	339	337	337	351
<i>F. tularensis</i> subsp. <i>tularensis</i> , (A.II)												
Rabbit, 1953, Nevada	054	Nevada 14	353	345	241	341	349	347	341	337	337	351
Human lymph node, 1920, Utah	230	ATCC6223	353	345	242	341	349	347	341	337	338	351
Foal, 1959, Montana	604	8859	353	346	241	341	349	346	341	337	338	351
<i>F. tularensis</i> subsp. <i>tularensis</i> , (A.I)												
Human ulcer, 1941, Ohio	237	SCHU S4	352	346	242	341	348	347	340	337	338	352
Tick, 1935, British Columbia, Canada	041	Vavenby	352	345	241	341	348	347	340	336	338	351
Human pleural fluid, 1940, Ohio	046	Fox Downs	353	345	242	341	349	348	341	337	338	352
<i>F. tularensis</i> subsp. <i>mediasiatica</i>												
Miday gerbil, 1965, Kazakhstan	147	GIEM 543	353	346	242	341	349	347	340	337	337	351
Tick, 1982, Central Asia	148	240	352	346	241	340	349	347	340	337	337	351

Hare, 1965, Central Asia	149	120	352	346	241	340	349	347	340	337	337	351
<i>F. tularensis</i> subsp. <i>holarctica</i> (Japanese)												
Human lymph node, 1926, Japan	017	S-2	342	312	228	336	349	347	340	337	338	351
Human, 1958, Japan	021	Tsuchiya	343	312	228	336	349	347	340	337	339	351
Human, 1950, Japan	022	Ebina	343	312	228	336	349	347	340	337	338	351
<i>F. tularensis</i> subsp. <i>holarctica</i> (non-Japanese)												
Beaver, 1976, Hamilton, Montana	035	B423A	342	312	230	336	345	284	315	310	331	352
Tick, 1941, Montana	012	425F4G	343	312	228	336	344	284	315	311	330	351
Human, 2004, Örebro, Sweden	519		343	312	228	336	343	284	315	310	330	351
Live vaccine strain, Russia	458	LVS, ATCC 29684	343	312	229	336	344	284	315	311	331	346
Human, 1995, Ljusdal, Sweden	171		343	313	228	336	344	284	315	310	331	346
Human, 2003, Örebro, Sweden	398		342	313	228	336	343	284	315	311	330	346
Human, 2003, Örebro, Sweden	412		343	312	228	336	344	284	315	310	330	346
Human, 2003, Örebro, Sweden	429		342	312	228	336	344	284	315	310	330	346
Tick, 1949, Moscow area, Russia	257	GIEM 503/840	342	312	228	336	343	284	315	311	330	346

Species (genetic clade), source	FSC no.	Other designation	Amplified DNA size (bp) at indel locus*						
			Ftind31	Ftind32	Ftind33	Ftind34	Ftind35	Ftind36	Ftind37
<i>F. tularensis</i> subsp. <i>novicida</i>									
Water, 1950, Utah	040	U112, ATCC 15482	328	334	315	308	388	244	398
Human, 2004, UK-Germany-Brazil	595	F58	327	335	315	308	386	244	397
<i>F. tularensis</i> subsp. <i>tularensis</i> (A.II)									
Rabbit, 1953, Nevada	054	Nevada 14	325	334	315	308	387	244	399
Human lymph node, 1920, Utah	230	ATCC6223	325	324 334	315	308	388	244	398
Foal, 1959, Montana	604	8859	325	335	314	308	387	244	398
<i>F. tularensis</i> subsp. <i>tularensis</i> (A.I)									
Human ulcer, 1941, Ohio	237	SCHU S4	325	335	314	305	388	244	399

Tick, 1935, British Columbia, Canada	041	Vavenby	325	334	315	308	387	244	397
Human pleural fluid, 1940, Ohio	046	Fox Downs	325	335	315	308	387	244	399
<i>F. tularensis</i> subsp. <i>mediasiatica</i>									
Miday gerbil, 1965, Kazakhstan	147	GIEM 543	324	334	315	308	388	243	398
Tick, 1982, Central Asia	148	240	325	334	315	308	387	244	399
Hare, 1965, Central Asia	149	120	325	334	315	308	387	244	398
<i>F. tularensis</i> subsp. <i>holarctica</i> (Japanese)									
Human lymph node, 1926, Japan	017	S-2	325	334	315	308	387	244	398
Human, 1958, Japan	021	Tsuchiya	325	334	315	308	388	245	399
Human, 1950, Japan	022	Ebina	325	334	315	308	387	244	398
<i>F. tularensis</i> subsp. <i>holarctica</i> (non-Japanese)									
Beaver, 1976, Hamilton, Montana	035	B423A	326	334	315	308	371	236	391
Tick, 1941, Montana	012	425F4G	325	334	315	308	388	244	398
Human, 2004, Örebro, Sweden	519		325	334	315	308	387	244	397
Live vaccine strain, Russia	458	LVS, ATCC 29684	321	324	296	268	388	244	398
Human, 1995, Ljusdal, Sweden	171		320	324	295	268	387	244	398
Human, 2003, Örebro, Sweden	398		319	325	295	268	387	244	399
Human, 2003, Örebro, Sweden	412		320	324	295	268	387	244	399
Human, 2003, Örebro, Sweden	429		320	324	294	266	387	244	397
Tick, 1949, Moscow area, Russia	257	GIEM 503/840	319	324	295	266	387	243	399

\*Size determinations include a 19-bp M-13 primer tail sequence used for fluorescent labeling. Ftindel, *F. tularensis* insertion-deletion loci; FSC, *Francisella* Strain Collection; ATCC, American Type Culture Collection.



Appendix Table 3. *Francisella tularensis* strains and repeat copy number at 25 MLVA loci\*

Species (genetic clade), source	FSC no.	Other designation	Repeat copy no. at MLVA locus																								
			Ft- M1	Ft- M2	Ft- M3	Ft- M4	Ft- M5	Ft- M6	Ft- M7	Ft- M8	Ft- M9	Ft- M10	Ft- M11	Ft- M12	Ft- M13	Ft- M14	Ft- M15	Ft- M16	Ft- M17	Ft- M18	Ft- M19	Ft- M20	Ft- M21	Ft- M22	Ft- M23	Ft- M24	Ft- M25
<i>F. tularensis</i> subsp. <i>novicida</i>																											
Water, 1950, Utah	040	U112, ATCC 15482	3	2	32	5	2	4	2	2	4	2	4	2	2	3	3	1	2	2	1	31	2	4	1	1	5
Human, 2004, UK- Germany-Brazil	595	F58	3	2	20	5	2	4	2	1	2	2	4	2	2	4	3	1	2	2	1	31	2	4	1	1	5
<i>F. tularensis</i> subsp. <i>tularensis</i> (A.II)																											
Rabbit, 1953, Nevada	054	Nevada 14	3	21	16	3	2	4	2	2	2	1	5	2	1	3	3	1	2	2	1	22	2	3	1	1	5
Human lymph node, 1920, Utah	230	ATCC6223	3	25	8	3	2	4	2	4	2	1	5	2	1	3	3	1	2	2	1	25	2	3	1	1	5
Foal, 1959, Montana	604	8859	3	7	11	4	2	4	2	2	2	1	5	2	1	4	3	1	2	2	1	21	3	3	1	1	5
<i>F. tularensis</i> subsp. <i>tularensis</i> , (A.I)																											
Human ulcer, 1941, Ohio	237	SCHU S4	3	4	25	3	3	4	4	4	4	20	5	2	2	3	2	2	3	4	1	3	3	2	2	1	5
Tick, 1935, British Columbia, Canada	041	Vavenby	3	24	19	3	2	4	3	2	5	10	5	2	2	3	2	2	3	4	1	3	2	2	1	1	5
Human pleural fluid, 1940, Ohio	046	Fox Downs	3	4	27	3	5	5	3	3	5	9	5	2	2	3	2	2	3	4	1	3	3	2	2	1	5
<i>F. tularensis</i> subsp. <i>mediasiatica</i>																											
Miday gerbil, 1965, Kazakhstan	147	GIEM 543	4	2	28	3	6	2	2	1	3	2	3	1	1	2	3	1	2	2	1	7	2	7	1	1	5
Tick, 1982, Central Asia	148	240	4	2	25	3	10	3	2	1	3	2	3	1	1	2	3	1	2	2	1	6	2	14	1	1	5
Hare, 1965, Central Asia	149	120	4	2	27	3	6	2	2	1	3	2	3	1	1	2	3	1	2	2	1	7	2	7	1	1	5
<i>F. tularensis</i> subsp. <i>holarctica</i> (Japanese)																											
Human lymph node, 1926, Japan	017	S-2	3	24	3	5	2	4	2	3	2	4	5	2	1	3	2	1	2	2	0	18	3	3	1	1	5
Human, 1958, Japan	021	Tsuchiya	3	8	3	5	2	5	2	1	2	8	5	2	1	3	3	1	2	2	0	11	2	3	1	1	5

Human, 1950, Japan	022	Ebina	3	12	3	5	2	4	1	4	2	4	5	2	1	3	3	1	2	2	0	11	2	3	1	1	5
<i>F. tularensis</i> subsp. <i>holarctica</i> (non-Japanese)																											
Beaver, 1976, Hamilton, Montana	035	B423A	3	2	19	6	2	5	2	2	2	2	5	2	1	3	3	1	2	2	0	3	2	3	1	2	4
Tick, 1941, Montana	012	425F4G	3	2	18	5	2	7	2	2	2	2	5	2	1	3	3	1	2	2	0	3	2	3	1	1	4
Human, 2004, Örebro, Sweden	519		3	2	12	5	4	4	2	2	2	2	5	2	1	4	3	1	2	2	0	3	2	3	1	1	4
Live vaccine strain, Russia	458	LVS, ATCC 29684	3	2	16	5	2	4	2	2	2	2	5	2	1	3	3	1	2	2	0	4	2	4	1	2	4
Human, 1995, Ljusdal, Sweden	171		3	2	10	5	2	5	2	2	2	2	5	2	1	4	3	1	2	2	0	3	2	4	1	2	4
Human, 2003, Örebro, Sweden	398		3	2	21	5	2	5	2	2	2	2	5	2	1	4	3	1	2	2	0	3	2	4	1	2	4
Human, 2003, Örebro, Sweden	412		3	2	21	5	2	5	2	2	2	2	5	2	1	4	3	1	2	2	0	3	2	4	1	2	4
Human, 2003, Örebro, Sweden	429		3	2	22	5	2	5	2	2	2	2	5	2	1	4	3	1	2	2	0	3	2	4	1	2	4
Tick, 1949, Moscow area, Russia	257	GIEM 503/840	3	2	17	5	2	4	2	2	2	2	5	2	1	3	3	1	2	2	0	3	4	4	1	2	4

\*MLVA, multilocus variable-number tandem repeat analysis; FSC, *Francisella* Strain Collection; ATCC, American Type Culture Collection.