

4. Vázquez A, Ruiz S, Herrero L, Moreno J, Molero F, Magallanes A, et al. West Nile and Usutu viruses in mosquitoes in Spain, 2008–2009. *Am J Trop Med Hyg.* 2011;85:178–81. <http://dx.doi.org/10.4269/ajtmh.2011.11-0042>
5. Sotelo E, Llorente F, Rebollo B, Camunas A, Venteo A, Gallardo C, et al. Development and evaluation of a new epitope-blocking ELISA for universal detection of antibodies to West Nile virus. *J Virol Methods.* 2011;174:35–41. <http://dx.doi.org/10.1016/j.jviromet.2011.03.015>
6. Figuerola J, Jimenez-Clavero MA, Rojo G, Gomez-Tejedor C, Soriguer R. Prevalence of West Nile virus neutralizing antibodies in colonial aquatic birds in southern Spain. *Avian Pathol.* 2007;36:209–12. <http://dx.doi.org/10.1080/03079450701332329>
7. Scaramozzino N, Crance JM, Jouan A, DeBriel DA, Stoll F, Garin D. Comparison of flavivirus universal primer pairs and development of a rapid, highly sensitive heminested reverse transcription-PCR assay for detection of flaviviruses targeted to a conserved region of the NS5 gene sequences. *J Clin Microbiol.* 2001;39:1922–7. <http://dx.doi.org/10.1128/JCM.39.5.1922-1927.2001>
8. Dupuis AP II, Marra PP, Kramer LD. Serologic evidence of West Nile virus transmission, Jamaica, West Indies. *Emerg Infect Dis.* 2003;9:860–3. <http://dx.doi.org/10.3201/eid0907.030249>
9. Monaco F, Savini G, Calistri P, Polci A, Pinoni C, Bruno R, et al. West Nile disease epidemic in Italy: first evidence of overwintering in Western Europe? *Res Vet Sci.* 2011;91:321–6.
10. Sotelo E, Fernandez-Pinero J, Llorente F, Vazquez A, Moreno A, Agüero M, et al. Phylogenetic relationships of western Mediterranean West Nile virus strains (1996–2010) using full-length genome sequences: single or multiple introductions? *J Gen Virol.* 2011;92:2512–22. <http://dx.doi.org/10.1099/vir.0.033829-0>

Address for correspondence: Miguel Ángel Jiménez-Clavero, Centro de Investigación en Sanidad Animal CISA (INIA), Ctra Algete—El Casar s/n, 28130, Valdeolmos, Spain; email: majimenez@inia.es

Books, Other Media

Reviews (250-500) words of new books or other media on emerging disease issues are welcome. Name, publisher, number of pages, and other pertinent details should be included.

Absence of Rift Valley Fever Virus in Wild Small Mammals, Madagascar

To the Editor: Rift Valley fever virus (RVFV) is a mosquito-borne zoonotic virus in the family *Bunyaviridae*, genus *Phlebovirus*, which affects mainly domestic ruminants and humans on continental Africa, Madagascar, and the Arabian Peninsula (1). RVFV is transmitted between ruminants mainly by bites of mosquitoes of several genera (1). Infection can lead to mild symptoms or can cause abortion in pregnant animals and high mortality rates among newborns. Humans are mostly infected by aerosol transmission when handling infected tissues (aborted fetuses or meat), which results in dengue-like illness. Some cases in humans can be in a severe form (hemorrhagic fever and meningoencephalitis), which can be fatal. Outbreaks in southern and eastern Africa are associated with periods of heavy rainfall (1). In eastern Africa, RVFV is believed to be maintained during interepizootic periods through vertical transmission in *Aedes* spp. mosquitoes (1). It has been suspected that wild mammals, especially rodents, play a role in the maintenance of RVFV during interepizootic periods (2). However, evidence of a wild mammal reservoir in the epidemiologic cycle of RVFV has yet to be demonstrated (2).

In Madagascar, the first RVFV isolate was obtained from mosquitoes captured in the Périnet Forest (Andasibe, Moramanga District) in 1979, outside an epizootic period (3). Two epizootic episodes occurred, during 1990–91 and 2008–09 (4). After the most recent episode, domestic ruminants were shown to be involved in RVFV circulation during interepizootic periods (5,6); together with the potential vertical transmission in *Aedes* spp. mosquitoes in Madagascar,

they might play a role in the maintenance of RVFV. However, genetic evidence indicates that RVFV outbreaks in Madagascar are not associated with emergence from enzootic cycles but that they are associated with recurrent virus introductions from mainland east Africa (7). Although these mechanisms for RVFV epidemiology on Madagascar are documented, the possibility of a wild mammal reservoir cannot be excluded. We therefore explored the role of wild terrestrial small mammals in the maintenance of RVFV in Madagascar, especially the nonnative, abundant, and ubiquitous black rats (*Rattus rattus*) (8), as has been suggested in rural Egypt (9,10).

For this study, 1,610 blood samples were obtained from different species of wild terrestrial small mammals in Madagascar (Figure). Permits to capture and collect animals were obtained from national authorities. Animals were sampled from October 2008 through March 2010 at a site in the Anjozorobe-Angavo (Anjozorobe District) forest corridor (18°18'41.9' S, 48°00'57.6' E), where RVFV was first detected in humans and cattle in February 2008 (4) and within 100 km from where the first RVFV was isolated in 1979 (3). We collected 378 serum samples from 11 native Tenrecidae (Afrosoricida) tenrecs, 114 samples from 6 native Nesomyidae (Rodentia) rodents, and 471 samples from introduced *R. rattus* (Muridae, Rodentia) rats (online Technical Appendix, wwwnc.cdc.gov/EID/article/19/6/12-1074-Techapp1.pdf). In addition, during 2008, we obtained serum samples from 647 *R. rattus* or *R. norvegicus* rats living near humans in areas where RVFV was reportedly circulating during 2008 and 2009: the districts of Ankazobe, Antsiranana, Betafo, Ihoisy, Marovoay, and Moramanga (4,5) (Figure).

Serum samples were tested for IgG against RVFV by ELISA, as described (4), by using peroxidase-labeled recombinant protein A/G (Pierce, Rockford, IL, USA) or




Figure. Collection sites of wild terrestrial small mammals on Madagascar and the number of mammals tested for Rift Valley fever virus (RVFV). At certain localities, the genus and species of sampled rats were *Rattus rattus* or *R. norvegicus*.

anti-mouse or rat IgG (H+L) according to the ability to recognize the immunoglobulin of species endemic to Madagascar (data not shown). The results were negative for all samples tested. Liver and spleen samples from 947 animals caught in the Anjozorobe-Angavo forest corridor were also tested. Approximately 50–100 mg of liver and spleen from each individual was mixed and homogenized at a dilution of 1:10 in culture medium containing 40% fetal bovine serum. After centrifugation, supernatants were collected and pooled by species (maximum 5 individuals/pool). RNA was extracted from pooled supernatants by using TRIzol LS reagent (Invitrogen,

Carlsbad, CA, USA) according to the manufacturers' instructions. Detection of RVFV RNA was attempted by using real-time reverse transcription PCR (4). The results were negative for the 220 monospecific pools tested.

Serologic and virologic results from rodent and tenrec samples collected during and after the epizootic 2008–2009 periods were negative for RVFV; 72.8% had been collected in municipalities where RVFV cases were reported. This finding does not indicate a role of native Rodentia and Afrosoricida mammals in the epidemiology of RVFV in Madagascar, nor does it indicate evidence of infection of *Rattus* spp. rats, as suggested in

Egypt (9,10). The absence of infection in *Rattus* spp. rats during a period of intense RVFV circulation does not support its potential role during the outbreak and, a fortiori, in the maintenance of RVFV during interepizootic periods. Among wild terrestrial mammals in Madagascar, animals of the orders Carnivora and Primata are not considered as candidates for the maintenance of RVFV; however, bats (order Chiroptera) and introduced bushpigs (order Artiodactyla, family Suidae, genus *Potamochoerus*) could be candidates, and their role in RVFV maintenance should be investigated (2). At present, no evidence is available for the maintenance of RVFV in wild terrestrial small mammals (native and introduced) in Madagascar.

Acknowledgments

We thank Lalaina Nomenjanahary and Gabriel Ravelojaona for their participation in the field work. We are grateful to Lila Rahalison for sharing some of the rodent serum.

This study was supported by a grant from the Groupement d'Intérêt Scientifique, Centre de Recherche et de Veille sur les Maladies Emergentes dans l'Océan (project RIFT-OI no. PRAO/AIRD/CRVOI/08/07).

**Marie-Marie Olive,
Nadia Razafindralambo,
Tony Andrianaivo Barivelo,
Jean-Théophile
Rafisandratantsoa,
Voahangy Soarimalala,
Steven M. Goodman,
Pierre E. Rollin,
Jean-Michel Héraud,
and Jean-Marc Reynes**

Author affiliations: Institut Pasteur, Antananarivo, Madagascar (M.-M. Olive, N. Razafindralambo, J.-T. Rafisandratantsoa, J.-M. Héraud); Association Vahatra, Antananarivo (T. Andrianaivo Barivelo, V. Soarimalala, S.M. Goodman); Université d'Antananarivo, Antananarivo (T. Andrianaivo Barivelo); Field Museum of Natural History, Chicago, Illinois, USA (S.M. Goodman); Centers for

Disease Control and Prevention, Atlanta, Georgia, USA (P.E. Rollin); and Institut Pasteur, Lyon, France (J.-M. Reynes)

DOI: <http://dx.doi.org/10.3201/eid1906.121074>

References

- Swanepoel R, Coetzer JAW. Rift Valley fever. In: Coetzer JAW, Thomson GR, Tustin RC, editors. Infectious diseases of livestock. Oxford (UK): Oxford University Press; 2004. p. 1037–70.
- Olive MM, Goodman SM, Reynes JM. The role of the wild mammals in the maintenance of Rift Valley fever virus. *J Wildl Dis*. 2012;48:241–66.
- Fontenille D. Arbovirus transmission cycles in Madagascar [in French]. *Arch Inst Pasteur Madagascar*. 1989;55:1–317.
- Andriamandimby SF, Randrianarivo-Solofoniaina AE, Jeanmaire EM, Ravalolomanana L, Razafimanantsoa LT, Rakotojoelinandrasana T, et al. Rift Valley fever during rainy seasons, Madagascar, 2008 and 2009. *Emerg Infect Dis*. 2010;16:963–70. <http://dx.doi.org/10.3201/eid1606.091266>
- Jeanmaire EM, Rabenarivahiny R, Biarmann M, Rabibisoa L, Ravaomanana F, Randriamparany T, et al. Prevalence of Rift Valley fever infection in ruminants in Madagascar after the 2008 outbreak. *Vector Borne Zoonotic Dis*. 2011;11:395–402. <http://dx.doi.org/10.1089/vbz.2009.0249>
- Chevalier V, Rakotondrafara T, Jourdan M, Heraud JM, Andriamanivo HR, Durand B, et al. An unexpected recurrent transmission of Rift Valley fever virus in cattle in a temperate and mountainous area of Madagascar. *PLoS Negl Trop Dis*. 2011;5:e1423. <http://dx.doi.org/10.1371/journal.pntd.0001423>
- Carroll SA, Reynes JM, Khristova ML, Andriamandimby SF, Rollin PE, Nichol ST. Genetic evidence for Rift Valley fever outbreaks in Madagascar resulting from virus introductions from the East African mainland rather than enzootic maintenance. *J Virol*. 2011;85:6162–7. <http://dx.doi.org/10.1128/JVI.00335-11>.
- Soarimalala V, Goodman SM. Les petits mammifères de Madagascar. Antananarivo (Madagascar): Association Vahatra; 2011.
- Youssef BZ, Donia HA. The potential role of *Rattus rattus* in enzootic cycle of Rift Valley fever in Egypt. 2—Application of reverse transcriptase polymerase chain reaction (RT-PCR) in blood samples of *Rattus rattus*. *J Egypt Public Health Assoc*. 2002;77:133–41.
- Youssef BZ, Donia HA. The potential role of *Rattus rattus* in enzootic cycle of Rift Valley fever in Egypt. 1—Detection of RVF

antibodies in *R. rattus* blood samples by both enzyme linked immuno sorbent assay (ELISA) and immuno-diffusion technique (ID). *J Egypt Public Health Assoc*. 2001;76:431–41.

Address for correspondence: Marie-Marie Olive, Unité de Virologie, Institut Pasteur de Madagascar, Antananarivo 101, Madagascar; email: mmolive@pasteur.mg

Colostrum Replacer and Bovine Leukemia Virus Seropositivity in Calves

To the Editor: Bovine leukemia virus (BLV), a deltaretrovirus in the family *Retroviridae*, is the causative agent of enzootic bovine leukosis (EBL). Although EBL is still endemic to the Americas and eastern Europe, most countries in western Europe are EBL free in accordance with European Union legislation; for example, Great Britain has held EBL-free status since 1999. EBL is notifiable to the World Organisation for Animal Health (1), with disease incursion affecting international trade. Infection with BLV is life-long and persistent; the presence of antibodies or integrated proviral DNA are indicators of virus exposure. Two tests prescribed for international trade are the agar gel immunodiffusion test (AGIDT) and ELISA (2); these tests are used widely for diagnosis (2).

We report 5 cases that occurred in the United Kingdom during 2009, in which calves became seropositive for BLV after consuming colostrum replacer. In Dumfriesshire, Scotland, routine serologic screening for BLV detected seropositivity in 2 calves, which were artificial insemination bull candidates. In Newport, Wales, a BLV-seropositive

calf was identified during pre-export testing. And in Yorkshire, England, 2 more BLV-seropositive calves, also artificial insemination bull candidates, were identified. All calves were home bred, and there was no evidence (as documented by serologic testing) or history of EBL within the herd. The farms were considered to have low risk for disease incursion because the introduction of new animals was limited.

Further inquiry revealed that the calves had each been exclusively fed a colostrum replacer from North America, where BLV is endemic. Antibodies to BLV might have been present in the colostrum replacer and thus passively acquired by the calves, resulting in seropositivity.

The hypothesis was tested by monthly blood sampling and ELISA analysis for antibodies against BLV (Institute Pourquier, Montpellier, France). Although the batch of colostrum replacer that had been fed to the calves from Dumfriesshire was not available for investigation, another colostrum sample was obtained from the same manufacturer for analysis. The reconstituted colostrum replacer was tested by AGIDT (IDEXX, Bern, Switzerland) at the following dilutions: neat (manufacturer's guidelines), 1:2, 1:4, and 1:8. In addition, 2 commercial ELISA tests (Institute Pourquier and IDEXX) were used over a series of dilutions to 1:125. All serologic tests were conducted according to manufacturer's recommendations. To examine the samples for proviral DNA, we conducted PCRs to amplify a 385-bp fragment of the envelope gene (3).

At the various dilutions of colostrum replacer, all serologic tests gave clearly positive reactions. Proviral PCR of the colostrum replacer also returned positive results, which were confirmed by sequencing. The resultant envelope sequence (GenBank accession no. HF545344) was aligned with 23 other sequences obtained from GenBank, which encompassed all known BLV genotypes. Phylogenetic analysis

Absence of Rift Valley Fever Virus in Wild Mammals, Madagascar

Technical Appendix

Technical Appendix Table. Taxonomy of small terrestrial animals sampled in Anjozorobe, Madagascar, October 2008 and 2009, March 2009 and 2010

Order	Family	Species	No.
Afrosoricida	Tenrecidae	<i>Hemicentetes semispinosus</i>	25
		<i>Microgale dobsoni</i>	219
		<i>M. fotsifotsy</i>	3
		<i>M. gymnorhyncha</i>	3
		<i>M. longicaudata/majori</i>	8
		<i>M. parvula</i>	2
		<i>M. thomasi</i>	15
		<i>M. soricooides</i>	61
		<i>Oryzorictes hova</i>	38
		<i>Tenrec ecaudatus</i>	2
		<i>Setifer setosus</i>	2
Rodentia	Nesomyidae	<i>Eliurus majori</i>	15
		<i>E. minor</i>	17
		<i>E. tanala</i>	15
		<i>E. grandidieri</i>	3
		<i>Gymnuromys roberti</i>	11
		<i>Nesomys rufus</i>	53
	Muridae	<i>Rattus rattus</i>	471