

Plasmodium falciparum kelch 13 Mutations, 9 Countries in Africa, 2014–2018

Sarah E. Schmedes,¹ Dhruviben Patel,¹ Simran Dhal, Julia Kelley, Samaly S. Svigel, Pedro Rafael Dimbu, Adicatou-Lai Adeothy, Gauthier Mesia Kahunu, Papy Mandoko Nkoli, Abdoul Habib Beavogui, Simon Kariuki, Don P. Mathanga, Ousmane Koita, Deus Ishengoma, Ally Mohamad, Moonga Hawela, Leah F. Moriarty, Aaron M. Samuels, Julie Gutman, Mateusz M. Plucinski, Venkatachalam Udhayakumar, Zhiyong Zhou, Naomi W. Lucchi, Meera Venkatesan, Eric S. Halsey, Eldin Talundzic

The spread of drug resistance to antimalarial treatments poses a serious public health risk globally. To combat this risk, molecular surveillance of drug resistance is imperative. We report the prevalence of mutations in the *Plasmodium falciparum* kelch 13 propeller domain associated with partial artemisinin resistance, which we determined by using Sanger sequencing samples from patients enrolled in therapeutic efficacy studies from 9 sub-Saharan countries during 2014–2018. Of the 2,865 samples successfully sequenced before treatment (day of enrollment) and on the day of treatment failure, 29 (1.0%) samples contained 11 unique nonsynonymous mutations and 83 (2.9%) samples contained 27 unique synonymous mutations. Two samples from Kenya contained the S522C mutation, which has been associated with delayed parasite clearance; however, no samples contained validated or candidate artemisinin-resistance mutations.

Malaria remains a serious global health concern, causing ≈405,000 deaths annually, mainly in young children in Africa (1). Although substantial progress has been made over the past decade to reduce the global burden of malaria, several factors threaten these gains, including the emergence and spread of antimalarial drug resistance (1). Artemisinin-based combination therapies (ACTs) are the first-line treatment for uncomplicated malaria caused by *Plasmodium falciparum* parasites, as recommended by the World Health Organization (WHO) (2). Unfortunately, resistance to ACTs (i.e., delayed parasite clearance and clinical treatment failures) has emerged in the Greater Mekong Subregion of Southeast Asia, posing a considerable risk to malaria control in the region (3). Even though clinical resistance to ACTs has not been reported in Africa (1), the threat of its emergence remains.

Author affiliations: Association of Public Health Laboratories, Silver Spring, Maryland, USA (S.E. Schmedes); Centers for Disease Control and Prevention, Atlanta, Georgia, USA (S.E. Schmedes, D. Patel, S.S. Svigel, A.M. Samuels, J. Gutman, M.M. Plucinski, V. Udhayakumar, Z. Zhou, N.W. Lucchi, E. Talundzic); Williams Consulting LLC, Baltimore, Maryland, USA (D. Patel); The Wallace H. Coulter Department of Biomedical Engineering, Georgia Institute of Technology, Atlanta (S. Dahl); CDC Foundation, Atlanta (J. Kelley); National Malaria Control Program, Ministry of Health, Luanda, Angola (P.R. Dimbu); National Malaria Control Program, Ministry of Health, Porto-Novo, Benin (A.-L. Adeothy); University of Kinshasa, Kinshasa, Democratic Republic of the Congo (G.M. Kahunu); National Institute of Biomedical Research, Kinshasa (P.M. Nkoli); Maferinyah Rural Health Research Center, Maferinyah, Guinea (A.H. Beavogui); Kenya Medical Research Institute, Centre for Global Health Research, Kisumu, Kenya (S. Kariuki); University

of Malawi College of Medicine, Blantyre, Malawi (D.P. Mathanga); University of Sciences, Techniques, and Technologies of Bamako, Bamako, Mali (O. Koita); Harvard T.H. Chan School of Public Health, Boston, Massachusetts, USA (D. Ishengoma); National Institute for Medical Research, Tanga Research Centre, Dar es Salaam, Tanzania (D. Ishengoma, A. Mohamad); Monash University Faculty of Pharmaceutical Sciences, Melbourne, Australia (D. Ishengoma); National Malaria Elimination Centre, Lusaka, Zambia (M. Hawela); US President's Malaria Initiative, Centers for Disease Control and Prevention, Atlanta (L.F. Moriarty, M.M. Plucinski, E.S. Halsey); US President's Malaria Initiative, US Agency for International Development, Washington, DC, USA (M. Venkatesan)

DOI: <https://doi.org/10.3201/eid2707.203230>

¹These authors contributed equally to this article.

As part of antimalarial therapeutic efficacy activities, WHO recommends molecular surveillance of the *P. falciparum* kelch 13 gene (*Pfk13*) (with focus on the propeller domain region), a molecular marker associated with delayed clearance of parasitemia after therapy with artemisinin monotherapy or an ACT (3–7). Because specific single-nucleotide polymorphisms (SNPs) within the propeller domain region of *Pfk13* continue to be discovered, WHO continues to update a list of these SNPs on the basis of association with delayed parasite clearance and reduced in vitro drug susceptibility (Table 1). Nine SNPs are currently considered validated by WHO to have delayed parasite clearance and in vitro data demonstrating partial resistance to artemisinin (3). WHO categorized 11 SNPs as candidate mutations, correlated with delayed parasite clearance but not validated with in vitro data (3). An additional 11 SNPs are listed by WHO as associated with delayed parasite clearance but without statistical significance because of limited data (3).

WHO recommends that malaria-endemic countries perform therapeutic efficacy studies (TESs) every 2 years to evaluate antimalarial treatments currently used in a particular region (8). Surveillance for molecular markers associated with antimalarial resistance is a recommended part of a TES to detect the presence of mutations associated with resistance (8). As part of the US President’s Malaria Initiative, the Centers for Disease Control and Prevention (CDC) and the US Agency for International Development provide support to countries in Africa to perform TESs, including molecular characterization of antimalarial-resistance markers, through the PMI-supported Antimalarial Resistance Monitoring in Africa (PARMA) network (9). Established in 2015, this endeavor involves laboratory trainees in Africa who bring TES samples from their home country to the CDC (Atlanta, Georgia, USA) to receive advanced laboratory training and perform molecular testing for antimalarial-resistance mutations (9). In this article, we report *Pfk13* mutation data generated from samples analyzed and collected from TESs conducted in 9 countries in Africa during 2014–2018.

Methods

Samples, Ethics Statement, and TES Protocols

Before initiation, all work described in this article was approved by the respective institutional ethics review committee in each country and the Office of the Associate Director of Science of CDC’s Center for Global Health and assigned the following tracking numbers: 2014–233a and 2014–233b (Angola), 2017–141 (Benin), 2018–035 (DRC), 2016–046 (Guinea), 6696.0 (Kenya), 6029.0 (Malawi), 2016–012a (Mali), 2015–073a (Tanzania), and 2016–200 (Zambia). Dried blood spots were collected from TESs conducted in 9 countries in Africa (Angola, Benin, the Democratic Republic of the Congo [DRC], Guinea, Kenya, Malawi, Mali, Tanzania, and Zambia; Table 2) during 2014–2018. The samples included those obtained pretreatment (at day of enrollment) and at day of treatment failure. Day of treatment failure samples came from patients experiencing a recrudescence or new infection during the follow-up period of (usually ending at 28 or 42 days) after administration of an ACT. TES and antimalarial molecular marker results for some of the data analyzed have been previously published for Angola (10–12), Kenya (13), and Tanzania (14). Results might differ slightly from previously published works because those works might not have reported results from all samples, might not have reported mutations in mixed infections, or might not have reported synonymous mutation results. Our study was a reanalysis of all available sequences using the same sequence data analysis quality filters, cut-offs, and quality scores for all countries.

Sequencing of *Pfk13* Propeller Domain Region

We extracted DNA from dried blood spots using the QIAamp Blood DNA Kit (QIAGEN, <https://www.qiagen.com>) according to the manufacturer’s instructions. We amplified the propeller domain region from codon positions 389–649 by PCR and Sanger sequenced according to methods previously described (15).

Table 1. Mutations in the *Pfk13* gene and WHO classification related to *Plasmodium falciparum* artemisinin resistance*

Validated <i>Pfk13</i> mutations	Candidate <i>Pfk13</i> mutations	Non–statistically significant associated <i>Pfk13</i> mutations
F446I	P441L	D452E
N458Y	G449A	C469Y
M476I	C469F	K479I
Y493H	A481V	R515K
R539T	P527H	S522C
I543T	N537I	N537D
P553L	G538V	R575K
R561H	V568G	M579I
C580Y	P574L, F673I, A675V	D584V, P667T, H719N

*Adapted from an August 2018 WHO status report on artemisinin resistance and artemisinin-based combination therapy efficacy (3). *Pfk13*, *P. falciparum* kelch 13; WHO, World Health Organization.

Table 2. Summary of antimalarial therapeutic efficacy studies, 9 countries in Africa, 2002–2007*

Country	Sites	Treatments studied	Age of patients enrolled	Year	Total no. samples			ACTs introduced
					D0 + DF	D0	DF	
Angola	Benguela, Zaire, Lunda Sul	AL, ASAQ, DP	6 mo–12 y	2015	379	379	0	2005
				2017	76	38	38	2005
Benin	Klouanmey, Djougou	AL	6–59 mo	2017	194	175	19	2004
DRC	Kabondo, Kapolowe, Rutshuru, Mikalayi, Kimpese	AL, ASAQ, DP	6–59 mo	2017–2018	633	317	316	2006
Guinea	Maferinyah, Labè	AL	6–59 mo	2016	432	409	23	2004–2005
Kenya	Siaya County	AL, DP	6–59 mo	2016–2017	417	325	92	2006
Malawi	Machinga, Nkhotakota, Karonga	AL, ASAQ	6–59 mo	2014	27	8	19	2007
Mali	Dioro, Sèlinguè	AL, ASAQ	2–59 mo	2015–2016	410	320	90	2006
Tanzania	Kibaha, Ujiji, Mkuzi, Mlimba	AL	6 mo–10 y	2016	417	345	72	2006
Zambia	Gwembe, Katete, Mansa	AL, DP	>6 mo	2016	263	263	0	2002
Total					3,248	2,579	669	

*ACTs, artemisinin-based combination therapies; AL, artemether/lumefantrine; ASAQ, artesunate/amodiaquine; D0, day of enrollment (pretreatment); DF, day of failure; DP, dihydroartemisinin/piperaquine; DRC, Democratic Republic of the Congo.

Data Analysis

We analyzed sequence data by using Geneious Prime (Biomatters, <https://www.geneious.com>). We trimmed and quality filtered forward and reverse sequence reads for each sample (error probability limit 0.05, maximum low-quality bases 30) from the 3' and 5' ends to remove low-quality bases. We aligned trimmed sequences to the *Pfk13* National Center for Biotechnology Information gene reference no. PF3D7_1343700 (<https://www.ncbi.nlm.nih.gov/gene/814205>) and assessed for SNPs. We only considered SNPs if they had a Phred quality score of ≥ 30 and were present in both forward and reverse strands. Mixed infections were detected by using the heterozygous caller plug-in tool in Geneious with a threshold of $\geq 30\%$. A second analyst confirmed all SNP and heterozygous calls by manual technical review. We submitted all *Pfk13* sequences with SNPs reported in this study to GenBank (accession nos. MN072940–3042). We used R software version 4.0.1 (R Foundation for Statistical Computing, <https://www.r-project.org>) to generate a map showing the distribution of mutations in the 9 countries (Figure).

Results

We attempted *Pfk13* sequencing on 3,248 samples (2,579 pretreatment and 669 day of failure samples) from the 9 countries (Table 2); 2,865 were successfully sequenced (Table 3). Of those, 2,753 samples were wild-type. A total of 11 unique nonsynonymous mutations and 27 unique synonymous mutations were detected in 2,865 successfully sequenced pretreatment and day of failure samples from Angola, Benin, DRC, Guinea, Kenya, Malawi, Mali, Tanzania, and Zambia collected during 2014–2018 (Figure, Table 4; Appendix 1, <https://wwwnc.cdc.gov/EID/article/27/7/20-3230-App1.pdf>).

Of the 2,303 sequenced pretreatment samples, 2,213 were wild-type and 90 (3.9%) contained mutations (Table 3). Of the 90 pretreatment samples with mutations, 10 unique nonsynonymous mutations were present in 25 samples from 8 of the 9 countries assessed (Table 4) and 25 unique synonymous mutations were present in 65 samples from 8 of the 9 countries assessed (Appendix 1 Table 1). Two samples from Kenya contained the S522C mutation, reported by WHO as a less-frequent mutation associated with delayed parasite clearance but without statistical significance because of limited data (3). Both of these patients cleared their initial infection. A578S, the most commonly found mutation in Africa (not associated with resistance) (3), was the most common nonsynonymous mutation we identified. The mutation was found in 14 pretreatment isolates: 4 in Angola, 1 in DRC, 1 in Mali, 6 in Kenya, 1 in Tanzania, and 1 in Zambia (Table 4). No mutations were identified in the samples from Malawi. Eight of the 10 unique nonsynonymous mutations in the pretreatment samples have been reported previously in other countries, whereas 2 mutations, P419S (Guinea) and Q613R (Angola), were newly identified in our study. No WHO-validated or candidate *Pfk13* mutations were identified.

Of the 669 day of failure samples, 562 were successfully sequenced; 107 (16.0%) samples failed to amplify, produced poor-quality sequences, or both (Table 3). A total of 540 samples were wild-type. Two nonsynonymous mutations were found in 4 day of failure samples (Table 4) and 10 synonymous mutations (Appendix 1 Table 2) were identified in 18 day of failure samples from 4 countries. Of the nonsynonymous mutations in day of failure samples, 2 samples from Kenya and 1 sample from DRC contained the A578S mutation, and 1 sample from DRC contained the S477Y mutation (Table 4). We compiled the complete results of the




Figure. Prevalence of *Plasmodium falciparum* kelch 13 mutations in pretreatment therapeutic efficacy study samples, 9 countries in Africa, 2014–2018. A total of 11 unique nonsynonymous and 27 unique synonymous mutations were detected in 2,865 successfully sequenced pretreatment and day of failure samples from Angola, Benin, Democratic Republic of the Congo, Guinea, Kenya, Malawi, Mali, Tanzania, and Zambia collected during 2014–2018. A total of 2,753 samples were wild-type. Data from Angola includes results from 2 therapeutic efficacy studies.

sequence data reanalysis (Appendix 2, <https://wwwnc.cdc.gov/EID/article/27/7/20-3230-App2.xlsx>).

Discussion

This work provides an update on *Pfk13* genetic markers in 9 countries in Africa with endemic malaria. Although clinical resistance to ACTs has yet to be confirmed in Africa (1), the early detection of *Pfk13* mutations through surveillance allows for swift action before resistance spreads widely. To

date, all WHO-validated SNPs detected in Africa have been the result of independent emergence as opposed to spreading through imported cases from Southeast Asia (21). More than 200 *Pfk13* mutations have been identified in global samples (3,18,21), and ≥74 *Pfk13* nonsynonymous mutations have been reported in Africa (22,23). In this study, we report the presence of S522C in Kenya, a less frequent mutation that has been previously reported to be associated with delayed parasite clearance but lacking

Table 3. Summary of *Pfk13* gene mutations detected in *Plasmodium falciparum* pretreatment and DF samples, 9 countries in Africa, 2014–2018*

Country (year)	Total with sequencing attempted	Poor quality or no amplification	No. samples pretreatment (DF)			
			Successfully sequenced	Wild-type samples pretreatment	Other nonsynonymous mutations	Synonymous mutations
Angola (2015)	379 (0)	77 (0)	302 (0)	291 (0)	5 (0)	6 (0)
Angola (2017)	38 (38)	0 (2)	38 (36)	37 (36)	1 (0)	0 (0)
Benin (2017)	175 (19)	20 (1)	155 (18)	151 (18)	1 (0)	3 (0)
DRC (2017–2018)	317 (316)	13 (34)	304 (282)	295 (269)	1 (2)	8 (11)
Guinea (2016)	409 (23)	20 (1)	389 (22)	380 (22)	1 (0)	8 (0)
Kenya (2016–2017)	325 (92)	7 (4)	318 (88)	302 (85)	8 (2)	8 (1)
Malawi (2014)	8 (19)	1 (5)	7 (14)	7 (14)	0 (0)	0 (0)
Mali (2015–2016)	320 (90)	68 (48)	252 (42)	244 (39)	1 (0)	7 (3)
Tanzania (2016)	345 (72)	20 (12)	325 (60)	306 (57)	6 (0)	13 (3)
Zambia (2016)	263 (0)	50 (0)	213 (0)	200 (0)	1 (0)	12 (0)
Total	2,579 (669)	276 (107)	2,303 (562)	2,213 (540)	25 (4)	65 (18)

*DF, day of failure; DRC, Democratic Republic of the Congo; *Pfk13*, *Plasmodium falciparum* kelch 13.

Table 4. Summary of *Pfk13* nonsynonymous mutations detected in *Plasmodium falciparum* pretreatment and DF samples, 9 countries in Africa, 2014–2018*

Mutation	Country	Codon change	No. samples pretreatment (DF)	Country or region where previously reported (reference)
I416V	Tanzania	ATA → GTA	1 (0)	Tanzania (14)
P419S	Guinea	CCA → TCA	1 (0)	NA
E433D	Tanzania	GAA → GAC	1 (0)	Tanzania (14)
R471S	Tanzania	CGT → AGT	1 (0)	Tanzania (14)
S477Y	DRC	TCT → TAT	0 (1)	Grande Comore Island (16)
A504V	Angola (2017)	GCT → GTT	1 (0)	Gabon (17)
S522C	Kenya	AGT → TGT	2 (0)	Africa (18)
A569G	Benin	GCA → GGA	1 (0)	Gambia (19) and Niger (20)
A578S	Angola (2015)	GCT → TCT	4 (0)	Africa (19)
A578S	DRC	GCT → TCT	1 (1)	Africa (19)
A578S	Mali	GCT → TCT	1 (0)	Africa (19)
A578S	Kenya	GCT → TCT	6 (2)	Africa (19)
A578S	Tanzania	GCT → TCT	1 (0)	Africa (19)
A578S	Zambia	GCT → TCT	1 (0)	Africa (19)
Q613R	Angola (2015)	CAA → CGA	1 (0)	NA
Q613E	Tanzania	CAA → GAA	2 (0)	Tanzania (14)
Total			25 (4)	

*DF, day of failure; DRC, Democratic Republic of the Congo; NA, not available; *Pfk13*, *Plasmodium falciparum* kelch 13.

sufficient evidence to be considered a WHO-validated or candidate mutation (3).

As more molecular surveillance data are collected, previous results should be reinterpreted to determine the presence of WHO-reportable mutations because the importance of these mutations in drug resistance might change based on new data (3,24). Although we report only 1 mutation identified by WHO to possibly play a role in resistance, other detected mutations, such as the other nonsynonymous mutations with unknown resistance status reported in this study, might be deemed important in the future as more data are collected and validated. In 2017, WHO categorized only 5 mutations as validated (N458Y, Y493H, R539T, I543T, and 580Y) (24), but in 2018 the validated list was updated to include an additional 4 mutations, including F446L, P553L, and R561H (formerly candidate markers) and M476I (formerly reported as a less frequent variant associated with in vivo or in vitro test results) (3). In addition, the Worldwide Antimalarial Resistance Network tracks *Pfk13* mutations worldwide and strives to detect new associations of mutations with delayed parasite clearance, which might inform WHO classifications (18).

We report the presence of 11 unique nonsynonymous mutations in Angola, Benin, Guinea, DRC, Kenya, Mali, Tanzania, and Zambia; all were previously reported in the literature (Table 4) except P419S and Q613R. The most common nonsynonymous mutation observed in our study was A578S, a nonsynonymous mutation frequently described in Africa (3) and, to a lesser extent, Asia (e.g., Thailand [19] and Bangladesh [25]). WHO has reported that A578S is not associated with partial artemisinin resistance (3). Most muta-

tions detected were synonymous mutations consistent with previous reports (21). Because synonymous mutations do not result in an amino acid change, they are not associated with resistance. Parasites from Africa have been shown to have a higher prevalence of synonymous mutations, which is not surprising given that *P. falciparum* originated in Africa and continues to have a high level of transmission in this region (19).

The results described in this article represent the collaborative output of the PARMA network, which originated in 2015 with the objectives of assisting countries in Africa in testing malaria samples from TESs for genetic markers associated with antimalarial resistance and supporting training and capacity building of collaborators in Africa (9). In 8 of the 9 countries included in this report (all but Angola), the *Pfk13* results were generated during a 6–8-week visit to CDC by trainees from a laboratory in the country where the TES was performed. Results were subsequently shared by the trainee's laboratory with their national malaria control program and other local stakeholders to make decisions related to antimalarial use. Although the *Pfk13* results we have described would not be cause for alarm or policy change, recent findings in Rwanda suggests a substantial presence of the *Pfk13* R561H mutation (26) that has evolved locally, highlighting the importance of molecular surveillance for early detection of emerging patterns of resistance. In this context, PARMA training visits generate a vast amount of data from TES samples, ranging from efficacy results to prevalence of other molecular markers (e.g., *P. falciparum* multidrug-resistant protein 1 and *P. falciparum* chloroquine-resistance transporter) to the

presence of *P. falciparum* histidine-rich protein 2 and 3 deletions (which might affect rapid diagnostic test performance). Generating phenotypic (i.e., efficacy) and genotypic data on the same sample provides an opportunity to identify novel mutations associated with resistance and enables detection of known mutations in samples with well-characterized efficacy outcomes. Because the PARMA network encourages standardization of laboratory methods and data reporting, such explorations might detect trends over time in a single country or produce insightful observations by using data from multiple countries. With the increased use of next-generation sequencing, the PARMA network has embarked on applying these principles of data generation, capacity building, networking, and standardization to this emerging technology (27). The ultimate goal of laboratories in Africa independently analyzing their own malaria samples.

This work was made possible with funding provided by the US President's Malaria Initiative (PMI) through PARMA. We also acknowledge partial support from the Advanced Molecular Detection Initiative at the CDC and partial support by the Bioinformatics Fellowship Program administered by the Association of Public Health Laboratories and funded by CDC. S.E.S. was supported by the Bioinformatics Fellowship Program. J.K. was supported in part by the CDC Foundation. D.P. was employed by Williams Consulting LLC, which provided support in the form of salary for D.P. D.S.I. was partly supported by the Developing Excellence in Leadership and Genetics Training for Malaria Elimination (DELGEME) in Sub-Saharan Africa program through the Developing Excellence in Leadership, Training and Science Africa Initiative (DELGEME grant no. 107740/Z/15/Z). The Developing Excellence in Leadership, Training and Science Africa Initiative is an independent funding scheme of the African Academy of Sciences' Alliance for Accelerating Excellence in Science in Africa and is supported by the New Partnership for Africa's Development Planning and Coordinating Agency with funding from the Wellcome Trust (DELGEME grant no. 107740/Z/15/Z) and government of the United Kingdom. The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

About the Author

Dr. Schmedes is lead bioinformatician at the Florida Department of Health; her primary interests include developing and implementing bioinformatics methods for studying bacterial and viral pathogens of public health significance in the state of Florida. Dr. Talundzic is an

informatics health scientist in the Center for Global Health at CDC; his primary interests include developing and implementing next generation sequencing and bioinformatics methods for studying *Plasmodium* parasites.

References

1. World Health Organization. World malaria report 2019 [cited 2020 May 15]. <https://www.who.int/publications/i/item/9789241565721>
2. World Health Organization. Guidelines for the treatment of malaria. 3rd edition. 2015 [cited 2020 May 15]. <https://apps.who.int/iris/handle/10665/162441>
3. World Health Organization. Artemisinin resistance and artemisinin-based combination therapy efficacy: status report. 2018 [cited 2020 May 15]. <https://apps.who.int/iris/handle/10665/274362>
4. Cheeseman IH, Miller BA, Nair S, Nkhoma S, Tan A, Tan JC, et al. A major genome region underlying artemisinin resistance in malaria. *Science*. 2012;336:79–82. <https://doi.org/10.1126/science.1215966>
5. Takala-Harrison S, Clark TG, Jacob CG, Cummings MP, Miotto O, Dondorp AM, et al. Genetic loci associated with delayed clearance of *Plasmodium falciparum* following artemisinin treatment in Southeast Asia. *Proc Natl Acad Sci U S A*. 2013;110:240–5. <https://doi.org/10.1073/pnas.1211205110>
6. Arey F, Witkowski B, Amaratunga C, Beghain J, Langlois A-C, Khim N, et al. A molecular marker of artemisinin-resistant *Plasmodium falciparum* malaria. *Nature*. 2014;505:50–5. <https://doi.org/10.1038/nature12876>
7. Ashley EA, Dhorda M, Fairhurst RM, Amaratunga C, Lim P, Suon S, et al.; Tracking Resistance to Artemisinin Collaboration (TRAC). Spread of artemisinin resistance in *Plasmodium falciparum* malaria. *N Engl J Med*. 2014;371:411–23. <https://doi.org/10.1056/NEJMoa1314981>
8. World Health Organization. Methods for surveillance of antimalarial drug efficacy. 2009 [cited 2020 May 15]. <https://www.who.int/malaria/publications/atoz/9789241597531/en>
9. Halsey ES, Venkatesan M, Plucinski MM, Talundzic E, Lucchi NW, Zhou Z, et al. Capacity development through the US President's Malaria Initiative-Supported Antimalarial Resistance Monitoring in Africa Network. *Emerg Infect Dis*. 2017;23:S53–S56. <https://doi.org/10.3201/eid2313.170366>
10. Plucinski MM, Dimbu PR, Macaia AP, Ferreira CM, Samutondo C, Quivinja J, et al. Efficacy of artemether-lumefantrine, artesunate-amodiaquine, and dihydroartemisinin-piperazine for treatment of uncomplicated *Plasmodium falciparum* malaria in Angola, 2015. *Malar J*. 2017;16:62. <https://doi.org/10.1186/s12936-017-1712-4>
11. Ljolje D, Dimbu PR, Kelley J, Goldman I, Nace D, Macaia A, et al. Prevalence of molecular markers of artemisinin and lumefantrine resistance among patients with uncomplicated *Plasmodium falciparum* malaria in three provinces in Angola, 2015. *Malar J*. 2018;17:84. <https://doi.org/10.1186/s12936-018-2233-5>
12. Davlantes E, Dimbu PR, Ferreira CM, Florinda Joao M, Pode D, Félix J, et al. Efficacy and safety of artemether-lumefantrine, artesunate-amodiaquine, and dihydroartemisinin-piperazine for the treatment of uncomplicated *Plasmodium falciparum* malaria in three provinces in Angola, 2017. *Malar J*. 2018;17:144. <https://doi.org/10.1186/s12936-018-2290-9>

13. Chebore W, Zhou Z, Westercamp N, Otieno K, Shi YP, Sergeant SB, et al. Assessment of molecular markers of anti-malarial drug resistance among children participating in a therapeutic efficacy study in western Kenya. *Malar J*. 2020;19:291. <https://doi.org/10.1186/s12936-020-03358-7>
14. Ishengoma DS, Mandara CI, Francis F, Talundzic E, Lucchi NW, Ngasala B, et al. Efficacy and safety of artemether-lumefantrine for the treatment of uncomplicated malaria and prevalence of Pfk13 and Pfmdr1 polymorphisms after a decade of using artemisinin-based combination therapy in mainland Tanzania. *Malar J*. 2019;18:88. <https://doi.org/10.1186/s12936-019-2730-1>
15. Talundzic E, Chenet SM, Goldman IF, Patel DS, Nelson JA, Plucinski MM, et al. Genetic analysis and species specific amplification of the artemisinin resistance-associated kelch propeller domain in *P. falciparum* and *P. vivax*. *PLoS One*. 2015;10:e0136099.
16. Huang B, Deng C, Yang T, Xue L, Wang Q, Huang S, et al. Polymorphisms of the artemisinin resistant marker (K13) in *Plasmodium falciparum* parasite populations of Grande Comore Island 10 years after artemisinin combination therapy. *Parasit Vectors*. 2015;8:634. <https://doi.org/10.1186/s13071-015-1253-z>
17. Voumbo-Matoumona DF, Kouma LC, Madamet M, Maghendji-Nzondo S, Pradines B, Lekana-Douki JB. Prevalence of *Plasmodium falciparum* antimalarial drug resistance genes in southeastern Gabon from 2011 to 2014. *Infect Drug Resist*. 2018;11:1329–38. <https://doi.org/10.2147/IDR.S160164>
18. WWARN K13 Genotype-Phenotype Study Group. Association of mutations in the *Plasmodium falciparum* Kelch13 gene (Pf3D7_1343700) with parasite clearance rates after artemisinin-based treatments – a WWARN individual patient data meta-analysis. *BMC Med*. 2019;17:1. <https://doi.org/10.1186/s12916-018-1207-3>
19. Ménard D, Khim N, Beghain J, Adegnika AA, Shafiul-Alam M, Amodu O, et al; KARMA Consortium. A worldwide map of *Plasmodium falciparum* K13-propeller polymorphisms. *N Engl J Med*. 2016;374:2453–64. <https://doi.org/10.1056/NEJMoa1513137>
20. Laminou I, Lamine M, Arzika I, Mahamadou B, Gora D, Dieye A. Detection of *Plasmodium falciparum* K13 propeller A569G mutation after artesunate-amodiaquine treatment failure in Niger. *J Adv Biol Biotechnol*. 2018;18:1–8. <https://doi.org/10.9734/JABB/2018/42872>
21. MalariaGEN *Plasmodium falciparum* Community Project. Genomic epidemiology of artemisinin resistant malaria. *eLife*. 2016;5:e08714. <https://doi.org/10.7554/eLife.08714>
22. Ocan M, Akena D, Nsobya S, Kanya MR, Senono R, Kinengyere AA, et al. K13-propeller gene polymorphisms in *Plasmodium falciparum* parasite population in malaria affected countries: a systematic review of prevalence and risk factors. *Malar J*. 2019;18:60. <https://doi.org/10.1186/s12936-019-2701-6>
23. Conrad MD, Rosenthal PJ. Antimalarial drug resistance in Africa: the calm before the storm? *Lancet Infect Dis*. 2019; 19:e338–51. [https://doi.org/10.1016/S1473-3099\(19\)30261-0](https://doi.org/10.1016/S1473-3099(19)30261-0)
24. World Health Organization. Artemisinin and artemisinin-based combination therapy resistance: status report. 2017 [cited 2020 May 15]. <https://apps.who.int/iris/handle/10665/255213>
25. Mohon AN, Alam MS, Bayih AG, Folefoc A, Shahinas D, Haque R, et al. Mutations in *Plasmodium falciparum* K13 propeller gene from Bangladesh (2009–2013). *Malar J*. 2014;13:431. <https://doi.org/10.1186/1475-2875-13-431>
26. Uwimana A, Legrand E, Stokes BH, Ndikumana JM, Warsame M, Umulisa N, et al. Emergence and clonal expansion of in vitro artemisinin-resistant *Plasmodium falciparum* kelch13 R561H mutant parasites in Rwanda. *Nat Med*. 2020;26:1602–8. <https://doi.org/10.1038/s41591-020-1005-2>
27. Talundzic E, Ravishankar S, Kelley J, Patel D, Plucinski M, Schmedes S, et al. Next-generation sequencing and bioinformatics protocol for malaria drug resistance marker surveillance. *Antimicrob Agents Chemother*. 2018;62:e02474–17. <https://doi.org/10.1128/AAC.02474-17>

Address for correspondence: Eldin Talundzic, Centers for Disease Control and Prevention, 1600 Clifton Rd NE, Mailstop H23-10, Atlanta, GA 30329-4018, USA; email: etalundzic@cdc.gov

Plasmodium falciparum kelch 13 Mutations, 9 Countries in Africa, 2014–2018

Appendix 1

Appendix 1 Table 1. Summary of *Pfk13* synonymous mutations detected in pre-treatment samples from 9 countries in Africa*

Mutation	Country	Codon change	No. samples by country	No. samples total
P413P	Zambia, Mali	CCG → CCA	1, 1	2
P417P	Angola 2015, DRC, Kenya, Tanzania, Zambia	CCC → CCT	2, 2, 2, 3, 2	11
L429L	Guinea	TTG → CTG	1	1
V454V	Kenya	GTA → GTG	1	1
C469C	Angola 2015, Benin, Guinea, Kenya, Mali, Tanzania	TGC → TGT	1, 1, 3, 2, 2, 3	12
R471R	Angola 2015, DRC, Kenya, Tanzania, Zambia	CGT → CGC	2, 1, 1, 2, 3	9
S477S	DRC, Zambia	TCT → TCG	1, 2	3
T478T	Benin, Mali, DRC	ACC → ACG	1, 1, 1	3
V487V	Tanzania	GTA → GTG	4	4
Y493Y	Mali	TAC → TAT	1	1
G496G	Guinea	GGT → GGC	1	1
K503K	Mali	AAG → AAA	1	1
A504A	DRC, Kenya	GCT → GCC	1, 1	2
E509E	Guinea	GAG → GAA	1	1
V510V	Guinea	GTG → GTA	1	1
I526I	Angola 2015	ATA → ATT	1	1
R529R	Zambia	AGA → AGG	1	1
G533G	Kenya	GGT → GGA	1	1
R539R	Tanzania	AGA → CGA	1	1
A557A	Zambia	GCA → GCG	1	1
R575R	Zambia	AGA → AGG	2	2
S576S	DRC	TCA → TCG	2	2
V589V	Benin	GTC → GTA	1	1
Q613Q	Mali	CAA → CAG	1	1
A621A	Guinea	GCT → GCA	1	1
Total				65

*DRC, Democratic Republic of the Congo; *Pfk13*, *Plasmodium falciparum* kelch 13.

Appendix 1 Table 2. Summary of *Pfk13* synonymous mutations detected in day of failure samples from 4 countries in Africa*

Mutation	Country	Codon change	No. samples by country	No. samples total
P417P	DRC	CCC → CCT	1	1
C469C	DRC, Tanzania	TGC → TGT	1, 3	4
R471R	DRC	CGT → CGC	5	5
Y493Y	Mali	TAC → TAT	2†	2
G496G	DRC	GGT → GGC	1	1
Y511Y	DRC	TAT → TAC	1	1
G538G	Kenya	GGT → GGA	1	1
R539R	DRC	AGA → CGA	1	1
Q613Q	Mali	CAA → CAG	1	1
A621A	DRC	GCT → GCA	1	1
Total				18

*DRC, Democratic Republic of the Congo; *Pfk13*, *Plasmodium falciparum* kelch 13.

†Both day of failure samples were collected from the same patient at different time intervals post-treatment.