

Address for correspondence: Kanta Subbarao, WHO Collaborating Centre for Reference and Research on Influenza, Department of Microbiology and Immunology, The University of Melbourne at The Peter Doherty Institute for Infection and Immunity, 792 Elizabeth St, Melbourne, VIC, 3000, Australia; email: kanta.subbarao@influenzacentre.org

## Serologic Evidence of Human Exposure to Ehrlichiosis Agents in Japan

Hongru Su, Kenji Kubo, Shigetoshi Sakabe, Shinsuke Mizuno, Nobuhiro Komiyama, Shigehiro Akachi, Hiromi Fujita,<sup>1</sup> Kozue Sato, Hiroki Kawabata, Hiromi Nagaoka, Shuji Ando, Norio Ohashi

Author affiliations: University of Shizuoka, Shizuoka, Japan (H. Su, N. Ohashi); Japanese Red Cross Wakayama Medical Center, Wakayama, Japan (K. Kubo, S. Mizuno, N. Komiyama); Ise Red Cross Hospital, Ise, Japan (S. Sakabe); Mie Prefecture Health and Environment Research Institute, Yokkaichi, Japan (S. Akachi); Mahara Institute of Medical Acarology, Anan, Japan (H. Fujita); National Institute of Infectious Diseases, Tokyo, Japan (K. Sato, H. Kawabata, S. Ando); Shizuoka Institute of Environment and Hygiene, Shizuoka (H. Nagaoka)

DOI: <https://doi.org/10.3201/eid2811.212566>

In retrospective analyses, we report 3 febrile patients in Japan who had seroconversion to antibodies against *Ehrlichia chaffeensis* antigens detected by using an immunofluorescence and Western blot. Our results provide evidence of autochthonous human ehrlichiosis cases and indicate ehrlichiosis should be considered a potential cause of febrile illness in Japan.

**H**uman ehrlichiosis is a tickborne infectious disease caused by *Ehrlichia* sp. that has primarily been detected in the United States. Common clinical manifestations of human ehrlichiosis are fever, headache, myalgia, and malaise. Leukopenia and thrombocytopenia often occur. Symptoms range from mild

fever to severe illness with multiple organ dysfunction, which is occasionally fatal (1). In a retrospective analysis, we show serologic evidence for human ehrlichiosis in 3 febrile patients in Japan.

In case 1, a male patient, who was 48 years of age and worked in the manufacturing industry, sought care at a primary care clinic in 2015 for high fever (>40°C) and headache ≈1 month after hiking in the mountains. The clinic physician prescribed levofloxacin and acetaminophen, but the treatment was not effective. Therefore, the patient was seen at the Japanese Red Cross Wakayama Medical Center. The day before onset of high fever, the patient found a small rash on the left side of his abdomen. This date was considered day 0, although there might have been symptoms that the patient was unaware of before that time. The rash was an erythema migrans–like lesion that expanded on day 5. The patient was hospitalized, and borreliosis or tick-associated rash illness, which is similar to Lyme borreliosis–like erythema migrans, was suspected (2); however, a tick bite or eschar was not observed. After intravenous administration of minocycline (200 mg/d), the patient's fever abated, but the lesion expanded and was accompanied by puritis. On day 10, the patient was discharged from the hospital, after which the rash gradually disappeared. Diagnostic tests for borreliosis were negative. We retrospectively performed immunofluorescence assays (IFAs) and Western blot (Appendix, <https://wwwnc.cdc.gov/EID/article/28/11/21-2566-App1.pdf>) using patient serum samples collected on days 2 and 17. We showed seroconversion to antibodies against *Ehrlichia chaffeensis* antigens by IFA and the presence of IgM and IgG against *Ehrlichia* sp. P28 protein by Western blot (Table; Figure). We suspected the patient had ehrlichiosis and tick-associated rash illness.

In case 2, a male patient, who was 66 years of age and worked as a truck driver, sought care at the Ise Red Cross Hospital in 2018 for fever (38°C), annular erythema, and malaise. The patient had renal impairment and jaundice. The principal physician suspected leptospirosis, but diagnostic tests for leptospirosis were negative. The physician suspected other bacterial infections, including Japanese spotted fever (JSF) or anaplasmosis. The patient was treated intravenously with minocycline (200 mg/d) and sulbactam/ampicillin (6 g/d) for 4 days. Subsequently, amoxicillin (1.5 g/d) was administered orally for 14 days, and the patient recovered. Diagnostic tests for JSF were negative. We retrospectively analyzed patient serum samples collected on days 14, 32, and 60 after onset of illness. We showed seroconversion to antibodies against *E. chaffeensis*

<sup>1</sup>Current affiliation: Northern Fukushima Medical Center, Date, Japan.

**Table.** Evaluation of immunofluorescence assay titers and Western blots of serum samples from 3 febrile patients demonstrating serologic evidence of human exposure to ehrlichiosis agents in Japan\*

Case no. (year)	No. days†	<i>Ehrlichia chaffeensis</i> antigens, IgM/IgG			<i>Anaplasma phagocytophilum</i> antigens, IgM/IgG		
		IFA, THP-1 cells	Western blot		IFA		Western blot, THP-1 cells
			DH82 cells	THP-1 cells	THP-1 cells	HL60 cells	
1 (2015)	2	20/160	-/+	-/-	<20/<20	<20/<20	-/-
	17	80/640	+/+	+/+	<20/<20	<20/<20	-/-
2 (2018)	14	20/20	+/+	+/+	<20/<20	<20/<20	-/-
	32	40/320	+/+	+/+	<20/<20	<20/<20	-/-
	60	20/20	+/+	+/+	<20/<20	<20/<20	-/-
3 (2018)	5	20/20	+/+	+/+	<20/40	<20/20	-/+
	58	80/80	+/+	+/+	<20/40	<20/40	-/+
	115	20/320	+/+	+/+	<20/40	<20/40	-/+

\*Serum samples were collected from 3 patients in Japan in 2015 and 2018 and assayed by using THP-1, DH82, or HL60 cells infected with *E. chaffeensis* or *A. phagocytophilum*. Western blots were categorized as positive or negative for IgM and IgG against antigens from each bacterial species. IFA, immunofluorescence assay.


†No. days after onset of illness.

antigens by IFA and the presence of IgM and IgG against *Ehrlichia* sp. P28 protein by Western blot (Table; Appendix Figure 1). The IFA titers for both IgM and IgG decreased on day 60.

In case 3, a female patient, who was 69 years of age and owned a Japanese-style accommodation, sought care at the Ise Red Cross Hospital in 2018 for mild fever, generalized edema and rash, headache, and malaise. The principal physician suspected JSF

and treated the patient with oral minocycline (200 mg/d) and levofloxacin (500 mg/d) for 10 days; the patient recovered. Diagnostic tests for JSF were negative. We retrospectively analyzed patient serum samples collected on days 5, 58, and 115 by IFA and Western blot and found seroconversion to antibodies against *E. chaffeensis* antigens by IFA and the presence of both IgM and IgG against *Ehrlichia* sp. P28 protein antigens by Western blot (Table;

Figure. Western blots using serum samples from a febrile patient (case 1) in Wakayama Prefecture in study showing serologic evidence of human exposure to ehrlichiosis agents in Japan. Serum samples were collected from the patient on day 2 and 17 after onset of illness. Human THP-1 and canine DH82 cells were uninfected or infected with *Ehrlichia chaffeensis*. THP-1 cells were also infected with *Anaplasma phagocytophilum*. Cell lysates were separated and Western blot was performed as described (Appendix, <https://wwwnc.cdc.gov/EID/article/28/11/21-2566-App1.pdf>). We used uninfected THP-1 and DH82 cells as negative lysate controls. We used rabbit serum against recombinant P44 antigens specific for *A. phagocytophilum* and recombinant P28 antigens specific for *E. chaffeensis* (1:10,000 dilution) as positive



serum controls. We used serum from a healthy donor as a negative control serum (Precision for Medicine, <https://www.precisionbiospecimens.com>). The patient's serum samples and negative control serum were diluted 1:250 and used to probe the blots. We used alkaline-phosphatase-conjugated goat anti-human IgM  $\mu$ -chain and anti-human IgG  $\gamma$ -chain (Thermo Fisher Scientific, <https://www.thermofisher.com>) as secondary antibodies. Arrows indicate *E. chaffeensis*-specific P28 antigens (encoded by a p28 multigene family). Arrowhead shows *A. phagocytophilum*-specific P44 antigen (encoded by a p44 multigene family).

Appendix Figure 2). In this case, the IgM titer increased in the convalescent-phase serum on day 58 but decreased on day 115. However, the IgG titer increased on days 58 and 115 after onset of illness. In addition, we detected antibodies against *Anaplasma phagocytophilum* by IFA and *A. phagocytophilum*-specific P44 surface antigen by Western blot. We detected only IgG antibodies against *A. phagocytophilum* in all 3 serum samples, suggesting a past infection with *A. phagocytophilum*.

The 3 patients lived on the Kii peninsula of Japan (Appendix Figure 3), which is known to be a JSF-endemic area, especially in Wakayama and Mie Prefectures (3,4). In addition, anaplasmosis exists in those areas (5). Previously, we revealed the presence of ticks infected with *A. phagocytophilum* and *Ehrlichia* sp. that could potentially infect humans in Mie prefecture (6,7). In particular, members of the *Ehrlichia* sp. genotype 2 group, including *Ehrlichia* sp. MieHI92 and MieHI94, were considered candidate organisms that might cause human ehrlichiosis in Japan (6).

In conclusion, we provide serologic evidence of autochthonous cases of human ehrlichiosis in Japan. We recommend that ehrlichiosis should be considered as a clinical cause of febrile illness in this country.

This work was supported by a Grant-in-Aid for Scientific Research (nos. 17K08835 and 20K07499) from the Japan Society for the Promotion of Science (to N.O.). The research was partially supported by the Research Program on Emerging and Re-emerging Infectious Diseases from the Japan Agency for Medical Research and Development (AMED no. 18fk0108068h0201 to N.O.).

## About the Author

Dr. Su is an assistant professor in the Graduate Program in Pharmaceutical and Nutritional Sciences, Graduate School of Integrated Pharmaceutical and Nutritional Sciences, University of Shizuoka, Japan. Her research interests include the molecular microbiology, ecology, and epidemiology of zoonotic parasites, especially those causing tickborne infectious diseases.

## References

1. Ismail N, McBride JW. Tick-borne emerging infections: ehrlichiosis and anaplasmosis. *Clin Lab Med*. 2017;37:317–40. <https://doi.org/10.1016/j.cll.2017.01.006>
2. Moriyama Y, Kutsuna S, Toda Y, Kawabata H, Sato K, Ohmagari N. Three cases diagnosed not Lyme disease but “tick-associated rash illness (TARI)” in Japan. *J Infect Chemother*. 2021;27:650–2. <https://doi.org/10.1016/j.jiac.2020.11.026>
3. Ministry of Health, Labour and Welfare, Japan. National Institute of Infectious Diseases. Japanese spotted fever 1999–2019. *Infectious Agents Surveillance Report*, vol. 41, 2020 Aug [cited 2021 Dec 21]. <https://www.niid.go.jp/niid/en/iasr-vol41-e/865-iasr/10416-486te.html>
4. Gaowa, Ohashi N, Aochi M, Wurutu D, Wu, Yoshikawa Y, et al. Rickettsiae in ticks, Japan, 2007–2011. *Emerg Infect Dis*. 2013;19:338–40. <https://doi.org/10.3201/eid1902.120856>
5. Su H, Ito K, Kawarasaki Y, Morita H, Nose H, Ikeda K, et al. Insight of diagnostic performance using B-cell epitope antigens derived from triple P44-related proteins of *Anaplasma phagocytophilum*. *Diagn Microbiol Infect Dis*. 2019;95:125–30. <https://doi.org/10.1016/j.diagmicrobio.2019.05.008>
6. Su H, Onoda E, Tai H, Fujita H, Sakabe S, Azuma K, et al. Diversity unearthed by the estimated molecular phylogeny and ecologically quantitative characteristics of uncultured *Ehrlichia* bacteria in *Haemaphysalis* ticks, Japan. *Sci Rep*. 2021;11:687. <https://doi.org/10.1038/s41598-020-80690-7>
7. Su H, Sato A, Onoda E, Fujita H, Sakabe S, Akachi S, et al. Molecular detection and characterization of *p44/msp2* multigene family of *Anaplasma phagocytophilum* from *Haemaphysalis longicornis* in Mie Prefecture, Japan. *Jpn J Infect Dis*. 2019;72:199–202. <https://doi.org/10.7883/yoken.JJID.2018.485>

Address for correspondence: Norio Ohashi, Laboratory of Microbiology, Department of Food Science and Biotechnology, School of Food and Nutritional Sciences, Graduate School of Integrated Pharmaceutical and Nutritional Sciences, University of Shizuoka, 52-1 Yada, Suruga-ku, Shizuoka 422-8526, Japan; email: ohashi@u-shizuoka-ken.ac.jp

## Environmental Investigation during Legionellosis Outbreak, Montérégie, Quebec, Canada, 2021

Laura Atikessé, Nabila Kadaoui, Vincent Lavallée, Éric Levac, Marie St-Amour, François Milord

Author affiliations: Centre intégré de santé et de services sociaux de la Montérégie-Centre, Longueuil, Quebec, Canada (L. Atikessé, N. Kadaoui, V. Lavallée, É. Levac, M. St-Amour, F. Milord); Université de Sherbrooke, Longueuil (N. Kadaoui, É. Levac, M. St-Amour, F. Milord); Public Health Agency of Canada, Ottawa, Ontario, Canada (V. Lavallée)

DOI: <https://doi.org/10.3201/eid2811.220151>

# Serologic Evidence of Human Exposure to Ehrlichiosis Agents in Japan

## Appendix

### Materials and Methods

We obtained serum samples from 34 patients with fever of unknown origin from 14 prefectures in Japan during 2008–2021. The serum samples were from acute and convalescent phases. The study was approved by the Ethics Committee of the University of Shizuoka (approval no. 1-27) and Japan National Institute of Infectious Diseases (approval no. 999). The samples were retrospectively analyzed for human ehrlichiosis and anaplasmosis by immunofluorescence assay (IFA) and Western blotting by using *Ehrlichia chaffeensis*-infected THP-1 cells, *E. chaffeensis*-infected DH82 cells, *Anaplasma phagocytophilum*-infected THP-1 cells, and *A. phagocytophilum*-infected HL60 cells as antigens. The procedures for IFA and Western blotting were described previously (1,2).

### Results of Laboratory Tests for 3 Patients

Laboratory test results (reference values) for patient 1 (case 1) on day 2 after onset of high fever were: leukocytes,  $7.8 \times 10^3$  cells/ $\mu\text{L}$  ( $3.8\text{--}9.0 \times 10^3$  cells/ $\mu\text{L}$ ); thrombocytes,  $18.2 \times 10^4$  cells/ $\mu\text{L}$  ( $15.0\text{--}40.0 \times 10^4$  cells/ $\mu\text{L}$ ); lactate dehydrogenase, 281 U/L (124–222 U/L); and C-reactive protein, 5.46 mg/dL (<0.10 mg/dL). Test results (reference values) for patient 2 (case 2) on day 14 after onset of illness were: leukocytes,  $9.4 \times 10^3$  cells/ $\mu\text{L}$  ( $3.8\text{--}9.0 \times 10^3$  cells/ $\mu\text{L}$ );

thrombocytes,  $18.5 \times 10^4$  cells/ $\mu$ L ( $15.0\text{--}40.0 \times 10^4$  cells/ $\mu$ L); aspartate aminotransferase, 147 U/L (8.0–38.0 U/L); alanine aminotransferase, 239 U/L (4.0–44.0 U/L); C-reactive protein, 11.56 mg/dL (<0.10 mg/dL); bilirubin, 4.3 mg/dL (0.2–1.2 mg/dL); creatinine, 1.5 mg/dL (0.60–1.10 mg/dL). Test results (reference values) for patient 3 (case 3) on day 5 after onset of illness were: leukocytes,  $4.5 \times 10^3$  cells/ $\mu$ L ( $3.8\text{--}9.0 \times 10^3$  cells/ $\mu$ L); thrombocytes,  $12.8 \times 10^4$  cells/ $\mu$ L ( $15.0\text{--}40.0 \times 10^4$  cells/ $\mu$ L); aspartate aminotransferase, 40 U/L (8.0–38.0 U/L); alanine aminotransferase, 25 U/L (4.0–44.0 U/L); lactate dehydrogenase, 451 U/L (124–222 U/L); and C-reactive protein, 0.69 mg/dL (<0.10 mg/dL).

## Discussion

In Asia, cases of human ehrlichiosis are extremely rare. Serologic evidence of human ehrlichiosis was reported in South Korea (2 cases), and serologic evidence and PCR detection of ehrlichiosis was reported in Taiwan (2 cases) (3–5). Epidemiologic surveillance has shown that antibodies against *Ehrlichia* antigens have been detected in healthy volunteers in Thailand, Japan, and China (6–8). Thus, human ehrlichiosis is potentially present in Asia, including Japan.


## References

1. Ohashi N, Gaowa, Wuritu, Kawamori F, Wu D, Yoshikawa Y, et al. Human granulocytic anaplasmosis, Japan. *Emerg Infect Dis.* 2013;19:289–92. [PubMed](https://doi.org/10.3201/eid1902.120855)  
<https://doi.org/10.3201/eid1902.120855>
2. Gaowa, Yoshikawa Y, Ohashi N, Wu D, Kawamori F, Ikegaya A, et al. *Anaplasma phagocytophilum* antibodies in humans, Japan, 2010–2011. *Emerg Infect Dis.* 2014;20:508–9. [PubMed](https://doi.org/10.3201/eid2003.131337)  
<https://doi.org/10.3201/eid2003.131337>


3. Heo EJ, Park JH, Koo JR, Park MS, Park MY, Dumler JS, et al. Serologic and molecular detection of *Ehrlichia chaffeensis* and *Anaplasma phagocytophila* (human granulocytic ehrlichiosis agent) in Korean patients. *J Clin Microbiol.* 2002;40:3082–5. [PubMed](#)  
<https://doi.org/10.1128/JCM.40.8.3082-3085.2002>
4. Peng SH, Yang SL, Ho YN, Chen HF, Shu PY. Human case of *Ehrlichia chaffeensis* infection, Taiwan. *Emerg Infect Dis.* 2019;25:2141–3. [PubMed](#) <https://doi.org/10.3201/eid2511.190665>
5. Yen TY, Tung YJ, Wang HC, Tsai KH. Detection of *Ehrlichia chaffeensis* in a febrile patient in Kinmen, an offshore island of Taiwan. *J Formos Med Assoc.* 2020;119:1329–30. [PubMed](#)  
<https://doi.org/10.1016/j.jfma.2019.11.019>
6. Heppner DG, Wongsrichanalai C, Walsh DS, McDaniel P, Eamsila C, Hanson B, et al. Human ehrlichiosis in Thailand. *Lancet.* 1997;350:785–6. [PubMed](#) [https://doi.org/10.1016/S0140-6736\(05\)62571-8](https://doi.org/10.1016/S0140-6736(05)62571-8)
7. Kawahara M, Ito T, Suto C, Shibata S, Rikihisa Y, Hata K, et al. Comparison of *Ehrlichia muris* strains isolated from wild mice and ticks and serologic survey of humans and animals with *E. muris* as antigen. *J Clin Microbiol.* 1999;37:1123–9. [PubMed](#) <https://doi.org/10.1128/JCM.37.4.1123-1129.1999>
8. Zhang L, Shan A, Mathew B, Yin J, Fu X, Zhang J, et al. Rickettsial seroepidemiology among farm workers, Tianjin, People’s Republic of China. *Emerg Infect Dis.* 2008;14:938–40. [PubMed](#)  
<https://doi.org/10.3201/eid1406.071502>



## Case 2




**Appendix Figure 1.** Western blots using acute-and convalescent-phase serum samples from a febrile patient (case 2) in Mie prefecture in study showing serologic evidence of human exposure to ehrlichiosis agents in Japan. Serum samples were collected from the patient on days 14, 32, and 60 after onset of illness. Human THP-1 and canine DH82 cells were uninfected or infected with *Ehrlichia chaffeensis* (Ech). THP-1 cells were also infected with *Anaplasma phagocytophilum* (Aph). We used uninfected THP-1 and DH82 cells as negative lysate controls. The patient's serum samples were diluted 1:250 and used to probe the blots. We used alkaline-phosphatase-conjugated goat anti-human IgM  $\mu$ -chain and anti-human IgG  $\gamma$ -chain antibodies (Thermo Fisher Scientific, <https://www.thermofisher.com>) as secondary antibodies. Arrows indicate *E. chaffeensis*-specific P28 antigens (encoded by a *p28* multigene family).



**Appendix Figure 2.** Western blots using acute-and convalescent-phase serum samples from a febrile patient (case 3) in Mie prefecture in study showing serologic evidence of human exposure to ehrlichiosis agents in Japan. Serum samples were collected from the patient on days 5, 58, and 115 after onset of illness. Human THP-1 and canine DH82 cells were uninfected or infected with *Ehrlichia chaffeensis* (Ech). THP-1 cells were also infected with *Anaplasma phagocytophilum* (Aph). We used uninfected THP-1 and DH82 cells as negative lysate controls. The patient's serum samples were diluted 1:250 and used to probe the blots. We used alkaline-phosphatase-conjugated goat anti-human IgM  $\mu$ -chain and anti-human IgG  $\gamma$ -chain antibodies (Thermo Fisher Scientific, <https://www.thermofisher.com>) as secondary antibodies. Arrows indicate *E. chaffeensis*-specific P28 antigens (encoded by a *p28* multigene family). Arrowheads show *A. phagocytophilum*-specific P44 antigens (encoded by a *p44* multigene family).





**Appendix Figure 3.** Map of Japan showing residential locations of 3 febrile patients who had seroconversion to antibody against *E. chaffeensis* antigens in study of serologic evidence of human exposure to ehrlichiosis agents. Serum samples were collected from these patients in 2015 and 2018. Kii peninsula is known to be highly endemic for Japanese spotted fever, especially Wakayama and Mie prefectures, and anaplasmosis is also present. Closed circles indicate where each patient lived at the time of serum collection.