Appendix Table 1. Seroprevalence of 5 bacterial zoonoses among farm workers, by age group, near Tianjin, People's Republic of China, May-July 2006*

	Anaplasma phagocytophilum			Ehrlichia chaffeensis			Bartonella henselae			Coxiella burnetii			Rickettsia typhi		
Age, y	М	F	Total	М	F	Total	M	F	Total	M	F	Total	M	F	Total
<15	14.3 (1/7)	0 (0/1)	12.5 (1/8)	0 (0/5)	NT	0 (0/5)	0 (0/7)	NT	0 (0/7)	0 (0/7)	NT	0 (0/7)	0 (0/7)	NT	0 (0/7)
15–19	12.5 (1/8)	11.1 (1/9)	11.8 (2/17)	0 (0/3)	0 (0/4)	0 (0/7)	0 (0/1)	0 (0/3)	0 (0/4)	0 (0/1)	0 (0/3)	0 (0/4)	0 (0/1)	0 (0/3)	0 (0/4)
20–29	15.8 (6/38)	3.5 (1/29)	10.5 (7/67)	0 (0/19)	0 (0/16)	0 (0/35)	5.6 (1/18)	20.0 (3/15)	12.1 (4/33)	0 (0/18)	0 (0/15)	0 (0/33)	5.6 (1/18)	0 (0/15)	3.0 (1/33)
30–39	5.7 (3/53)	2.3 (1/44)	4.1 (4/97)	0 (0/38)	0 (0/30)	0 (0/68)	7.9 (3/38)	12.9 (4/31)	10.1 (7/69)	2.6 (1/38)	9.7 (3/31)	5.8 (4/69)	5.3 (2/38)	6.5 (2/31)	5.8 (4/69)
40–49	10.5 (6/57)	10.0 (4/40)	10.3 (10/97)	0 (0/39)	3.2 (1/32)	1.4 (1/71)	16.7 (6/36)	6.5 (2/31)	11.9 (8/67)	16.7 (6/36)	9.7 (3/31)	13.4 (9/67)	5.6 (2/36)	3.2 (1/31)	4.5 (3/67)
50–59	12.5 (4/32)	10.0 (3/30)	11.3 (7/62)	0 (0/16)	0 (0/19)	0 (0/35)	7.1 (1/14)	5.3 (1/19)	6.1 (2/33)	0 (0/14)	5.3 (1/19)	3.0 (1/33)	0 (0/14)	5.3 (1/19)	3.0 (1/33)
<u>≥</u> 60	11.1 (1/9)	0 (0/5)	7.1 (1/14)	0 (0/4)	0 (0/3)	0 (0/7)	0 (0/4)	0 (0/3)	0 (0/7)	0 (0/4)	0 (0/3)	0 (0/7)	0 (0/4)	0 (0/3)	0 (0/7)
Total	10.8	6.3	8.8	0	1.0	0.4	9.4	9.8	9.6	5.9	6.9	6.4	4.2	3.9	4.1
	(22/204)	(10/158)	(32/362)	(0/124)	(1/104)	(1/228)	(11/118)	(10/102)	(21/220)	(7/118)	(7/102)	(14/220)	(5/118)	(4/102)	(9/220)

^{*}Values are % (no. positive/no. tested); NT, none tested.