

Human Genome Epidemiology (HuGE) Review

A Meta-Analysis of the Association of *N*-Acetyltransferase 2 Gene (*NAT2*) Variants with Breast Cancer

Heather M. Ochs-Balcom^{1,2}, Georgia Wiesner^{2,3}, and Robert C. Elston^{1,2}

¹ Department of Epidemiology and Biostatistics, Case Western Reserve University, Cleveland, OH.

² Ireland Comprehensive Cancer Center at University Hospitals of Cleveland and Case Western Reserve University, Cleveland, OH.

³ Department of Genetics, Case Western Reserve University, Cleveland, OH.

Received for publication September 15, 2006; accepted for publication January 25, 2007.

The *N*-acetyltransferase 2 gene (*NAT2*) product is an enzyme important in carcinogen metabolism via activation and detoxification pathways. Therefore, *NAT2* variants may represent underlying susceptibility to breast cancer. Because a number of studies of the association of *NAT2* with breast cancer have been published, the authors performed a meta-analysis. They extracted all relevant data to examine evidence for a main effect (i.e., the effect in a model that does not include any interactions) of *NAT2* phenotype and genotype on breast cancer risk. They summarized the evidence for modification by smoking and meat intake, sources of exposure to aromatic and heterocyclic amines, respectively, which are metabolized by *NAT2*. The authors identified seven studies that measured *NAT2* phenotype and 20 studies that deduced phenotype via genotyping. They found no evidence for heterogeneity (Cochran's *Q* statistic $p = 0.74$) and no statistically significant increased risk from *NAT2* acetylation (slow/rapid) for breast cancer (summary odds ratio = 1.02, 95% confidence interval: 0.95, 1.08). These results suggest that there is no overall association between the *NAT2* slow- or rapid-acetylation phenotype and breast cancer risk. However, some evidence suggests that smoking may modify this association.

acetyltransferases; breast neoplasms; epidemiology; genotype; meat; *NAT2*; polymorphism, genetic

Abbreviations: *NAT2*, *N*-acetyltransferase 2; PhIP, 2-amino-1-methyl-6-phenylimidazo[4,5-b]-pyridine.

Editor's note: This paper is also available on the website of the Human Genome Epidemiology Network (<http://www.cdc.gov/genomics/hugenet/>).

important role in the metabolism of both aromatic and heterocyclic amines, via *N*- and *O*-acetylation, which are pathways responsible for deactivation and activation, respectively (2).

GENE

The *N*-acetyltransferase 2 gene (*NAT2*) is located on chromosome 8p21.3-23.1 and codes for a phase II xenobiotic metabolizing enzyme (1). The *NAT2* enzyme plays an

GENE VARIANTS AND FREQUENCY

A previous *NAT2* human genome epidemiology (HuGE) review detailed gene variants and their population frequencies (3). Briefly, *NAT2* phenotypes are characterized as being slow, intermediate, or rapid acetylators, which refers to their ability to metabolize or activate xenobiotics.

Correspondence to Dr. Robert C. Elston, Department of Epidemiology and Biostatistics, Case Western Reserve University, Wolstein Research Building, Room 1300, 2103 Cornell Road, Cleveland, OH 44106-7281 (e-mail: rce@darwin.epbi.cwru.edu).

DISEASE(S) OR OTHER OUTCOMES

Previous epidemiologic work implicates *NAT2* variants in carcinogenesis, especially for bladder, colon, and breast cancer (3–5). This finding is biologically plausible owing to the ability of the *NAT2* enzyme to *N*-acetylate or *O*-acetylate carcinogens to inert or noninert compounds capable of forming DNA adducts. Cigarette smoking and intake of well-done meat involve exposure to aryl and heterocyclic amines, substrates of *NAT2*; therefore, it is plausible that underlying *NAT2* genotype may modify risk of cancer based on the ability to activate or detoxify heterocyclic and aromatic amines (6, 7).

OBJECTIVES OF THE CURRENT REVIEW

Since there have been a number of published reports on *NAT2* and breast cancer, we conducted a meta-analysis to summarize the results of the effect on breast cancer of the *NAT2* acetylation phenotype and *NAT2* genotype and also to discuss evidence for an interaction of *NAT2* with exposure to smoking, meat intake, and meat cooking method.

METHODS

Study selection and inclusion criteria

We searched for relevant papers published before August 2006 by using the Human Genome Epidemiology Network (HuGENet) and the MEDLINE database (National Library of Medicine, Bethesda, Maryland). We searched MEDLINE by using the following terms: “arylamine *N*-acetyltransferase,” “acetyltransferases,” “*NAT2*,” “genetic polymorphism,” “restriction fragment polymorphism,” “single nucleotide polymorphism,” “breast cancer,” and “breast neoplasms.” We included all articles involving case-control or nested case-control studies that investigated *NAT2* (determined by using phenotyping or genotyping methods) and breast cancer risk. We also searched the reference lists of the published studies found this way.

Data extraction

We extracted the following information from each manuscript: year of publication, country, number of cases and controls, matching, phenotyping technique, genotyping technique, alleles measured, allele and genotype frequencies, method for phenotype classification, phenotype frequencies, mean age of case and controls, menopausal status, covariates, and results by smoking or meat intake. We also extracted risk estimates for gene-environment interactions.

Classification of phenotype

Because of the inherent differences in methods for measuring *NAT2* status—measuring phenotypes by using metabolic response to a particular compound or measuring alleles directly—we did not combine these studies for analysis. Respectively, these two types of studies are herein referred to as “phenotype” or “genotype” studies. Subse-

quently, we summarized phenotype frequencies (slow and rapid status). In the published genotype studies, individuals were classified as having a slow phenotype if homozygous for any of the slow-activity alleles and as having a rapid phenotype if homozygous or heterozygous for wild-type alleles. In some cases, we reclassified intermediate acetylators as rapid when the authors provided frequencies for an intermediate category (having one slow and one rapid allele) (6).

Statistical analysis

We performed a meta-analysis separately for phenotype and genotype studies. For each analysis, we investigated among-study heterogeneity by using Cochran’s *Q* statistic, and we examined fixed- and random-effect models based on the method of DerSimonian and Laird (8) in Number Cruncher Statistical software (9). We constructed a funnel plot to examine the influence of publication bias.

RESULTS

Phenotype studies

Seven studies were identified that analyzed differences in the proportion of cases and controls having a slow-acetylator phenotype (10–16). Table 1 summarizes these studies. Six were conducted in Europe, and the samples were primarily Caucasian. The sample size ranged from 79 to 515, totaling 1,330 women. Six studies measured acetylation phenotype via administration of arylamine drugs (four sulfamethazine, two dapsone); the other used isoniazid. We pooled the numbers across these studies and calculated the prevalence of the slow-acetylator phenotype to be 56 percent overall: 52 percent in cases and 58 percent in controls.

Three of the seven studies reported a higher prevalence of the rapid-acetylator phenotype in the breast cancer cases (10, 11, 15), but only two of the studies were statistically significant (10, 15). For the meta-analysis, the test for heterogeneity was not statistically significant at $p < 0.05$ (Q statistic = 10.9, $p = 0.09$). Figure 1 shows the individual study odds as well as the summary odds ratio (0.77, 95 percent confidence interval: 0.61, 0.96) for the combined phenotype studies (where the reference group is rapid acetylators) using the fixed-effect model.

Genotype studies

Two duplication studies and two abstracts were identified that were later published as full manuscripts (17–20). These two particular full manuscripts were included in the meta-analysis. Another study, a case-cohort study (21), was brought to our attention by the reviewers of our paper. Excluding overlapping papers and including the case-cohort study, we identified 15 case-control studies, four nested case-control studies, and one case-cohort study, 20 in total (21–40). Table 2 summarizes these studies. Ten studies were population based, and eight were hospital based. Most studies were conducted among Caucasian populations. The only study that included a larger sample of African-American women was published by Millikan et al. (35). Three studies

TABLE 1. Description of phenotype-only studies included in a meta-analysis of the association of *N*-acetyltransferase 2 variants with breast cancer

First author, year (reference no.)	Geographic region	Source/type of cases	Source/type of controls	No. of cases	No. of controls	Method for phenotype determination	% with the slow phenotype	
							Cases	Controls
Bulovskaya et al., 1978 (10)	Russia	Advanced breast cancer	Healthy, but some with age-associated cardiovascular disturbances	41	38	Sulfamethazine (not given)	31.7	63.2
Cartwright et al., 1984 (11)	United Kingdom	Invasive breast cancer	Not given	93	112	Dapsone (slow: MADDS/DDS* <0.33)	45.2	58.0
Ladero et al., 1987 (13)	Spain	Histologically confirmed	Healthy or with disease not related to acetylator phenotype	81	75	Sulfamethazine (slow: ACMZ/SMZ* <0.45 in plasma or <0.77 in urine)	60.5	60.0
Philip et al., 1987 (14)	United Kingdom	Malignant breast disease	Cardiovascular disease and normal	181	337	Dapsone (slow: MADDS/DDS <0.30)	54.7	55.2
Webster et al., 1989 (16)	United Kingdom	Histologically confirmed	68 healthy volunteers, 32 with breast abnormalities	100	100	Isoniazid (slow: ACIHN/INH* <1.5)	57.0	59.0
Sardas et al., 1990 (15)	Turkey	Advanced breast cancer	Healthy volunteers	28	51	Sulfamethazine (slow: not given)	39.3	64.7
Ilett et al., 1990 (12)	Australia	Surgical resection of primary breast carcinoma	No breast disease	45	48	Sulfamethazine (slow: ACMZ/SMZ <0.6)	55.6	64.6

* MADDS/DDS, monoacetyldapsone/dapsone ratio; ACMZ/SMZ, *N*-acetylmethazine/methazine ratio; ACIHN/INH, acetylisoniazid/isoniazid ratio.




FIGURE 1. Summary plot for phenotype studies (first author, date of publication (reference no.)) and combined estimate of odds ratio and 95% confidence interval. Reference group is rapid acetylators. Studies are sorted according to odds ratio. The sizes of the symbols are proportional to the study sizes.

were conducted among Asian populations (two in China and one in Korea) (29, 33, 39). Two recruited women based on menopausal status; the Iowa Women's Health Study recruited postmenopausal women only (26), while Matheson et al. (34) included only premenopausal women. Of the 16,091 women included in the 20 studies, there were 7,479 cases and 8,612 controls.

Web table 1 shows the *NAT2* alleles typed for each study as well as the frequency of slow- and rapid-acetylation phenotypes deduced from genotyping results by each author. (This information is described in the first of two supplementary tables; both are referred to as "Web table" in the text and are posted on the website of the Human Genome Epidemiology Network (<http://www.cdc.gov/genomics/hugenet/reviews.htm>) as well as on the *Journal's* website (<http://aje.oupjournals.org/>.) In spite of the differences in nomenclature across the 20 genotype studies included in the meta-analysis, there were few actual differences in the particular alleles measured. In Caucasians, the majority of variation in slow-acetylator phenotype is due to three polymorphic sites (41, 42). Each study similarly measured the three alleles responsible for the majority of the slow-acetylator phenotype and classified women as having slow acetylators if they had two slow alleles. Therefore, carrying one copy of the *4 allele resulted in classification as a rapid acetylator. Seven studies measured additional rapid alleles (*12 or *13) (21, 26, 31–33, 38, 40), while six measured the *14 allele as an additional slow allele (23, 26, 27, 29, 35, 37). Intermediate acetylators were reclassified as rapid for the purpose of this review and, in each case the definition of intermediate was the same, as having only one rapid allele.

There is a confusion regarding differences in nomenclature for *NAT2* alleles. An excellent review (6) as well as a website (www.louisville.edu/medschool/pharmacology/NAT.html) help to clarify the allele name with the specific nucleotide changes.

We reclassified intermediate acetylators as fast acetylators for the analysis in those studies in which the authors classified women into three categories (slow, intermediate, and rapid). Following this reclassification of intermediate phenotypes, we found that, overall, 54.4 percent of cases and 54.3 percent of controls were slow acetylators. From the 16 studies that included primarily Caucasian women and the Caucasian sample provided by Millikan et al. (35), we calculated prevalence of the slow genotype in Caucasians to be 56.9 percent: 57.0 percent in cases and 56.9 percent in controls. For the sample of African-American women, prevalence of the slow phenotype was 40 percent overall: 41 percent for cases and 40 percent for controls (35). In the three Asian studies, the pooled prevalence of the slow phenotype was 20 percent: 21 percent in cases and 18 percent in controls.

Associations

Web table 2 shows that three of the 20 studies reported a significant association between *NAT2* acetylation and breast cancer. Alberg et al. (23) showed that rapid-acetylation status increased risk both in the whole sample and among postmenopausal women. Conversely, Sillanpaa et al. (37) in the Finland sample and Huang et al. (29) in the Taiwanese population reported an increased risk for slow acetylators, which may also be highest for postmenopausal women.

For the meta-analysis of the genotype studies, we found no evidence for among-study heterogeneity (Q statistic = 14.58, $p = 0.75$). Figure 2 shows the individual odds ratios, as well as the summary odds ratio (1.02) and 95 percent confidence interval (0.95, 1.08) for the main effect of *NAT2* acetylator genotype obtained by using a fixed-effect model (8).

Figure 3 is a funnel plot showing the association of *NAT2* with breast cancer by study type (phenotype or genotype) and study size. There does not appear to be an obvious publication bias toward positive or negative findings among the genotype studies. However, in the phenotype studies, there may be a publication bias toward manuscripts showing risk associated with rapid phenotypes because the plot is asymmetrical.

Smoking

In terms of a main effect of smoking, 15 of the 20 genotype studies analyzed smoking and breast cancer risk by using a variety of exposure variables: smoking status (active, former, passive smoking), cumulative exposure, duration, and intensity via number of cigarettes smoked per day (which may have been measured as many as 20 years prior to diagnosis). Of these 15 studies, nine found no significant association between smoking and breast cancer risk. Hunter et al. (30) reported increased risk for women who smoked 15 or more cigarettes per day (10 years prior to diagnosis)

and women with 20–30 pack-years of exposure, a finding replicated by Millikan et al. (35), who reported increased risk of breast cancer with more than a 20-year duration of active smoking. Morabia et al. (36) reported increased risk for current and former smokers. Egan et al. (28) found increased risk for ever smokers and increasing pack-years; however, the association with increasing pack-years was significant for postmenopausal women only. Lissowska et al. (40) reported increased risk for women less than age 45 years in terms of smoking status, increasing number of cigarettes per day and duration, and age at which smoking started. None of the studies showed a decreased risk of breast cancer with active or passive smoking.

Interactions

Smoking. Of the 15 genotype studies that measured smoking, 13 investigated the hypothesized interaction of smoking and *NAT2* acetylator genotype. Some studies further stratified by menopausal status.

Among the studies that examined the *NAT2* × smoking interaction according to menopausal status, Ambrosone et al. (24) was the first to show increased risk of breast cancer for slow-acetylator, postmenopausal women who were current or former smokers (using cigarettes/day 2 years ago, cigarettes/day 20 years ago, and lifetime pack-years). These findings were later replicated by Egan et al. (28) (pack-years) and Alberg et al. (23) (active smoking), who reported the strongest association among postmenopausal women. Sillanpaa et al. (37), van der Hel et al. (38), and Matheson et al. (34) reported significant increased risk for smokers who were slow acetylators.

Krajinovic et al. (32), Delfino et al. (27), Hunter et al. (30), and Millikan et al. (35) showed no significant evidence for modification of *NAT2* and breast cancer by smoking. Krajinovic et al. reported a nonsignificant increase in risk among rapid acetylators who were current or former smokers, which became significant in the case-only analysis. Delfino et al. reported no trend in risk for rapid or slow acetylators, while Hunter et al.'s weak, nonsignificant trend was among slow acetylators. Millikan et al. reported an increased, yet nonsignificant trend in risk for duration of smoking among postmenopausal rapid acetylators. Interestingly, Chang-Claude et al. (25) reported increased risk of breast cancer in rapid acetylators exposed to passive smoke, a finding later replicated by Morabia et al. (36).

Two of the studies conducted a case-only analysis. Results from Ambrosone et al. (24) suggest that, among cases, the odds of being a slow acetylator were greatest for women who smoked more than 20 cigarettes per day (20 years ago), smoked for more than 20 pack-years, or initiated smoking at an earlier age (≤ 16 years). Conversely, Krajinovic et al. (32) reported increased odds for being a rapid acetylator for ever smokers.

Meat intake. There is much published literature on the association of breast cancer with red meat intake and meat doneness (43). Five of the studies included in this review investigated potential modification of the association of *NAT2* with breast cancer according to charred or well-done meat intake. The only study that found evidence for

TABLE 2. Description of genotype studies included in a meta-analysis of the association of *N*-acetyltransferase 2 variants with breast cancer

First author, year (reference no.)	Geographic region	Study type	Source of cases	Source of controls	No. of cases	No. of controls	Matching
Agundez et al., 1995 (22)	Spain	Case-control	Not given	Healthy women from the same region	160	132	None
Ambrosone et al., 1996 (24)	New York	Case-control	Erie and Niagara Counties, New York	Randomly selected using the DMV* and HCFA*	304	327	Age and county
Hunter et al., 1997 (30)	United States	Nested case-control (Nurses' Health Study)	From cohort	Cancer free from cohort	465	466	Year of birth, menopausal status, month and time of blood draw, fasting status at blood draw, hormone use
Millikan et al., 1998 (35)	North Carolina	Case-control	Primary, invasive, North Carolina Central Cancer Registry	Randomly selected using the DMV and HCFA	488	472	None
Huang et al., 1999 (29)	Taiwan, Republic of China	Case-control	One hospital in Taiwan	Hospital based, randomly selected	139	133	None
Delfino et al., 2000 (27)	California	Case-control	Breast centers in Orange County, California	Subjects with benign masses recruited from the same breast centers	113	107	None
Morabia et al., 2000 (36)	Geneva, Switzerland	Case-control	Surviving cases who remained residents of Geneva (from an earlier study)	Surviving controls who remained residents of Geneva (from an earlier study)	160	162	Age (within 10 years)
Deitz et al., 2000 (26)	Iowa	Nested case-control (Iowa Women's Health Study)	Postmenopausal, from cohort	Postmenopausal and cancer free from cohort	174	387	None
Krajcinovic et al., 2001 (32)	Montreal, Canada	Case-control	Three hospitals in Montreal	Randomly selected from Montreal and a hospital DNA bank	149	203	Neighborhood
Wu et al., 2002 (39)	Taiwan, Republic of China	Case-control	One hospital in Taiwan	Cancer free from the same area	60	60	Age, smoking, family history (first-degree relative)

Table continues

interaction between meat doneness and *NAT2* status was that of Deitz et al. (26), who reported a significant increasing trend in risk among rapid acetylators with increasing meat doneness score.

Ambrosone et al. (44) reported no significant association between meat consumption and breast cancer, nor did they find evidence of a significant interaction between meat intake and *NAT2* acetylation status. Similarly, Gertig et al. (45) found no association of red meat intake or meat cooking method with breast cancer risk in the Nurses' Health Study, and no significant interaction with *NAT2*. Delfino et al. (46) also found no evidence of increased risk for intake of well-cooked meat nor an interaction with *NAT2*. The last of the studies, by Krajcinovic et al. (32), did not report a significant elevated risk for rapid acetylators who consumed well-done meat. In each of these studies, the trends in risk according to increasing meat intake or charred meat intake

were not strong enough to suggest whether *NAT2* slow or rapid acetylators are at increased or decreased risk.

DISCUSSION

Main findings

The results of this meta-analysis show no evidence for an overall effect of *NAT2* acetylation capacity on the risk of breast cancer. There is suggestive evidence that acetylation may underlie a susceptibility to breast cancer if there is also exposure to tobacco smoke, but there is little evidence for effect modification by intake of well-done meat. Future studies evaluating gene variants in entire pathways may enable better evaluation of susceptibility to breast cancer in the presence of other factors.

TABLE 2. Continued

First author, year (reference no.)	Geographic region	Study type	Source of cases	Source of controls	No. of cases	No. of controls	Matching
Matheson et al., 2002 (34)	Melbourne, Australia	Case-control	Pre-menopausal, recruited from Victoria and New South Wales cancer registries	Pre-menopausal, randomly selected from electoral rolls	157	157	Age range
Chang-Claude et al., 2002 (25)	Southern Germany	Case-control	All hospitals in two regions	Randomly selected from the same regions	422	887	Age (within 5 years), study region
Egan et al., 2003 (28)	Massachusetts, New Hampshire, and Wisconsin	Case-control	State registries	Randomly selected using the DMV and HCFA	791	797	None
Lee et al., 2003 (33)	Seoul, Korea	Case-control	Three hospitals in Seoul	Hospital based, cancer free	251	288	None
van der Hel et al., 2003 (38)	The Netherlands	Nested case-control	Netherlands Cancer Registry and regional registries	Randomly selected from cohort	229	264	Age (within 5 years), menopausal status, residence
Kocabas et al., 2004 (31)	Turkey (central and western)	Case-control	Three hospitals in Ankara	University staff, students, and others from hospital, no family history of cancer	84	103	Age (within 10 years), time of blood draw, menopausal status
Alberg et al., 2004 (23)	Maryland	Nested case-control (Clue II cohort)	Washington County cancer registry	Cancer free from cohort	110	113	Age (within 1 year), race, menopausal status, day of menstrual cycle (premenopausal only), date of blood collection
Sillanpaa et al., 2005 (37)	Finland	Case-control	One hospital in Finland	Recruited from the National Population Register	478	479	None
van der Hel et al., 2005 (21)	The Netherlands	Case-cohort	Regional and the Netherlands cancer registry	Population-based screening cohort	845	875	None
Lissowska et al., 2006 (40)	Warsaw and Łódź, Poland	Case-control	Five hospitals and the Cancer Registry	Randomly selected residents of the same cities	1,900	2,200	Age (within 5 years), city

* DMV, Department of Motor Vehicles; HCFA, Health Care Financing Administration.

There are a number of challenges in summarizing studies of *NAT2* variants; there are not only differences in single nucleotide polymorphism selection and technology available at the time of the study but also a reliance on phenotypic characterization of *NAT2* variants. Any misclassification of individuals as slow, intermediate, or rapid acetylators would more likely result in bias toward the null and therefore may partly explain the lack of an overall significant association between *NAT2* acetylation status and breast cancer.

Following construction of a funnel plot (figure 3) of both phenotype and genotype studies, we conclude that there is some degree of publication bias in the phenotype studies. These studies were more likely to be published if they found a protective effect for slow acetylators. The distribution of odds ratio estimates in the genotype studies suggests that, in these publications, bias is less likely to have occurred.

Our original goal was to also conduct a quantitative review of the potential factors implicated in modifying the

association of *NAT2* with breast cancer, specifically smoking and meat intake. However, once we examined the studies and discovered the many different exposure assessments used to classify individuals, we concluded that the difficulty associated with summarizing the effect of different exposure variables would be better overcome by using a pooled design in which primary data are obtained from authors of published manuscripts. At the same time, we found that a pooled analysis, in which the exposure measurements can be better summarized and a more comprehensive analysis performed, is already under way.

Biology

Approximately 26 *NAT2* variants are known to exist in humans, some silent and some functional variants responsible for changes in enzymatic activity. An excellent review




FIGURE 2. Forest plot of genotype studies (first author, date of publication (reference no.)) and combined odds ratio and 95% confidence interval. Reference group is rapid acetylators: those with at least one rapid *N*-acetyltransferase 2 allele. Studies are sorted according to odds ratio. The sizes of the symbols are proportional to the study sizes.

and explanation of genotype-phenotype correlation was published by Hein et al. (6). This paper helps to clarify the confusion regarding *NAT2* allelic nomenclature as well as the functional significance and classification of the different alleles.

For the phenotype studies, our meta-analysis suggests evidence of decreased breast cancer risk for women with a slow phenotype (odds ratio = 0.77, 95 percent confidence interval: 0.62, 0.96) or, conversely, increased breast cancer risk for women with a rapid phenotype (odds ratio = 1.29, 95 percent confidence interval: 1.04, 1.62) when slow is the reference group. The published reports also suggest increased risk for rapid acetylators; however, the majority of these studies were unable to report a significant association. Evans (47) combined the results of Bulovskaya et al. (10) and Cartwright (11) and found an odds ratio of 2.05, suggesting an increased risk for carriers of the rapid allele. These studies were of limited size, however, which may have increased sampling error. In addition, misclassification may also be important because different xenobiotics and methods were used to assess *NAT2* activity. The *NAT2* phenotype may be influenced by disease status; therefore, the results of the genotype studies may be more reliable because genotype is not affected by disease status.




FIGURE 3. Funnel plot of the association of the *N*-acetyltransferase 2 (*NAT2*) genotype with breast cancer, by study size. Odds ratios for the main effect of *NAT2* are shown. Reference group is rapid acetylators.

The genotype studies purport a modifying effect of smoking in the association of *NAT2* with breast cancer, especially in postmenopausal women. Postmenopausal women are more likely to have a greater duration of smoking simply because of their age, which may explain these findings. These studies therefore suggest a role for *NAT2* in breast cancer carcinogenesis, especially for women who also smoke. This hypothesis can be supported by evidence from Pfau et al. (48), who showed that slow acetylators have a higher quantity of DNA adducts than rapid acetylators. A recent review and meta-analysis of the potential modifying effect of smoking suggests that, among slow acetylators, smoking may increase breast cancer risk, an effect that is strongest among postmenopausal women (odds ratio = 2.4, 95 percent confidence interval: 1.7, 3.3) (49), which agrees with our assessment of the studies included in our review.

The increased risk of breast cancer for women exposed to passive smoke who are also rapid acetylators is very interesting (25, 36) in that passive smoke contains 2-amino- α -carboline, a heterocyclic amine (50). If exposed to passive or sidestream smoke, rapid acetylators are at increased risk since they are more likely to *O*-acetylate the *N*-hydroxy heterocyclic amines that have been processed in the liver by *CYP1A2* and subsequently form compounds capable of creating DNA adducts (50). Further studies are needed to confirm this association.

Biologically, slow acetylators who have lower concentrations of active *NAT2* enzyme and are exposed to aromatic amines are more likely to undergo metabolism via the hydroxylation pathway. This process may then lead to electrophilic intermediates capable of binding DNA and initiating carcinogenesis, as compared with the process in rapid acetylators, whose aromatic amine exposure is metabolized via *N*-acetylation to become an inactive metabolite (51). Firozi et al. (52) reported a higher frequency of smoking-related

adducts in breast cancer cases with the slow-acetylation phenotype, which suggests that the hypothesized increased risk of breast cancer and smoking is especially important for slow acetylators.

The varying action of *NAT2* under different exposure scenarios may partially explain the null findings for the main effect of *NAT2* and breast cancer. While smoking involves exposure to both aromatic amines and heterocyclic amines, *N*- and *O*-acetylation may be taking place. Thus, the competing or heterogeneous nature of the pathway depending on exposure may attenuate the association. In spite of this heterogeneity, many of the studies found evidence of increased risk for slow-acetylating women who also smoke.

Another exposure that has been investigated as a potential modifier of the *NAT2* and breast cancer association is meat intake. Meat intake, especially intake of charred meat, is associated with exposure to heterocyclic amines because high meat intake results in increased exposure to 2-amino-1-methyl-6-phenylimidazo[4,5-b]pyridine (PhIP) (53). PhIP is the most abundant heterocyclic amine in cooked meat (53). Women with the rapid-acetylation phenotype and high intake of well-done or charred meat may especially be at increased breast cancer risk, since *O*-acetylation by *NAT2* predominates, resulting in reactive metabolites capable of forming DNA adducts and mutations. Zhu et al. (54) detected PhIP-DNA adducts in breast tissue and reported that rapid acetylators, compared with slow acetylators, had a higher level of PhIP-DNA adducts, conferring increased risk of breast cancer in rapid acetylators who have high exposure to PhIP.

Regarding *NAT2*, the molecular explanation for why particular exposures more readily result in *N*- or *O*-acetylation may be an important cause of the lack of significant main effect of the *NAT2* genotype and breast cancer risk. The results of the meta-analysis of the genotype studies are therefore not unexpected. Aryl and heterocyclic amines differ in how they are metabolized, namely, the propensity toward *N*- or *O*-acetylation pathways for these respective exposures. Without specific knowledge of which pathway may predominate and whether other genes are involved, it is difficult to discern epidemiologically the exact estimate of risk conferred via one particular gene. Future studies should also measure and analyze the interaction of *NAT2* and other genes, such as *CYP1A2*, in the carcinogen metabolism pathway since knowledge of underlying variation in both genes may more fully characterize an individual's carcinogen metabolism phenotype (55).

Future studies that investigate smoking and meat intake should attempt to collect and use standardized exposure measures. Doing so would greatly help summarize the results of published studies.

ACKNOWLEDGMENTS

This work was supported in part by US Public Health Service National Institutes of Health training grant R25 CA094186 (H. O.-B.), research grants RR03655 and

GM28356 (R. E.), and Cancer Center support grant P30CAD43703 (R. E.).

The authors thank Dr. Christine Ambrosone from the Roswell Park Cancer Institute for her constructive comments regarding the manuscript.

Conflict of interest: none declared.

REFERENCES

1. Smith CA, Smith G, Wolf CR. Genetic polymorphisms in xenobiotic metabolism. *Eur J Cancer* 1994;30A:1921-35.
2. Hein DW. Acetylator genotype and arylamine-induced carcinogenesis. *Biochim Biophys Acta* 1988;948:37-66.
3. Brockton N, Little J, Sharp L, et al. N-acetyltransferase polymorphisms and colorectal cancer: a HuGE review. *Am J Epidemiol* 2000;151:846-61.
4. Marcus PM, Hayes RB, Vineis P, et al. Cigarette smoking, N-acetyltransferase 2 acetylation status, and bladder cancer risk: a case-series meta-analysis of a gene-environment interaction. *Cancer Epidemiol Biomarkers Prev* 2000;9:461-7.
5. Marcus PM, Vineis P, Rothman N. *NAT2* slow acetylation and bladder cancer risk: a meta-analysis of 22 case-control studies conducted in the general population. *Pharmacogenetics* 2000;10:115-22.
6. Hein DW, Doll MA, Fretland AJ, et al. Molecular genetics and epidemiology of the *NAT1* and *NAT2* acetylation polymorphisms. *Cancer Epidemiol Biomarkers Prev* 2000;9:29-42.
7. Weber W. The acetylator genes and drug response. New York, NY: Oxford University Press, 1987.
8. DerSimonian R, Laird N. Meta-analysis in clinical trials. *Control Clin Trials* 1986;7:177-88.
9. Number Cruncher Statistical Systems. Kaysville, UT. (www.ncss.com).
10. Bulovskaya LN, Krupkin RG, Bochina TA, et al. Acetylator phenotype in patients with breast cancer. *Oncology* 1978;35:185-8.
11. Cartwright RA. Epidemiological studies on N-acetylation and C-center ring oxidation in neoplasia. Banbury report 16: Genetic variability in responses to chemical exposure. New York, NY: Cold Spring Harbor Laboratory, 1984.
12. Ilett KF, Detchon P, Ingram DM, et al. Acetylation phenotype is not associated with breast cancer. *Cancer Res* 1990;50:6649-51.
13. Ladero JM, Fernandez MJ, Palmeiro R, et al. Hepatic acetylator polymorphism in breast cancer patients. *Oncology* 1987;44:341-4.
14. Philip PA, Rogers HJ, Millis RR, et al. Acetylator status and its relationship to breast cancer and other diseases of the breast. *Eur J Cancer Clin Oncol* 1987;23:1701-6.
15. Sardas S, Cok I, Sardas OS, et al. Polymorphic N-acetylation capacity in breast cancer patients. *Int J Cancer* 1990;46:1138-9.
16. Webster DJ, Flook D, Jenkins J, et al. Drug acetylation in breast cancer. *Br J Cancer* 1989;60:236-7.
17. Deitz AC, Zheng W, Leff MA, et al. N-Acetyltransferase-2 (*NAT2*) acetylation polymorphism, well-done meat intake and breast cancer risk among post-menopausal women. (Abstract). *Proc Am Assoc Cancer Res* 1999;40:148.
18. Ambrosone CB, Freudenheim JL, Marshall JR, et al. The association of polymorphic N-acetyltransferase (*NAT2*) with breast cancer risk. *Ann N Y Acad Sci* 1995;768:250-2.
19. Chern HD, Huang CS, Wang CY, et al. Association of N-acetyltransferase polymorphism with the susceptibility of

- breast cancer in Taiwan. (Abstract). *Proc Am Assoc Cancer Res* 1997;38:214.
20. Lilla C, Risch A, Kropp S, et al. SULT1A1 genotype, active and passive smoking, and breast cancer risk by age 50 years in a German case-control study. *Breast Cancer Res* 2005;7:R229–37.
 21. van der Hel OL, Bueno-de-Mesquita HB, van Gils CH, et al. Cumulative genetic defects in carcinogen metabolism may increase breast cancer risk (The Netherlands). *Cancer Causes Control* 2005;16:675–81.
 22. Agundez JA, Ladero JM, Olivera M, et al. Genetic analysis of the arylamine N-acetyltransferase polymorphism in breast cancer patients. *Oncology* 1995;52:7–11.
 23. Alberg AJ, Daudt A, Huang HY, et al. N-acetyltransferase 2 (NAT2) genotypes, cigarette smoking, and the risk of breast cancer. *Cancer Detect Prev* 2004;28:187–93.
 24. Ambrosone CB, Freudenheim JL, Graham S, et al. Cigarette smoking, N-acetyltransferase 2 genetic polymorphisms, and breast cancer risk. *JAMA* 1996;276:1494–501.
 25. Chang-Claude J, Kropp S, Jager B, et al. Differential effect of NAT2 on the association between active and passive smoke exposure and breast cancer risk. *Cancer Epidemiol Biomarkers Prev* 2002;11:698–704.
 26. Deitz AC, Zheng W, Leff MA, et al. N-Acetyltransferase-2 genetic polymorphism, well-done meat intake, and breast cancer risk among postmenopausal women. *Cancer Epidemiol Biomarkers Prev* 2000;9:905–10.
 27. Delfino RJ, Smith C, West JG, et al. Breast cancer, passive and active cigarette smoking and N-acetyltransferase 2 genotype. *Pharmacogenetics* 2000;10:461–9.
 28. Egan KM, Newcomb PA, Titus-Ernstoff L, et al. Association of NAT2 and smoking in relation to breast cancer incidence in a population-based case-control study (United States). *Cancer Causes Control* 2003;14:43–51.
 29. Huang CS, Chern HD, Shen CY, et al. Association between N-acetyltransferase 2 (NAT2) genetic polymorphism and development of breast cancer in post-menopausal Chinese women in Taiwan, an area of great increase in breast cancer incidence. *Int J Cancer* 1999;82:175–9.
 30. Hunter DJ, Hankinson SE, Hough H, et al. A prospective study of NAT2 acetylation genotype, cigarette smoking, and risk of breast cancer. *Carcinogenesis* 1997;18:2127–32.
 31. Kocabas NA, Sardas S, Cholerton S, et al. N-acetyltransferase (NAT2) polymorphism and breast cancer susceptibility: a lack of association in a case-control study of Turkish population. *Int J Toxicol* 2004;23:25–31.
 32. Krajinovic M, Ghadirian P, Richer C, et al. Genetic susceptibility to breast cancer in French-Canadians: role of carcinogen-metabolizing enzymes and gene-environment interactions. *Int J Cancer* 2001;92:220–5.
 33. Lee KM, Park SK, Kim SU, et al. N-acetyltransferase (NAT1, NAT2) and glutathione S-transferase (GSTM1, GSTT1) polymorphisms in breast cancer. *Cancer Lett* 2003;196:179–86.
 34. Matheson MC, Stevenson T, Akbarzadeh S, et al. GSTT1 null genotype increases risk of premenopausal breast cancer. *Cancer Lett* 2002;181:73–9.
 35. Millikan RC, Pittman GS, Newman B, et al. Cigarette smoking, N-acetyltransferases 1 and 2, and breast cancer risk. *Cancer Epidemiol Biomarkers Prev* 1998;7:371–8.
 36. Morabia A, Bernstein MS, Bouchardy I, et al. Breast cancer and active and passive smoking: the role of the N-acetyltransferase 2 genotype. *Am J Epidemiol* 2000;152:226–32.
 37. Sillanpaa P, Hirvonen A, Kataja V, et al. NAT2 slow acetylator genotype as an important modifier of breast cancer risk. *Int J Cancer* 2005;114:579–84.
 38. van der Hel OL, Peeters PH, Hein DW, et al. NAT2 slow acetylation and GSTM1 null genotypes may increase postmenopausal breast cancer risk in long-term smoking women. *Pharmacogenetics* 2003;13:399–407.
 39. Wu FY, Lee YJ, Chen DR, et al. Association of DNA-protein crosslinks and breast cancer. *Mutat Res* 2002;501:69–78.
 40. Lissowska J, Brinton LA, Zatonski W, et al. Tobacco smoking, NAT2 acetylation genotype and breast cancer risk. *Int J Cancer* 2006;119:1961–9.
 41. Bell DA, Taylor JA, Butler MA, et al. Genotype/phenotype discordance for human arylamine N-acetyltransferase (NAT2) reveals a new slow-acetylator allele common in African-Americans. *Carcinogenesis* 1993;14:1689–92.
 42. Lin HJ, Han CY, Lin BK, et al. Slow acetylator mutations in the human polymorphic N-acetyltransferase gene in 786 Asians, blacks, Hispanics, and whites: application to metabolic epidemiology. *Am J Hum Genet* 1993;52:827–34.
 43. Boyd NF, Stone J, Vogt KN, et al. Dietary fat and breast cancer risk revisited: a meta-analysis of the published literature. *Br J Cancer* 2003;89:1672–85.
 44. Ambrosone CB, Freudenheim JL, Sinha R, et al. Breast cancer risk, meat consumption and N-acetyltransferase (NAT2) genetic polymorphisms. *Int J Cancer* 1998;75:825–30.
 45. Gertig DM, Hankinson SE, Hough H, et al. N-acetyltransferase 2 genotypes, meat intake and breast cancer risk. *Int J Cancer* 1999;80:13–17.
 46. Delfino RJ, Sinha R, Smith C, et al. Breast cancer, heterocyclic aromatic amines from meat and N-acetyltransferase 2 genotype. *Carcinogenesis* 2000;21:607–15.
 47. Evans D. Acetylation. In: Kalow W, Goedde H, Agarwal D, eds. *Ethnic differences in reactions to drugs and xenobiotics*. New York, NY: Alan R Liss, Inc, 1986.
 48. Pfau W, Stone EM, Brockstedt U, et al. DNA adducts in human breast tissue: association with N-acetyltransferase-2 (NAT2) and NAT1 genotypes. *Cancer Epidemiol Biomarkers Prev* 1998;7:1019–25.
 49. Terry PD, Goodman M. Is the association between cigarette smoking and breast cancer modified by genotype? A review of epidemiologic studies and meta-analysis. *Cancer Epidemiol Biomarkers Prev* 2006;15:602–11.
 50. King RS, Teitel CH, Kadlubar FF. In vitro bioactivation of N-hydroxy-2-amino-alpha-carboline. *Carcinogenesis* 2000;21:1347–54.
 51. Ambrosone CB. Impact of genetics on the relationship between smoking and breast cancer risk. *J Women's Cancer* 2001;3:17–22.
 52. Firozi PF, Bondy ML, Sahin AA, et al. Aromatic DNA adducts and polymorphisms of CYP1A1, NAT2, and GSTM1 in breast cancer. *Carcinogenesis* 2002;23:301–6.
 53. Sinha R, Gustafson DR, Kulldorff M, et al. 2-amino-1-methyl-6-phenylimidazo[4,5-b]pyridine, a carcinogen in high-temperature-cooked meat, and breast cancer risk. *J Natl Cancer Inst* 2000;92:1352–4.
 54. Zhu J, Chang P, Bondy ML, et al. Detection of 2-amino-1-methyl-6-phenylimidazo[4,5-b]-pyridine-DNA adducts in normal breast tissues and risk of breast cancer. *Cancer Epidemiol Biomarkers Prev* 2003;12:830–7.
 55. Murtaugh MA, Sweeney C, Ma KN, et al. The CYP1A1 genotype may alter the association of meat consumption patterns and preparation with the risk of colorectal cancer in men and women. *J Nutr* 2005;135:179–86.