

TABLES

TABLE 1. Sample Sizes and Response Rates, Selected U.S. Sites: School Health Profiles, Principal and Lead Health Education Teacher Surveys, 2006

Site	Principal surveys		Teacher surveys	
	Sample size	Response rate (%)	Sample size	Response rate (%)
STATE SURVEYS				
Alabama	260	77	252	74
Alaska	223	76	206	71
Arizona	337	85	307	78
Arkansas	254	79	249	78
Connecticut	244	77	238	75
Delaware	68	91	68	91
Florida	299	72	298	71
Georgia	279	72	271	70
Hawaii	78	74	78	74
Idaho	213	89	201	84
Illinois	318	70	NA [†]	NA
Iowa	273	78	275	79
Kansas	253	78	243	75
Maine	277	85	255	78
Massachusetts	661	88	659	88
Michigan	346	86	323	80
Mississippi	180	76	173	73
Missouri	333	85	321	82
Montana	263	81	253	78
Nebraska	236	82	222	77
New Hampshire	182	85	165	77
New York*	367	79	356	77
North Carolina	299	71	299	71
North Dakota	166	76	168	77
Oregon	272	79	255	74
Pennsylvania	349	78	342	76
Rhode Island	88	77	88	77
South Carolina	248	78	245	77
South Dakota	206	72	202	71
Tennessee	318	80	303	77
Texas	362	71	358	71
Utah	209	82	182	71
Vermont	136	88	114	74
Virginia	280	74	277	74
Washington	269	71	NA	NA
West Virginia	196	86	196	86

TABLE 1. Sample Sizes and Response Rates, Selected U.S. Sites: School Health Profiles, Principal and Lead Health Education Teacher Surveys, 2006 (*continued*)

Site	Principal surveys		Teacher surveys	
	Sample size	Response rate (%)	Sample size	Response rate (%)
LOCAL SURVEYS				
Charlotte-Mecklenburg County	42	86	40	82
Chicago	234	77	212	70
Dallas	51	98	52	100
District of Columbia	31	72	32	74
Hillsborough County	55	83	58	88
Los Angeles	87	73	90	76
Memphis	58	85	56	82
Miami	90	93	83	86
Orange County	35	71	35	71
Philadelphia	106	73	104	71
San Diego	57	98	56	97
San Francisco	34	79	34	79

* Survey did not include schools from the New York City Department of Education.

† Data not available.

TABLE 2. Percentage of Schools That Required Health Education in Any of Grades 6–12, the Percentage That Required Students to Take Only One Course or Two or More Courses, and Among Schools That Required Health Education, the Percentage That Taught Required Health Education in a Combined Course or in Another Course and the Percentage That Required Students Who Fail a Required Health Education Course to Repeat It, Selected U.S. Sites: School Health Profiles, Principal Surveys, 2006

Site	Required health education	Required only one health education course	Required two or more health education courses	Taught required health education in a combined health education and physical education course*	Taught required health education in a course mainly about another subject*†	Required students who fail a required health education* course to repeat it
STATE SURVEYS						
Alabama	86.4	57.1	18.8	59.3	33.0	76.6
Alaska	82.2	35.5	44.4	64.3	25.5	85.1
Arizona	55.6	29.4	16.2	67.1	37.4	47.0
Arkansas	97.0	52.0	43.3	60.2	21.8	74.0
Connecticut	89.7	29.3	53.0	52.7	19.6	46.8
Delaware	95.3	37.9	50.8	81.1	9.8	52.6
Florida	70.8	42.9	19.0	65.3	33.6	68.6
Georgia	90.9	52.1	37.4	81.3	15.4	53.4
Hawaii	93.5	62.5	30.6	62.0	18.6	65.5
Idaho	98.1	55.3	42.3	50.2	8.1	56.5
Illinois	97.3	46.2	46.4	62.2	15.3	54.5
Iowa	76.1	33.0	37.4	40.8	31.7	56.0
Kansas	88.3	49.6	29.8	86.8	22.2	64.8
Maine	93.4	39.0	49.2	49.2	17.6	50.4
Massachusetts	85.6	28.9	52.0	54.0	10.2	39.2
Michigan	82.1	45.9	29.4	53.0	21.5	57.0
Mississippi	91.5	82.4	8.1	42.3	15.9	95.4
Missouri	88.4	43.3	42.9	66.0	17.7	57.3
Montana	98.6	15.5	79.3	93.9	24.5	63.5
Nebraska	93.5	39.3	50.1	77.4	25.4	57.2
New Hampshire	88.5	39.0	42.0	40.7	15.3	52.2
New York‡	99.4	39.5	59.9	32.8	10.6	58.0
North Carolina	95.2	42.5	46.4	93.7	15.0	51.7
North Dakota	92.9	26.4	63.6	64.0	24.7	58.9
Oregon	98.0	21.5	73.2	59.5	21.3	54.7
Pennsylvania	97.0	30.6	65.1	70.0	13.9	52.2
Rhode Island	97.7	16.2	74.6	81.2	7.6	41.8
South Carolina	85.5	42.0	34.7	73.4	26.4	35.0
South Dakota	71.3	23.2	41.8	80.4	21.1	38.8
Tennessee	77.6	42.7	24.1	87.4	30.0	59.0
Texas	81.0	62.0	14.6	51.4	19.6	74.2
Utah	97.7	64.9	31.8	41.0	10.5	49.3
Vermont	87.0	32.3	41.5	47.6	29.7	54.0
Virginia	91.4	12.5	68.4	98.0	10.5	48.3
Washington	92.7	43.5	43.0	68.0	27.7	57.6
West Virginia	98.9	38.6	58.6	47.6	7.2	46.3
State Median	91.5	39.4	43.0	63.1	19.6	55.4
State Range	55.6 – 99.4	12.5 – 82.4	8.1 – 79.3	32.8 – 98.0	7.2 – 37.4	35.0 – 95.4

TABLE 2. Percentage of Schools That Required Health Education in Any of Grades 6–12, the Percentage That Required Students to Take Only One Course or Two or More Courses, and Among Schools That Required Health Education, the Percentage That Taught Required Health Education in a Combined Course or in Another Course and the Percentage That Required Students Who Fail a Required Health Education Course to Repeat It, Selected U.S. Sites: School Health Profiles, Principal Surveys, 2006 (continued)

Site	Required health education	Required only one health education course	Required two or more health education courses	Taught required health education in a combined health education and physical education course*	Taught required health education in a course mainly about another subject*†	Required students who fail a required health education course to repeat it*
LOCAL SURVEYS						
Charlotte-Mecklenburg County	100.0	33.1	66.9	100.0	0.0	45.6
Chicago	76.6	33.2	20.2	88.7	45.2	35.1
Dallas	60.8	44.8	12.0	23.9	14.1	86.2
District of Columbia	97.0	29.0	54.4	96.9	14.3	66.6
Hillsborough County	56.0	44.2	7.5	49.5	42.1	64.3
Los Angeles	100.0	74.7	25.3	12.4	41.2	49.8
Memphis	85.6	55.5	25.4	95.2	3.4	57.4
Miami	59.2	46.3	10.3	77.6	36.6	77.1
Orange County	59.6	32.8	8.1	56.1	51.7	84.8
Philadelphia	88.7	50.0	35.2	92.6	19.5	35.1
San Diego [§]	100.0	0.0	0.0	59.2	90.9	0.0
San Francisco	91.3	45.5	11.5	46.6	54.4	60.2
Local Median	87.2	44.5	16.1	68.4	38.9	60.2
Local Range	56.0 – 100.0	0.0 – 74.7	0.0 – 66.9	12.4 – 100.0	0.0 – 90.9	35.1 – 86.2

* Among schools that required health education.

† For example, science, social studies, home economics, or English.

‡ Survey did not include schools from the New York City Department of Education.

§ San Diego does not have a required health education course, but requires that health education be taught in science and physical education classes.

TABLE 3. Percentage of Schools That Taught a Required Health Education Course in Each Grade,* Selected U.S. Sites: School Health Profiles, Principal Surveys, 2006

Site	Grade 6	Grade 7	Grade 8	Grade 9	Grade 10	Grade 11	Grade 12
STATE SURVEYS							
Alabama	39.0	33.0	31.5	30.1	53.4	13.1	12.6
Alaska	39.0	42.8	44.0	65.9	44.5	42.3	34.9
Arizona	25.9	29.2	28.3	22.2	16.6	8.1	8.5
Arkansas	48.1	67.7	37.7	83.3	58.4	48.3	49.3
Connecticut	61.7	62.9	62.6	60.0	52.3	37.2	32.7
Delaware	61.4	79.8	73.6	63.0	32.2	14.6	11.7
Florida	29.9	28.9	29.9	39.0	25.9	19.0	17.2
Georgia	73.0	73.6	74.1	74.7	19.8	20.3	20.2
Hawaii	47.4	77.5	33.9	46.8	61.8	22.4	15.7
Idaho	35.9	62.0	63.0	16.9	56.1	35.5	11.2
Illinois	55.0	75.4	61.8	52.7	47.9	12.3	11.4
Iowa	33.3	40.9	40.2	35.5	30.3	16.6	14.4
Kansas	33.6	45.4	41.6	64.9	9.0	1.9	1.9
Maine	72.4	77.3	72.6	49.5	43.3	8.4	7.0
Massachusetts	58.2	65.0	66.9	52.1	40.2	19.9	15.2
Michigan	30.7	47.5	41.1	51.9	24.1	13.4	14.2
Mississippi	11.8	11.4	15.0	84.5	60.6	53.9	52.9
Missouri	52.0	68.4	67.3	67.9	16.6	11.1	11.0
Montana	71.1	85.9	88.1	86.1	82.4	8.6	7.3
Nebraska	44.4	58.5	58.8	55.8	30.3	8.5	7.2
New Hampshire	50.6	61.1	64.3	44.9	40.1	11.8	11.7
New York†	53.7	73.6	57.8	32.3	70.3	42.8	35.2
North Carolina	78.5	80.2	78.5	75.3	17.1	11.1	11.8
North Dakota	66.9	74.2	67.4	52.5	32.3	9.0	10.1
Oregon	77.0	81.7	83.4	67.0	64.8	40.6	21.3
Pennsylvania	71.2	72.8	73.0	54.3	51.1	53.5	23.4
Rhode Island	79.0	79.9	82.3	80.9	74.3	77.2	73.7
South Carolina	63.7	65.6	65.6	55.3	23.4	19.0	18.4
South Dakota	34.5	54.3	50.6	28.2	10.5	5.5	4.9
Tennessee	39.7	45.2	44.3	46.0	18.9	8.5	8.5
Texas	27.2	30.0	36.2	55.3	41.4	35.9	34.1
Utah	56.4	47.1	60.3	7.9	83.9	28.6	22.2
Vermont	48.5	55.7	51.9	44.0	24.1	15.3	14.9
Virginia	57.4	61.2	56.3	68.4	63.3	2.8	2.0
Washington	50.3	56.3	62.8	56.9	39.5	21.8	20.8
West Virginia	93.2	93.3	93.3	63.6	61.6	16.8	16.8
State Median	51.3	62.5	61.1	54.8	40.8	16.7	14.7
State Range	11.8 – 93.2	11.4 – 93.3	15.0 – 93.3	7.9 – 86.1	9.0 – 83.9	1.9 – 77.2	1.9 – 73.7

TABLE 3. Percentage of Schools That Taught a Required Health Education Course in Each Grade,* Selected U.S. Sites: School Health Profiles, Principal Surveys, 2006 (continued)

Site	Grade 6	Grade 7	Grade 8	Grade 9	Grade 10	Grade 11	Grade 12
LOCAL SURVEYS							
Charlotte-Mecklenburg County	100.0	100.0	100.0	100.0	39.7	11.1	11.1
Chicago	39.5	40.7	41.8	25.8	12.2	4.5	3.7
Dallas	4.2	18.5	14.8	31.6	42.8	35.6	33.3
District of Columbia	43.6	53.0	53.0	61.0	58.8	9.2	9.2
Hillsborough County	22.3	16.9	16.9	37.5	24.6	22.1	22.1
Los Angeles	48.5	93.5	4.6	97.3	10.0	6.9	6.9
Memphis	49.3	55.7	55.7	70.2	21.1	19.2	19.2
Miami	15.7	13.7	15.7	19.4	45.0	11.5	11.7
Orange County	0.0	4.8	0.0	35.3	11.2	8.4	8.4
Philadelphia	72.9	74.6	74.6	46.2	50.4	42.3	37.3
San Diego [‡]	0.0	0.0	0.0	0.0	0.0	0.0	0.0
San Francisco	34.6	33.2	33.2	29.3	18.5	13.3	9.3
Local Median	37.1	37.0	25.1	36.4	22.9	11.3	10.2
Local Range	0.0 – 100.0	0.0 – 100.0	0.0 – 100.0	0.0 – 100.0	0.0 – 58.8	0.0 – 42.3	0.0 – 37.3

* Among schools with students in that grade.

† Survey did not include schools from the New York City Department of Education.

‡ San Diego does not have a required health education course, but requires that health education be taught in science and physical education classes.

TABLE 4. Percentage of Schools That Required Teachers to Use Specific Materials in a Required Health Education Course, Selected U.S. Sites: School Health Profiles, Lead Health Education Teacher Surveys, 2006

Site	National Health Education Standards	State, district, or school developed curriculum	Commercially developed curriculum	Commercially developed student textbook	Commercially developed teacher's guide	Health education performance assessment materials	Materials from health organizations
STATE SURVEYS							
Alabama	39.7	75.4	33.3	54.7	45.2	37.5	45.6
Alaska	30.3	60.0	19.8	34.0	27.3	28.6	14.5
Arizona	24.7	36.3	15.0	20.7	21.4	23.1	20.5
Arkansas	65.6	91.6	38.6	76.5	62.4	54.1	41.8
Connecticut	60.8	75.1	18.4	18.9	19.2	47.0	29.6
Delaware	73.4	82.0	17.2	25.1	22.5	31.6	20.3
Florida	26.2	52.1	20.4	35.1	28.7	30.2	28.1
Georgia	39.9	86.8	42.7	67.6	62.8	46.0	41.8
Hawaii	64.4	56.4	18.0	20.2	15.8	46.1	21.5
Idaho	51.9	87.7	26.8	63.8	52.6	47.5	31.0
Iowa	33.6	61.0	20.1	35.9	30.2	25.5	27.0
Kansas	32.0	67.1	20.9	42.5	36.7	31.4	28.8
Maine	30.7	76.1	21.3	28.4	23.1	70.5	20.3
Massachusetts	46.2	71.8	33.7	24.4	29.3	39.1	36.2
Michigan	34.7	67.4	22.0	34.6	29.6	27.0	24.8
Mississippi	47.3	95.9	39.6	68.4	63.8	60.7	40.7
Missouri	44.8	80.3	20.9	53.3	43.6	54.6	34.2
Montana	57.6	86.5	26.6	43.4	38.4	42.6	38.2
Nebraska	35.8	65.4	27.3	50.9	42.4	28.9	33.6
New Hampshire	56.7	69.2	17.5	30.4	30.4	32.4	27.1
New York*	71.6	86.3	16.9	34.1	24.3	45.9	35.1
North Carolina	38.6	83.1	27.0	49.9	45.8	32.4	33.8
North Dakota	46.8	65.5	31.2	50.0	46.2	28.3	31.8
Oregon	57.4	76.9	36.0	46.9	39.7	30.5	24.6
Pennsylvania	63.0	84.6	19.2	50.9	37.4	38.5	37.5
Rhode Island	76.9	85.7	19.5	30.7	30.6	44.2	39.0
South Carolina	53.5	63.2	29.1	47.4	43.3	35.0	27.5
South Dakota	33.6	48.1	22.2	35.0	33.2	23.0	24.1
Tennessee	34.0	61.0	23.6	44.4	39.8	34.2	23.2
Texas	29.0	68.0	34.5	57.3	49.4	42.7	34.0
Utah	30.5	87.4	20.0	48.4	34.5	26.9	20.6
Vermont	43.3	49.5	27.4	17.6	25.0	24.4	17.6
Virginia	34.3	78.5	24.1	57.1	48.1	39.4	36.9
West Virginia	59.3	91.6	43.9	80.7	71.0	58.1	35.9
State Median	44.1	75.3	22.9	43.9	37.1	36.3	30.3
State Range	24.7 – 76.9	36.3 – 95.9	15.0 – 43.9	17.6 – 80.7	15.8 – 71.0	23.0 – 70.5	14.5 – 45.6

TABLE 4. Percentage of Schools That Required Teachers to Use Specific Materials in a Required Health Education Course, Selected U.S. Sites: School Health Profiles, Lead Health Education Teacher Surveys, 2006 (continued)

Site	National Health Education Standards	State, district, or school developed curriculum	Commercially developed curriculum	Commercially developed student textbook	Commercially developed teacher's guide	Health education performance assessment materials	Materials from health organizations
LOCAL SURVEYS							
Charlotte-Mecklenburg County	63.7	97.1	20.4	34.7	39.3	49.6	55.1
Chicago	29.8	39.4	20.8	28.6	31.6	23.6	24.0
Dallas	34.0	55.1	19.1	36.2	37.5	56.3	46.9
District of Columbia	49.1	47.2	25.2	53.9	51.8	55.8	31.2
Hillsborough County	16.6	39.3	12.3	30.6	26.5	20.0	21.7
Los Angeles	57.7	91.4	58.4	83.9	67.3	28.7	35.1
Memphis	62.8	77.8	40.4	71.1	62.6	59.2	30.1
Miami	32.5	47.5	14.1	30.4	22.8	22.9	36.8
Orange County	30.6	49.5	19.8	27.3	27.3	24.6	19.9
Philadelphia	58.3	77.5	29.3	43.5	39.5	46.6	42.7
San Diego [†]	0.0	0.0	0.0	0.0	0.0	0.0	0.0
San Francisco	25.9	48.2	35.7	28.6	32.1	17.9	17.9
Local Median	33.3	48.9	20.6	32.7	34.8	26.7	30.7
Local Range	0.0 – 63.7	0.0 – 97.1	0.0 – 58.4	0.0 – 83.9	0.0 – 67.3	0.0 – 59.2	0.0 – 55.1

* Survey did not include schools from the New York City Department of Education.

† San Diego does not have a required health education course, but requires that health education be taught in science and physical education classes.

TABLE 5a. Percentage of All Schools That Tried to Increase Student Knowledge on a Specific Health-Related Topic in a Required Health Education Course During the 2005–2006 School Year, Selected U.S. Sites: School Health Profiles, Lead Health Education Teacher Surveys, 2006

Site	Alcohol-use or other drug-use prevention	Asthma awareness	Consumer health	CPR*	Dental and oral health	Emotional and mental health	Environmental health
STATE SURVEYS							
Alabama	77.3	56.1	73.4	68.6	67.2	74.8	65.7
Alaska	73.8	35.3	61.9	42.7	58.9	67.2	51.4
Arizona	43.9	25.3	34.8	23.6	34.1	40.9	33.5
Arkansas	96.4	66.0	91.5	72.3	85.6	93.6	79.8
Connecticut	87.3	38.8	74.6	44.6	39.3	83.6	47.3
Delaware	90.5	51.8	77.1	46.8	49.1	85.3	52.8
Florida	55.9	28.3	48.2	41.8	37.1	52.3	37.7
Georgia	89.8	60.7	81.5	66.5	70.0	85.5	68.3
Hawaii	96.1	48.0	91.8	51.5	58.4	94.8	69.1
Idaho	94.9	61.3	88.4	69.0	70.5	92.9	72.0
Iowa	74.9	39.5	66.7	45.6	53.1	72.3	54.3
Kansas	75.9	36.1	64.1	44.3	50.0	68.9	50.8
Maine	90.3	39.1	83.7	53.4	48.3	85.4	60.8
Massachusetts	82.3	36.2	67.7	42.1	46.4	79.8	46.2
Michigan	76.3	35.4	65.2	33.2	42.9	73.0	43.2
Mississippi	97.9	64.1	95.5	78.6	87.4	96.0	80.8
Missouri	86.2	55.1	81.9	59.8	66.9	82.6	67.7
Montana	96.1	50.6	83.0	66.0	67.8	89.9	70.9
Nebraska	84.8	55.2	71.7	53.4	54.1	79.3	58.4
New Hampshire	85.4	38.1	78.9	44.1	53.9	81.4	60.1
New York [†]	99.0	61.3	93.8	66.2	64.6	98.4	74.3
North Carolina	87.3	63.1	77.7	65.0	54.5	84.4	67.4
North Dakota	89.5	54.4	81.2	62.5	69.8	84.3	66.6
Oregon	93.5	47.5	81.9	55.3	60.8	90.7	64.9
Pennsylvania	94.2	58.5	82.8	71.4	59.6	89.0	57.8
Rhode Island	96.5	48.5	86.8	49.5	46.0	92.9	60.0
South Carolina	70.0	41.6	57.9	40.1	50.1	65.4	50.7
South Dakota	68.2	35.4	58.8	47.1	49.5	60.6	44.2
Tennessee	70.3	46.1	58.9	49.2	51.9	66.0	50.7
Texas	78.6	49.9	70.7	65.5	59.9	75.5	63.7
Utah	94.8	45.0	80.1	62.1	55.9	92.6	59.8
Vermont	80.8	28.8	74.2	29.0	44.1	76.4	46.2
Virginia	83.5	51.7	73.3	58.5	59.7	81.0	57.6
West Virginia	98.3	71.6	90.9	67.1	81.4	96.7	83.2
State Median	86.8	48.3	77.4	53.4	55.2	83.1	59.9
State Range	43.9 – 99.0	25.3 – 71.6	34.8 – 95.5	23.6 – 78.6	34.1 – 87.4	40.9 – 98.4	33.5 – 83.2

TABLE 5a. Percentage of All Schools That Tried to Increase Student Knowledge on a Specific Health-Related Topic in a Required Health Education Course During the 2005–2006 School Year, Selected U.S. Sites: School Health Profiles, Lead Health Education Teacher Surveys, 2006 (continued)

Site	Alcohol-use or other drug-use prevention	Asthma awareness	Consumer health	CPR*	Dental and oral health	Emotional and mental health	Environmental health
LOCAL SURVEYS							
Charlotte-Mecklenburg County	100.0	67.2	91.9	68.3	40.3	97.4	62.1
Chicago	51.2	38.1	37.0	30.4	42.2	42.1	37.3
Dallas	58.0	29.2	56.0	58.0	42.9	57.1	57.1
District of Columbia	76.4	43.1	62.4	66.1	45.1	65.8	41.2
Hillsborough County	47.4	17.8	37.9	37.9	24.0	37.5	35.5
Los Angeles	100.0	76.8	95.0	57.6	91.7	98.9	84.8
Memphis	84.6	76.7	77.3	66.3	70.8	83.0	66.1
Miami	52.5	26.2	43.7	42.5	40.0	50.0	44.3
Orange County	49.5	22.2	41.2	40.6	24.9	49.5	35.9
Philadelphia	83.9	70.2	71.4	50.7	60.8	77.7	52.7
San Diego [‡]	0.0	0.0	0.0	0.0	0.0	0.0	0.0
San Francisco	57.2	35.7	48.2	25.0	33.3	57.2	50.0
Local Median	57.6	36.9	52.1	46.6	41.3	57.2	47.2
Local Range	0.0 – 100.0	0.0 – 76.8	0.0 – 95.0	0.0 – 68.3	0.0 – 91.7	0.0 – 98.9	0.0 – 84.8

* Cardiopulmonary resuscitation.

† Survey did not include schools from the New York City Department of Education.

‡ San Diego does not have a required health education course, but requires that health education be taught in science and physical education classes.

TABLE 5b. Percentage of All Schools That Tried to Increase Student Knowledge on a Specific Health-Related Topic in a Required Health Education Course During the 2005–2006 School Year, Selected U.S. Sites: School Health Profiles, Lead Health Education Teacher Surveys, 2006

Site	First aid	Foodborne illness prevention	Growth and development	HIV* prevention	Human sexuality	Immunizations	Injury prevention and safety
STATE SURVEYS							
Alabama	71.3	62.4	72.7	76.9	68.5	61.5	73.7
Alaska	51.9	51.4	66.4	69.3	61.9	51.7	63.6
Arizona	28.6	29.2	40.3	35.6	28.9	24.1	35.0
Arkansas	85.4	70.8	91.0	92.0	76.0	73.6	92.8
Connecticut	58.6	50.2	83.1	87.6	78.8	43.3	76.8
Delaware	68.8	60.2	84.4	88.1	83.8	42.6	83.5
Florida	39.4	42.6	53.6	55.2	52.1	33.3	46.2
Georgia	75.5	63.8	86.5	86.8	78.9	63.7	83.0
Hawaii	61.6	60.2	86.1	94.7	89.9	45.2	91.5
Idaho	81.6	76.6	85.9	92.8	77.5	69.2	86.6
Iowa	52.9	53.7	66.9	71.5	65.0	45.7	61.3
Kansas	49.5	53.0	71.2	74.7	70.3	39.3	63.1
Maine	61.0	57.4	83.4	86.6	79.2	45.9	78.4
Massachusetts	49.9	48.7	76.0	77.6	73.7	39.5	68.2
Michigan	44.9	46.9	70.1	76.4	67.4	33.5	62.2
Mississippi	88.2	84.5	94.1	97.9	84.8	79.4	90.1
Missouri	73.5	67.9	80.4	83.8	74.2	58.2	78.8
Montana	77.0	69.4	89.5	92.4	79.7	61.1	87.0
Nebraska	58.9	62.8	76.4	83.9	76.7	49.9	71.7
New Hampshire	55.0	61.3	80.8	84.3	78.5	52.8	73.3
New York [†]	77.1	74.1	96.5	99.3	96.9	73.9	88.8
North Carolina	71.9	64.5	75.3	84.0	70.5	46.4	82.2
North Dakota	73.6	71.4	82.4	84.4	73.9	66.9	79.1
Oregon	62.2	69.3	90.6	94.8	89.2	62.4	82.5
Pennsylvania	77.9	59.6	86.7	92.8	84.1	54.9	84.7
Rhode Island	60.5	61.7	89.1	96.3	89.2	55.4	84.5
South Carolina	53.0	51.0	64.9	69.5	65.7	45.1	63.5
South Dakota	52.1	43.5	55.2	61.0	51.1	36.6	57.5
Tennessee	59.0	52.2	62.6	65.7	62.0	47.4	65.0
Texas	68.9	64.2	75.3	73.6	64.5	54.2	73.1
Utah	70.0	62.6	88.1	92.2	84.0	53.7	84.0
Vermont	41.8	45.5	76.8	77.4	76.5	39.7	65.5
Virginia	69.5	61.4	78.2	78.1	72.7	50.2	74.0
West Virginia	81.5	83.3	90.8	94.7	86.0	76.8	93.9
State Median	61.9	61.4	80.6	84.2	76.3	51.0	77.6
State Range	28.6 – 88.2	29.2 – 84.5	40.3 – 96.5	35.6 – 99.3	28.9 – 96.9	24.1 – 79.4	35.0 – 93.9

TABLE 5b. Percentage of All Schools That Tried to Increase Student Knowledge on a Specific Health-Related Topic in a Required Health Education Course During the 2005–2006 School Year, Selected U.S. Sites: School Health Profiles, Lead Health Education Teacher Surveys, 2006 (continued)

Site	First aid	Foodborne illness prevention	Growth and development	HIV* prevention	Human sexuality	Immunizations	Injury prevention and safety
LOCAL SURVEYS							
Charlotte-Mecklenburg County	70.4	53.8	97.2	100.0	97.3	35.8	88.2
Chicago	38.0	33.2	48.6	45.2	42.3	33.9	45.7
Dallas	53.2	53.1	57.1	57.1	55.1	51.0	57.1
District of Columbia	58.5	48.8	68.5	75.5	66.1	54.4	65.8
Hillsborough County	34.1	35.6	40.8	45.7	42.2	25.7	37.6
Los Angeles	75.1	88.9	98.7	100.0	95.0	78.7	88.7
Memphis	80.8	65.4	83.0	84.6	82.8	62.9	84.6
Miami	42.5	41.7	51.2	52.5	48.7	38.7	47.5
Orange County	38.0	30.0	46.6	48.0	46.6	24.6	46.9
Philadelphia	70.9	50.0	80.8	81.3	76.5	51.0	74.3
San Diego‡	0.0	0.0	0.0	0.0	0.0	0.0	0.0
San Francisco	25.0	32.1	46.4	57.2	57.2	25.0	42.9
Local Median	47.9	45.3	54.2	57.2	56.2	37.3	52.3
Local Range	0.0 – 80.8	0.0 – 88.9	0.0 – 98.7	0.0 – 100.0	0.0 – 97.3	0.0 – 78.7	0.0 – 88.7

* Human immunodeficiency virus.

† Survey did not include schools from the New York City Department of Education.

‡ San Diego does not have a required health education course, but requires that health education be taught in science and physical education classes.

TABLE 5c. Percentage of All Schools That Tried to Increase Student Knowledge on a Specific Health-Related Topic in a Required Health Education Course During the 2005–2006 School Year, Selected U.S. Sites: School Health Profiles, Lead Health Education Teacher Surveys, 2006

Site	Nutrition and dietary behavior	Physical activity and fitness	Pregnancy prevention	STD* prevention	Suicide prevention	Sun safety or skin cancer prevention	Tobacco-use prevention	Violence prevention
STATE SURVEYS								
Alabama	76.6	77.1	75.1	74.6	69.3	66.2	77.3	74.8
Alaska	73.5	74.0	62.2	62.9	60.3	42.2	73.0	69.3
Arizona	43.1	44.0	29.6	30.8	32.6	37.3	42.9	39.7
Arkansas	95.9	96.0	87.1	86.4	78.8	78.9	95.5	91.3
Connecticut	85.4	86.4	84.3	83.2	69.3	56.9	87.1	80.8
Delaware	91.9	91.7	83.4	85.1	68.8	58.6	90.1	81.8
Florida	54.2	53.0	53.0	54.0	44.2	45.7	55.7	49.6
Georgia	88.7	89.2	83.8	84.3	66.4	73.6	89.8	79.5
Hawaii	96.1	96.1	92.0	92.3	72.9	64.9	96.1	91.1
Idaho	95.3	94.6	89.3	87.9	85.7	82.7	93.4	86.1
Iowa	73.4	73.5	68.6	69.8	58.4	63.0	74.2	61.7
Kansas	75.5	76.2	71.4	72.4	55.8	60.0	75.1	64.5
Maine	90.8	90.1	81.7	81.1	62.3	60.7	90.0	81.6
Massachusetts	81.4	81.1	74.1	74.1	61.0	66.0	81.2	77.9
Michigan	76.5	77.8	71.6	73.8	50.9	48.2	76.2	68.4
Mississippi	97.9	97.3	96.6	96.1	86.9	76.4	97.9	87.7
Missouri	86.1	85.7	80.0	81.1	65.6	73.3	85.9	75.1
Montana	95.4	95.5	87.4	83.2	72.4	69.7	95.5	88.7
Nebraska	85.4	86.7	78.2	77.0	62.9	65.0	86.2	65.8
New Hampshire	83.8	83.1	79.9	78.6	66.5	65.3	83.0	70.7
New York [†]	98.6	98.9	99.3	98.9	89.5	85.7	99.0	92.8
North Carolina	86.6	87.2	80.8	81.2	70.4	66.6	86.9	82.5
North Dakota	88.4	89.1	78.6	78.4	70.3	73.1	88.9	84.1
Oregon	93.2	94.2	93.2	90.2	75.3	69.5	94.9	85.6
Pennsylvania	92.8	94.1	89.8	89.3	64.4	70.2	93.8	80.7
Rhode Island	96.5	96.4	92.0	92.9	69.9	65.8	93.9	92.7
South Carolina	69.6	70.5	66.8	66.5	47.0	51.0	68.6	61.0
South Dakota	66.8	68.7	53.3	56.5	46.1	53.0	68.4	59.3
Tennessee	68.9	70.5	63.3	62.5	53.1	51.6	69.6	63.4
Texas	78.3	78.7	70.5	70.2	64.9	66.3	78.6	70.9
Utah	94.7	94.7	85.3	85.7	88.3	69.2	94.8	86.3
Vermont	79.4	76.6	75.2	75.2	53.9	49.4	80.8	73.7
Virginia	83.2	83.4	74.1	74.8	59.6	66.9	82.2	77.1
West Virginia	97.7	97.7	92.6	93.2	80.5	84.2	98.3	95.3
State Median	85.8	86.5	80.0	79.9	66.0	65.9	86.6	78.7
State Range	43.1 – 98.6	44.0 – 98.9	29.6 – 99.3	30.8 – 98.9	32.6 – 89.5	37.3 – 85.7	42.9 – 99.0	39.7 – 95.3

TABLE 5c. Percentage of All Schools That Tried to Increase Student Knowledge on a Specific Health-Related Topic in a Required Health Education Course During the 2005–2006 School Year, Selected U.S. Sites: School Health Profiles, Lead Health Education Teacher Surveys, 2006 (continued)

Site	Nutrition and dietary behavior	Physical activity and fitness	Pregnancy prevention	STD* prevention	Suicide prevention	Sun safety or skin cancer prevention	Tobacco-use prevention	Violence prevention
LOCAL SURVEYS								
Charlotte-Mecklenburg County	97.3	100.0	100.0	100.0	91.9	69.7	97.3	94.4
Chicago	50.9	52.2	39.8	41.4	29.4	29.2	48.3	45.4
Dallas	57.1	55.3	55.1	57.1	55.1	46.9	57.1	57.1
District of Columbia	71.7	75.5	69.7	68.5	48.5	32.4	71.0	62.4
Hillsborough County	42.8	42.0	39.2	44.0	36.7	41.4	45.8	42.8
Los Angeles	100.0	100.0	100.0	96.2	81.1	87.9	100.0	95.2
Memphis	82.2	84.6	84.4	82.4	67.5	53.8	83.8	82.4
Miami	50.6	50.6	52.5	52.5	46.3	42.6	51.9	46.2
Orange County	46.6	46.4	46.6	46.6	40.6	43.9	49.5	44.2
Philadelphia	84.4	82.8	75.4	76.2	50.6	46.3	82.5	76.9
San Diego‡	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
San Francisco	57.2	57.2	57.2	57.2	50.0	32.2	57.2	57.2
Local Median	57.2	56.3	56.2	57.2	49.3	43.3	57.2	57.2
Local Range	0.0 – 100.0	0.0 – 100.0	0.0 – 100.0	0.0 – 100.0	0.0 – 91.9	0.0 – 87.9	0.0 – 100.0	0.0 – 95.2

* Sexually transmitted disease.

† Survey did not include schools from the New York City Department of Education.

‡ San Diego does not have a required health education course, but requires that health education be taught in science and physical education classes.

TABLE 6. Percentage of All Schools That Tried To Improve Specific Student Skills in a Required Health Education Course During the 2005–2006 School Year, Selected U.S. Sites: School Health Profiles, Lead Health Education Teacher Surveys, 2006

Site	Finding valid information or services related to personal health and wellness	Influence of media on personal health and wellness	Communication skills	Decision-making skills	Goal-setting skills	Conflict resolution skills	Resisting peer pressure to engage in unhealthy behaviors
STATE SURVEYS							
Alabama	72.9	69.7	71.9	74.8	75.1	74.3	77.0
Alaska	63.6	58.8	58.3	56.7	67.3	66.5	70.7
Arizona	35.7	37.6	36.8	34.6	40.3	39.1	42.0
Arkansas	89.2	91.4	87.2	87.9	94.2	88.5	96.0
Connecticut	79.7	83.3	80.7	79.3	80.4	81.2	87.7
Delaware	83.8	86.9	83.2	88.3	88.3	86.6	91.7
Florida	49.7	52.0	51.5	51.3	52.8	52.9	55.1
Georgia	83.7	83.4	81.1	81.3	88.7	81.8	89.4
Hawaii	90.0	93.1	91.3	89.4	96.1	88.2	96.1
Idaho	85.4	89.0	86.6	87.2	92.0	89.1	92.5
Iowa	65.0	67.3	62.5	63.2	69.2	63.3	71.9
Kansas	65.4	67.0	59.4	63.8	70.7	65.1	75.2
Maine	83.2	86.7	82.5	79.8	87.0	81.2	89.3
Massachusetts	75.8	78.4	75.6	75.7	78.2	75.6	81.6
Michigan	67.3	72.3	64.7	65.7	70.8	69.3	75.3
Mississippi	95.3	87.4	93.7	91.0	92.7	91.1	97.3
Missouri	79.8	80.4	75.5	77.6	83.0	78.1	85.2
Montana	81.4	86.9	82.0	79.8	92.4	87.4	95.0
Nebraska	75.7	78.6	67.1	76.7	81.4	68.4	86.2
New Hampshire	79.5	80.8	80.8	76.5	79.0	70.9	84.1
New York*	96.7	96.7	94.1	94.9	98.0	90.3	98.8
North Carolina	78.1	79.5	77.2	78.1	84.4	83.3	85.8
North Dakota	78.8	80.7	77.2	80.5	81.5	81.5	86.2
Oregon	85.5	90.9	80.6	82.9	90.4	88.0	93.3
Pennsylvania	85.8	87.4	83.4	83.3	88.5	84.8	93.0
Rhode Island	89.3	94.1	92.9	92.9	95.3	94.0	94.1
South Carolina	62.8	61.7	65.2	63.5	68.0	63.6	68.3
South Dakota	54.7	56.7	48.7	47.1	60.5	55.3	66.5
Tennessee	60.5	59.6	61.7	60.8	65.6	65.3	68.5
Texas	70.8	69.6	69.7	70.6	77.0	71.0	76.8
Utah	81.2	93.3	84.5	87.1	93.5	87.4	94.3
Vermont	71.8	78.1	75.3	71.0	77.4	70.1	78.0
Virginia	73.1	75.2	71.5	70.5	80.5	76.1	82.1
West Virginia	94.6	96.5	92.4	94.9	96.1	95.6	97.9
State Median	79.2	80.6	77.2	77.9	81.5	79.7	86.0
State Range	35.7 – 96.7	37.6 – 96.7	36.8 – 94.1	34.6 – 94.9	40.3 – 98.0	39.1 – 95.6	42.0 – 98.8

TABLE 6. Percentage of All Schools That Tried To Improve Specific Student Skills in a Required Health Education Course During the 2005–2006 School Year, Selected U.S. Sites: School Health Profiles, Lead Health Education Teacher Surveys, 2006 (*continued*)

Site	Finding valid information or services related to personal health and wellness	Influence of media on personal health and wellness	Communication skills	Decision-making skills	Goal-setting skills	Conflict resolution skills	Resisting peer pressure to engage in unhealthy behaviors
LOCAL SURVEYS							
Charlotte-Mecklenburg County	97.2	94.5	100.0	92.1	97.4	94.5	100.0
Chicago	43.0	38.0	43.7	40.5	46.6	47.4	48.3
Dallas	55.1	55.1	55.1	57.1	55.1	57.1	57.1
District of Columbia	76.4	58.8	69.4	69.7	76.4	69.1	72.7
Hillsborough County	37.5	39.6	37.5	37.5	44.7	46.5	44.7
Los Angeles	91.4	97.5	90.3	91.2	90.4	92.9	98.9
Memphis	78.7	83.0	81.9	79.0	80.4	82.7	84.4
Miami	48.8	47.5	50.0	51.3	50.0	48.7	51.2
Orange County	49.5	49.5	46.6	49.5	49.5	46.9	49.5
Philadelphia	67.5	73.3	68.7	69.8	76.7	77.1	83.0
San Diego [†]	0.0	0.0	0.0	0.0	0.0	0.0	0.0
San Francisco	53.6	53.6	53.6	57.2	53.6	57.2	53.6
Local Median	54.4	54.4	54.4	57.2	54.4	57.2	55.4
Local Range	0.0 – 97.2	0.0 – 97.5	0.0 – 100.0	0.0 – 92.1	0.0 – 97.4	0.0 – 94.5	0.0 – 100.0

* Survey did not include schools from the New York City Department of Education.

† San Diego does not have a required health education course, but requires that health education be taught in science and physical education classes.

TABLE 7a. Percentage of All Schools That Taught About Health Outcomes and Risks of Tobacco Use in a Required Health Education Course During the 2005–2006 School Year, Selected U.S. Sites: School Health Profiles, Lead Health Education Teacher Surveys, 2006

Site	Addictive effects of nicotine	Benefits of not smoking cigarettes	Benefits of not using smokeless tobacco	Short-term and long-term health consequences of cigarette smoking	Short-term and long-term health consequences of using smokeless tobacco	Health effects of ETS* or second-hand smoke	Short-term and long-term health consequences of cigar smoking
STATE SURVEYS							
Alabama	76.6	76.6	73.5	77.1	74.3	75.0	72.5
Alaska	70.7	71.7	68.3	71.4	67.9	66.3	50.7
Arizona	40.8	42.0	37.6	41.3	38.5	40.2	35.9
Arkansas	94.7	95.6	94.8	95.6	95.1	93.3	88.5
Connecticut	86.3	86.0	81.3	86.7	83.8	85.6	68.8
Delaware	90.1	88.5	86.9	88.7	87.1	87.0	80.0
Florida	53.6	54.7	50.4	54.7	50.9	53.5	47.3
Georgia	89.7	89.7	88.5	89.7	88.6	89.3	83.4
Hawaii	92.5	96.1	84.6	96.1	88.8	89.9	76.2
Idaho	93.4	94.0	92.3	94.0	92.8	93.3	85.6
Iowa	71.6	73.4	68.4	73.8	68.6	70.7	57.7
Kansas	71.8	73.5	72.7	73.6	74.0	71.7	63.0
Maine	88.5	88.5	81.5	88.5	82.7	86.6	65.3
Massachusetts	79.6	79.0	73.8	79.5	75.4	78.5	65.6
Michigan	73.7	74.6	72.3	75.0	73.0	73.7	65.7
Mississippi	96.7	97.9	96.8	97.3	96.8	96.0	90.5
Missouri	83.3	85.0	80.6	85.0	81.0	82.8	72.0
Montana	93.8	94.5	92.4	93.8	93.8	93.0	79.0
Nebraska	84.9	85.7	81.9	85.7	84.4	82.3	68.3
New Hampshire	80.8	81.9	79.6	82.5	79.5	80.0	72.4
New York†	99.0	98.7	95.9	99.0	98.1	97.9	91.0
North Carolina	86.7	86.3	84.3	86.7	84.9	85.6	77.4
North Dakota	87.6	87.6	85.6	87.6	86.3	87.4	75.3
Oregon	93.2	93.4	87.3	93.5	91.0	92.1	71.3
Pennsylvania	91.9	92.7	89.9	92.8	91.3	91.7	82.0
Rhode Island	92.8	92.8	88.2	94.0	89.4	90.5	74.4
South Carolina	66.5	66.9	64.4	66.5	65.2	66.5	58.1
South Dakota	65.4	66.4	64.2	66.9	65.5	64.5	52.4
Tennessee	67.1	69.1	65.9	68.6	67.7	66.8	61.9
Texas	76.8	77.5	76.9	77.5	77.2	76.0	71.4
Utah	92.3	94.2	89.7	94.7	90.7	92.4	75.6
Vermont	78.1	78.1	75.2	79.1	76.4	75.5	64.0
Virginia	80.3	81.5	77.9	81.5	78.0	79.4	67.9
West Virginia	97.7	98.3	98.3	98.3	98.3	98.3	90.3
State Median	85.6	85.9	81.4	86.2	83.3	84.2	71.7
State Range	40.8 – 99.0	42.0 – 98.7	37.6 – 98.3	41.3 – 99.0	38.5 – 98.3	40.2 – 98.3	35.9 – 91.0

TABLE 7a. Percentage of All Schools That Taught About Health Outcomes and Risks of Tobacco Use in a Required Health Education Course During the 2005–2006 School Year, Selected U.S. Sites: School Health Profiles, Lead Health Education Teacher Surveys, 2006 (continued)

Site	Addictive effects of nicotine	Benefits of not smoking cigarettes	Benefits of not using smokeless tobacco	Short-term and long-term health consequences of cigarette smoking	Short-term and long-term health consequences of using smokeless tobacco	Health effects of ETS* or second-hand smoke	Short-term and long-term health consequences of cigar smoking
LOCAL SURVEYS							
Charlotte-Mecklenburg County	97.4	97.4	89.3	97.4	91.9	94.7	81.0
Chicago	45.2	47.6	33.8	47.6	34.9	47.1	36.5
Dallas	56.3	56.3	56.3	56.3	54.2	55.3	52.1
District of Columbia	68.5	68.5	48.5	68.5	55.8	64.8	52.4
Hillsborough County	43.1	44.0	43.1	44.0	43.1	43.1	39.0
Los Angeles	100.0	100.0	98.7	100.0	97.5	98.9	93.9
Memphis	80.3	81.9	73.8	79.5	73.8	77.3	70.0
Miami	48.1	50.0	44.3	50.0	44.3	49.4	45.6
Orange County	46.6	46.6	46.6	46.6	46.6	46.6	41.2
Philadelphia	77.8	79.6	64.2	80.9	69.2	77.6	67.0
San Diego‡	0.0	0.0	0.0	0.0	0.0	0.0	0.0
San Francisco	57.2	57.2	50.0	57.2	53.6	53.6	32.1
Local Median	56.8	56.8	49.3	56.8	53.9	54.5	48.9
Local Range	0.0 – 100.0	0.0 – 100.0	0.0 – 98.7	0.0 – 100.0	0.0 – 97.5	0.0 – 98.9	0.0 – 93.9

* Environmental tobacco smoke.

† Survey did not include schools from the New York City Department of Education.

‡ San Diego does not have a required health education course, but requires that health education be taught in science and physical education classes.

TABLE 7b. Percentage of All Schools That Taught About External Influences on Tobacco Use in a Required Health Education Course During the 2005–2006 School Year, Selected U.S. Sites: School Health Profiles, Lead Health Education Teacher Surveys, 2006

Site	Influence of families on tobacco use	Influence of the media on tobacco use	Social or cultural influences on tobacco use	How many young people use tobacco
STATE SURVEYS				
Alabama	74.3	74.4	73.6	73.8
Alaska	64.8	66.1	63.5	67.8
Arizona	35.7	38.1	36.5	37.2
Arkansas	93.4	93.0	91.4	93.8
Connecticut	82.2	84.2	78.2	80.6
Delaware	90.1	90.1	83.5	85.1
Florida	51.2	53.9	51.0	51.7
Georgia	88.2	89.0	85.1	88.0
Hawaii	88.8	94.8	89.1	91.7
Idaho	91.2	91.2	88.6	90.8
Iowa	67.2	69.0	65.7	68.0
Kansas	70.3	70.9	66.8	69.5
Maine	83.7	85.5	76.4	79.6
Massachusetts	74.6	78.7	72.8	75.3
Michigan	71.2	73.7	69.0	70.1
Mississippi	95.2	94.0	91.5	94.1
Missouri	79.2	81.3	79.8	78.7
Montana	91.3	91.7	91.0	89.7
Nebraska	82.2	83.5	79.3	81.8
New Hampshire	78.2	80.5	73.6	76.8
New York*	96.4	98.7	94.9	96.2
North Carolina	82.9	86.4	81.9	83.9
North Dakota	82.6	85.2	80.2	83.3
Oregon	89.2	92.7	86.3	87.0
Pennsylvania	89.4	91.7	85.1	90.2
Rhode Island	86.9	89.4	87.0	89.2
South Carolina	63.6	63.7	62.5	64.1
South Dakota	63.7	65.2	60.4	62.2
Tennessee	63.3	64.3	63.1	63.8
Texas	72.8	76.0	71.3	73.7
Utah	89.1	93.6	85.1	86.1
Vermont	69.6	75.4	72.0	70.0
Virginia	77.9	78.5	74.4	76.4
West Virginia	97.1	97.7	95.0	96.6
State Median	82.2	83.9	78.8	80.1
State Range	35.7 – 97.1	38.1 – 98.7	36.5 – 95.0	37.2 – 96.6

TABLE 7b. Percentage of All Schools That Taught About External Influences on Tobacco Use in a Required Health Education Course During the 2005–2006 School Year, Selected U.S. Sites: School Health Profiles, Lead Health Education Teacher Surveys, 2006 (continued)

Site	Influence of families on tobacco use	Influence of the media on tobacco use	Social or cultural influences on tobacco use	How many young people use tobacco
LOCAL SURVEYS				
Charlotte-Mecklenburg County	94.5	91.7	89.1	94.7
Chicago	40.4	41.0	39.2	40.7
Dallas	56.3	56.3	56.3	54.2
District of Columbia	67.1	67.1	59.1	67.1
Hillsborough County	43.1	43.1	41.0	43.1
Los Angeles	97.6	100.0	96.3	96.6
Memphis	80.1	77.8	78.4	74.9
Miami	48.1	49.4	48.2	48.1
Orange County	43.9	46.6	43.9	46.6
Philadelphia	76.0	77.8	73.4	74.9
San Diego†	0.0	0.0	0.0	0.0
San Francisco	50.0	57.2	57.2	57.2
Local Median	53.2	56.8	56.8	55.7
Local Range	0.0 – 97.6	0.0 – 100.0	0.0 – 96.3	0.0 – 96.6

* Survey did not include schools from the New York City Department of Education.

† San Diego does not have a required health education course, but requires that health education be taught in science and physical education classes.

TABLE 7c. Percentage of All Schools That Taught Skills to Avoid and to Stop Using Tobacco in a Required Health Education Course During the 2005–2006 School Year, Selected U.S. Sites: School Health Profiles, Lead Health Education Teacher Surveys, 2006

Site	Resisting peer pressure to use tobacco	Making a personal commitment not to use tobacco	How students can influence or support others to prevent tobacco use	How to find valid information or services related to tobacco-use prevention or cessation	How students can influence or support others in efforts to quit using tobacco
STATE SURVEYS					
Alabama	76.8	64.3	75.2	70.4	74.0
Alaska	69.0	51.1	61.8	57.3	62.1
Arizona	40.0	32.2	36.7	33.1	35.5
Arkansas	95.6	75.5	90.8	80.8	90.5
Connecticut	83.9	57.9	77.1	68.2	74.2
Delaware	90.1	64.6	79.8	81.7	76.3
Florida	54.0	42.7	50.3	48.2	50.5
Georgia	88.9	75.0	87.4	79.1	88.1
Hawaii	96.1	68.5	88.4	77.8	88.4
Idaho	92.8	73.9	86.9	80.5	89.0
Iowa	69.6	51.7	65.9	59.4	63.7
Kansas	73.1	47.1	65.9	55.5	64.4
Maine	87.6	57.4	77.3	67.5	78.2
Massachusetts	78.5	60.1	72.1	64.6	70.1
Michigan	73.9	50.1	68.9	58.4	67.5
Mississippi	96.7	80.6	93.5	89.3	92.7
Missouri	83.1	64.3	79.2	70.3	78.4
Montana	93.2	75.8	87.0	77.7	86.8
Nebraska	83.5	63.0	78.2	64.5	74.3
New Hampshire	77.4	60.1	70.3	67.1	68.9
New York*	97.5	79.7	93.6	89.0	93.4
North Carolina	85.6	70.8	81.3	72.8	79.8
North Dakota	86.8	68.4	84.8	72.3	78.5
Oregon	92.2	66.0	87.0	71.7	84.6
Pennsylvania	92.9	71.7	84.9	77.2	83.0
Rhode Island	89.4	75.7	86.9	79.4	83.4
South Carolina	68.6	55.3	64.0	56.7	63.1
South Dakota	65.6	47.9	59.9	51.3	61.0
Tennessee	67.2	53.4	59.7	53.0	61.5
Texas	77.2	58.6	73.8	66.1	72.6
Utah	93.7	77.2	89.7	74.4	88.6
Vermont	76.7	48.7	66.9	62.9	62.2
Virginia	80.5	63.9	74.1	66.7	74.1
West Virginia	97.9	84.5	96.8	93.2	95.1
State Median	83.7	64.1	77.8	69.3	75.3
State Range	40.0 – 97.9	32.2 – 84.5	36.7 – 96.8	33.1 – 93.2	35.5 – 95.1

TABLE 7c. Percentage of All Schools That Taught Skills to Avoid and to Stop Using Tobacco in a Required Health Education Course During the 2005–2006 School Year, Selected U.S. Sites: School Health Profiles, Lead Health Education Teacher Surveys, 2006 (continued)

Site	Resisting peer pressure to use tobacco	Making a personal commitment not to use tobacco	How students can influence or support others to prevent tobacco use	How to find valid information or services related to tobacco-use prevention or cessation	How students can influence or support others in efforts to quit using tobacco
LOCAL SURVEYS					
Charlotte-Mecklenburg County	97.4	67.4	89.3	75.9	86.6
Chicago	45.2	36.2	41.6	33.2	39.3
Dallas	56.3	47.9	52.1	47.9	52.1
District of Columbia	68.5	57.7	67.1	51.8	71.0
Hillsborough County	43.1	37.0	39.0	43.1	41.0
Los Angeles	98.9	86.9	97.7	92.7	95.3
Memphis	77.6	70.2	77.9	58.6	78.4
Miami	49.4	43.0	49.4	48.1	50.7
Orange County	46.6	38.6	41.2	39.6	46.6
Philadelphia	79.8	59.8	68.1	62.4	69.0
San Diego†	0.0	0.0	0.0	0.0	0.0
San Francisco	57.2	50.0	57.2	53.6	57.2
Local Median	56.8	49.0	54.7	50.0	54.7
Local Range	0.0 – 98.9	0.0 – 86.9	0.0 – 97.7	0.0 – 92.7	0.0 – 95.3

* Survey did not include schools from the New York City Department of Education.

† San Diego does not have a required health education course, but requires that health education be taught in science and physical education classes.

TABLE 8a. Percentage of All Schools That Taught About HIV* Transmission and Prevention in a Required Health Education Course During the 2005–2006 School Year, Selected U.S. Sites: School Health Profiles, Lead Health Education Teacher Surveys, 2006

Site	Abstinence as the most effective method to avoid pregnancy, HIV, and STDs [†]	How HIV is transmitted	How HIV affects the human body	Condom efficacy	Risks associated with having multiple sexual partners
STATE SURVEYS					
Alabama	74.0	74.7	74.4	60.3	70.9
Alaska	58.3	60.4	58.6	44.0	56.2
Arizona	28.5	28.0	28.3	20.0	26.5
Arkansas	85.9	84.9	83.1	56.3	79.7
Connecticut	82.3	84.1	82.8	68.5	75.1
Delaware	87.9	83.1	83.1	71.2	83.2
Florida	52.0	51.8	51.3	40.9	50.2
Georgia	84.0	84.5	83.3	62.6	80.1
Hawaii	93.1	91.8	89.9	76.2	88.0
Idaho	88.9	87.6	85.1	51.6	78.8
Iowa	67.4	67.9	67.5	55.6	65.7
Kansas	69.1	69.7	67.7	49.1	64.7
Maine	80.9	81.2	79.7	65.5	74.5
Massachusetts	72.0	73.1	72.0	58.7	66.8
Michigan	69.8	70.5	70.1	49.3	64.8
Mississippi	97.0	97.2	95.8	74.2	93.8
Missouri	78.7	80.2	79.9	51.1	71.9
Montana	84.9	84.0	83.0	59.3	78.3
Nebraska	76.8	77.1	73.8	54.1	70.3
New Hampshire	77.3	79.0	78.3	61.9	74.4
New York [‡]	99.3	99.3	99.3	90.0	96.2
North Carolina	81.0	78.4	76.4	50.5	67.6
North Dakota	76.9	77.8	77.2	47.2	74.1
Oregon	91.5	91.6	91.4	76.7	83.6
Pennsylvania	88.4	89.0	89.0	67.0	84.7
Rhode Island	92.0	93.4	93.4	68.2	85.8
South Carolina	65.9	65.8	64.9	45.6	59.4
South Dakota	50.3	50.1	48.9	28.4	43.3
Tennessee	61.5	62.4	61.2	47.2	59.7
Texas	69.2	67.4	66.8	44.0	62.6
Utah	87.5	87.0	85.5	31.5	75.4
Vermont	72.2	74.0	71.5	65.4	73.2
Virginia	72.4	73.1	72.9	11.7	66.0
West Virginia	92.4	91.1	91.1	72.6	86.8
State Median	78.0	78.7	77.8	56.0	73.7
State Range	28.5 – 99.3	28.0 – 99.3	28.3 – 99.3	11.7 – 90.0	26.5 – 96.2

TABLE 8a. Percentage of All Schools That Taught About HIV* Transmission and Prevention in a Required Health Education Course During the 2005–2006 School Year, Selected U.S. Sites: School Health Profiles, Lead Health Education Teacher Surveys, 2006 (continued)

Site	Abstinence as the most effective method to avoid pregnancy, HIV, and STDs†	How HIV is transmitted	How HIV affects the human body	Condom efficacy	Risks associated with having multiple sexual partners
LOCAL SURVEYS					
Charlotte-Mecklenburg County	100.0	100.0	97.3	83.4	91.7
Chicago	37.4	39.4	39.5	31.0	35.6
Dallas	56.3	54.2	54.2	35.4	54.2
District of Columbia	71.0	71.0	71.0	60.2	71.0
Hillsborough County	39.2	39.2	39.2	34.0	38.5
Los Angeles	97.4	98.7	98.7	91.1	95.0
Memphis	84.4	84.4	84.4	65.6	84.4
Miami	52.5	52.5	51.2	50.0	52.5
Orange County	48.0	46.6	46.6	43.6	46.6
Philadelphia	75.4	74.1	72.0	56.0	73.4
San Diego§	0.0	0.0	0.0	0.0	0.0
San Francisco	57.2	57.2	57.2	57.2	57.2
Local Median	56.8	55.7	55.7	53.0	55.7
Local Range	0.0 – 100.0	0.0 – 100.0	0.0 – 98.7	0.0 – 91.1	0.0 – 95.0

* Human immunodeficiency virus.

† Sexually transmitted disease.

‡ Survey did not include schools from the New York City Department of Education.

§ San Diego does not have a required health education course, but requires that health education be taught in science and physical education classes.

TABLE 8b. Percentage of All Schools That Taught About External Influences on Behaviors Related to HIV* Infection, and Skills to Avoid HIV Infection, STDs,† and Pregnancy in a Required Health Education Course During the 2005–2006 School Year, Selected U.S. Sites: School Health Profiles, Lead Health Education Teacher Surveys, 2006

Site	Influence of alcohol and other drugs on HIV-related risk behaviors	Social or cultural influences on sexual behavior	How to prevent HIV infection	How to find valid information or services related to HIV or HIV testing	How to correctly use a condom	Compassion for persons living with HIV or AIDS‡
STATE SURVEYS						
Alabama	73.5	67.3	74.2	68.0	21.5	65.3
Alaska	55.7	52.7	60.0	49.6	22.3	44.3
Arizona	26.8	24.1	27.6	22.9	9.6	23.3
Arkansas	83.3	71.3	83.4	72.8	19.5	71.4
Connecticut	82.6	65.6	83.8	62.9	39.7	73.9
Delaware	82.8	74.4	83.1	76.3	43.8	71.6
Florida	50.8	46.0	51.9	46.3	25.5	45.2
Georgia	81.6	72.5	84.0	74.8	19.0	73.2
Hawaii	89.3	82.2	91.8	82.8	51.6	84.4
Idaho	85.9	65.4	87.6	69.3	18.4	69.2
Iowa	63.8	59.3	67.4	52.8	28.6	54.7
Kansas	66.9	55.1	69.0	51.8	20.4	54.2
Maine	77.6	63.5	80.1	68.8	45.4	65.2
Massachusetts	70.7	61.0	72.5	60.8	36.4	65.3
Michigan	67.7	57.2	70.4	55.9	23.1	60.4
Mississippi	95.1	86.9	97.2	87.8	28.9	84.4
Missouri	77.4	65.3	79.9	65.2	19.0	64.6
Montana	87.3	67.7	84.4	75.3	29.2	71.2
Nebraska	76.5	62.6	75.8	55.4	19.4	59.6
New Hampshire	79.1	68.4	78.4	67.7	42.5	71.0
New York [§]	99.3	87.7	98.9	91.9	59.1	91.6
North Carolina	76.4	62.9	78.0	60.0	12.8	63.7
North Dakota	76.2	67.6	77.8	65.5	15.8	68.7
Oregon	88.8	79.5	91.6	70.4	44.5	75.0
Pennsylvania	87.1	70.1	88.7	72.8	36.2	75.9
Rhode Island	90.8	75.7	89.7	82.2	43.6	82.0
South Carolina	62.5	55.8	65.0	56.8	26.3	55.9
South Dakota	48.3	42.7	50.4	35.7	11.6	32.9
Tennessee	59.0	55.7	62.0	55.2	17.7	51.8
Texas	64.5	57.0	66.5	56.1	16.5	55.1
Utah	80.2	69.3	85.7	56.4	1.0	74.1
Vermont	70.1	60.2	74.0	60.8	46.0	65.7
Virginia	71.5	56.1	72.5	62.8	7.7	61.1
West Virginia	89.2	82.9	92.4	81.3	33.6	81.5
State Median	77.0	65.4	78.2	64.1	24.3	65.5
State Range	26.8 – 99.3	24.1 – 87.7	27.6 – 98.9	22.9 – 91.9	1.0 – 59.1	23.3 – 91.6

TABLE 8b. Percentage of All Schools That Taught About External Influences on Behaviors Related to HIV* Infection, and Skills to Avoid HIV Infection, STDs,† and Pregnancy in a Required Health Education Course During the 2005–2006 School Year, Selected U.S. Sites: School Health Profiles, Lead Health Education Teacher Surveys, 2006 (continued)

Site	Influence of alcohol and other drugs on HIV-related risk behaviors	Social or cultural influences on sexual behavior	How to prevent HIV infection	How to find valid information or services related to HIV or HIV testing	How to correctly use a condom	Compassion for persons living with HIV or AIDS‡
LOCAL SURVEYS						
Charlotte-Mecklenburg County	97.3	86.1	97.3	72.8	22.0	75.1
Chicago	37.1	32.4	39.0	33.6	18.5	37.1
Dallas	52.1	50.0	54.2	52.1	12.5	50.0
District of Columbia	71.0	64.8	71.0	67.1	44.0	64.8
Hillsborough County	39.2	37.6	39.2	38.4	18.2	36.2
Los Angeles	97.4	90.1	98.7	92.3	74.8	88.8
Memphis	84.4	83.0	84.4	81.9	33.7	80.1
Miami	51.3	48.7	52.5	51.2	42.4	51.2
Orange County	43.9	40.9	46.6	46.6	34.7	43.9
Philadelphia	68.1	60.6	73.2	61.0	37.5	63.9
San Diego	0.0	0.0	0.0	0.0	0.0	0.0
San Francisco	57.2	46.4	57.2	53.6	42.9	53.6
Local Median	54.7	49.4	55.7	52.9	34.2	52.4
Local Range	0.0 – 97.4	0.0 – 90.1	0.0 – 98.7	0.0 – 92.3	0.0 – 74.8	0.0 – 88.8

* Human immunodeficiency virus.

† Sexually transmitted disease.

‡ Acquired immunodeficiency syndrome.

§ Survey did not include schools from the New York City Department of Education.

|| San Diego does not have a required health education course, but requires that health education be taught in science and physical education classes.

TABLE 9. Percentage of All Schools That Taught Required HIV* Prevention Units or Lessons in Specific Courses, Selected U.S. Sites: School Health Profiles, Lead Health Education Teacher Surveys, 2006

Site	Science	Home economics or family and consumer education	Physical education	Family life education or life skills	Special education	Social studies
STATE SURVEYS						
Alabama	66.3	45.3	53.2	45.1	30.6	18.3
Alaska	29.1	11.0	22.6	31.7	9.8	12.1
Arizona	26.0	7.2	19.7	16.9	7.9	7.2
Arkansas	44.1	53.4	31.5	55.9	28.2	12.5
Connecticut	32.5	12.4	20.3	41.4	15.0	6.8
Delaware	22.8	15.9	22.4	29.1	20.0	5.4
Florida	54.2	28.1	27.7	52.0	26.1	12.4
Georgia	35.2	25.5	45.8	36.5	24.4	7.7
Hawaii	12.9	6.4	14.6	14.6	13.3	2.0
Idaho	29.7	26.1	19.7	27.3	9.5	5.2
Iowa	38.1	48.1	13.5	45.0	11.9	6.0
Kansas	34.4	39.6	58.4	42.4	19.3	4.7
Maine	25.4	10.4	9.7	35.0	11.4	4.9
Massachusetts	24.5	11.2	14.6	23.0	9.6	3.5
Michigan	28.9	21.4	28.2	35.7	18.7	4.7
Mississippi	55.3	66.9	28.5	59.3	47.3	19.9
Missouri	31.5	47.1	36.4	39.5	18.7	8.5
Montana	32.4	31.5	71.8	34.2	19.8	8.2
Nebraska	30.1	50.0	33.5	45.4	14.9	7.8
New Hampshire	30.7	22.2	9.3	26.9	12.8	7.0
New York [†]	44.5	22.8	9.6	40.3	26.6	7.6
North Carolina	31.4	21.1	62.0	45.9	15.8	8.2
North Dakota	41.2	44.8	24.5	52.6	16.8	7.8
Oregon	35.4	13.0	20.1	31.5	20.0	12.9
Pennsylvania	27.4	19.4	34.4	28.4	18.8	6.4
Rhode Island	21.7	10.3	40.5	28.7	14.1	5.0
South Carolina	32.4	21.8	49.2	33.6	14.2	6.5
South Dakota	40.1	31.3	25.4	30.0	8.4	10.4
Tennessee	36.0	28.7	45.8	47.8	19.5	10.2
Texas	45.2	34.4	26.7	39.0	16.5	9.4
Utah	31.4	42.4	9.4	43.5	9.3	6.8
Vermont	34.3	23.0	15.4	37.6	5.4	8.9
Virginia	24.2	21.9	55.8	86.3	17.5	8.9
West Virginia	36.8	31.4	27.1	49.6	24.1	10.8
State Median	32.4	24.3	26.9	38.3	16.7	7.8
State Range	12.9 – 66.3	6.4 – 66.9	9.3 – 71.8	14.6 – 86.3	5.4–47.3	2.0–19.9

TABLE 9. Percentage of All Schools That Taught Required HIV* Prevention Units or Lessons in Specific Courses, Selected U.S. Sites: School Health Profiles, Lead Health Education Teacher Surveys, 2006 (continued)

Site	Science	Home economics or family and consumer education	Physical education	Family life education or life skills	Special education	Social studies
LOCAL SURVEYS						
Charlotte-Mecklenburg County	26.1	15.3	55.8	34.5	21.2	9.1
Chicago	44.5	8.0	37.2	46.9	27.9	9.5
Dallas	54.0	38.8	25.0	40.0	24.5	10.4
District of Columbia	37.3	3.7	81.6	38.1	13.9	3.5
Hillsborough County	75.8	22.1	13.7	33.7	14.4	0.0
Los Angeles	53.8	8.4	6.4	28.7	56.3	4.7
Memphis	44.9	46.2	83.4	71.2	45.4	24.2
Miami	69.3	27.7	36.1	47.3	30.9	26.4
Orange County	69.1	14.3	11.3	40.6	34.9	5.8
Philadelphia	25.4	9.7	50.0	29.2	27.8	14.5
San Diego	76.4	0.0	39.2	44.3	40.7	31.5
San Francisco	65.6	6.9	30.0	38.0	32.1	13.8
Local Median	53.9	12.0	36.7	39.1	29.4	10.0
Local Range	25.4 – 76.4	0.0 – 46.2	6.4 – 83.4	28.7 – 71.2	13.9–56.3	0.0–31.5

* Human immunodeficiency virus.

† Survey did not include schools from the New York City Department of Education.

TABLE 10a. Percentage of All Schools That Taught About Choosing Healthful Foods in a Required Health Education Course During the 2005–2006 School Year, Selected U.S. Sites: School Health Profiles, Lead Health Education Teacher Surveys, 2006

Site	Benefits of healthy eating	Using food labels	Food guidance using MyPyramid*	Eating more fruits, vegetables, and grain products	Choosing foods that are low in fat, saturated fat, and cholesterol	Using sugars in moderation	Using salt and sodium in moderation	Eating more calcium-rich foods
STATE SURVEYS								
Alabama	76.0	73.0	69.6	75.1	73.3	72.5	71.7	69.0
Alaska	72.5	67.6	61.8	70.5	68.4	69.1	63.2	61.3
Arizona	42.4	37.9	35.7	41.7	40.1	39.3	36.0	37.4
Arkansas	95.9	90.7	89.5	93.7	92.8	92.2	89.8	86.2
Connecticut	83.9	76.7	72.9	81.9	81.6	80.1	77.6	73.6
Delaware	90.3	83.5	81.8	88.5	86.9	85.3	83.6	75.2
Florida	53.6	51.2	49.9	53.0	52.3	52.2	51.6	52.3
Georgia	88.5	84.8	82.6	87.0	86.7	85.0	85.0	79.8
Hawaii	96.1	89.0	85.2	90.3	91.6	90.8	82.8	82.0
Idaho	93.8	91.4	88.3	94.1	93.0	91.4	90.4	91.0
Iowa	72.1	66.6	63.9	69.8	67.9	65.7	65.1	65.2
Kansas	74.3	66.6	66.8	72.2	70.7	71.8	68.0	66.8
Maine	89.6	82.1	77.8	84.9	87.3	85.7	84.9	79.8
Massachusetts	79.0	71.7	70.5	77.5	75.3	73.4	69.1	70.0
Michigan	75.1	68.7	67.2	74.2	72.5	71.1	68.1	65.3
Mississippi	97.9	95.0	89.7	96.6	95.4	94.7	92.9	92.4
Missouri	85.5	82.1	80.9	84.4	83.8	81.6	80.8	81.1
Montana	93.4	84.5	86.2	91.6	90.9	90.9	87.3	87.4
Nebraska	81.9	70.8	72.1	77.8	78.6	72.6	71.1	70.7
New Hampshire	81.7	78.8	77.8	79.9	79.3	79.4	77.8	75.9
New York†	98.2	89.1	86.5	97.0	96.7	94.5	92.1	90.1
North Carolina	85.7	83.4	78.6	84.8	84.2	82.8	81.0	80.1
North Dakota	87.7	80.0	81.7	84.4	80.1	80.5	80.4	81.3
Oregon	91.4	85.3	82.0	89.6	89.5	89.6	84.7	80.9
Pennsylvania	89.8	82.6	82.1	87.6	87.5	85.6	82.8	79.4
Rhode Island	93.0	87.0	84.7	90.7	91.8	84.6	82.2	84.3
South Carolina	68.1	61.8	62.3	68.1	66.2	63.7	62.5	60.1
South Dakota	65.4	56.1	57.0	61.9	60.2	59.2	56.1	54.5
Tennessee	68.0	62.5	59.8	66.5	65.7	63.2	62.5	61.1
Texas	77.8	74.4	72.6	76.1	75.3	74.1	73.4	72.3
Utah	94.7	90.4	90.2	94.2	92.1	90.4	87.1	87.4
Vermont	78.8	66.4	63.4	75.7	73.2	72.9	68.7	63.7
Virginia	82.2	77.8	76.8	82.2	81.0	79.7	78.2	76.6
West Virginia	97.7	95.6	92.5	95.0	96.0	95.0	92.2	88.7
State Median	84.7	79.4	77.8	83.3	81.3	80.3	79.3	76.3
State Range	42.4 – 98.2	37.9 – 95.6	35.7 – 92.5	41.7 – 97.0	40.1 – 96.7	39.3 – 95.0	36.0 – 92.9	37.4 – 92.4

TABLE 10a. Percentage of All Schools That Taught About Choosing Healthful Foods in a Required Health Education Course During the 2005–2006 School Year, Selected U.S. Sites: School Health Profiles, Lead Health Education Teacher Surveys, 2006 (continued)

Site	Benefits of healthy eating	Using food labels	Food guidance using MyPyramid*	Eating more fruits, vegetables, and grain products	Choosing foods that are low in fat, saturated fat, and cholesterol	Using sugars in moderation	Using salt and sodium in moderation	Eating more calcium-rich foods
LOCAL SURVEYS								
Charlotte-Mecklenburg County	97.4	94.5	89.1	94.5	91.9	94.5	91.9	83.4
Chicago	49.6	43.8	43.6	48.3	44.9	45.0	43.6	41.6
Dallas	56.3	56.3	56.3	54.2	56.3	52.1	54.2	50.0
District of Columbia	69.1	62.1	65.8	69.1	69.1	62.4	58.8	62.4
Hillsborough County	41.0	37.4	37.1	41.0	41.0	39.2	41.0	41.0
Los Angeles	100.0	98.7	98.7	100.0	98.7	98.7	95.1	94.0
Memphis	82.2	74.5	71.6	82.2	82.2	73.3	74.2	76.4
Miami	50.0	47.5	46.3	50.0	50.0	50.0	47.5	48.8
Orange County	46.6	40.9	46.6	46.6	43.6	43.6	43.6	43.6
Philadelphia	83.5	77.3	75.5	81.6	80.5	75.3	75.2	69.1
San Diego‡	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
San Francisco	57.2	53.6	53.6	53.6	53.6	53.6	53.6	53.6
Local Median	56.8	55.0	55.0	53.9	55.0	52.9	53.9	51.8
Local Range	0.0 – 100.0	0.0 – 98.7	0.0 – 98.7	0.0 – 100.0	0.0 – 98.7	0.0 – 98.7	0.0 – 95.1	0.0 – 94.0

* Visit the MyPyramid Web site at <http://www.mypyramid.gov>.

† Survey did not include schools from the New York City Department of Education.

‡ San Diego does not have a required health education course, but requires that health education be taught in science and physical education classes.

TABLE 10b. Percentage of All Schools That Taught About Food Safety and Behaviors That Contribute to Maintaining a Healthy Weight in a Required Health Education Course During the 2005–2006 School Year, Selected U.S. Sites: School Health Profiles, Lead Health Education Teacher Surveys, 2006

Site	Food safety	Balancing food intake and physical activity	Preparing healthy meals and snacks	Risks of unhealthy weight control practices	Accepting body size differences	Eating disorders
STATE SURVEYS						
Alabama	69.2	74.4	71.8	74.1	73.8	73.9
Alaska	59.6	69.5	63.7	64.8	61.1	57.9
Arizona	33.3	41.2	36.0	38.6	37.1	36.1
Arkansas	83.7	95.6	89.2	92.4	88.9	90.5
Connecticut	59.5	80.3	65.0	80.7	76.2	76.5
Delaware	74.8	90.2	80.0	90.0	83.6	80.2
Florida	47.8	53.0	50.4	51.3	49.5	52.0
Georgia	78.4	87.4	80.7	86.2	84.8	86.3
Hawaii	73.3	94.7	82.4	87.9	85.8	85.3
Idaho	82.6	94.1	84.4	93.0	90.4	91.3
Iowa	58.5	70.9	58.9	69.5	64.1	69.3
Kansas	60.6	73.2	62.2	70.2	70.0	71.6
Maine	68.2	87.8	73.0	83.8	82.0	80.6
Massachusetts	57.3	77.2	63.6	74.3	74.4	74.2
Michigan	58.4	73.9	66.4	72.7	69.3	66.7
Mississippi	93.1	97.3	93.5	96.7	94.4	94.4
Missouri	75.6	84.5	77.8	82.2	79.4	81.8
Montana	79.2	93.9	83.9	92.5	89.7	92.0
Nebraska	67.5	81.9	68.1	81.0	76.4	76.5
New Hampshire	66.8	79.7	69.3	78.2	77.3	76.7
New York*	76.0	96.5	87.6	97.3	94.1	95.2
North Carolina	75.1	85.6	81.7	84.4	81.4	82.6
North Dakota	77.5	84.3	77.3	82.4	79.9	81.7
Oregon	73.7	90.5	80.0	87.3	87.3	87.4
Pennsylvania	68.7	90.1	80.1	87.7	84.7	85.9
Rhode Island	77.2	89.5	78.5	95.4	89.3	93.0
South Carolina	56.8	67.4	60.6	65.1	62.3	63.2
South Dakota	48.4	61.3	51.8	55.7	54.6	58.3
Tennessee	58.5	66.4	61.9	66.7	60.2	62.8
Texas	69.9	76.2	74.5	75.9	74.6	74.7
Utah	75.2	92.6	80.7	92.2	92.0	93.6
Vermont	52.7	73.9	61.1	69.2	68.9	67.6
Virginia	70.1	81.2	73.9	78.5	73.1	76.3
West Virginia	84.9	96.1	93.4	94.3	93.9	95.4
State Median	69.6	83.1	74.2	81.6	78.4	78.5
State Range	33.3 – 93.1	41.2 – 97.3	36.0 – 93.5	38.6 – 97.3	37.1 – 94.4	36.1 – 95.4

TABLE 10b. Percentage of All Schools That Taught About Food Safety and Behaviors That Contribute to Maintaining a Healthy Weight in a Required Health Education Course During the 2005–2006 School Year, Selected U.S. Sites: School Health Profiles, Lead Health Education Teacher Surveys, 2006 (*continued*)

Site	Food safety	Balancing food intake and physical activity	Preparing healthy meals and snacks	Risks of unhealthy weight control practices	Accepting body size difference	Eating disorders
LOCAL SURVEYS						
Charlotte-Mecklenburg County	65.2	94.7	94.7	97.4	91.9	89.3
Chicago	39.4	48.8	43.9	43.4	45.0	39.8
Dallas	52.1	54.2	47.9	54.2	54.2	56.3
District of Columbia	58.5	67.9	65.8	61.0	62.1	55.5
Hillsborough County	34.4	41.0	38.9	41.0	40.3	41.0
Los Angeles	89.2	100.0	91.5	98.7	93.9	95.0
Memphis	76.2	82.2	72.9	80.6	71.8	74.7
Miami	47.5	50.0	48.7	48.8	48.7	48.1
Orange County	40.9	46.6	43.6	43.6	43.6	43.6
Philadelphia	71.7	81.6	80.5	76.9	72.1	73.0
San Diego [†]	0.0	0.0	0.0	0.0	0.0	0.0
San Francisco	42.9	50.0	53.6	53.6	57.2	57.2
Local Median	49.8	52.1	51.2	53.9	55.7	55.9
Local Range	0.0 – 89.2	0.0 – 100.0	0.0 – 94.7	0.0 – 98.7	0.0 – 93.9	0.0 – 95.0

* Survey did not include schools from the New York City Department of Education.

† San Diego does not have a required health education course, but requires that health education be taught in science and physical education classes.

TABLE 11a. Percentage of All Schools That Taught About the Benefits of Physical Activity and Guidance for Engaging in Physical Activity in a Required Health Education Course During the 2005–2006 School Year, Selected U.S. Sites: School Health Profiles, Lead Health Education Teacher Surveys, 2006

Site	Physical, psychological, or social benefits	Health-related fitness	Difference between physical activity, exercise, and fitness	Phases of a workout	How much physical activity is enough	Decreasing sedentary activities
STATE SURVEYS						
Alabama	76.2	74.5	72.1	73.4	72.3	74.2
Alaska	70.3	65.9	55.8	62.3	59.3	60.7
Arizona	41.4	41.3	36.8	36.8	35.1	40.2
Arkansas	94.8	93.8	90.8	91.8	90.1	91.2
Connecticut	77.7	67.2	57.8	58.5	60.0	68.6
Delaware	86.9	85.3	80.2	80.1	78.9	86.9
Florida	51.4	47.8	45.6	43.8	44.1	48.6
Georgia	86.7	85.1	81.4	80.4	80.9	82.7
Hawaii	90.8	79.3	70.7	71.4	72.9	81.2
Idaho	92.6	88.3	82.0	84.6	86.2	88.7
Iowa	71.1	70.0	66.5	61.1	61.5	66.1
Kansas	72.1	72.6	66.3	67.9	67.9	67.5
Maine	82.8	76.0	67.4	68.5	67.3	75.0
Massachusetts	76.7	67.3	58.2	58.1	58.7	69.4
Michigan	74.0	68.6	63.3	64.0	64.4	65.6
Mississippi	95.5	91.7	86.6	84.2	84.0	90.9
Missouri	83.2	80.9	77.2	77.6	77.8	79.4
Montana	94.4	93.1	88.3	93.2	88.1	90.9
Nebraska	81.5	77.0	72.1	73.5	73.7	73.4
New Hampshire	79.9	71.1	56.9	57.8	62.5	73.5
New York*	94.9	85.5	75.3	68.5	79.3	92.0
North Carolina	86.5	85.2	77.8	82.3	76.4	83.2
North Dakota	87.3	83.7	79.1	80.1	74.3	82.9
Oregon	90.3	84.2	78.1	79.2	79.1	82.8
Pennsylvania	89.9	87.6	78.9	82.4	79.0	86.4
Rhode Island	91.6	85.8	83.1	83.4	83.5	83.4
South Carolina	67.3	65.2	61.8	63.4	62.2	63.0
South Dakota	66.0	65.0	58.5	56.7	54.9	61.1
Tennessee	68.5	68.2	64.6	67.2	64.5	66.6
Texas	76.3	76.7	74.5	68.6	71.1	74.1
Utah	92.9	88.7	82.9	80.4	81.5	88.0
Vermont	69.3	59.3	49.0	48.4	55.5	60.5
Virginia	82.4	81.3	75.0	76.9	77.5	79.3
West Virginia	96.7	94.5	86.8	90.0	92.2	93.6
State Median	82.6	78.2	73.3	72.4	73.3	77.2
State Range	41.4 – 96.7	41.3 – 94.5	36.8 – 90.8	36.8 – 93.2	35.1 – 92.2	40.2 – 93.6

TABLE 11a. Percentage of All Schools That Taught About the Benefits of Physical Activity and Guidance for Engaging in Physical Activity in a Required Health Education Course During the 2005–2006 School Year, Selected U.S. Sites: School Health Profiles, Lead Health Education Teacher Surveys, 2006 (*continued*)

Site	Physical, psychological, or social benefits	Health-related fitness	Difference between physical activity, exercise, and fitness	Phases of a workout	How much physical activity is enough	Decreasing sedentary activities
LOCAL SURVEYS						
Charlotte-Mecklenburg County	91.6	86.0	72.1	83.1	85.8	91.4
Chicago	50.8	47.4	44.4	45.2	41.3	45.0
Dallas	51.1	45.7	45.8	39.6	37.5	46.8
District of Columbia	72.1	75.5	64.3	75.5	63.6	68.3
Hillsborough County	39.1	38.0	35.9	32.1	32.8	37.2
Los Angeles	92.6	90.1	80.5	77.4	86.8	92.8
Memphis	84.6	84.6	81.1	84.6	82.8	82.2
Miami	47.5	41.8	38.5	38.5	40.5	45.6
Orange County	41.9	35.6	30.6	38.3	33.5	39.1
Philadelphia	76.6	76.6	69.7	74.4	65.8	65.4
San Diego [†]	0.0	0.0	0.0	0.0	0.0	0.0
San Francisco	53.6	53.6	50.0	46.4	50.0	53.6
Local Median	52.4	50.5	47.9	45.8	45.7	50.2
Local Range	0.0 – 92.6	0.0 – 90.1	0.0 – 81.1	0.0 – 84.6	0.0 – 86.8	0.0 – 92.8

* Survey did not include schools from the New York City Department of Education.

† San Diego does not have a required health education course, but requires that health education be taught in science and physical education classes.

TABLE 11b. Percentage of All Schools That Taught About the Challenges to Engaging in Physical Activity in a Required Health Education Course During the 2005–2006 School Year, Selected U.S. Sites: School Health Profiles, Lead Health Education Teacher Surveys, 2006

Site	Overcoming barriers to physical activity	Developing an individualized physical activity plan	Monitoring progress toward reaching goals	Opportunities for physical activity in the community	Preventing injury during physical activity	Weather-related safety	Dangers of using performance-enhancing drugs
STATE SURVEYS							
Alabama	67.6	63.4	64.3	69.8	74.0	72.0	75.0
Alaska	52.4	49.7	48.3	59.4	57.6	56.0	58.2
Arizona	32.6	29.1	28.9	33.0	36.3	37.4	34.3
Arkansas	79.3	79.6	74.8	81.8	92.8	89.1	92.4
Connecticut	54.5	49.0	48.7	61.2	63.0	55.2	77.9
Delaware	78.6	65.3	66.3	81.9	81.8	73.6	83.4
Florida	43.9	39.5	38.1	41.7	44.0	47.0	47.3
Georgia	77.2	72.4	72.4	78.3	84.8	83.7	86.0
Hawaii	70.2	67.0	61.6	74.8	68.0	74.2	87.3
Idaho	75.2	76.1	67.2	75.8	85.0	82.9	89.4
Iowa	55.5	51.7	48.4	56.0	61.0	61.1	68.2
Kansas	54.9	50.8	47.9	54.1	64.9	63.2	71.2
Maine	61.4	61.0	55.1	64.7	70.6	68.3	76.6
Massachusetts	52.3	46.6	45.1	56.0	59.6	60.5	72.3
Michigan	58.0	54.2	49.9	56.4	62.0	58.7	69.0
Mississippi	77.4	67.2	63.9	81.9	87.9	88.9	95.3
Missouri	67.2	65.9	60.6	68.0	77.3	76.9	79.9
Montana	81.4	70.6	68.6	79.7	92.4	84.4	89.0
Nebraska	59.6	57.2	54.4	64.1	71.7	70.6	80.9
New Hampshire	59.6	52.2	49.8	63.2	63.8	65.6	76.9
New York*	73.4	55.9	50.0	73.0	73.8	80.4	96.3
North Carolina	73.2	71.1	69.6	79.3	81.9	77.4	81.6
North Dakota	69.2	58.3	50.4	70.5	74.9	76.0	83.6
Oregon	68.5	66.4	60.1	70.6	77.9	70.5	83.1
Pennsylvania	69.4	66.2	64.2	73.3	80.7	73.7	83.5
Rhode Island	76.2	59.7	59.6	73.6	79.7	71.2	88.0
South Carolina	60.0	54.5	51.7	62.5	64.3	59.2	64.3
South Dakota	48.6	41.7	40.5	52.0	59.4	50.6	58.3
Tennessee	57.7	53.0	51.4	57.5	66.6	59.2	62.6
Texas	64.4	63.6	59.8	66.2	72.0	72.6	75.4
Utah	74.3	68.1	64.2	74.4	80.8	75.2	89.7
Vermont	46.4	41.7	41.6	52.1	49.7	49.8	65.3
Virginia	70.1	67.9	66.7	70.6	76.7	72.0	77.2
West Virginia	84.2	74.7	70.9	85.9	92.9	89.3	92.7
State Median	67.4	60.4	57.4	68.9	72.9	71.6	78.9
State Range	32.6 – 84.2	29.1 – 79.6	28.9 – 74.8	33.0 – 85.9	36.3 – 92.9	37.4 – 89.3	34.3 – 96.3

TABLE 11b. Percentage of All Schools That Taught About the Challenges to Engaging in Physical Activity in a Required Health Education Course During the 2005–2006 School Year, Selected U.S. Sites: School Health Profiles, Lead Health Education Teacher Surveys, 2006 (continued)

Site	Overcoming barriers to physical activity	Developing an individualized physical activity plan	Monitoring progress toward reaching goals	Opportunities for physical activity in the community	Preventing injury during physical activity	Weather-related safety	Dangers of using performance-enhancing drugs
LOCAL SURVEYS							
Charlotte-Mecklenburg County	69.4	69.6	69.4	80.0	82.9	74.0	88.8
Chicago	41.0	33.1	32.7	37.9	47.4	41.3	41.4
Dallas	37.5	33.3	31.3	44.7	41.7	45.7	52.2
District of Columbia	65.7	61.4	63.3	68.3	72.1	61.1	57.7
Hillsborough County	29.5	28.8	26.7	24.6	32.1	34.9	37.0
Los Angeles	80.6	77.0	69.8	72.1	78.3	79.1	94.8
Memphis	79.5	79.3	79.5	77.1	84.6	78.0	80.4
Miami	43.6	39.8	37.2	42.4	41.1	43.7	45.5
Orange County	33.6	30.0	36.4	29.7	33.9	44.9	43.2
Philadelphia	61.6	61.4	59.1	63.6	69.4	59.3	64.7
San Diego [†]	0.0	0.0	0.0	0.0	0.0	0.0	0.0
San Francisco	50.0	42.9	39.3	35.7	46.4	48.2	57.2
Local Median	46.8	41.4	38.3	43.6	46.9	47.0	54.7
Local Range	0.0 – 80.6	0.0 – 79.3	0.0 – 79.5	0.0 – 80.0	0.0 – 84.6	0.0 – 79.1	0.0 – 94.8

* Survey did not include schools from the New York City Department of Education.

† San Diego does not have a required health education course, but requires that health education be taught in science and physical education classes.

TABLE 12. Percentage of All Schools That Taught All 16 Tobacco-Use Prevention Topics; All 11 Pregnancy, HIV* or STD† Prevention Topics; All 14 Nutrition and Dietary Behavior Topics; or All 13 Physical Activity Topics in a Required Health Education Course During the 2005–2006 School Year, Selected U.S. Sites: School Health Profiles, Lead Health Education Teacher Surveys, 2006

Site	Taught all 16 tobacco-use prevention topics	Taught all 11 pregnancy, HIV, or STD prevention topics	Taught all 14 nutrition and dietary behavior topics	Taught all 13 physical activity topics
STATE SURVEYS				
Alabama	58.1	21.0	57.0	54.2
Alaska	32.8	18.8	42.9	27.9
Arizona	23.1	9.0	23.8	20.1
Arkansas	66.3	19.1	71.8	62.0
Connecticut	43.7	32.6	45.9	24.8
Delaware	51.9	37.8	59.6	56.3
Florida	35.0	21.2	40.2	28.2
Georgia	64.8	18.3	67.2	60.1
Hawaii	50.0	44.9	60.0	37.6
Idaho	58.6	16.1	69.1	52.0
Iowa	37.7	23.7	43.2	35.8
Kansas	34.4	15.2	44.3	31.6
Maine	39.3	34.9	53.1	39.4
Massachusetts	41.9	30.8	45.3	31.5
Michigan	37.6	19.7	43.8	32.5
Mississippi	70.8	28.5	78.0	52.1
Missouri	51.1	16.8	64.8	46.6
Montana	59.2	26.5	64.5	54.8
Nebraska	44.4	16.8	52.5	37.4
New Hampshire	47.6	37.4	56.2	37.9
New York‡	67.8	53.1	64.5	34.4
North Carolina	58.5	12.1	66.0	52.8
North Dakota	51.9	15.3	65.7	40.8
Oregon	42.2	35.0	54.6	39.8
Pennsylvania	56.2	29.7	57.7	44.1
Rhode Island	59.6	39.9	63.5	45.3
South Carolina	43.0	23.0	46.5	42.7
South Dakota	34.5	9.5	35.7	27.5
Tennessee	39.4	15.5	46.0	39.0
Texas	50.5	15.1	61.3	48.8
Utah	51.5	1.0	63.1	43.4
Vermont	30.8	33.3	40.5	25.4
Virginia	48.6	1.5	58.8	50.2
West Virginia	74.5	31.0	75.9	58.2
State Median	49.3	21.1	57.4	40.3
State Range	23.1 – 74.5	1.0 – 53.1	23.8 – 78.0	20.1 – 62.0

TABLE 12. Percentage of All Schools That Taught All 16 Tobacco-Use Prevention Topics; All 11 Pregnancy, HIV* or STD† Prevention Topics; All 14 Nutrition and Dietary Behavior Topics; or All 13 Physical Activity Topics in a Required Health Education Course During the 2005–2006 School Year, Selected U.S. Sites: School Health Profiles, Lead Health Education Teacher Surveys, 2006 (continued)

Site	Taught all 16 tobacco-use prevention topics	Taught all 11 pregnancy, HIV, or STD prevention topics	Taught all 14 nutrition and dietary behavior topics	Taught all 13 physical activity topics
LOCAL SURVEYS				
Charlotte-Mecklenburg County	59.5	13.9	59.7	50.2
Chicago	23.0	16.8	28.8	22.3
Dallas	38.3	10.4	43.8	22.9
District of Columbia	38.5	44.0	48.5	36.0
Hillsborough County	33.0	18.2	30.6	20.0
Los Angeles	76.8	66.5	79.2	50.9
Memphis	51.1	33.7	56.0	64.8
Miami	37.9	40.0	39.9	33.8
Orange County	33.2	32.0	40.9	24.3
Philadelphia	44.8	28.5	55.1	33.8
San Diego [§]	0.0	0.0	0.0	0.0
San Francisco	32.1	28.6	42.9	28.6
Local Median	38.1	28.6	43.4	31.2
Local Range	0.0 – 76.8	0.0 – 66.5	0.0 – 79.2	0.0 – 64.8

* Human immunodeficiency virus.

† Sexually transmitted disease.

‡ Survey did not include schools from the New York City Department of Education.

§ San Diego does not have a required health education course, but requires that health education be taught in science and physical education classes.

TABLE 13a. Percentage of All Schools That Sometimes, Almost Always, or Always Used Specific Teaching Methods in a Required Health Education Course During the 2005–2006 School Year, Selected U.S. Sites: School Health Profiles, Lead Health Education Teacher Surveys, 2006

Site	Audio-visual media, (e.g., videos)	Group discussions	Cooperative group activities	Role play simulations, or practice	Language, performing, or visual arts	Pledges or contracts for changing behavior or abstaining from a behavior
STATE SURVEYS						
Alabama	74.1	76.8	70.6	49.2	35.2	40.1
Alaska	59.8	72.0	65.6	46.1	35.7	23.0
Arizona	39.1	42.6	39.9	32.0	23.0	21.9
Arkansas	83.1	95.6	84.5	53.2	35.4	29.5
Connecticut	79.3	86.8	84.3	65.1	42.6	22.7
Delaware	82.5	92.0	91.8	73.1	45.2	29.8
Florida	49.9	53.4	50.1	39.2	29.7	25.0
Georgia	85.3	87.4	81.1	68.1	49.4	41.3
Hawaii	77.5	96.1	90.5	78.5	66.1	43.3
Idaho	86.7	92.7	85.7	57.7	42.2	30.3
Iowa	66.2	72.1	68.4	47.3	34.8	19.0
Kansas	64.5	73.6	68.5	42.4	30.0	18.7
Maine	80.7	91.2	86.4	65.9	42.6	21.3
Massachusetts	78.4	82.4	80.2	64.3	44.6	24.2
Michigan	72.7	77.9	75.6	54.6	37.9	24.6
Mississippi	90.8	94.1	87.2	60.5	46.1	31.9
Missouri	78.5	84.6	77.6	52.4	40.5	32.5
Montana	87.1	95.2	88.2	55.0	36.6	24.8
Nebraska	81.1	84.5	78.5	49.4	39.1	26.5
New Hampshire	74.5	83.3	80.1	66.7	44.1	20.0
New York*	92.8	97.8	94.3	78.0	55.6	36.7
North Carolina	80.1	84.7	80.1	60.1	44.7	35.3
North Dakota	83.0	88.4	83.6	61.6	45.7	26.9
Oregon	85.9	93.8	91.7	67.9	51.8	27.8
Pennsylvania	89.1	92.4	87.1	59.8	41.4	27.7
Rhode Island	84.6	95.4	90.7	71.3	51.5	31.4
South Carolina	66.0	71.1	65.6	51.7	37.7	38.7
South Dakota	58.9	67.6	63.0	40.3	28.5	17.5
Tennessee	61.8	70.0	64.1	40.0	33.5	31.2
Texas	70.2	76.7	72.6	45.4	37.4	29.5
Utah	89.4	94.3	88.3	65.0	48.4	32.9
Vermont	71.9	80.4	77.0	61.0	38.7	19.9
Virginia	78.9	82.0	76.8	53.9	39.0	29.1
West Virginia	91.7	97.3	90.7	61.9	53.8	34.6
State Median	79.1	84.7	80.2	58.8	41.0	28.5
State Range	39.1 – 92.8	42.6 – 97.8	39.9 – 94.3	32.0 – 78.5	23.0 – 66.1	17.5 – 43.3

TABLE 13a. Percentage of All Schools That Sometimes, Almost Always, or Always Used Specific Teaching Methods in a Required Health Education Course During the 2005–2006 School Year, Selected U.S. Sites: School Health Profiles, Lead Health Education Teacher Surveys, 2006 (continued)

Site	Audio-visual media, (e.g., videos)	Group discussions	Cooperative group activities	Role play simulations, or practice	Language, performing, or visual arts	Pledges or contracts for changing behavior or abstaining from a behavior
LOCAL SURVEYS						
Charlotte-Mecklenburg County	86.4	100.0	94.7	83.8	66.7	49.0
Chicago	46.5	51.2	48.8	37.0	33.6	31.6
Dallas	56.3	56.3	52.1	37.5	39.6	18.8
District of Columbia	62.4	76.4	69.1	62.4	48.4	42.8
Hillsborough County	43.7	44.7	40.7	30.9	23.8	16.7
Los Angeles	94.9	95.0	97.7	81.1	76.1	54.8
Memphis	76.4	84.6	80.6	68.3	63.5	38.7
Miami	44.8	51.3	46.1	37.6	32.0	24.3
Orange County	49.5	49.5	49.5	32.9	24.3	16.6
Philadelphia	75.8	83.7	80.7	56.1	49.7	39.1
San Diego [†]	0.0	0.0	0.0	0.0	0.0	0.0
San Francisco	50.0	53.6	53.6	48.2	48.2	17.8
Local Median	53.2	55.0	52.9	42.9	43.9	28.0
Local Range	0.0 – 94.9	0.0 – 100.0	0.0 – 97.7	0.0 – 83.8	0.0 – 76.1	0.0 – 54.8

* Survey did not include schools from the New York City Department of Education.

† San Diego does not have a required health education course, but requires that health education be taught in science and physical education classes.

TABLE 13b. Percentage of All Schools That Sometimes, Almost Always, or Always Used Specific Teaching Methods in a Required Health Education Course During the 2005–2006 School Year, Selected U.S. Sites: School Health Profiles, Lead Health Education Teacher Surveys, 2006

Site	Peer teaching	Internet	Computer-assisted instruction	Guest speakers	Health education programs through available distance learning methods
STATE SURVEYS					
Alabama	47.8	58.6	40.5	57.8	14.9
Alaska	49.4	64.4	37.9	49.1	14.0
Arizona	28.7	31.5	22.5	29.8	5.0
Arkansas	51.3	61.2	37.3	51.8	9.9
Connecticut	54.0	70.7	48.7	51.0	5.4
Delaware	61.1	68.0	41.8	66.5	1.7
Florida	32.2	40.3	25.3	42.6	9.0
Georgia	56.5	67.0	51.5	54.7	12.3
Hawaii	82.7	70.5	41.6	77.2	5.3
Idaho	61.3	71.3	44.5	61.4	5.6
Iowa	46.4	58.3	37.3	45.0	8.5
Kansas	43.6	57.3	37.8	36.8	7.0
Maine	48.6	79.6	53.0	56.9	3.6
Massachusetts	48.1	60.6	39.6	44.6	4.8
Michigan	42.8	55.4	37.3	47.0	6.9
Mississippi	57.5	75.9	57.3	62.4	16.7
Missouri	49.9	64.2	44.3	53.0	10.2
Montana	55.2	68.0	51.0	67.2	8.9
Nebraska	44.4	64.5	42.1	48.6	6.0
New Hampshire	49.6	62.1	36.6	52.7	5.9
New York*	63.5	85.4	62.4	69.6	5.9
North Carolina	54.3	58.5	39.5	59.0	9.8
North Dakota	54.3	79.4	52.0	52.9	7.5
Oregon	61.9	73.3	42.5	59.0	8.8
Pennsylvania	50.1	70.6	42.9	57.9	5.9
Rhode Island	54.9	73.6	53.8	60.2	9.8
South Carolina	44.0	48.6	34.8	41.7	8.7
South Dakota	36.0	58.7	40.1	36.2	7.1
Tennessee	40.1	48.7	36.1	54.6	9.5
Texas	53.3	55.1	43.7	41.1	10.6
Utah	57.7	61.0	39.3	72.7	8.0
Vermont	46.4	53.2	33.6	49.7	3.5
Virginia	48.2	64.4	54.1	47.0	6.5
West Virginia	58.1	76.3	59.1	66.4	14.8
State Median	50.0	64.3	41.7	53.0	7.8
State Range	28.7 – 82.7	31.5 – 85.4	22.5 – 62.4	29.8 – 77.2	1.7 – 16.7

TABLE 13b. Percentage of All Schools That Sometimes, Almost Always, or Always Used Specific Teaching Methods in a Required Health Education Course During the 2005–2006 School Year, Selected U.S. Sites: School Health Profiles, Lead Health Education Teacher Surveys, 2006 (continued)

Site	Peer teaching	Internet	Computer-assisted instruction	Guest speakers	Health education programs through available distance learning methods
LOCAL SURVEYS					
Charlotte-Mecklenburg County	55.7	51.6	40.7	64.8	10.7
Chicago	33.1	39.5	27.4	35.5	9.6
Dallas	41.7	35.4	27.1	38.3	10.4
District of Columbia	45.1	42.1	36.4	62.0	14.2
Hillsborough County	21.6	25.7	9.8	44.0	3.4
Los Angeles	75.2	82.9	59.1	64.7	12.1
Memphis	63.5	72.1	48.9	57.2	23.5
Miami	34.6	39.8	20.5	41.5	6.4
Orange County	27.9	38.6	14.0	35.9	2.7
Philadelphia	48.9	46.7	30.4	38.0	10.4
San Diego [†]	0.0	0.0	0.0	0.0	0.0
San Francisco	42.9	37.1	14.3	46.5	0.0
Local Median	42.3	39.7	27.3	42.8	10.0
Local Range	0.0 – 75.2	0.0 – 82.9	0.0 – 59.1	0.0 – 64.8	0.0 – 23.5

* Survey did not include schools from the New York City Department of Education.

† San Diego does not have a required health education course, but requires that health education be taught in science and physical education classes.

TABLE 14. Percentage of All Schools That Used Specific Methods to Highlight Diversity or the Values of Various Cultures in a Required Health Education Course During the 2005–2006 School Year, Selected U.S. Sites: School Health Profiles, Lead Health Education Teacher Surveys, 2006

Site	Using textbooks or curricular materials reflective of various cultures	Using textbooks or curricular materials designed for students with limited English proficiency	Asking students or families to share their own cultural experiences related to health topics	Teaching about cultural differences and similarities	Modifying teaching methods to match students' learning styles, health beliefs, or cultural values
STATE SURVEYS					
Alabama	54.8	26.2	55.7	64.1	69.8
Alaska	53.6	21.8	54.3	63.2	68.1
Arizona	31.2	21.6	32.4	36.4	40.6
Arkansas	64.7	34.1	61.0	82.6	85.2
Connecticut	46.9	16.9	53.3	61.3	81.3
Delaware	64.6	30.4	57.8	74.6	88.6
Florida	41.8	29.1	37.5	42.6	51.1
Georgia	57.0	30.5	63.6	72.6	80.6
Hawaii	49.7	28.4	72.3	78.7	89.3
Idaho	63.7	32.5	62.4	78.7	83.9
Iowa	53.4	18.4	43.2	57.7	66.8
Kansas	40.6	15.0	34.2	52.1	53.2
Maine	44.4	13.5	50.8	62.8	79.2
Massachusetts	50.8	18.7	54.3	64.9	73.2
Michigan	50.5	15.4	41.2	52.9	67.8
Mississippi	72.2	26.2	68.1	84.4	87.7
Missouri	58.1	22.5	48.0	67.2	72.8
Montana	54.2	11.2	57.8	75.3	79.7
Nebraska	66.1	20.5	54.5	65.6	75.1
New Hampshire	51.0	11.8	48.9	62.6	75.4
New York*	61.5	21.2	69.5	77.6	87.9
North Carolina	56.7	38.2	55.4	69.0	79.7
North Dakota	60.9	15.7	54.8	71.3	73.4
Oregon	65.5	35.9	57.8	76.1	86.1
Pennsylvania	53.3	17.3	56.2	66.0	82.1
Rhode Island	48.3	25.9	63.8	69.5	84.2
South Carolina	48.4	22.9	47.1	54.0	63.1
South Dakota	31.1	4.7	27.7	38.8	49.1
Tennessee	48.3	20.8	41.9	56.5	59.7
Texas	60.5	40.6	54.2	65.6	72.0
Utah	61.5	41.3	48.2	73.8	84.1
Vermont	42.4	9.4	37.7	53.3	74.7
Virginia	44.8	27.0	45.6	60.3	73.1
West Virginia	72.8	26.5	67.4	85.5	92.1
State Median	53.5	22.2	54.3	65.6	75.3
State Range	31.1 – 72.8	4.7 – 41.3	27.7 – 72.3	36.4 – 85.5	40.6 – 92.1

TABLE 14. Percentage of All Schools That Used Specific Methods to Highlight Diversity or the Values of Various Cultures in a Required Health Education Course During the 2005–2006 School Year, Selected U.S. Sites: School Health Profiles, Lead Health Education Teacher Surveys, 2006 (continued)

Site	Using textbooks or curricular materials reflective of various cultures	Using textbooks or curricular materials designed for students with limited English proficiency	Asking students or families to share their own cultural experiences related to health topics	Teaching about cultural differences and similarities	Modifying teaching methods to match students' learning styles, health beliefs, or cultural values
LOCAL SURVEYS					
Charlotte-Mecklenburg County	65.2	44.4	67.8	84.0	91.9
Chicago	37.6	21.6	37.1	47.0	47.0
Dallas	47.9	31.3	45.8	54.2	56.3
District of Columbia	45.1	20.6	55.8	59.1	66.1
Hillsborough County	31.7	25.7	25.9	37.3	41.0
Los Angeles	87.6	85.4	86.7	91.5	97.5
Memphis	62.4	43.0	63.5	73.2	78.7
Miami	39.7	33.8	39.8	46.1	44.8
Orange County	35.6	24.9	43.9	43.9	49.5
Philadelphia	59.6	19.6	51.3	67.2	71.6
San Diego [†]	0.0	0.0	0.0	0.0	0.0
San Francisco	46.5	28.6	35.7	53.6	55.6
Local Median	45.8	27.2	44.9	53.9	56.0
Local Range	0.0 – 87.6	0.0 – 85.4	0.0 – 86.7	0.0 – 91.5	0.0 – 97.5

* Survey did not include schools from the New York City Department of Education.

† San Diego does not have a required health education course, but requires that health education be taught in science and physical education classes.

Table 15. Percentage of All Schools with a Health Education Coordinator and, Among those Schools, the Percentage in Which Specific Staff Served as the Health Education Coordinator, Selected U.S. Sites: School Health Profiles, Principal Surveys, 2006

Site	Had a health education coordinator	District administrator or district health education or curriculum coordinator	School administrator	Health education teacher	School nurse	Someone else
STATE SURVEYS						
Alabama	95.8	21.6	33.9	37.2	2.7	4.6
Alaska	89.4	34.7	31.7	25.0	2.5	6.1
Arizona	80.4	21.8	30.6	24.8	8.6	14.2
Arkansas	96.8	16.8	27.4	53.6	0.9	1.4
Connecticut	94.3	47.1	16.6	29.6	3.5	3.2
Delaware	98.5	19.7	27.3	49.9	1.5	1.5
Florida	89.6	24.6	29.0	29.9	3.7	12.8
Georgia	98.1	24.5	23.8	48.6	0.4	2.7
Hawaii	93.2	5.9	15.4	75.1	0.0	3.7
Idaho	98.4	27.9	12.4	57.1	0.0	2.6
Illinois	98.3	19.9	20.7	55.1	0.6	3.7
Iowa	93.7	29.2	16.5	46.9	2.0	5.4
Kansas	93.2	28.3	13.4	48.6	2.6	7.1
Maine	94.9	37.6	11.0	43.1	3.9	4.5
Massachusetts	94.7	49.0	14.3	29.3	2.5	4.9
Michigan	95.1	38.0	21.1	34.6	1.8	4.5
Mississippi	94.6	16.1	32.8	45.3	4.5	1.3
Missouri	96.6	27.4	16.6	46.9	5.3	3.9
Montana	99.2	19.1	15.8	62.8	0.0	2.3
Nebraska	95.9	26.8	21.4	47.3	2.4	2.2
New Hampshire	93.0	18.0	23.4	44.2	6.9	7.5
New York*	98.1	47.5	18.5	31.9	0.0	2.2
North Carolina	97.5	24.8	14.7	56.2	0.4	3.9
North Dakota	95.8	13.4	26.9	54.5	2.2	2.9
Oregon	95.8	18.2	20.8	54.3	0.0	6.7
Pennsylvania	97.2	34.4	20.3	41.6	1.5	2.2
Rhode Island	96.3	31.9	8.5	46.2	2.4	11.0
South Carolina	96.1	27.8	19.8	46.2	1.3	4.9
South Dakota	87.4	21.7	23.7	51.7	0.0	2.8
Tennessee	91.7	28.1	22.3	37.0	5.6	7.0
Texas	93.7	26.4	25.0	41.6	5.3	1.7
Utah	99.5	19.2	18.1	59.8	1.0	1.9
Vermont	91.1	11.2	21.4	49.0	8.7	9.7
Virginia	98.3	39.1	14.9	43.5	0.7	1.8
Washington	92.8	25.6	14.7	51.4	3.4	4.9
West Virginia	97.4	10.5	28.1	60.4	0.0	1.1
State Median	95.8	25.2	20.8	46.9	2.1	3.8
State Range	80.4 – 99.5	5.9 – 49.0	8.5 – 33.9	24.8 – 75.1	0.0 – 8.7	1.1 – 14.2

Table 15. Percentage of All Schools with a Health Education Coordinator and, Among those Schools, the Percentage in Which Specific Staff Served as the Health Education Coordinator, Selected U.S. Sites: School Health Profiles, Principal Surveys, 2006 (continued)

Site	Had a health education coordinator	District administrator or district health education or curriculum coordinator	School administrator	Health education teacher	School nurse	Someone else
LOCAL SURVEYS						
Charlotte-Mecklenburg County	100.0	33.5	9.7	56.8	0.0	0.0
Chicago	81.4	5.8	14.3	44.8	9.1	25.9
Dallas	74.5	38.1	8.1	40.6	7.9	5.3
District of Columbia	100.0	10.3	12.6	63.7	3.1	10.3
Hillsborough County	74.2	37.8	32.6	14.6	5.2	9.8
Los Angeles	97.5	12.9	22.3	62.5	0.0	2.3
Memphis	92.7	31.0	12.0	55.1	0.0	1.9
Miami	81.0	21.3	22.4	35.5	0.0	20.8
Orange County	96.8	9.2	36.9	30.9	3.3	19.8
Philadelphia	92.7	17.8	19.1	53.7	2.3	7.1
San Diego	92.6	37.3	5.9	21.5	5.8	29.5
San Francisco	97.2	6.1	6.5	57.6	0.0	29.8
Local Median	92.7	19.6	13.5	49.3	2.7	10.1
Local Range	74.2 – 100.0	5.8 – 38.1	5.9 – 36.9	14.6 – 63.7	0.0 – 9.1	0.0 – 29.8

* Survey did not include schools from the New York City Department of Education.

TABLE 16. Percentage of All Schools in Which Health Education Staff Worked on Health Education Activities with Other School Staff During the 2005–2006 School Year, Selected U.S. Sites: School Health Profiles, Lead Health Education Teacher Surveys, 2006

Site	Physical education staff	School health services staff	School mental health or social services staff	Nutrition or food service staff
STATE SURVEYS				
Alabama	74.1	78.3	59.8	39.2
Alaska	54.6	42.7	45.9	24.6
Arizona	56.4	47.1	45.9	31.7
Arkansas	79.9	67.7	51.3	37.0
Connecticut	76.5	73.0	65.6	34.1
Delaware	91.1	86.3	74.5	37.8
Florida	56.4	55.2	51.6	30.3
Georgia	90.9	64.4	51.4	33.2
Hawaii	79.9	34.9	38.3	13.5
Idaho	82.2	59.1	63.7	37.2
Iowa	63.8	70.0	44.7	39.0
Kansas	73.9	65.0	43.4	46.2
Maine	78.6	78.7	66.7	43.9
Massachusetts	78.7	74.2	66.7	43.7
Michigan	66.6	28.2	44.2	31.3
Mississippi	58.9	59.2	53.3	42.9
Missouri	83.8	80.6	59.3	41.2
Montana	85.4	56.8	64.6	39.4
Nebraska	71.8	71.5	51.0	37.1
New Hampshire	74.2	76.0	73.8	49.4
New York*	75.9	64.7	66.2	40.2
North Carolina	87.3	74.0	57.2	37.9
North Dakota	68.8	36.4	52.5	45.6
Oregon	76.8	48.7	59.1	30.6
Pennsylvania	89.4	81.0	53.7	42.6
Rhode Island	87.5	72.9	63.7	26.5
South Carolina	76.4	64.0	46.5	31.2
South Dakota	61.5	48.5	40.1	40.1
Tennessee	75.1	69.4	57.3	39.1
Texas	71.8	63.5	40.8	28.4
Utah	76.9	41.0	54.9	20.0
Vermont	77.1	87.8	82.4	48.1
Virginia	88.7	77.2	56.7	31.7
West Virginia	79.3	80.0	67.3	40.2
State Median	76.7	66.4	55.8	37.9
State Range	54.6 – 91.1	28.2 – 87.8	38.3 – 82.4	13.5 – 49.4

TABLE 16. Percentage of All Schools in Which Health Education Staff Worked on Health Education Activities with Other School Staff During the 2005–2006 School Year, Selected U.S. Sites: School Health Profiles, Lead Health Education Teacher Surveys, 2006 (continued)

Site	Physical education staff	School health services staff	School mental health or social services staff	Nutrition or food service staff
LOCAL SURVEYS				
Charlotte-Mecklenburg County	95.1	67.4	59.9	27.3
Chicago	61.3	55.1	56.3	27.7
Dallas	51.0	62.7	37.3	24.0
District of Columbia	75.5	74.8	60.4	39.9
Hillsborough County	53.5	70.4	22.9	19.0
Los Angeles	45.1	67.8	56.2	30.8
Memphis	96.8	76.0	58.5	54.3
Miami	46.8	27.1	46.9	21.3
Orange County	58.5	26.2	38.6	8.5
Philadelphia	83.8	84.8	50.5	53.5
San Diego	51.5	79.6	87.5	9.1
San Francisco	81.2	53.4	67.9	36.7
Local Median	59.9	67.6	56.3	27.5
Local Range	45.1 – 96.8	26.2 – 84.8	22.9 – 87.5	8.5 – 54.3

* Survey did not include schools from the New York City Department of Education.

TABLE 17. Percentage of All Schools in Which the Major Emphasis of the Lead Health Education Teacher’s Professional Preparation Was in a Specific Discipline, Selected U.S. Sites: School Health Profiles, Lead Health Education Teacher Surveys, 2006

Site	Health and physical education combined	Health education only	Physical education only	Other education degree	Kinesiology, exercise science or exercise physiology, home economics or family and consumer science, biology, or other science	Nursing or counseling	Public health, nutrition, or another discipline
STATE SURVEYS*							
Alabama	53.2	10.4	18.7	2.0	11.1	3.8	0.9
Alaska	9.5	4.3	5.8	33.7	21.9	10.3	14.4
Arizona	18.6	2.3	18.9	16.7	14.4	15.1	14.1
Arkansas	67.5	4.6	16.2	1.7	5.7	1.2	3.0
Connecticut	44.1	16.0	18.4	5.2	7.9	3.6	4.8
Delaware	74.2	4.7	12.0	0.0	4.5	1.6	3.0
Florida	26.6	14.2	12.0	14.2	19.7	4.1	9.1
Georgia	78.7	4.0	6.2	2.9	4.3	2.0	2.0
Hawaii	45.6	4.7	16.5	10.3	15.8	0.0	7.1
Idaho	57.5	8.1	12.8	8.7	7.6	1.7	3.7
Iowa	37.3	3.3	13.0	6.5	28.6	4.6	6.8
Kansas	50.8	2.3	26.6	6.3	4.5	7.6	2.0
Maine	40.3	14.9	8.2	10.6	15.3	7.3	3.6
Massachusetts	40.9	21.4	11.4	5.4	6.6	10.6	3.6
Michigan	39.5	9.6	15.3	7.8	23.7	1.7	2.5
Mississippi	47.3	9.7	7.5	2.8	27.1	2.1	3.5
Missouri	56.3	4.1	15.0	3.5	16.4	3.7	1.0
Montana	64.2	1.2	10.5	18.6	3.6	0.6	1.3
Nebraska	38.9	2.3	22.6	2.8	26.9	4.4	2.1
New Hampshire	20.4	21.0	19.7	4.7	12.6	18.2	3.4
New York [†]	40.8	42.9	6.5	0.7	6.8	1.1	1.2
North Carolina	60.2	3.4	23.3	4.3	5.1	2.6	1.1
North Dakota	40.2	1.7	11.6	12.8	26.3	2.3	5.0
Oregon	45.0	14.3	9.0	12.5	12.0	0.0	7.2
Pennsylvania	88.9	3.0	2.0	3.6	1.2	1.0	0.4
Rhode Island	74.9	7.5	5.0	0.0	0.0	11.2	1.3
South Carolina	45.4	4.2	36.4	3.0	5.7	2.5	2.7
South Dakota	46.2	1.5	6.5	20.3	17.9	2.7	4.8
Tennessee	61.1	3.0	9.5	8.6	6.3	10.1	1.5
Texas	45.2	8.6	9.1	3.8	26.1	3.9	3.3
Utah	45.4	16.0	20.6	5.2	10.3	0.0	2.5
Vermont	36.4	12.9	7.2	5.4	20.8	17.3	0.0
Virginia	81.9	2.4	9.3	3.1	1.9	0.7	0.7
West Virginia	79.2	6.6	6.7	4.7	0.6	0.5	1.8
State Median	45.5	4.7	11.8	5.2	10.7	2.7	2.9
State Range	9.5 – 88.9	1.2 – 42.9	2.0 – 36.4	0.0 – 33.7	0.0 – 28.6	0.0 – 18.2	0.0 – 14.4

TABLE 17. Percentage of All Schools in Which the Major Emphasis of the Lead Health Education Teacher's Professional Preparation Was in a Specific Discipline, Selected U.S. Sites: School Health Profiles, Lead Health Education Teacher Surveys, 2006 (continued)

Site	Health and physical education combined	Health education only	Physical education only	Other education degree	Kinesiology, exercise science or exercise physiology, home economics or family and consumer science, biology, or other science	Nursing or counseling	Public health, nutrition, or another discipline
LOCAL SURVEYS*							
Charlotte-Mecklenburg County	69.7	5.4	22.0	0.0	2.9	0.0	0.0
Chicago	28.2	1.7	24.5	14.0	10.6	15.0	5.6
Dallas	22.4	16.3	0.0	6.1	55.1	0.0	0.0
District of Columbia	82.4	0.0	0.0	0.0	3.4	10.7	3.4
Hillsborough County	32.3	10.6	14.6	10.8	11.4	1.8	18.4
Los Angeles	12.3	29.3	4.7	7.8	32.5	1.2	11.3
Memphis	75.7	4.8	12.5	1.6	5.3	0.0	0.0
Miami	14.5	22.4	22.3	1.4	26.4	9.2	3.9
Orange County	24.6	37.1	14.8	3.0	17.5	0.0	3.0
Philadelphia	85.0	2.3	3.5	2.3	2.3	3.5	1.2
San Diego	1.8	0.0	1.8	10.9	0.0	85.4	0.0
San Francisco	10.0	16.7	3.3	26.7	26.7	6.7	10.0
Local Median	26.4	8.0	8.6	4.6	11.0	2.7	3.2
Local Range	1.8 – 85.0	0.0 – 37.1	0.0 – 24.5	0.0 – 26.7	0.0 – 55.1	0.0 – 85.4	0.0 – 18.4

* Percentages for each row might not add up to 100.0 because of rounding.

† Survey did not include schools from the New York City Department of Education.

TABLE 18. Percentage of All Schools in Which a Newly Hired Health Education Teacher Was Required to Be Certified* in Health Education, the Lead Health Education Teacher Was Certified in Health Education, or the Lead Health Education Teacher Had Experience Teaching Health Education Classes or Topics for a Specific Number of Years, Selected U.S. Sites: School Health Profiles, Principal Surveys and Lead Health Education Teacher Surveys, 2006

Site	Newly hired health education teacher required to be certified in health education	Lead health education teacher certified in health education	Number of years lead health education teacher taught health education classes or topics				
			1 year	2–5 years	6–9 years	10–14 years	≥15 years
STATE SURVEYS							
Alabama	79.0	78.2	10.3	24.3	17.4	13.6	34.4
Alaska	23.8	26.7	23.1	30.3	19.0	11.4	16.2
Arizona	36.2	31.0	21.3	30.4	13.9	11.8	22.6
Arkansas	94.7	93.7	8.8	26.9	12.3	13.3	38.7
Connecticut	84.3	80.6	5.7	22.3	21.4	12.8	37.8
Delaware	93.9	85.5	7.4	22.3	23.8	22.5	24.0
Florida	65.3	51.1	12.8	23.4	16.1	12.1	35.6
Georgia	88.1	97.2	3.9	19.6	18.6	11.6	46.3
Hawaii	64.1	63.1	6.9	38.4	12.1	20.1	22.5
Idaho	92.9	91.2	5.3	20.2	15.8	21.0	37.7
Illinois	87.6	NA [†]	NA	NA	NA	NA	NA
Iowa	79.0	75.4	10.7	22.3	15.1	14.8	37.1
Kansas	76.8	71.2	8.6	23.3	17.0	9.7	41.4
Maine	80.1	74.2	9.9	26.5	13.1	10.5	40.1
Massachusetts	84.6	76.9	6.6	20.8	15.7	17.0	39.9
Michigan	90.0	82.2	10.4	28.0	20.8	14.5	26.2
Mississippi	92.1	91.9	8.2	23.2	20.8	14.0	33.7
Missouri	85.1	83.2	8.3	28.9	22.5	15.6	24.6
Montana	85.4	90.8	6.1	20.2	17.5	18.3	37.9
Nebraska	56.7	65.7	8.2	19.4	18.8	15.6	38.0
New Hampshire	68.9	54.2	3.7	25.6	23.8	16.2	30.7
New York [‡]	97.4	94.6	3.1	22.6	18.8	13.1	42.4
North Carolina	80.2	79.1	6.4	20.9	14.1	14.5	44.0
North Dakota	88.9	93.1	5.3	16.2	16.3	19.6	42.6
Oregon	77.7	78.1	7.2	24.0	14.0	13.5	41.3
Pennsylvania	89.9	95.3	1.7	18.7	14.4	14.3	50.9
Rhode Island	92.8	97.7	3.5	18.9	19.8	19.5	38.3
South Carolina	69.9	74.9	8.5	23.1	16.4	11.0	41.0
South Dakota	81.5	75.5	7.6	30.6	13.0	11.9	36.9
Tennessee	80.1	80.1	7.8	22.5	16.7	14.8	38.2
Texas	87.6	77.7	14.5	24.9	16.8	10.2	33.6
Utah	95.7	89.0	6.9	25.4	19.1	13.4	35.1
Vermont	75.0	62.7	10.2	19.8	14.6	17.8	37.6
Virginia	90.8	93.3	2.9	12.4	13.9	11.4	59.4
Washington	69.2	NA	NA	NA	NA	NA	NA
West Virginia	99.0	94.3	5.2	13.2	13.0	11.3	57.2
State Median	84.5	79.6	7.5	22.9	16.6	13.8	37.9
State Range	23.8 – 99.0	26.7 – 97.7	1.7 – 23.1	12.4 – 38.4	12.1 – 23.8	9.7 – 22.5	16.2 – 59.4

TABLE 18. Percentage of All Schools in Which a Newly Hired Health Education Teacher Was Required to Be Certified* in Health Education, the Lead Health Education Teacher Was Certified in Health Education, or the Lead Health Education Teacher Had Experience Teaching Health Education Classes or Topics for a Specific Number of Years, Selected U.S. Sites: School Health Profiles, Principal Surveys and Lead Health Education Teacher Surveys, 2006 (continued)

Site	Newly hired health education teacher required to be certified in health education	Lead health education teacher certified in health education	Number of years lead health education teacher taught health education classes or topics				
			1 year	2–5 years	6–9 years	10–14 years	≥15 years
LOCAL SURVEYS							
Charlotte-Mecklenburg County	100.0	74.4	2.4	20.2	15.0	27.5	34.8
Chicago	44.3	48.5	10.9	25.2	9.9	14.5	39.5
Dallas	82.6	65.4	12.2	14.3	16.3	12.2	44.9
District of Columbia	93.2	87.4	3.4	19.0	15.9	3.4	58.4
Hillsborough County	76.0	48.4	12.4	21.3	27.5	13.4	25.4
Los Angeles	91.7	77.3	3.6	24.7	32.7	11.1	27.9
Memphis	96.4	84.9	4.7	23.2	14.3	20.7	37.2
Miami	73.4	54.1	12.9	22.1	18.1	14.4	32.4
Orange County	54.2	73.3	8.9	28.8	5.9	17.8	38.6
Philadelphia	75.3	91.7	3.1	12.6	6.1	15.1	63.1
San Diego	0.0	57.8	22.3	48.5	1.9	0.0	27.4
San Francisco	52.2	35.3	29.4	35.3	17.7	8.8	8.8
Local Median	75.7	69.4	9.9	22.7	15.5	13.9	36.0
Local Range	0.0 – 100.0	35.3 – 91.7	2.4 – 29.4	12.6 – 48.5	1.9 – 32.7	0.0 – 27.5	8.8 – 63.1

* Certification, licensure, or endorsement by the state.

† Data not available.

‡ Survey did not include schools from the New York City Department of Education.

TABLE 19a. Percentage of All Schools in Which the Lead Health Education Teacher Received Staff Development During the 2 Years Preceding the Survey on Specific Health Topics, Selected U.S. Sites: School Health Profiles, Lead Health Education Teacher Surveys, 2006

Site	Alcohol-use or other drug-use prevention	Asthma awareness	Consumer health	CPR*	Dental and oral health	Emotional and mental health	Environmental health
STATE SURVEYS							
Alabama	56.0	30.8	27.5	85.4	14.6	36.9	21.4
Alaska	47.9	14.9	15.4	40.9	13.5	33.1	15.5
Arizona	46.6	24.9	21.4	61.1	19.5	32.8	19.5
Arkansas	52.3	21.5	27.8	70.7	16.8	37.0	16.6
Connecticut	36.3	15.4	16.2	55.0	6.8	34.4	14.1
Delaware	55.9	12.0	18.1	66.8	4.5	34.9	7.7
Florida	51.5	22.1	24.7	63.4	15.5	38.5	18.8
Georgia	49.8	24.3	21.9	72.8	15.1	32.8	15.3
Hawaii	61.8	17.9	42.2	29.6	13.7	47.5	16.0
Idaho	64.6	15.5	35.8	70.1	12.1	48.5	20.5
Iowa	36.3	11.0	17.6	51.2	6.9	25.3	10.9
Kansas	40.7	16.0	20.2	63.7	11.8	27.9	13.2
Maine	41.5	16.6	22.6	53.5	12.7	41.3	14.2
Massachusetts	47.6	15.1	15.6	63.7	11.0	41.6	11.4
Michigan	40.8	24.9	18.0	55.2	6.6	30.1	11.3
Mississippi	61.2	25.1	40.9	60.9	22.7	39.3	20.5
Missouri	49.3	26.2	22.3	68.8	13.8	32.1	11.9
Montana	70.0	13.8	24.1	71.4	9.5	38.9	13.4
Nebraska	49.1	57.4	16.2	65.3	7.7	27.1	9.8
New Hampshire	68.1	17.4	35.8	68.2	13.2	62.3	19.9
New York [†]	55.2	13.0	24.0	73.0	6.2	47.0	15.1
North Carolina	42.5	30.6	18.9	67.7	9.6	25.6	12.7
North Dakota	57.7	17.6	27.8	59.3	10.2	36.3	13.5
Oregon	56.1	21.1	17.9	74.8	13.5	38.0	15.9
Pennsylvania	52.3	23.6	19.8	78.6	7.4	40.0	10.2
Rhode Island	37.7	14.8	14.0	67.1	3.5	26.6	9.3
South Carolina	37.5	17.5	22.8	66.5	14.4	25.3	16.0
South Dakota	45.3	17.8	20.1	69.4	11.4	23.8	9.6
Tennessee	53.6	28.6	26.8	69.0	20.1	41.2	18.5
Texas	46.6	21.7	23.8	82.6	13.6	27.4	17.4
Utah	63.0	19.0	23.1	68.2	7.8	43.4	7.8
Vermont	67.2	22.5	19.2	53.5	12.5	42.6	13.3
Virginia	47.8	24.0	23.4	78.0	8.4	31.9	13.5
West Virginia	50.9	19.4	22.0	55.0	14.7	28.8	15.2
State Median	50.4	19.2	22.2	67.0	12.3	35.6	14.2
State Range	36.3 – 70.0	11.0 – 57.4	14.0 – 42.2	29.6 – 85.4	3.5 – 22.7	23.8 – 62.3	7.7 – 21.4

TABLE 19a. Percentage of All Schools in Which the Lead Health Education Teacher Received Staff Development During the 2 Years Preceding the Survey on Specific Health Topics, Selected U.S. Sites: School Health Profiles, Lead Health Education Teacher Surveys, 2006 (continued)

Site	Alcohol-use or other drug-use prevention	Asthma awareness	Consumer health	CPR*	Dental and oral health	Emotional and mental health	Environmental health
LOCAL SURVEYS							
Charlotte-Mecklenburg County	67.2	54.9	43.7	77.1	15.0	50.4	22.9
Chicago	39.1	48.0	25.8	40.6	29.8	35.5	24.3
Dallas	66.0	13.7	28.0	62.7	13.5	33.3	24.0
District of Columbia	44.5	34.9	35.5	38.2	29.3	48.6	38.9
Hillsborough County	39.0	32.7	17.2	76.6	5.0	18.4	14.5
Los Angeles	78.2	35.1	24.7	48.1	15.1	48.1	29.1
Memphis	71.7	67.4	53.3	88.0	37.3	54.5	38.5
Miami	42.0	11.3	29.7	59.7	13.8	27.7	17.8
Orange County	62.9	5.9	17.4	56.4	14.2	42.4	14.5
Philadelphia	41.7	68.0	19.4	87.9	14.6	32.8	11.8
San Diego	100.0	11.2	40.9	43.0	5.4	86.7	7.5
San Francisco	87.9	21.2	27.3	39.4	12.5	56.2	9.1
Local Median	64.5	33.8	27.7	58.1	14.4	45.3	20.4
Local Range	39.0 – 100.0	5.9 – 68.0	17.2 – 53.3	38.2 – 88.0	5.0 – 37.3	18.4 – 86.7	7.5 – 38.9

* Cardiopulmonary resuscitation.

† Survey did not include schools from the New York City Department of Education.

TABLE 19b. Percentage of All Schools in Which the Lead Health Education Teacher Received Staff Development During the 2 Years Preceding the Survey on Specific Health Topics, Selected U.S. Sites: School Health Profiles, Lead Health Education Teacher Surveys, 2006

Site	First aid	Foodborne illness prevention	Growth and development	HIV* prevention	Human sexuality	Immunizations	Injury prevention and safety
STATE SURVEYS							
Alabama	77.1	24.3	25.8	54.7	32.3	21.3	58.6
Alaska	46.6	20.1	20.7	25.7	15.5	22.8	30.5
Arizona	55.5	22.1	27.5	32.4	19.3	30.7	43.0
Arkansas	66.8	15.3	21.9	30.6	19.8	16.9	52.5
Connecticut	50.7	13.6	23.4	39.0	29.5	10.8	34.3
Delaware	48.6	9.2	24.2	39.5	39.6	13.6	25.9
Florida	59.4	18.9	31.4	56.3	36.9	21.5	43.7
Georgia	60.4	15.1	31.9	50.0	38.4	15.6	42.6
Hawaii	29.1	19.5	25.6	53.4	46.2	7.4	36.8
Idaho	65.8	16.3	21.3	48.3	35.9	15.2	44.4
Iowa	41.6	21.1	17.9	32.6	18.1	20.9	26.3
Kansas	55.3	19.5	25.1	43.9	42.0	16.9	33.0
Maine	45.7	15.4	29.1	43.4	35.9	11.7	36.1
Massachusetts	45.8	13.6	26.1	28.9	29.5	12.0	33.4
Michigan	47.5	17.0	29.8	57.2	48.9	8.2	29.2
Mississippi	52.8	25.6	27.1	41.0	30.9	24.7	48.2
Missouri	66.0	21.9	27.9	34.3	23.3	18.4	49.9
Montana	72.2	21.3	23.6	44.8	27.1	13.5	47.9
Nebraska	50.6	17.4	20.8	29.5	24.6	18.2	29.2
New Hampshire	57.9	21.1	49.8	63.9	65.6	13.2	43.9
New York [†]	61.2	13.1	28.2	52.5	39.0	10.9	38.3
North Carolina	64.2	18.9	24.1	48.2	37.1	12.9	44.8
North Dakota	48.3	30.2	22.6	32.5	20.4	21.1	31.3
Oregon	75.1	21.8	22.7	48.7	39.9	12.2	41.8
Pennsylvania	59.1	9.2	20.7	37.8	26.6	9.2	41.4
Rhode Island	53.5	20.9	19.4	24.3	12.7	16.2	36.7
South Carolina	64.2	19.1	30.8	51.3	41.9	18.1	42.3
South Dakota	47.5	14.3	18.1	21.3	15.8	17.0	25.0
Tennessee	66.0	19.3	26.6	51.0	25.1	22.8	52.7
Texas	71.7	18.3	25.9	35.0	29.9	20.5	52.5
Utah	61.4	14.9	29.5	53.6	60.5	11.4	35.9
Vermont	41.6	19.3	33.3	44.2	40.6	18.0	28.3
Virginia	64.9	16.5	30.8	45.0	41.2	18.1	44.4
West Virginia	46.5	14.8	22.4	24.0	18.9	12.5	30.2
State Median	56.7	18.9	25.7	43.7	31.6	16.6	39.9
State Range	29.1 – 77.1	9.2 – 30.2	17.9 – 49.8	21.3 – 63.9	12.7 – 65.6	7.4 – 30.7	25.0 – 58.6

TABLE 19b. Percentage of All Schools in Which the Lead Health Education Teacher Received Staff Development During the 2 Years Preceding the Survey on Specific Health Topics, Selected U.S. Sites: School Health Profiles, Lead Health Education Teacher Surveys, 2006 (continued)

Site	First aid	Foodborne illness prevention	Growth and development	HIV* prevention	Human sexuality	Immunizations	Injury prevention and safety
LOCAL SURVEYS							
Charlotte-Mecklenburg County	79.8	20.0	57.3	62.4	79.6	17.6	67.4
Chicago	44.0	21.2	35.6	42.9	33.4	33.9	41.3
Dallas	51.9	26.0	65.4	61.5	65.4	19.6	53.8
District of Columbia	47.2	23.7	48.9	65.9	50.3	29.3	38.9
Hillsborough County	63.1	14.6	26.3	58.9	32.2	16.9	34.4
Los Angeles	41.7	22.1	37.6	82.5	62.1	25.6	43.3
Memphis	90.0	38.5	54.0	89.2	71.0	36.9	75.8
Miami	49.5	17.6	49.5	69.2	43.2	22.5	37.6
Orange County	53.4	14.2	45.4	88.1	76.6	22.8	42.4
Philadelphia	71.3	13.6	34.5	52.4	44.1	14.0	50.4
San Diego	24.3	14.6	63.0	100.0	100.0	36.6	24.5
San Francisco	36.3	6.1	21.2	60.6	66.7	6.1	34.4
Local Median	50.7	18.8	47.2	64.2	63.8	22.7	41.9
Local Range	24.3 – 90.0	6.1 – 38.5	21.2 – 65.4	42.9 – 100.0	32.2 – 100.0	6.1 – 36.9	24.5 – 75.8

* Human immunodeficiency virus.

† Survey did not include schools from the New York City Department of Education.

TABLE 19c. Percentage of All Schools in Which the Lead Health Education Teacher Received Staff Development During the 2 Years Preceding the Survey on Specific Health Topics, Selected U.S. Sites: School Health Profiles, Lead Health Education Teacher Surveys, 2006

Site	Nutrition and dietary behavior	Physical activity and fitness	Pregnancy prevention	STD* prevention	Suicide prevention	Sun safety or skin cancer prevention	Tobacco-use prevention	Violence prevention
STATE SURVEYS								
Alabama	35.4	52.4	34.4	44.0	24.0	17.4	35.1	59.7
Alaska	21.3	29.9	12.6	21.5	34.5	6.8	26.0	50.8
Arizona	30.5	39.6	16.3	23.2	25.9	26.8	37.9	59.4
Arkansas	39.9	56.7	22.7	26.0	28.3	14.2	49.0	60.9
Connecticut	35.1	44.0	22.8	32.2	20.4	8.9	21.1	45.9
Delaware	27.4	51.7	36.5	39.6	15.3	7.6	40.7	51.5
Florida	46.0	48.1	34.9	46.3	31.6	26.9	36.1	57.0
Georgia	30.4	49.7	39.6	48.4	25.7	14.1	34.2	45.4
Hawaii	52.6	58.9	46.0	47.1	25.2	15.6	46.8	56.3
Idaho	46.9	48.1	26.9	42.6	39.6	16.0	41.6	54.0
Iowa	30.3	27.5	15.0	23.0	15.5	8.5	16.7	46.7
Kansas	39.1	54.2	29.8	39.4	20.6	10.5	26.3	50.1
Maine	40.9	45.7	30.1	36.7	39.1	20.9	32.9	49.0
Massachusetts	43.9	45.1	22.0	26.8	28.4	43.1	25.7	61.2
Michigan	35.3	42.6	37.0	50.3	20.5	9.1	26.4	47.7
Mississippi	47.4	42.6	33.4	37.4	27.3	19.6	46.4	45.5
Missouri	40.2	53.2	21.3	30.8	24.6	14.2	32.0	52.9
Montana	34.5	51.9	23.4	33.2	30.7	10.3	39.4	58.6
Nebraska	30.3	37.7	23.0	28.8	22.4	11.0	27.7	51.6
New Hampshire	72.8	64.6	55.7	64.8	25.8	21.1	45.0	49.8
New York†	44.9	41.4	32.1	41.9	27.0	13.1	38.9	64.5
North Carolina	33.0	58.4	35.6	43.9	17.9	17.3	37.7	51.5
North Dakota	40.8	44.9	13.6	29.6	33.9	12.0	42.4	57.6
Oregon	27.5	35.2	35.0	45.7	30.3	10.0	35.0	57.5
Pennsylvania	43.5	58.0	21.2	30.7	22.1	8.4	31.5	56.3
Rhode Island	27.5	39.8	10.3	17.4	14.8	8.1	19.4	45.1
South Carolina	27.5	56.8	36.3	41.6	11.4	12.2	26.2	41.5
South Dakota	34.6	34.4	11.3	18.0	21.7	9.4	25.6	40.7
Tennessee	34.8	51.0	24.5	35.9	31.4	14.3	31.4	62.8
Texas	28.8	48.4	25.7	31.8	21.0	18.2	30.2	49.0
Utah	38.1	51.7	35.2	47.6	30.0	15.8	49.7	47.6
Vermont	46.4	45.6	28.0	37.9	24.2	11.6	43.0	70.1
Virginia	36.1	64.4	28.8	36.6	38.5	13.1	30.8	67.2
West Virginia	29.4	60.4	22.4	22.8	14.2	13.7	49.0	54.0
State Median	35.4	48.3	27.5	36.7	25.5	13.4	34.6	52.3
State Range	21.3 – 72.8	27.5 – 64.6	10.3 – 55.7	17.4 – 64.8	11.4 – 39.6	6.8 – 43.1	16.7 – 49.7	40.7 – 70.1

TABLE 19c. Percentage of All Schools in Which the Lead Health Education Teacher Received Staff Development During the 2 Years Preceding the Survey on Specific Health Topics, Selected U.S. Sites: School Health Profiles, Lead Health Education Teacher Surveys, 2006 (continued)

Site	Nutrition and dietary behavior	Physical activity and fitness	Pregnancy prevention	STD* prevention	Suicide prevention	Sun safety or skin cancer prevention	Tobacco-use prevention	Violence prevention
LOCAL SURVEYS								
Charlotte-Mecklenburg County	50.4	77.3	62.5	60.1	39.0	46.1	37.6	77.3
Chicago	38.6	54.2	30.4	35.4	21.3	11.8	34.7	58.7
Dallas	41.2	32.7	42.0	61.5	28.8	19.6	50.0	61.5
District of Columbia	45.5	48.6	39.2	56.9	29.3	22.4	41.2	55.1
Hillsborough County	32.4	47.7	22.9	37.6	16.3	12.6	28.1	55.0
Los Angeles	40.8	34.6	53.8	70.8	32.6	17.2	85.5	69.9
Memphis	51.1	81.2	64.0	89.1	42.9	29.8	51.1	76.6
Miami	35.1	46.1	43.2	57.3	25.0	26.3	29.8	44.0
Orange County	48.1	40.1	65.6	79.8	36.5	23.1	40.1	60.2
Philadelphia	63.2	80.3	27.2	42.0	20.3	11.7	35.8	68.0
San Diego	21.8	23.6	92.8	100.0	82.8	10.7	100.0	81.4
San Francisco	63.6	39.4	41.9	54.6	34.4	6.1	87.9	72.7
Local Median	43.4	46.9	42.6	58.7	31.0	18.4	40.7	64.8
Local Range	21.8 – 63.6	23.6 – 81.2	22.9 – 92.8	35.4 – 100.0	16.3 – 82.8	6.1 – 46.1	28.1 – 100.0	44.0 – 81.4

* Sexually transmitted disease.

† Survey did not include schools from the New York City Department of Education.

TABLE 20a. Percentage of All Schools in Which the Lead Health Education Teacher Wanted to Receive Staff Development on Specific Health Topics, Selected U.S. Sites: School Health Profiles, Lead Health Education Teacher Surveys, 2006

Site	Alcohol-use or other drug-use prevention	Asthma awareness	Consumer health	CPR*	Dental and oral health	Emotional and mental health	Environmental health	First aid
STATE SURVEYS								
Alabama	76.5	68.3	53.5	68.4	55.1	70.6	50.3	67.6
Alaska	67.8	54.1	48.6	64.9	45.3	71.6	52.7	64.9
Arizona	70.5	60.3	48.5	66.9	42.8	64.8	52.3	67.4
Arkansas	79.7	61.7	54.9	71.4	49.8	66.4	52.4	79.6
Connecticut	73.3	56.6	62.2	55.9	41.7	78.7	55.7	56.3
Delaware	78.4	54.3	44.6	59.5	35.6	71.4	44.5	67.0
Florida	74.3	59.5	57.0	70.1	49.9	71.9	55.4	70.1
Georgia	79.2	63.6	50.1	71.1	49.0	67.3	55.7	76.0
Hawaii	69.3	58.9	64.6	69.0	53.9	64.5	61.5	67.8
Idaho	73.1	53.8	47.6	65.3	43.4	69.7	52.5	66.6
Iowa	60.5	44.2	41.9	60.4	28.6	62.2	42.5	57.0
Kansas	70.3	50.4	42.9	63.9	34.6	62.6	39.5	60.7
Maine	60.9	42.9	47.1	53.8	32.1	63.7	42.8	59.1
Massachusetts	78.8	55.8	59.3	65.2	43.8	78.8	57.4	67.7
Michigan	67.8	53.6	51.2	62.5	39.1	67.4	48.1	63.8
Mississippi	79.7	64.4	58.8	78.4	57.4	71.7	58.6	78.8
Missouri	68.8	51.3	48.0	67.1	42.0	65.5	45.3	67.8
Montana	71.8	61.2	44.7	61.0	43.4	70.7	46.9	63.2
Nebraska	61.5	43.5	36.1	55.3	33.7	55.6	37.7	54.6
New Hampshire	76.4	56.6	62.3	51.8	46.4	78.5	65.9	54.5
New York [†]	81.4	54.1	62.8	60.4	41.7	79.4	59.8	60.1
North Carolina	70.0	64.4	47.2	69.5	42.2	62.3	45.5	72.8
North Dakota	55.9	51.8	44.2	63.5	30.4	63.3	37.1	61.4
Oregon	65.0	44.7	44.7	48.9	38.5	67.4	54.2	52.0
Pennsylvania	78.2	61.7	62.4	68.6	46.0	76.1	58.1	71.0
Rhode Island	74.6	56.4	65.5	54.0	35.0	76.6	57.6	56.5
South Carolina	70.5	57.6	45.7	67.2	45.3	64.8	48.4	61.3
South Dakota	54.7	47.9	48.3	61.4	34.4	57.7	37.2	63.8
Tennessee	76.5	61.2	54.4	72.2	50.4	72.1	54.1	76.1
Texas	74.0	57.3	51.0	67.2	46.9	69.4	55.6	69.1
Utah	73.5	51.0	55.6	63.2	39.6	80.2	53.3	65.2
Vermont	53.7	33.0	46.3	49.0	27.0	61.4	42.0	50.2
Virginia	63.1	62.3	44.3	61.1	37.4	61.0	43.6	59.9
West Virginia	73.4	65.9	53.0	74.3	45.7	63.3	56.2	76.7
State Median	72.5	56.5	49.4	64.4	42.5	67.4	52.5	65.1
State Range	53.7 – 81.4	33.0 – 68.3	36.1 – 65.5	48.9 – 78.4	27.0 – 57.4	55.6 – 80.2	37.1 – 65.9	50.2 – 79.6

TABLE 20a. Percentage of All Schools in Which the Lead Health Education Teacher Wanted to Receive Staff Development on Specific Health Topics, Selected U.S. Sites: School Health Profiles, Lead Health Education Teacher Surveys, 2006 (continued)

Site	Alcohol-use or other drug-use prevention	Asthma awareness	Consumer health	CPR*	Dental and oral health	Emotional and mental health	Environmental health	First aid
LOCAL SURVEYS								
Charlotte-Mecklenburg County	74.5	72.0	61.8	74.5	51.1	77.4	61.6	67.0
Chicago	67.9	71.4	59.7	83.5	57.0	77.9	56.7	82.9
Dallas	80.4	66.7	56.3	75.0	51.0	76.9	62.7	79.6
District of Columbia	81.2	84.1	74.1	84.1	70.7	83.5	80.6	84.4
Hillsborough County	48.3	35.3	35.2	66.4	33.6	62.9	39.4	67.8
Los Angeles	73.3	65.6	66.6	70.4	56.1	77.0	68.5	72.8
Memphis	85.9	85.2	74.5	84.6	76.3	82.9	71.1	81.5
Miami	73.1	62.7	57.5	76.5	53.7	73.2	58.5	74.4
Orange County	73.9	46.3	46.0	62.9	48.4	68.5	51.3	63.5
Philadelphia	76.3	80.2	67.0	78.6	62.0	79.5	72.6	76.5
San Diego	81.8	50.1	60.9	60.6	39.1	87.4	38.6	56.8
San Francisco	77.4	46.9	56.3	60.6	45.4	87.5	60.6	71.9
Local Median	75.4	66.2	60.3	74.8	52.4	77.7	61.1	73.6
Local Range	48.3 – 85.9	35.3 – 85.2	35.2 – 74.5	60.6 – 84.6	33.6 – 76.3	62.9 – 87.5	38.6 – 80.6	56.8 – 84.4

* Cardiopulmonary resuscitation.

† Survey did not include schools from the New York City Department of Education.

TABLE 20b. Percentage of All Schools in Which the Lead Health Education Teacher Wanted to Receive Staff Development on Specific Health Topics, Selected U.S. Sites: School Health Profiles, Lead Health Education Teacher Surveys, 2006

Site	Foodborne illness prevention	Growth and development	HIV* prevention	Human sexuality	Immunizations	Injury prevention and safety	Nutrition and dietary behavior
STATE SURVEYS							
Alabama	53.4	56.8	68.3	59.4	53.1	69.1	74.8
Alaska	48.1	55.4	57.3	53.5	44.5	58.0	66.0
Arizona	45.0	51.8	55.9	48.2	42.1	61.9	64.7
Arkansas	55.7	59.1	66.2	56.9	47.3	72.4	75.9
Connecticut	54.6	64.6	69.7	68.4	45.6	55.1	76.7
Delaware	44.7	59.6	66.3	71.4	41.6	56.6	72.6
Florida	59.8	61.4	69.3	66.2	53.9	65.2	73.2
Georgia	57.9	60.9	70.9	61.5	46.4	70.6	74.4
Hawaii	61.4	57.3	54.6	55.1	59.5	53.0	68.3
Idaho	49.7	54.8	64.3	59.2	47.4	58.5	74.9
Iowa	38.5	45.9	55.0	50.3	38.2	47.7	62.2
Kansas	40.4	55.9	60.0	58.3	40.8	53.3	71.7
Maine	37.0	43.7	46.1	46.3	31.4	50.0	62.2
Massachusetts	53.3	63.1	70.7	75.0	45.9	62.5	76.5
Michigan	47.0	50.0	57.6	57.6	43.3	57.3	67.3
Mississippi	61.0	59.3	74.9	60.3	57.5	70.7	68.0
Missouri	44.7	52.8	63.4	55.1	44.1	62.3	69.2
Montana	49.8	56.6	62.4	56.1	50.3	62.4	73.9
Nebraska	35.8	40.7	54.1	49.3	34.9	46.2	57.3
New Hampshire	53.7	60.7	62.1	66.0	50.8	67.3	78.0
New York [†]	51.1	66.0	74.7	73.0	52.0	55.8	76.1
North Carolina	48.2	52.9	61.7	55.5	43.4	66.3	75.2
North Dakota	42.7	45.0	54.4	53.0	37.3	48.2	60.3
Oregon	42.5	54.1	57.8	56.1	40.4	47.1	67.1
Pennsylvania	56.8	67.0	77.4	75.4	53.0	65.0	78.5
Rhode Island	51.8	55.9	70.2	66.7	44.4	62.0	77.9
South Carolina	46.2	51.6	64.2	55.7	42.7	56.1	74.2
South Dakota	37.1	39.4	48.4	40.9	33.8	51.1	66.4
Tennessee	55.4	59.5	63.5	54.6	48.3	69.5	73.2
Texas	58.3	58.1	65.0	59.6	49.1	63.5	74.0
Utah	50.5	57.4	66.7	66.0	47.1	60.7	78.5
Vermont	36.4	41.5	46.2	47.5	30.7	42.0	64.9
Virginia	45.7	47.3	47.2	45.2	41.1	55.1	72.7
West Virginia	59.4	53.2	64.7	60.0	49.0	65.7	76.6
State Median	49.8	55.9	63.5	57.3	45.1	59.6	73.2
State Range	35.8 – 61.4	39.4 – 67.0	46.1 – 77.4	40.9 – 75.4	30.7 – 59.5	42.0 – 72.4	57.3 – 78.5

TABLE 20b. Percentage of All Schools in Which the Lead Health Education Teacher Wanted to Receive Staff Development on Specific Health Topics, Selected U.S. Sites: School Health Profiles, Lead Health Education Teacher Surveys, 2006 (continued)

Site	Foodborne illness prevention	Growth and development	HIV* prevention	Human sexuality	Immunizations	Injury prevention and safety	Nutrition and dietary behavior
LOCAL SURVEYS							
Charlotte-Mecklenburg County	61.4	64.3	67.0	56.4	51.2	64.1	77.2
Chicago	61.1	65.4	66.9	62.8	57.8	67.9	73.4
Dallas	68.6	65.4	76.0	72.0	60.8	59.2	76.9
District of Columbia	80.6	77.3	83.8	80.6	73.8	77.3	81.2
Hillsborough County	41.9	47.8	60.5	50.5	41.5	48.3	58.6
Los Angeles	68.2	66.8	72.5	72.6	67.8	56.5	71.8
Memphis	72.6	83.8	84.6	79.5	80.3	78.6	83.1
Miami	59.8	61.0	70.7	67.2	53.7	71.5	79.3
Orange County	51.6	47.3	62.6	62.6	33.8	49.0	71.2
Philadelphia	72.7	76.1	78.7	78.3	66.7	75.7	77.0
San Diego	42.5	51.6	69.4	63.9	44.0	42.5	58.5
San Francisco	39.4	56.2	63.6	71.0	48.4	64.5	75.7
Local Median	61.3	64.8	70.1	69.1	55.8	64.3	76.3
Local Range	39.4 – 80.6	47.3 – 83.8	60.5 – 84.6	50.5 – 80.6	33.8 – 80.3	42.5 – 78.6	58.5 – 83.1

* Human immunodeficiency virus.

† Survey did not include schools from the New York City Department of Education.

TABLE 20c. Percentage of All Schools in Which the Lead Health Education Teacher Wanted to Receive Staff Development on Specific Health Topics, Selected U.S. Sites: School Health Profiles, Lead Health Education Teacher Surveys, 2006

Site	Physical activity and fitness	Pregnancy prevention	STD* prevention	Suicide prevention	Sun safety or skin cancer prevention	Tobacco-use prevention	Violence prevention
STATE SURVEYS							
Alabama	71.1	62.7	71.2	75.5	60.3	70.2	79.6
Alaska	60.3	53.5	55.8	72.3	42.2	63.1	68.2
Arizona	57.6	53.0	54.7	66.1	59.8	60.3	72.5
Arkansas	78.5	59.6	67.4	77.9	58.6	74.3	77.9
Connecticut	62.3	62.6	68.4	77.6	58.0	65.5	76.1
Delaware	68.4	61.9	66.7	69.9	47.5	66.9	83.5
Florida	68.7	65.6	68.0	72.9	66.1	65.6	75.3
Georgia	72.1	62.1	68.3	74.8	60.9	71.5	82.3
Hawaii	58.7	59.2	56.4	72.4	62.8	64.6	81.3
Idaho	69.4	55.9	60.7	74.6	53.0	62.9	79.3
Iowa	60.1	52.0	54.4	69.3	47.9	53.2	67.0
Kansas	68.8	57.3	64.1	68.9	49.7	63.9	70.7
Maine	52.7	44.6	52.6	65.9	44.0	50.3	66.8
Massachusetts	69.7	66.6	74.0	82.4	57.3	64.0	80.9
Michigan	64.8	53.0	57.1	68.8	51.3	57.3	70.7
Mississippi	70.3	71.3	74.4	80.5	65.0	70.8	83.5
Missouri	66.6	58.1	62.5	69.9	49.9	59.4	74.9
Montana	71.5	57.1	61.5	71.7	50.0	59.9	72.1
Nebraska	53.8	49.6	53.3	63.2	40.0	57.3	66.8
New Hampshire	68.6	58.8	68.3	73.9	64.9	64.8	76.7
New York [†]	69.9	69.2	75.0	85.2	60.3	65.7	83.6
North Carolina	75.4	57.6	62.2	68.8	56.7	64.2	76.4
North Dakota	60.4	42.8	55.8	71.0	46.7	51.9	69.0
Oregon	58.8	51.9	55.8	67.6	50.8	53.5	67.2
Pennsylvania	75.8	72.3	77.7	81.8	60.1	73.3	78.6
Rhode Island	64.4	63.4	63.2	72.3	55.1	51.9	81.4
South Carolina	71.1	57.4	61.9	68.3	50.8	59.9	76.6
South Dakota	60.9	39.5	47.9	62.6	41.7	46.6	65.0
Tennessee	75.0	59.3	63.6	74.5	58.4	67.0	81.3
Texas	65.7	58.3	63.0	76.5	57.8	65.8	77.7
Utah	67.9	54.8	65.2	74.2	57.9	61.0	79.5
Vermont	50.9	46.6	47.9	64.4	42.6	45.1	58.7
Virginia	67.7	46.2	46.2	63.8	53.5	56.2	73.9
West Virginia	67.3	65.3	65.2	75.9	60.9	64.4	74.0
State Median	67.8	57.9	62.8	72.3	55.9	63.5	76.3
State Range	50.9 – 78.5	39.5 – 72.3	46.2 – 77.7	62.6 – 85.2	40.0 – 66.1	45.1 – 74.3	58.7 – 83.6

TABLE 20c. Percentage of All Schools in Which the Lead Health Education Teacher Wanted to Receive Staff Development on Specific Health Topics, Selected U.S. Sites: School Health Profiles, Lead Health Education Teacher Surveys, 2006 (continued)

Site	Physical activity and fitness	Pregnancy prevention	STD* prevention	Suicide prevention	Sun safety or skin cancer prevention	Tobacco-use prevention	Violence prevention
LOCAL SURVEYS							
Charlotte-Mecklenburg County	61.6	56.4	64.3	71.8	61.6	66.6	81.9
Chicago	68.9	62.1	68.1	74.4	57.9	65.9	85.7
Dallas	65.3	68.6	74.0	78.8	52.9	66.0	78.4
District of Columbia	83.8	83.8	83.8	78.6	73.8	83.8	93.7
Hillsborough County	46.8	49.3	55.1	62.9	61.3	52.4	63.4
Los Angeles	65.7	69.0	71.4	82.9	58.6	58.8	83.2
Memphis	81.8	84.4	88.0	86.3	69.0	74.1	83.1
Miami	75.6	63.9	64.7	70.8	64.5	58.5	69.8
Orange County	51.3	65.3	68.5	73.6	49.7	61.6	73.6
Philadelphia	73.6	78.3	75.2	85.9	69.6	71.4	80.2
San Diego	55.0	72.7	72.6	96.3	42.8	68.8	94.8
San Francisco	71.0	65.6	66.7	75.8	43.7	51.5	87.9
Local Median	67.3	67.1	70.0	77.2	60.0	66.0	82.5
Local Range	46.8 – 83.8	49.3 – 84.4	55.1 – 88.0	62.9 – 96.3	42.8 – 73.8	51.5 – 83.8	63.4 – 94.8

* Sexually transmitted disease.

† Survey did not include schools from the New York City Department of Education.

TABLE 21. Percentage of All Schools in Which the Lead Health Education Teacher Received Staff Development During the 2 Years Preceding the Survey on Specific Topics, Selected U.S. Sites: School Health Profiles, Lead Health Education Teacher Surveys, 2006

Site	Teaching students with physical or cognitive disabilities	Teaching students of various cultural backgrounds	Teaching students with limited English proficiency	Using interactive teaching methods	Encouraging family or community involvement	Teaching skills for behavior change	Classroom management techniques	Assessing or evaluating students in health education
STATE SURVEYS								
Alabama	63.0	46.4	42.5	50.5	50.6	56.0	72.9	30.6
Alaska	55.3	65.9	40.2	50.8	50.3	43.2	50.4	28.2
Arizona	59.1	63.8	77.9	61.3	47.5	51.7	65.1	27.0
Arkansas	54.3	49.5	33.4	56.1	72.0	59.1	75.7	39.6
Connecticut	53.4	33.2	18.6	47.1	26.1	46.2	55.2	49.2
Delaware	46.3	47.8	25.4	50.8	32.8	46.1	59.7	46.4
Florida	52.6	60.6	58.8	66.1	47.1	58.9	70.9	30.1
Georgia	38.8	38.8	28.8	53.4	33.6	39.1	53.1	27.4
Hawaii	36.6	42.4	32.5	54.2	31.0	43.0	53.0	66.7
Idaho	43.3	38.4	39.5	46.3	35.1	41.8	45.8	26.5
Iowa	52.9	43.9	26.3	55.6	38.7	50.5	56.3	21.0
Kansas	39.0	35.9	21.4	49.7	38.0	40.9	48.7	24.9
Maine	39.2	12.5	7.6	46.1	27.5	37.8	39.4	58.3
Massachusetts	50.9	37.8	34.1	44.9	30.1	44.4	54.2	36.7
Michigan	41.2	30.3	17.4	52.2	33.8	43.3	53.5	24.9
Mississippi	52.9	40.4	22.3	56.5	49.3	54.3	73.0	36.6
Missouri	52.0	42.9	21.1	63.1	53.4	56.3	72.8	37.4
Montana	43.6	33.0	8.8	45.2	35.8	52.7	57.2	34.6
Nebraska	35.1	30.6	20.2	40.1	29.1	39.5	50.3	21.4
New Hampshire	64.9	20.6	8.8	68.0	43.1	61.1	63.4	60.7
New York*	52.7	34.5	16.6	56.2	36.5	52.7	56.7	39.6
North Carolina	47.0	47.1	31.3	52.0	35.2	45.7	63.7	32.2
North Dakota	49.5	25.3	13.4	43.5	33.9	43.2	56.6	15.2
Oregon	49.7	54.8	53.8	54.0	37.2	52.2	57.0	33.2
Pennsylvania	54.0	35.3	33.0	46.0	30.7	45.0	58.1	35.3
Rhode Island	37.0	26.9	18.6	44.5	39.4	32.8	47.9	31.2
South Carolina	39.5	44.9	29.1	51.2	42.2	46.1	64.4	30.0
South Dakota	43.4	22.7	9.5	35.7	32.2	31.6	49.3	22.7
Tennessee	48.3	41.8	25.4	59.6	53.6	49.6	66.1	30.3
Texas	58.9	52.1	51.0	54.9	44.5	56.3	70.9	30.7
Utah	43.4	46.9	49.8	55.7	35.0	42.3	53.2	30.2
Vermont	52.2	24.4	13.0	54.8	38.2	54.9	58.2	55.0
Virginia	51.8	42.5	29.3	61.3	38.2	46.2	62.8	31.6
West Virginia	48.4	37.1	11.0	52.0	35.4	48.4	65.9	39.1
State Median	49.6	39.6	25.9	52.1	36.9	46.2	57.1	31.4
State Range	35.1 – 64.9	12.5 – 65.9	7.6 – 77.9	35.7 – 68.0	26.1 – 72.0	31.6 – 61.1	39.4 – 75.7	15.2 – 66.7

TABLE 21. Percentage of All Schools in Which the Lead Health Education Teacher Received Staff Development During the 2 Years Preceding the Survey on Specific Topics, Selected U.S. Sites: School Health Profiles, Lead Health Education Teacher Surveys, 2006 (continued)

Site	Teaching students with physical or cognitive disabilities	Teaching students of various cultural backgrounds	Teaching students with limited English proficiency	Using interactive teaching methods	Encouraging family or community involvement	Teaching skills for behavior change	Classroom management techniques	Assessing or evaluating students in health education
LOCAL SURVEYS								
Charlotte-Mecklenburg County	57.2	69.3	48.9	89.8	61.6	61.4	74.5	82.2
Chicago	66.2	50.7	39.3	67.6	54.9	62.6	67.8	29.0
Dallas	57.7	59.6	50.0	73.1	51.9	61.5	90.2	34.6
District of Columbia	40.7	21.7	18.7	59.6	37.4	48.4	49.7	40.4
Hillsborough County	41.5	63.6	63.5	52.3	42.7	42.9	59.0	30.0
Los Angeles	73.6	77.5	85.6	68.7	54.0	62.0	67.0	37.8
Memphis	39.0	41.4	27.0	69.9	61.8	64.0	77.6	53.9
Miami	39.0	45.2	64.7	62.3	41.7	35.8	56.2	29.2
Orange County	35.7	62.6	71.8	54.6	37.7	60.5	55.5	25.8
Philadelphia	24.9	33.8	17.5	52.7	32.6	40.6	54.8	60.5
San Diego	30.0	60.5	48.4	87.4	73.1	78.0	62.9	30.2
San Francisco	67.6	64.7	76.5	60.6	60.6	44.1	60.6	29.4
Local Median	41.1	60.1	49.5	65.0	53.0	61.0	61.8	32.4
Local Range	24.9 – 73.6	21.7 – 77.5	17.5 – 85.6	52.3 – 89.8	32.6 – 73.1	35.8 – 78.0	49.7 – 90.2	25.8 – 82.2

* Survey did not include schools from the New York City Department of Education.

TABLE 22. Percentage of All Schools in Which the Lead Health Education Teacher Wanted to Receive Staff Development on Specific Topics, Selected U.S. Sites: School Health Profiles, Lead Health Education Teacher Surveys, 2006

Site	Teaching students with physical, medical, or cognitive disabilities	Teaching students of various cultural backgrounds	Teaching students with limited English proficiency	Using interactive teaching methods	Encouraging family or community involvement	Teaching skills for behavior change	Classroom management techniques	Assessing or evaluating students in health education
STATE SURVEYS								
Alabama	65.9	55.7	54.8	63.4	65.2	72.1	68.8	68.5
Alaska	60.2	54.7	49.9	58.2	62.5	71.3	65.9	62.5
Arizona	61.7	59.6	56.7	56.5	64.9	69.3	66.8	62.5
Arkansas	69.5	60.6	60.3	64.2	70.0	74.1	73.7	78.0
Connecticut	65.5	62.5	54.7	71.5	70.5	77.9	71.4	77.3
Delaware	72.9	64.1	64.8	67.1	67.0	80.3	68.3	81.8
Florida	64.6	60.3	55.2	64.1	66.5	77.1	68.9	68.0
Georgia	65.1	58.3	59.1	63.2	68.7	73.0	67.3	74.4
Hawaii	72.5	66.8	65.0	66.7	69.0	76.8	66.0	78.0
Idaho	67.2	58.5	55.5	57.8	61.0	72.4	70.5	70.2
Iowa	57.4	46.9	42.7	46.9	60.8	68.6	62.1	60.3
Kansas	56.3	42.3	38.7	55.1	63.9	72.8	64.5	68.6
Maine	55.5	36.5	35.0	54.9	55.9	70.8	64.8	58.6
Massachusetts	69.5	64.6	56.5	72.1	70.8	81.6	73.4	80.7
Michigan	62.1	59.7	46.3	60.7	63.0	72.8	63.2	64.9
Mississippi	63.9	63.8	60.6	63.5	67.1	73.6	70.5	72.6
Missouri	61.3	50.8	44.7	61.4	66.6	71.2	66.2	69.0
Montana	61.5	43.7	34.9	60.9	62.3	79.4	74.0	71.1
Nebraska	49.8	47.3	38.0	44.7	53.6	65.2	64.6	60.8
New Hampshire	64.9	53.4	47.4	67.1	68.2	79.6	70.7	77.1
New York*	68.7	58.0	53.5	69.9	69.9	85.7	73.7	83.2
North Carolina	69.6	63.0	63.3	59.9	63.6	72.7	67.3	68.3
North Dakota	56.0	41.4	30.7	46.3	56.9	73.3	58.9	60.9
Oregon	52.2	50.5	50.4	55.5	58.6	70.1	58.6	63.0
Pennsylvania	71.2	59.3	52.0	70.6	72.4	81.9	75.7	75.8
Rhode Island	68.7	52.8	49.4	66.6	61.6	81.3	68.8	82.4
South Carolina	64.0	59.4	60.4	59.1	64.1	71.8	63.3	72.5
South Dakota	50.1	41.8	31.1	51.8	57.0	68.5	65.2	60.8
Tennessee	72.3	62.3	60.4	62.1	65.9	77.4	73.1	70.2
Texas	61.8	58.4	56.9	63.3	63.2	69.3	63.8	62.2
Utah	63.6	59.9	57.6	66.7	60.4	79.1	69.2	68.4
Vermont	54.1	44.7	34.3	54.8	61.7	68.8	52.4	65.3
Virginia	62.3	53.8	54.1	52.3	53.8	69.3	62.3	66.4
West Virginia	59.9	45.3	42.2	61.1	64.6	72.5	66.8	68.9
State Median	63.8	58.2	53.8	61.3	64.0	72.8	67.1	68.8
State Range	49.8 – 72.9	36.5 – 66.8	30.7 – 65.0	44.7 – 72.1	53.6 – 72.4	65.2 – 85.7	52.4 – 75.7	58.6 – 83.2

TABLE 22. Percentage of All Schools in Which the Lead Health Education Teacher Wanted to Receive Staff Development on Specific Topics, Selected U.S. Sites: School Health Profiles, Lead Health Education Teacher Surveys, 2006 (continued)

Site	Teaching students with physical, medical, or cognitive disabilities	Teaching students of various cultural backgrounds	Teaching students with limited English proficiency	Using interactive teaching methods	Encouraging family or community involvement	Teaching skills for behavior change	Classroom management techniques	Assessing or evaluating students in health education
LOCAL SURVEYS								
Charlotte-Mecklenburg County	71.8	74.3	69.1	64.1	72.0	78.9	74.5	77.0
Chicago	74.3	66.6	60.2	70.1	76.1	78.7	81.3	71.9
Dallas	53.8	71.2	63.5	65.4	72.5	87.8	69.2	55.8
District of Columbia	77.3	77.3	65.7	73.2	76.8	80.7	80.0	74.1
Hillsborough County	45.2	43.6	43.4	46.5	53.0	63.2	58.0	48.9
Los Angeles	62.7	61.5	65.7	66.1	71.7	78.3	74.1	80.0
Memphis	90.2	82.0	81.4	81.9	87.7	90.7	86.5	91.8
Miami	72.9	67.1	54.4	71.3	72.9	77.9	76.7	67.1
Orange County	59.9	51.3	42.7	59.6	50.7	74.5	72.7	62.6
Philadelphia	81.6	78.0	75.5	78.2	75.9	91.0	82.5	73.5
San Diego	80.3	85.8	71.6	77.0	75.6	87.8	89.3	64.7
San Francisco	59.4	65.6	54.8	61.3	71.9	67.8	67.7	78.8
Local Median	72.4	69.2	64.6	68.1	72.7	78.8	75.6	72.7
Local Range	45.2 – 90.2	43.6 – 85.8	42.7 – 81.4	46.5 – 81.9	50.7 – 87.7	63.2 – 91.0	58.0 – 89.3	48.9 – 91.8

* Survey did not include schools from the New York City Department of Education.

TABLE 23. Percentage of All Schools That Required Physical Education in Any of Grades 6–12; the Percentage That Required Students to Take Only One Course or Two or More Courses; Among Schools That Required a Physical Education Course, the Percentage That Required Students Who Fail a Required Physical Education Course to Repeat It; and the Percentage of All Schools in Which a Newly Hired Physical Education Teacher is Required to be Certified* in Physical Education, Selected U.S. Sites: School Health Profiles, Principal Surveys, 2006

Site	Required physical education	Required only one physical education course	Required two or more physical education courses	Required students who fail a required physical education course to repeat it [†]	Newly hired physical education teacher required to be certified in physical education
STATE SURVEYS					
Alabama	98.0	36.6	60.1	62.1	99.2
Alaska	86.1	19.9	65.3	80.6	35.0
Arizona	73.1	31.8	39.1	39.6	67.9
Arkansas	99.2	36.8	62.0	73.1	97.7
Connecticut	100.0	16.7	82.4	48.7	97.9
Delaware	94.1	18.2	75.7	51.4	95.5
Florida	79.1	26.6	51.4	61.9	93.1
Georgia	85.5	41.9	42.9	56.6	98.3
Hawaii	93.1	35.3	57.6	66.5	85.3
Idaho	95.8	17.7	77.5	53.1	94.3
Illinois	100.0	11.7	88.0	40.8	98.2
Iowa	93.8	8.7	84.9	62.2	96.7
Kansas	98.8	37.6	60.6	58.5	98.8
Maine	98.8	23.0	75.0	46.2	95.9
Massachusetts	95.2	15.4	79.3	37.9	95.7
Michigan	90.0	35.2	54.1	55.3	97.3
Mississippi	36.8	19.0	14.4	64.6	97.8
Missouri	99.0	27.3	70.6	60.2	98.0
Montana	99.3	10.0	89.3	63.5	94.1
Nebraska	99.6	16.0	83.6	66.1	95.9
New Hampshire	97.7	18.0	78.4	43.4	99.5
New York [‡]	100.0	10.3	89.7	60.5	100.0
North Carolina	97.3	42.7	54.6	46.2	96.5
North Dakota	100.0	9.7	90.3	77.2	94.8
Oregon	97.3	18.7	78.1	51.2	85.4
Pennsylvania	98.4	11.5	86.9	56.2	97.7
Rhode Island	100.0	11.9	88.1	53.0	98.8
South Carolina	98.0	47.4	50.1	53.1	99.2
South Dakota	78.1	14.3	61.5	38.5	92.4
Tennessee	83.0	35.6	44.4	42.1	96.5
Texas	95.2	16.3	78.5	60.8	98.9
Utah	99.4	4.8	94.6	60.3	98.5
Vermont	97.1	10.5	85.9	49.6	98.6
Virginia	93.2	6.5	83.7	52.0	96.1
Washington	96.3	6.2	88.8	55.8	89.6
West Virginia	99.4	36.2	62.7	46.4	99.5
State Median	97.3	18.1	76.6	55.6	97.0
State Range	36.8 – 100.0	4.8 – 47.4	14.4 – 94.6	37.9 – 80.6	35.0 – 100.0

TABLE 23. Percentage of All Schools That Required Physical Education in Any of Grades 6–12; the Percentage That Required Students to Take Only One Course or Two or More Courses; Among Schools That Required a Physical Education Course, the Percentage That Required Students Who Fail a Required Physical Education Course to Repeat It; and the Percentage of All Schools in Which a Newly Hired Physical Education Teacher is Required to be Certified* in Physical Education, Selected U.S. Sites: School Health Profiles, Principal Surveys, 2006 (continued)

Site	Required physical education	Required only one physical education course	Required two or more physical education courses	Required students who fail a required physical education course to repeat it [†]	Newly hired physical education teacher required to be certified in physical education
LOCAL SURVEYS					
Charlotte-Mecklenburg County	100.0	34.0	66.0	43.8	97.7
Chicago	94.5	41.0	49.1	26.4	97.8
Dallas	98.0	16.1	81.9	59.1	100.0
District of Columbia	91.1	34.7	56.1	73.1	96.0
Hillsborough County	96.6	15.3	79.5	49.3	100.0
Los Angeles	98.7	4.7	93.9	45.3	98.7
Memphis	81.2	47.3	30.9	54.8	100.0
Miami	65.7	32.7	31.9	82.7	98.9
Orange County	44.0	12.4	31.6	78.8	96.9
Philadelphia	96.9	44.6	51.4	34.5	96.9
San Diego	96.6	8.9	87.7	57.3	100.0
San Francisco	93.7	22.3	68.2	37.9	90.6
Local Median	95.6	27.5	61.1	52.1	98.3
Local Range	44.0 – 100.0	4.7 – 47.3	30.9 – 93.9	26.4 – 82.7	90.6 – 100.0

* Certification, licensure, or endorsement by the state.

† Among schools that required physical education.

‡ Survey did not include schools from the New York City Department of Education.

TABLE 24. Percentage of Schools That Taught a Required Physical Education Course in Each Grade,* Selected U.S. Sites: School Health Profiles, Principal Surveys, 2006

Site	Grade 6	Grade 7	Grade 8	Grade 9	Grade 10	Grade 11	Grade 12
STATE SURVEYS							
Alabama	91.8	94.6	94.0	83.3	48.4	46.8	45.9
Alaska	60.3	64.1	63.4	71.3	65.0	47.5	40.8
Arizona	58.8	60.8	57.0	44.8	25.8	21.1	19.4
Arkansas	91.0	89.4	81.9	91.7	63.7	59.9	59.7
Connecticut	95.6	95.0	94.9	94.3	91.3	76.3	65.0
Delaware	80.0	81.5	82.9	76.3	58.7	21.2	18.7
Florida	60.0	59.4	58.8	59.5	47.0	36.1	34.1
Georgia	69.6	70.0	70.3	70.7	25.0	21.9	21.3
Hawaii	71.3	76.3	69.2	80.9	40.4	18.5	15.6
Idaho	83.2	88.0	82.8	54.9	58.9	39.2	36.2
Illinois	98.9	99.1	99.0	97.9	97.2	96.5	94.3
Iowa	87.1	90.0	89.9	87.8	88.0	87.5	87.5
Kansas	91.3	87.7	85.4	90.6	17.0	7.1	7.1
Maine	95.9	95.4	94.3	88.7	74.3	28.0	26.5
Massachusetts	89.7	91.0	91.2	85.0	77.6	57.3	52.4
Michigan	75.7	76.8	68.6	73.3	37.3	26.5	25.9
Mississippi	9.9	7.8	9.2	23.3	20.9	17.9	17.3
Missouri	92.6	95.3	95.3	90.4	46.9	27.9	29.1
Montana	94.8	95.1	96.8	98.8	94.5	12.8	11.5
Nebraska	93.8	96.1	96.1	87.7	46.7	18.6	16.0
New Hampshire	94.1	94.0	94.0	88.7	66.8	43.8	39.3
New York [†]	100.0	100.0	100.0	98.9	98.9	98.9	98.9
North Carolina	94.3	95.1	93.6	90.7	20.7	14.6	14.7
North Dakota	98.8	100.0	100.0	94.7	72.5	28.6	22.7
Oregon	92.5	91.5	91.6	87.3	66.2	35.8	31.1
Pennsylvania	95.4	97.0	97.0	92.8	92.3	88.7	82.4
Rhode Island	100.0	100.0	100.0	100.0	97.6	100.0	97.5
South Carolina	94.7	94.4	94.4	93.1	41.9	41.0	41.4
South Dakota	58.4	70.1	68.6	37.4	12.8	7.7	11.0
Tennessee	70.0	70.9	69.7	56.3	20.7	12.5	13.9
Texas	87.9	83.5	71.5	88.5	87.8	61.0	59.7
Utah	96.7	98.2	97.3	90.8	90.7	69.7	48.6
Vermont	92.2	94.8	94.8	88.6	80.3	53.8	50.1
Virginia	80.9	81.1	68.7	82.4	81.2	5.4	5.4
Washington	87.6	90.7	89.3	79.9	74.5	46.8	41.6
West Virginia	96.3	96.1	96.1	86.2	46.4	20.4	19.2
State Median	91.6	91.3	91.4	87.8	64.4	36.0	32.6
State Range	9.9 – 100.0	7.8 – 100.0	9.2 – 100.0	23.3 – 100.0	12.8 – 98.9	5.4 – 100.0	5.4 – 98.9

TABLE 24. Percentage of Schools That Taught a Required Physical Education Course in Each Grade,* Selected U.S. Sites: School Health Profiles, Principal Surveys, 2006 (continued)

Site	Grade 6	Grade 7	Grade 8	Grade 9	Grade 10	Grade 11	Grade 12
LOCAL SURVEYS							
Charlotte-Mecklenburg County	100.0	100.0	100.0	100.0	27.3	18.2	18.2
Chicago	87.1	88.0	87.9	64.2	63.2	19.3	16.0
Dallas	90.0	92.6	96.3	95.8	91.5	77.4	77.4
District of Columbia	64.7	56.8	54.4	87.4	82.1	19.6	19.6
Hillsborough County	83.4	87.1	87.1	72.3	72.3	47.8	47.8
Los Angeles	97.8	97.8	97.8	93.7	93.7	23.2	19.9
Memphis	53.9	60.3	63.4	67.1	21.2	19.3	19.3
Miami	30.9	26.0	21.7	50.2	42.8	17.3	17.6
Orange County	14.3	14.3	14.3	38.3	33.7	20.2	20.2
Philadelphia	91.0	90.1	90.1	63.9	83.1	68.2	60.2
San Diego	92.8	93.5	93.5	90.9	80.6	20.8	20.8
San Francisco	85.6	86.2	86.2	78.4	76.9	41.7	41.7
Local Median	86.4	87.6	87.5	75.4	74.6	20.5	20.1
Local Range	14.3 – 100.0	14.3 – 100.0	14.3 – 100.0	38.3 – 100.0	21.2 – 93.7	17.3 – 77.4	16.0 – 77.4

* Among schools with students in that grade.

† Survey did not include schools from the New York City Department of Education.

TABLE 25a. Among Schools That Required a Physical Education Course for Students in Any of Grades 6–12, the Percentage That Allowed Students to be Exempted from Taking a Required Physical Education Course for Specific Reasons, Selected U.S. Sites: School Health Profiles, Principal Surveys, 2006

Site	Religious reasons	Long-term physical or medical disability	Cognitive disability	Enrollment in other courses	Participation in school sports	Participation in other school activities*
STATE SURVEYS						
Alabama	23.0	63.1	18.1	7.6	17.5	38.2
Alaska	34.4	78.4	27.5	22.4	35.0	9.6
Arizona	52.8	86.1	34.8	18.8	13.6	25.8
Arkansas	32.8	75.3	28.6	16.5	44.4	15.7
Connecticut	33.0	87.2	12.9	5.5	2.9	0.4
Delaware	30.6	77.5	20.0	8.2	4.9	8.2
Florida	44.7	75.3	36.7	43.1	35.2	56.9
Georgia	38.0	77.1	31.2	30.7	3.8	36.0
Hawaii	40.0	85.0	36.5	18.4	1.9	10.4
Idaho	38.7	81.6	38.1	22.2	9.6	8.0
Illinois	47.1	85.7	25.8	24.1	32.1	22.4
Iowa	53.6	79.4	27.5	40.0	17.5	10.7
Kansas	47.7	79.8	34.6	9.9	5.8	3.9
Maine	34.0	83.3	21.8	15.7	3.9	2.6
Massachusetts	42.9	89.2	24.3	13.8	4.5	7.1
Michigan	42.8	87.9	35.0	15.1	15.2	28.0
Mississippi	64.5	79.7	59.8	48.7	57.4	51.6
Missouri	37.7	76.9	26.8	15.2	2.5	6.8
Montana	28.7	85.3	23.5	10.7	0.7	2.8
Nebraska	32.5	77.3	29.9	13.3	3.6	3.4
New Hampshire	33.2	81.1	18.0	16.0	12.0	7.3
New York [†]	16.6	65.5	11.5	1.4	10.9	0.3
North Carolina	34.1	68.4	18.7	11.2	1.4	9.6
North Dakota	20.4	64.9	21.1	10.6	0.6	1.2
Oregon	55.6	89.4	36.5	25.3	12.1	10.9
Pennsylvania	36.4	83.9	21.2	6.7	2.8	3.9
Rhode Island	29.7	88.1	15.6	0.0	2.5	0.0
South Carolina	47.9	82.2	31.4	21.4	3.1	52.5
South Dakota	22.4	69.9	30.4	11.2	5.0	2.4
Tennessee	45.2	79.7	37.9	12.7	4.6	36.7
Texas	35.7	79.2	37.1	16.9	75.4	56.8
Utah	44.7	92.7	51.6	17.3	29.3	10.4
Vermont	33.5	77.4	19.9	9.5	20.2	4.0
Virginia	40.9	79.5	28.7	11.4	0.5	6.8
Washington	61.5	89.7	36.8	34.3	35.6	20.8
West Virginia	16.3	70.2	15.0	8.0	0.5	4.5
State Median	37.1	79.7	28.1	15.2	5.4	8.9
State Range	16.3 – 64.5	63.1 – 92.7	11.5 – 59.8	0.0 – 48.7	0.5 – 75.4	0.0 – 56.9

TABLE 25a. Among Schools That Required a Physical Education Course for Students in Any of Grades 6–12, the Percentage That Allowed Students to be Exempted from Taking a Required Physical Education Course for Specific Reasons, Selected U.S. Sites: School Health Profiles, Principal Surveys, 2006 (*continued*)

Site	Religious reasons	Long-term physical or medical disability	Cognitive disability	Enrollment in other courses	Participation in school sports	Participation in other school activities*
LOCAL SURVEYS						
Charlotte-Mecklenburg County	37.3	69.2	12.7	15.0	5.0	19.5
Chicago	53.5	75.4	19.0	5.1	2.5	10.6
Dallas	33.3	79.3	47.2	12.8	53.9	64.3
District of Columbia	40.1	79.4	22.7	21.9	0.0	17.3
Hillsborough County	47.6	83.4	38.2	49.0	34.6	61.1
Los Angeles	27.5	73.8	16.3	5.8	42.8	33.4
Memphis	50.6	65.7	33.8	10.2	0.0	56.2
Miami	35.7	75.2	47.1	32.9	32.4	45.7
Orange County	55.0	47.9	42.3	31.0	73.2	64.8
Philadelphia	67.9	90.0	31.2	14.1	3.3	7.7
San Diego	32.2	92.6	40.1	16.8	49.3	50.2
San Francisco	48.3	87.3	26.3	7.0	13.9	20.9
Local Median	43.9	77.4	32.5	14.6	23.2	39.6
Local Range	27.5 – 67.9	47.9 – 92.6	12.7 – 47.2	5.1 – 49.0	0.0 – 73.2	7.7 – 64.8

* Such as ROTC, marching band, chorus, or cheerleading.

† Survey did not include schools from the New York City Department of Education.

TABLE 25b. Among Schools That Required a Physical Education Course for Students in Any of Grades 6–12, the Percentage That Allowed Students to be Exempted from Taking a Required Physical Education Course for Specific Reasons, Selected U.S. Sites: School Health Profiles, Principal Surveys, 2006

Site	Participation in community sports activities	High physical fitness competency test score	Participation in vocational training	Participation in community service activities	Could not be exempted from a required physical education course for certain reasons
STATE SURVEYS					
Alabama	0.4	1.1	7.2	0.8	55.9
Alaska	13.4	7.1	6.2	4.5	49.5
Arizona	5.4	1.3	5.1	4.1	59.3
Arkansas	2.9	2.5	4.5	2.1	46.9
Connecticut	1.4	0.5	1.2	0.4	90.2
Delaware	1.7	0.0	1.6	1.6	79.7
Florida	3.6	12.4	11.3	5.0	21.1
Georgia	0.8	1.0	5.1	2.5	50.4
Hawaii	1.9	0.0	0.0	0.0	70.7
Idaho	4.5	1.0	3.7	1.5	64.8
Illinois	3.5	0.6	5.7	0.4	54.5
Iowa	1.3	0.8	8.3	1.7	50.5
Kansas	0.9	0.0	2.1	1.7	80.5
Maine	1.8	0.4	2.3	1.2	79.8
Massachusetts	2.7	0.2	2.5	1.2	77.0
Michigan	2.0	1.0	2.8	0.7	58.0
Mississippi	7.2	14.3	25.7	11.0	21.7
Missouri	0.3	0.0	1.7	0.3	80.5
Montana	1.1	0.7	1.1	1.1	88.5
Nebraska	0.0	0.0	1.0	0.9	81.1
New Hampshire	3.6	0.0	1.2	1.3	71.9
New York [†]	2.1	0.3	0.5	0.3	86.3
North Carolina	1.0	0.0	0.7	0.4	82.8
North Dakota	0.0	0.8	0.0	1.3	87.3
Oregon	11.6	3.2	5.1	3.9	63.1
Pennsylvania	2.3	2.1	6.3	1.0	82.7
Rhode Island	1.2	0.0	2.4	0.0	95.2
South Carolina	1.7	0.8	2.1	0.9	38.7
South Dakota	0.6	0.6	1.2	1.2	82.3
Tennessee	0.4	0.7	2.9	1.6	54.9
Texas	16.8	2.1	18.1	1.9	17.0
Utah	3.8	17.4	1.4	0.5	44.7
Vermont	7.5	0.0	3.2	3.1	72.1
Virginia	4.2	0.0	0.9	0.4	80.5
Washington	16.3	2.0	4.3	0.8	35.8
West Virginia	0.5	1.1	1.8	0.5	90.7
State Median	2.0	0.8	2.5	1.2	71.3
State Range	0.0 – 16.8	0.0 – 17.4	0.0 – 25.7	0.0 – 11.0	17.0 – 95.2

TABLE 25b. Among Schools That Required a Physical Education Course for Students in Any of Grades 6–12, the Percentage That Allowed Students to be Exempted from Taking a Required Physical Education Course for Specific Reasons, Selected U.S. Sites: School Health Profiles, Principal Surveys, 2006 (continued)

Site	Participation in community sports activities	High physical fitness competency test score	Participation in vocational training	Participation in community service activities	Could not be exempted from a required physical education course for certain reasons
LOCAL SURVEYS					
Charlotte-Mecklenburg County	5.0	0.0	2.6	2.6	70.7
Chicago	3.1	0.5	1.0	2.0	83.6
Dallas	10.6	6.3	23.8	2.1	31.4
District of Columbia	0.0	0.0	0.0	3.8	68.0
Hillsborough County	2.1	14.7	2.5	0.0	16.8
Los Angeles	3.5	2.6	2.3	1.2	48.6
Memphis	0.0	2.7	2.4	0.0	34.3
Miami	2.0	13.4	7.4	1.8	29.4
Orange County	0.0	0.0	0.0	0.0	12.7
Philadelphia	3.6	3.5	1.2	2.1	76.2
San Diego	15.3	0.0	0.0	0.0	39.6
San Francisco	9.7	0.0	0.0	0.0	59.0
Local Median	3.3	1.6	1.8	1.5	44.1
Local Range	0.0 – 15.3	0.0 – 14.7	0.0 – 23.8	0.0 – 3.8	12.7 – 83.6

* These reasons included enrollment in other courses, participation in school sports, participation in other school activities, participation in community sports activities, high physical fitness competency test score, participation in vocational training, and participation in community service activities.

† Survey did not include schools from the New York City Department of Education.

TABLE 26. Percentage of All Schools That Supported or Promoted Walking or Biking to and from School, the Percentage of All Schools That Allowed Use of Their Physical Activity or Athletic Facilities* or Offered Opportunities for Students to Participate in Intramural Activities or Physical Activity Clubs, and Among Schools That Offered Opportunities for Students to Participate in Intramural Activities or Physical Activity Clubs, the Percentage That Provided Transportation Home,† Selected U.S. Sites: School Health Profiles, Principal Surveys, 2006

Site	Supported or promoted walking or biking to and from school	Allowed use of physical activity or athletic facilities	Offered intramural activities or physical activity clubs	Provided transportation home
STATE SURVEYS				
Alabama	24.8	75.5	44.6	10.3
Alaska	46.8	84.1	68.6	12.1
Arizona	62.9	64.0	71.3	51.7
Arkansas	37.7	82.7	38.1	12.1
Connecticut	31.1	89.7	80.9	52.2
Delaware	31.0	91.1	69.2	54.2
Florida	52.2	68.7	66.9	39.2
Georgia	28.1	82.6	60.7	23.9
Hawaii	42.9	73.4	90.1	14.1
Idaho	45.9	95.1	63.5	16.8
Illinois	55.3	89.1	58.3	35.8
Iowa	53.2	90.2	44.7	32.0
Kansas	48.2	93.2	35.4	31.8
Maine	48.6	94.6	82.1	46.9
Massachusetts	45.2	88.6	87.0	44.8
Michigan	46.3	86.4	72.7	13.6
Mississippi	10.3	66.8	43.3	18.7
Missouri	41.4	92.0	58.2	25.5
Montana	58.5	93.9	60.4	16.1
Nebraska	53.2	95.9	40.8	30.9
New Hampshire	53.0	94.4	84.7	39.3
New York‡	46.8	97.0	88.7	76.5
North Carolina	26.7	88.9	65.5	22.7
North Dakota	49.9	86.5	42.9	31.6
Oregon	59.9	91.0	60.7	27.1
Pennsylvania	36.2	84.9	82.9	38.4
Rhode Island	36.3	80.3	77.3	67.4
South Carolina	35.6	81.3	59.8	18.2
South Dakota	57.4	91.4	35.8	17.1
Tennessee	26.1	78.5	60.0	12.1
Texas	38.3	85.9	41.4	37.5
Utah	59.9	96.8	74.2	27.3
Vermont	58.5	97.6	83.3	30.6
Virginia	29.2	95.2	68.6	56.6
Washington	50.9	90.7	65.2	48.2
West Virginia	23.4	93.9	65.7	24.2
State Median	46.1	89.4	65.4	30.8
State Range	10.3 – 62.9	64.0 – 97.6	35.4 – 90.1	10.3 – 76.5

TABLE 26. Percentage of All Schools That Supported or Promoted Walking or Biking to and from School, the Percentage of All Schools That Allowed Use of Their Physical Activity or Athletic Facilities* or Offered Opportunities for Students to Participate in Intramural Activities or Physical Activity Clubs, and Among Schools That Offered Opportunities for Students to Participate in Intramural Activities or Physical Activity Clubs, the Percentage That Provided Transportation Home,† Selected U.S. Sites: School Health Profiles, Principal Surveys, 2006 (continued)

Site	Supported or promoted walking or biking to and from school	Allowed use of physical activity or athletic facilities	Offered intramural activities or physical activity clubs	Provided transportation home
LOCAL SURVEYS				
Charlotte-Mecklenburg County	30.8	97.7	61.9	23.4
Chicago	51.7	58.6	93.6	19.0
Dallas	18.1	58.0	67.0	51.2
District of Columbia	39.3	63.7	93.2	20.8
Hillsborough County	44.9	82.5	81.4	15.5
Los Angeles	70.6	98.9	75.5	32.4
Memphis	37.6	67.2	71.0	14.9
Miami	51.6	66.6	92.0	52.0
Orange County	80.8	90.7	95.0	18.3
Philadelphia	35.1	64.4	92.3	14.7
San Diego	70.1	77.0	71.9	68.0
San Francisco	38.1	60.7	87.7	6.8
Local Median	42.1	66.9	84.6	19.9
Local Range	18.1 – 80.8	58.0 – 98.9	61.9 – 95.0	6.8 – 68.0

* For community-sponsored sports teams, classes, or lessons outside of school hours or when school is not in session.

† For students who participate in after-school intramural activities or physical activity clubs.

‡ Survey did not include schools from the New York City Department of Education.

TABLE 27. Percentage of All Schools That Served Lunch to Students, and Among Those Schools, the Percentage in Which Students Usually Had 20 or More Minutes to Eat Lunch Once They Were Seated, Selected U.S. Sites: School Health Profiles, Principal Surveys, 2006

Site	Served lunch to students	≥20 minutes to eat lunch*
STATE SURVEYS		
Alabama	100.0	81.3
Alaska	91.1	90.4
Arizona	88.9	80.7
Arkansas	100.0	88.2
Connecticut	99.5	77.1
Delaware	100.0	82.2
Florida	98.9	85.5
Georgia	99.6	84.8
Hawaii	96.8	81.3
Idaho	100.0	84.5
Illinois	99.7	85.5
Iowa	97.7	66.8
Kansas	99.6	78.3
Maine	99.3	67.4
Massachusetts	99.8	76.7
Michigan	99.2	84.1
Mississippi	100.0	73.4
Missouri	100.0	64.7
Montana	99.2	85.2
Nebraska	99.2	83.6
New Hampshire	99.5	67.8
New York [†]	99.7	87.9
North Carolina	100.0	80.6
North Dakota	97.6	78.4
Oregon	99.6	83.3
Pennsylvania	98.4	89.5
Rhode Island	100.0	82.1
South Carolina	100.0	83.7
South Dakota	98.9	67.9
Tennessee	99.3	85.7
Texas	99.8	86.6
Utah	99.5	89.0
Vermont	100.0	68.2
Virginia	99.7	76.5
Washington	96.5	86.7
West Virginia	99.4	95.9
State Median	99.6	82.8
State Range	88.9 – 100.0	64.7 – 95.9

TABLE 27. Percentage of All Schools That Served Lunch to Students, and Among Those Schools, the Percentage in Which Students Usually Had 20 or More Minutes to Eat Lunch Once They Were Seated, Selected U.S. Sites: School Health Profiles, Principal Surveys, 2006 (continued)

Site	Served lunch to students	≥20 minutes to eat lunch*
LOCAL SURVEYS		
Charlotte-Mecklenburg County	100.0	64.2
Chicago	99.6	58.5
Dallas	98.0	67.4
District of Columbia	96.0	94.0
Hillsborough County	100.0	62.2
Los Angeles	100.0	81.8
Memphis	100.0	76.5
Miami	100.0	92.3
Orange County	100.0	74.5
Philadelphia	99.2	84.0
San Diego	100.0	80.5
San Francisco	100.0	94.4
Local Median	100.0	78.5
Local Range	96.0 – 100.0	58.5 – 94.4

* Among schools that served lunch to students.

† Survey did not include schools from the New York City Department of Education.

TABLE 28a. Percentage of All Schools That Allowed Students to Purchase Snack Foods or Beverages from One or More Vending Machines or at the School Store, Canteen, or Snack Bar, and That Allowed Students to Purchase Less Nutritious and More Nutritious Snack Foods and Beverages, Selected U.S. Sites: School Health Profiles, Principal Surveys, 2006

Site	Allowed students to purchase snack foods or beverages	Less Nutritious Foods and Beverages					
		2% or whole milk (plain or flavored)	Chocolate candy	Other kinds of candy	Salty snacks that are not low in fat*	Soda pop or fruit drinks that are not 100% juice	Sports drinks
STATE SURVEYS							
Alabama	86.7	32.3	32.3	37.4	45.4	69.7	81.9
Alaska	62.7	15.9	41.2	42.3	44.0	50.4	53.3
Arizona	68.6	27.1	32.8	36.2	40.0	43.1	58.8
Arkansas	70.7	33.5	23.5	26.3	26.2	64.2	58.5
Connecticut	71.8	49.5	21.2	25.8	41.2	39.5	57.3
Delaware	79.0	40.8	34.3	37.2	44.7	45.4	67.6
Florida	72.3	44.4	28.9	32.9	38.1	57.4	66.0
Georgia	87.1	40.3	53.9	56.9	59.4	73.3	82.6
Hawaii	61.9	17.3	12.6	14.2	11.0	39.5	30.5
Idaho	93.4	44.8	65.5	67.4	63.7	82.5	90.2
Illinois	77.2	50.2	43.2	46.5	52.2	63.7	67.5
Iowa	87.9	45.3	46.6	54.4	48.3	74.9	81.3
Kansas	85.7	32.5	62.2	63.0	60.4	79.1	78.9
Maine	77.6	42.4	8.4	11.2	23.0	25.3	59.5
Massachusetts	77.5	50.9	18.2	23.8	38.7	37.4	59.1
Michigan	87.6	55.2	58.6	64.2	68.2	67.7	78.9
Mississippi	87.9	34.0	71.0	72.0	75.0	78.2	78.5
Missouri	87.1	50.2	50.8	54.9	60.9	74.2	76.2
Montana	87.3	23.9	52.2	55.2	49.9	71.3	85.3
Nebraska	86.0	37.8	44.9	46.1	46.4	78.3	81.3
New Hampshire	90.5	60.2	22.2	24.5	44.6	43.4	73.1
New York [†]	93.3	60.9	34.5	44.8	61.7	62.5	81.5
North Carolina	84.3	40.1	35.0	40.3	50.0	56.0	72.2
North Dakota	78.4	23.3	45.7	44.1	38.2	69.1	73.4
Oregon	78.6	35.9	49.9	55.1	55.6	62.0	70.9
Pennsylvania	76.9	48.8	39.0	43.0	46.9	50.7	62.3
Rhode Island	89.5	67.9	26.4	28.8	49.8	44.0	71.0
South Carolina	94.0	49.2	56.4	66.0	69.9	76.0	86.6
South Dakota	80.5	35.4	28.3	29.7	27.5	66.6	77.1
Tennessee	88.0	45.6	58.4	61.9	62.5	73.3	81.9
Texas	81.0	49.6	46.9	39.9	47.9	56.3	70.9
Utah	93.0	58.6	82.9	82.6	75.9	86.0	87.9
Vermont	75.5	54.4	13.4	15.7	36.5	39.3	56.3
Virginia	80.2	47.2	47.2	51.5	60.0	62.4	67.0
Washington	88.2	41.1	39.4	46.5	39.6	57.8	75.1
West Virginia	82.3	32.9	10.1	18.2	28.3	37.3	48.6
State Median	83.3	43.4	40.3	43.6	47.4	62.5	72.7
State Range	61.9 – 94.0	15.9 – 67.9	8.4 – 82.9	11.2 – 82.6	11.0 – 75.9	25.3 – 86.0	30.5 – 90.2

TABLE 28a. Percentage of All Schools That Allowed Students to Purchase Snack Foods or Beverages from One or More Vending Machines or at the School Store, Canteen, or Snack Bar, and That Allowed Students to Purchase Less Nutritious and More Nutritious Snack Foods and Beverages, Selected U.S. Sites: School Health Profiles, Principal Surveys, 2006 (*continued*)

Site	Allowed students to purchase snack foods or beverages	Less Nutritious Foods and Beverages					
		2% or whole milk (plain or flavored)	Chocolate candy	Other kinds of candy	Salty snacks that are not low in fat*	Soda pop or fruit drinks that are not 100% juice	Sports drinks
LOCAL SURVEYS							
Charlotte-Mecklenburg County	85.7	48.0	47.6	57.3	81.0	66.1	73.3
Chicago	31.5	16.1	4.0	5.7	4.4	9.8	18.0
Dallas	76.9	35.0	59.1	56.9	65.6	71.4	69.8
District of Columbia	64.0	16.1	18.3	22.3	18.3	37.1	35.1
Hillsborough County	88.6	53.4	27.1	32.3	51.1	69.4	84.3
Los Angeles	88.0	55.3	8.2	16.0	15.5	9.6	76.5
Memphis	77.7	30.2	53.3	56.8	52.5	67.5	67.0
Miami	86.1	57.8	52.2	59.3	63.4	71.9	80.1
Orange County	80.7	49.9	21.1	24.3	33.3	47.4	74.3
Philadelphia	61.4	28.5	9.9	12.6	24.3	12.9	25.8
San Diego	84.5	64.6	43.8	43.8	67.0	57.3	78.6
San Francisco	62.3	30.2	5.9	12.3	6.5	15.1	28.1
Local Median	79.2	41.5	24.1	28.3	42.2	52.4	71.6
Local Range	31.5 – 88.6	16.1 – 64.6	4.0 – 59.1	5.7 – 59.3	4.4 – 81.0	9.6 – 71.9	18.0 – 84.3

* Such as regular potato chips.

† Survey did not include schools from the New York City Department of Education.

TABLE 28b. Percentage of All Schools That Allowed Students to Purchase Snack Foods or Beverages from One or More Vending Machines or at the School Store, Canteen, or Snack Bar, and That Allowed Students to Purchase Less Nutritious and More Nutritious Snack Foods and Beverages, Selected U.S. Sites: School Health Profiles, Principal Surveys, 2006

Site	More Nutritious Foods and Beverages					
	1% or skim milk	100% fruit juice or vegetable juice	Bottled water	Fruits or vegetables, not juice	Low-fat baked goods*	Salty snacks that are low in fat†
STATE SURVEYS						
Alabama	33.3	71.5	84.4	17.5	71.6	76.7
Alaska	10.7	50.6	55.6	14.7	36.2	42.7
Arizona	26.3	44.7	64.4	24.9	39.7	50.2
Arkansas	27.6	48.7	66.6	14.6	30.8	35.1
Connecticut	50.1	57.8	69.1	39.4	47.7	58.5
Delaware	40.2	62.1	74.5	23.9	47.6	56.6
Florida	43.7	55.4	70.2	29.6	42.5	49.5
Georgia	36.9	65.3	85.9	15.2	49.7	58.1
Hawaii	16.3	41.0	60.3	6.6	9.8	12.5
Idaho	36.0	77.0	90.8	28.6	57.6	71.2
Illinois	40.4	62.0	73.7	34.0	49.2	57.8
Iowa	44.1	72.5	85.9	28.2	49.3	57.6
Kansas	26.2	62.0	80.0	19.5	56.9	66.2
Maine	47.5	68.6	74.9	32.2	46.1	53.2
Massachusetts	52.1	64.7	75.2	34.9	53.9	62.4
Michigan	48.7	68.9	84.4	43.4	56.2	70.8
Mississippi	27.1	54.3	83.3	18.2	53.2	72.1
Missouri	45.6	65.5	81.9	23.6	52.4	62.4
Montana	22.0	69.6	83.1	25.1	41.9	52.1
Nebraska	35.6	62.6	78.8	17.1	45.4	52.2
New Hampshire	60.1	78.6	89.7	43.7	65.6	73.2
New York‡	59.8	77.0	89.6	46.8	65.2	79.2
North Carolina	41.2	63.8	79.9	30.6	55.7	62.0
North Dakota	21.7	64.0	75.9	14.5	33.7	43.2
Oregon	35.4	64.4	76.3	30.3	51.2	63.4
Pennsylvania	47.6	65.2	74.8	32.8	53.8	61.8
Rhode Island	66.2	77.7	84.6	46.8	55.8	68.7
South Carolina	42.7	66.9	90.2	25.9	66.7	75.9
South Dakota	33.2	66.1	79.7	19.0	36.7	39.9
Tennessee	35.7	63.1	85.1	22.0	57.1	67.4
Texas	44.9	67.2	77.6	41.2	59.6	68.5
Utah	45.8	74.4	89.7	36.8	73.4	82.9
Vermont	55.1	64.6	71.2	38.4	43.8	55.9
Virginia	40.6	62.9	77.6	25.2	58.1	69.1
Washington	36.0	73.7	85.1	33.9	57.9	62.9
West Virginia	33.1	67.0	79.3	7.6	62.3	67.8
State Median	40.3	65.0	79.5	27.1	52.8	62.2
State Range	10.7 – 66.2	41.0 – 78.6	55.6 – 90.8	6.6 – 46.8	9.8 – 73.4	12.5 – 82.9

TABLE 28b. Percentage of All Schools That Allowed Students to Purchase Snack Foods or Beverages from One or More Vending Machines or at the School Store, Canteen, or Snack Bar, and That Allowed Students to Purchase Less Nutritious and More Nutritious Snack Foods and Beverages, Selected U.S. Sites: School Health Profiles, Principal Surveys, 2006 (*continued*)

Site	More Nutritious Foods and Beverages					
	1% or skim milk	100% fruit juice or vegetable juice	Bottled water	Fruits or vegetables, not juice	Low-fat baked goods*	Salty snacks that are low in fat†
LOCAL SURVEYS						
Charlotte-Mecklenburg County	50.4	71.7	81.0	41.0	69.2	81.0
Chicago	14.7	25.0	29.0	10.3	13.4	14.2
Dallas	21.8	43.9	72.1	19.7	45.8	56.7
District of Columbia	24.3	42.7	55.8	14.2	22.3	25.5
Hillsborough County	53.6	66.9	86.5	38.7	46.6	65.1
Los Angeles	56.7	75.9	86.6	43.6	66.5	67.8
Memphis	25.1	61.6	56.6	17.9	32.0	42.7
Miami	60.1	62.7	80.3	39.3	62.5	72.0
Orange County	44.2	53.2	78.2	32.6	51.9	55.7
Philadelphia	29.2	53.5	54.7	23.1	40.4	42.7
San Diego	46.1	67.1	78.6	58.8	56.6	72.4
San Francisco	36.7	52.2	61.2	39.6	45.7	46.1
Local Median	40.5	57.6	75.2	35.7	46.2	56.2
Local Range	14.7 – 60.1	25.0 – 75.9	29.0 – 86.6	10.3 – 58.8	13.4 – 69.2	14.2 – 81.0

* Such as cookies, crackers, cakes, pastries, or other low-fat baked goods.

† Such as pretzels, baked chips, or other low-fat chips.

‡ Survey did not include schools from the New York City Department of Education.

TABLE 29. Percentage of All Schools That Allowed Students to Purchase Candy; Snacks That are Not Low in Fat; Soda Pop, Sports Drinks, or Fruit Drinks That Are Not 100% Juice; or 2% or Whole Milk During Specific Times and the Percentage of Schools That Had a Policy Stating That Fruits and Vegetables Will Be Among the Foods Offered at School Settings,* Selected U.S. Sites: School Health Profiles, Principal Surveys, 2006

Site	Before classes begin in the morning	During any school hours when meals are not being served	During school lunch periods	Offered fruits or vegetables
STATE SURVEYS				
Alabama	28.9	45.3	13.9	38.0
Alaska	29.2	25.3	29.8	22.2
Arizona	38.8	28.2	34.9	24.6
Arkansas	30.8	25.5	34.6	35.5
Connecticut	21.6	20.3	32.6	27.0
Delaware	23.9	11.9	26.9	15.6
Florida	34.2	33.7	33.8	19.5
Georgia	41.0	36.6	44.4	11.1
Hawaii	20.2	16.7	3.9	21.0
Idaho	67.5	52.5	69.1	16.5
Illinois	42.3	23.0	45.6	9.9
Iowa	59.4	44.3	35.5	11.9
Kansas	60.9	39.5	39.9	11.2
Maine	21.3	13.0	18.5	35.5
Massachusetts	24.1	13.7	44.0	18.1
Michigan	52.3	28.9	64.3	11.9
Mississippi	28.9	50.0	11.9	17.6
Missouri	56.1	25.7	57.5	13.8
Montana	57.3	41.6	58.4	18.1
Nebraska	55.6	40.9	14.4	10.8
New Hampshire	38.6	29.2	37.6	19.9
New York [†]	35.3	30.1	45.7	14.4
North Carolina	27.2	22.8	22.4	17.2
North Dakota	52.7	32.9	28.8	21.6
Oregon	41.8	33.4	46.2	17.5
Pennsylvania	31.3	15.3	40.1	29.3
Rhode Island	34.7	16.9	58.4	24.4
South Carolina	38.6	30.3	63.2	14.9
South Dakota	54.3	46.8	22.7	16.8
Tennessee	32.3	46.2	22.2	14.6
Texas	31.3	29.1	34.8	29.6
Utah	72.5	56.6	81.3	20.2
Vermont	32.2	29.4	32.1	15.2
Virginia	33.9	23.4	31.3	11.1
Washington	43.4	33.1	43.8	27.6
West Virginia	25.4	26.2	11.6	29.8
State Median	35.0	29.3	34.9	17.9
State Range	20.2 – 72.5	11.9 – 56.6	3.9 – 81.3	9.9 – 38.0

TABLE 29. Percentage of All Schools That Allowed Students to Purchase Candy; Snacks That are Not Low in Fat; Soda Pop, Sports Drinks, or Fruit Drinks That Are Not 100% Juice; or 2% or Whole Milk During Specific Times and the Percentage of Schools That Had a Policy Stating That Fruits and Vegetables Will Be Among the Foods Offered at School Settings,* Selected U.S. Sites: School Health Profiles, Principal Surveys, 2006 (continued)

Site	Before classes begin in the morning	During any school hours when meals are not being served	During school lunch periods	Offered fruits or vegetables
LOCAL SURVEYS				
Charlotte-Mecklenburg County	31.8	16.9	48.6	21.4
Chicago	7.2	3.1	15.7	31.9
Dallas	38.8	28.1	47.4	19.9
District of Columbia	24.7	3.1	45.7	30.9
Hillsborough County	30.5	37.1	29.2	11.7
Los Angeles	18.0	5.7	19.4	28.2
Memphis	14.1	5.5	35.9	15.7
Miami	58.0	33.7	52.6	24.6
Orange County	39.1	39.1	37.8	8.7
Philadelphia	10.5	7.1	30.8	41.5
San Diego	34.3	19.3	72.3	21.6
San Francisco	11.9	2.9	18.4	62.3
Local Median	27.6	12.0	36.9	23.1
Local Range	7.2 – 58.0	2.9 – 39.1	15.7 – 72.3	8.7 – 62.3

* Such as student parties, after-school or extended day programs, or concession stands.

† Survey did not include schools from the New York City Department of Education.

TABLE 30. Percentage of All Schools That Had a Policy Prohibiting Tobacco Use and the Percentage That Prohibited All Tobacco Use in All Locations,* Selected U.S. Sites: School Health Profiles, Principal Surveys, 2006

Site	Had a policy prohibiting tobacco use	Prohibited all tobacco use in all locations
STATE SURVEYS		
Alabama	97.9	57.3
Alaska	94.3	26.8
Arizona	97.4	56.2
Arkansas	99.6	57.4
Connecticut	97.9	52.1
Delaware	100.0	56.4
Florida	95.4	40.4
Georgia	99.6	56.7
Hawaii	98.4	62.2
Idaho	99.1	44.2
Illinois	99.0	44.9
Iowa	98.0	25.3
Kansas	98.4	26.2
Maine	99.3	58.0
Massachusetts	98.4	59.2
Michigan	95.9	39.2
Mississippi	99.4	62.2
Missouri	98.3	24.0
Montana	100.0	53.8
Nebraska	99.6	31.6
New Hampshire	99.5	49.6
New York†	98.0	64.4
North Carolina	97.2	45.3
North Dakota	100.0	39.7
Oregon	98.9	58.4
Pennsylvania	98.8	53.9
Rhode Island	97.7	61.7
South Carolina	99.3	47.0
South Dakota	96.0	22.8
Tennessee	99.0	30.3
Texas	98.5	62.2
Utah	99.5	53.8
Vermont	100.0	63.1
Virginia	98.9	43.7
Washington	99.0	64.9
West Virginia	98.9	76.3
State Median	98.9	53.8
State Range	94.3 – 100.0	22.8 – 76.3

TABLE 30. Percentage of All Schools That Had a Policy Prohibiting Tobacco Use and the Percentage That Prohibited All Tobacco Use in All Locations,* Selected U.S. Sites: School Health Profiles, Principal Surveys, 2006 (continued)

Site	Had a policy prohibiting tobacco use	Prohibited all tobacco use in all locations
LOCAL SURVEYS		
Charlotte-Mecklenburg County	100.0	74.8
Chicago	84.0	36.4
Dallas	98.0	62.6
District of Columbia	94.2	41.0
Hillsborough County	92.6	15.5
Los Angeles	100.0	67.9
Memphis	98.3	50.3
Miami	96.6	53.3
Orange County	100.0	79.5
Philadelphia	95.5	34.1
San Diego	100.0	74.0
San Francisco	100.0	58.7
Local Median	98.2	56.0
Local Range	84.0 – 100.0	15.5 – 79.5

* Prohibited the use of all tobacco, including cigarettes, smokeless tobacco (i.e., chewing tobacco, snuff, or dip), cigars, and pipes; by students, faculty and school staff, and visitors; in school buildings, outside on school grounds (including parking lots and playing fields), on school buses or other vehicles used to transport students, and at off-campus, school-sponsored events.

† Survey did not include schools from the New York City Department of Education.

TABLE 31a. Among Schools with a Policy Prohibiting Tobacco Use, the Percentage of Schools That Sometimes, Almost Always, or Always Took Specific Actions When Students Were Caught Smoking Cigarettes, Selected U.S. Sites: School Health Profiles, Principal Surveys, 2006

Site	Informed parents or guardians	Referred to a school counselor	Referred to a school administrator	Encouraged to participate in an assistance, education, or cessation program	Required to participate in an assistance, education, or cessation program
STATE SURVEYS					
Alabama	96.8	69.0	97.5	39.3	14.8
Alaska	93.2	61.1	92.6	51.1	26.8
Arizona	96.8	68.8	95.9	58.2	29.6
Arkansas	99.2	67.8	98.8	44.0	23.5
Connecticut	97.3	83.2	96.5	64.6	29.2
Delaware	100.0	77.4	98.4	68.7	32.7
Florida	95.3	73.1	93.6	58.0	43.1
Georgia	99.2	69.6	98.9	41.7	19.9
Hawaii	98.3	88.6	96.6	75.1	40.0
Idaho	99.1	87.4	98.1	69.5	58.9
Illinois	98.4	71.5	97.1	49.2	23.6
Iowa	97.9	76.6	97.9	61.9	37.7
Kansas	98.4	76.4	97.0	48.5	25.0
Maine	98.8	86.1	98.4	76.0	46.4
Massachusetts	97.4	78.6	97.5	68.0	31.2
Michigan	95.0	72.1	95.5	59.9	30.6
Mississippi	98.9	57.1	98.9	30.4	10.3
Missouri	97.7	63.9	97.9	32.5	15.3
Montana	100.0	83.0	100.0	68.4	48.4
Nebraska	99.2	76.9	99.2	56.6	26.8
New Hampshire	99.4	85.0	98.9	73.2	39.9
New York*	97.7	85.8	97.7	71.1	32.6
North Carolina	96.7	73.4	96.7	55.9	42.6
North Dakota	99.2	77.5	98.3	60.4	28.2
Oregon	98.1	77.1	96.8	74.7	50.2
Pennsylvania	98.1	78.6	97.4	71.6	41.0
Rhode Island	96.5	90.5	97.6	73.6	47.1
South Carolina	98.8	71.8	99.3	49.0	26.6
South Dakota	94.8	74.0	95.3	46.7	28.3
Tennessee	98.0	68.9	98.7	50.2	34.4
Texas	98.2	70.1	97.9	45.5	22.9
Utah	99.5	78.7	99.5	81.2	70.4
Vermont	98.6	92.1	98.6	76.5	56.9
Virginia	98.9	76.8	98.9	57.4	39.5
Washington	97.9	85.5	98.7	76.3	58.1
West Virginia	98.8	81.3	98.8	69.8	64.0
State Median	98.3	76.9	97.9	60.2	32.7
State Range	93.2 – 100.0	57.1 – 92.1	92.6 – 100.0	30.3 – 81.2	10.3 – 70.4

TABLE 31a. Among Schools with a Policy Prohibiting Tobacco Use, the Percentage of Schools That Sometimes, Almost Always, or Always Took Specific Actions When Students Were Caught Smoking Cigarettes, Selected U.S. Sites: School Health Profiles, Principal Surveys, 2006 (continued)

Site	Informed parents or guardians	Referred to a school counselor	Referred to a school administrator	Encouraged to participate in an assistance, education, or cessation program	Required to participate in an assistance, education, or cessation program
LOCAL SURVEYS					
Charlotte-Mecklenburg County	97.4	90.1	97.5	79.6	84.7
Chicago	82.5	66.2	81.1	38.7	18.2
Dallas	97.9	66.4	97.8	44.5	24.4
District of Columbia	93.7	74.7	93.7	51.4	46.1
Hillsborough County	92.3	65.6	92.2	48.9	31.0
Los Angeles	98.9	92.1	87.6	74.2	72.2
Memphis	98.3	84.4	98.3	65.1	54.6
Miami	96.5	93.0	95.4	72.3	38.2
Orange County	100.0	85.4	100.0	66.2	51.0
Philadelphia	94.1	78.0	88.4	50.7	23.2
San Diego	100.0	98.2	100.0	90.7	72.8
San Francisco	90.3	96.8	93.7	80.5	78.9
Local Median	97.0	84.9	94.6	65.7	48.6
Local Range	82.5 – 100.0	65.6 – 98.2	81.1 – 100.0	38.7 – 90.7	18.2 – 84.7

* Survey did not include schools from the New York City Department of Education.

TABLE 31b. Among Schools with a Policy Prohibiting Tobacco Use, the Percentage of Schools That Sometimes, Almost Always, or Always Took Specific Actions When Students Were Caught Smoking Cigarettes, Selected U.S. Sites: School Health Profiles, Principal Surveys, 2006

Site	Referred to legal authorities	Placed in detention	Given in-school suspension	Not allowed to participate in extracurricular activities or interscholastic sports	Suspended from school	Expelled from school	Reassigned to an alternative school
STATE SURVEYS							
Alabama	19.0	61.5	67.7	58.5	83.5	7.0	37.7
Alaska	51.8	55.8	54.7	76.8	72.2	6.5	1.6
Arizona	51.2	66.4	69.2	64.2	80.0	12.2	13.7
Arkansas	31.2	62.4	79.1	64.2	75.6	11.3	20.0
Connecticut	28.4	67.0	70.1	63.4	78.3	7.5	3.6
Delaware	18.9	62.5	75.8	54.2	82.9	9.4	10.8
Florida	59.8	59.2	66.8	49.9	72.2	5.3	13.7
Georgia	19.9	58.4	83.5	56.9	72.9	9.4	18.0
Hawaii	17.3	77.8	62.2	53.3	78.8	0.0	2.2
Idaho	95.5	67.8	71.9	92.4	82.2	15.5	11.2
Illinois	40.2	73.3	73.8	87.8	80.5	8.3	8.2
Iowa	78.8	59.9	74.8	94.2	66.4	4.8	2.9
Kansas	69.1	64.5	67.1	86.1	71.5	13.4	3.8
Maine	73.9	56.3	59.2	89.5	85.8	8.2	2.6
Massachusetts	22.2	70.4	60.9	65.3	75.1	1.6	1.6
Michigan	63.2	51.5	54.7	79.2	88.5	8.9	5.5
Mississippi	25.0	57.7	74.3	62.6	79.4	4.5	20.6
Missouri	41.7	70.1	80.6	74.8	72.6	7.9	9.0
Montana	83.9	70.8	71.7	92.3	78.1	9.2	2.7
Nebraska	56.1	75.6	81.1	93.4	73.9	11.4	5.0
New Hampshire	83.7	57.9	61.5	77.6	80.6	3.7	0.6
New York*	19.5	74.1	83.4	82.8	71.6	5.2	4.9
North Carolina	16.6	65.6	75.3	54.5	74.9	7.2	10.3
North Dakota	68.7	67.4	67.4	97.7	60.8	3.7	1.4
Oregon	66.7	62.8	68.8	76.3	79.8	17.1	14.8
Pennsylvania	74.2	53.1	67.3	69.5	67.6	5.9	6.3
Rhode Island	32.7	84.4	55.6	59.8	67.1	2.3	3.5
South Carolina	15.4	62.1	70.1	58.7	89.0	18.9	26.6
South Dakota	63.8	72.8	71.3	93.7	61.8	8.5	2.7
Tennessee	77.2	55.3	66.7	57.6	70.8	14.9	32.1
Texas	65.6	78.3	89.7	74.0	53.2	10.9	29.4
Utah	97.2	59.4	62.7	74.7	86.5	15.2	14.2
Vermont	65.7	54.7	74.8	80.1	73.6	5.0	4.4
Virginia	68.9	50.9	62.5	71.4	88.1	5.6	11.5
Washington	48.6	59.1	60.9	85.0	85.9	11.2	6.4
West Virginia	69.3	63.6	57.3	61.6	73.3	9.0	16.4
State Median	58.0	62.7	69.0	74.4	75.4	8.3	7.3
State Range	15.4 – 97.2	50.9 – 84.4	54.7 – 89.7	49.9 – 97.7	53.2 – 89.0	0.6 – 18.9	0.6 – 37.7

TABLE 31b. Among Schools with a Policy Prohibiting Tobacco Use, the Percentage of Schools That Sometimes, Almost Always, or Always Took Specific Actions When Students Were Caught Smoking Cigarettes, Selected U.S. Sites: School Health Profiles, Principal Surveys, 2006 (continued)

Site	Referred to legal authorities	Placed in detention	Given in-school suspension	Not allowed to participate in extracurricular activities or interscholastic sports	Suspended from school	Expelled from school	Reassigned to an alternative school
LOCAL SURVEYS							
Charlotte-Mecklenburg County	45.0	87.0	67.3	56.1	67.3	9.8	17.8
Chicago	25.3	65.4	64.9	51.5	70.0	7.0	4.7
Dallas	55.1	77.3	93.3	54.5	84.1	17.7	26.5
District of Columbia	35.1	68.2	61.6	57.1	76.9	17.6	31.6
Hillsborough County	75.4	57.4	72.1	36.9	75.2	1.9	3.4
Los Angeles	52.3	65.5	56.5	45.0	74.6	3.9	5.3
Memphis	53.3	64.3	75.9	60.0	92.8	22.3	22.7
Miami	37.3	67.8	85.8	51.7	70.2	10.5	17.4
Orange County	51.9	63.1	88.5	57.1	84.8	9.6	9.6
Philadelphia	19.3	79.5	73.2	64.0	74.9	5.6	10.9
San Diego	66.4	71.8	54.7	61.4	90.9	24.2	24.1
San Francisco	30.7	52.2	61.4	51.3	56.4	0.0	0.0
Local Median	48.5	66.7	69.7	55.3	75.1	9.7	14.2
Local Range	19.3 – 75.4	52.2 – 87.0	54.7 – 93.3	36.9 – 64.0	56.4 – 92.8	0.0 – 24.2	0.0 – 31.6

* Survey did not include schools from the New York City Department of Education.

TABLE 32. Among Schools with a Policy Prohibiting Tobacco Use, the Percentage of Schools That Had Procedures to Inform Specific Groups About the Tobacco Prevention Policy That Prohibits Their Use of Tobacco and a Policy to Inform Students' Families About the Rules Related to Tobacco Use by Students, Selected U.S. Sites: School Health Profiles, Principal Surveys, 2006

Site	Groups informed about policy prohibiting their use of tobacco			Informed students' families of rules related to tobacco use
	Students	Faculty and staff	Visitors	
STATE SURVEYS				
Alabama	99.6	98.8	92.2	98.9
Alaska	99.0	94.2	83.8	97.9
Arizona	98.4	95.4	87.3	97.2
Arkansas	99.7	99.3	96.2	100.0
Connecticut	98.8	93.4	76.4	98.8
Delaware	98.4	93.7	87.1	100.0
Florida	99.7	95.6	89.2	96.9
Georgia	99.7	97.6	83.8	97.8
Hawaii	100.0	97.6	89.8	98.7
Idaho	100.0	94.5	76.9	95.4
Illinois	98.1	94.3	82.5	96.9
Iowa	100.0	94.7	78.1	97.7
Kansas	98.4	93.5	77.1	97.9
Maine	99.3	98.2	87.3	98.5
Massachusetts	99.5	98.3	85.0	99.1
Michigan	99.0	95.4	81.0	97.6
Mississippi	100.0	97.6	92.5	98.3
Missouri	99.4	95.5	78.8	98.6
Montana	99.1	98.3	96.1	99.6
Nebraska	99.5	94.4	74.0	99.0
New Hampshire	98.9	95.9	85.6	98.2
New York*	99.7	98.6	93.3	99.4
North Carolina	99.7	98.6	89.1	99.0
North Dakota	100.0	95.9	87.9	98.7
Oregon	100.0	98.6	90.2	100.0
Pennsylvania	99.7	97.4	88.7	98.2
Rhode Island	97.6	96.5	83.9	95.2
South Carolina	99.1	97.5	87.2	98.7
South Dakota	98.9	90.6	77.8	96.7
Tennessee	99.7	97.5	91.0	99.3
Texas	99.5	97.6	89.6	96.8
Utah	100.0	96.0	74.7	98.6
Vermont	99.3	96.3	90.3	100.0
Virginia	100.0	98.9	92.7	99.6
Washington	99.3	95.6	90.0	98.9
West Virginia	100.0	99.4	96.6	100.0
State Median	99.5	96.4	87.3	98.7
State Range	97.6 – 100.0	90.6 – 99.4	74.0 – 96.6	95.2 – 100.0

TABLE 32. Among Schools with a Policy Prohibiting Tobacco Use, the Percentage of Schools That Had Procedures to Inform Specific Groups About the Tobacco Prevention Policy That Prohibits Their Use of Tobacco and a Policy to Inform Students' Families About the Rules Related to Tobacco Use by Students, Selected U.S. Sites: School Health Profiles, Principal Surveys, 2006 (continued)

Site	Groups informed about policy prohibiting their use of tobacco			Informed students' families of rules related to tobacco use
	Students	Faculty and staff	Visitors	
LOCAL SURVEYS				
Charlotte-Mecklenburg County	100.0	100.0	94.8	100.0
Chicago	96.9	96.4	85.2	92.8
Dallas	97.9	100.0	90.8	91.5
District of Columbia	95.8	81.3	75.9	92.5
Hillsborough County	97.8	81.6	69.5	100.0
Los Angeles	100.0	100.0	96.0	100.0
Memphis	98.0	98.0	89.4	94.6
Miami	98.8	96.4	86.7	93.0
Orange County	100.0	100.0	93.8	100.0
Philadelphia	98.0	93.8	86.6	96.0
San Diego	100.0	85.9	85.9	96.1
San Francisco	97.2	91.4	78.8	90.9
Local Median	98.0	96.4	86.7	95.3
Local Range	95.8 – 100.0	81.3 – 100.0	69.5 – 96.0	90.9 – 100.0

* Survey did not include schools from the New York City Department of Education.

TABLE 33. Percentage of All Schools That Prohibited Tobacco Advertising in Specific Locations, Through Sponsorship of School Events, or on Student Apparel or Merchandise, or Prohibited All Tobacco Advertising, Selected U.S. Sites: School Health Profiles, Principal Surveys, 2006

Site	In school building	On school grounds*	On school buses, or other vehicles†	In school publications	Through sponsorship of school events	On tobacco brand-name apparel or merchandise‡	All tobacco advertising§
STATE SURVEYS							
Alabama	96.6	95.8	95.7	96.3	94.9	98.3	89.8
Alaska	96.4	94.8	95.4	94.1	90.4	84.7	75.8
Arizona	96.4	96.4	96.1	93.7	93.3	98.8	88.7
Arkansas	95.4	94.6	95.4	93.2	91.2	95.8	81.8
Connecticut	95.7	95.7	95.3	95.3	92.9	87.5	77.9
Delaware	100.0	100.0	98.5	98.5	92.7	85.3	78.0
Florida	96.7	95.4	95.0	96.0	95.2	97.0	87.7
Georgia	94.2	94.2	92.8	93.0	89.3	97.2	84.0
Hawaii	94.5	95.1	92.1	93.4	94.1	96.3	82.5
Idaho	96.6	95.1	96.1	94.1	95.5	98.6	89.4
Illinois	95.6	95.2	95.6	94.9	95.5	95.7	86.8
Iowa	95.3	94.0	93.6	93.7	92.2	96.5	86.5
Kansas	93.6	93.5	93.2	93.1	92.2	95.1	84.2
Maine	95.3	95.6	94.6	95.0	93.8	94.8	84.4
Massachusetts	94.2	93.6	93.3	93.1	91.5	79.1	69.5
Michigan	94.7	93.2	92.9	92.6	92.6	94.8	84.2
Mississippi	95.5	95.5	95.5	93.2	90.3	96.7	84.3
Missouri	92.1	91.4	91.3	91.5	91.5	95.3	80.5
Montana	98.4	98.1	98.0	98.0	95.9	96.5	91.6
Nebraska	93.6	92.3	91.8	93.2	92.5	98.4	84.6
New Hampshire	93.9	93.9	93.3	93.3	94.5	95.7	85.7
New York¶	93.3	92.4	92.7	93.0	95.0	82.6	71.8
North Carolina	93.2	92.2	92.2	92.2	89.8	84.7	71.5
North Dakota	93.7	93.2	92.3	93.7	94.6	96.7	87.3
Oregon	92.6	92.7	93.1	92.6	92.3	98.3	85.4
Pennsylvania	93.0	91.8	92.4	92.4	91.7	93.6	80.2
Rhode Island	95.4	96.5	95.3	94.1	89.5	78.2	68.8
South Carolina	97.2	97.2	96.4	95.4	93.9	90.7	82.1
South Dakota	92.2	90.8	92.2	90.6	91.5	98.5	85.5
Tennessee	95.6	93.9	94.6	93.1	92.0	95.4	83.0
Texas	95.4	94.9	94.8	93.8	92.1	98.4	86.6
Utah	95.9	94.0	95.0	93.9	93.7	98.9	87.0
Vermont	98.6	98.6	96.5	98.6	95.5	96.9	88.9
Virginia	95.4	94.3	94.6	94.9	93.1	91.2	81.3
Washington	98.4	97.8	96.3	95.4	94.0	94.1	84.8
West Virginia	95.8	95.3	95.4	93.8	91.5	95.5	84.9
State Median	95.4	94.7	94.7	93.7	92.7	95.7	84.4
State Range	92.1 – 100.0	90.8 – 100.0	91.3 – 98.5	90.6 – 98.6	89.3 – 95.9	78.2 – 98.9	68.8 – 91.6

TABLE 33. Percentage of All Schools That Prohibited Tobacco Advertising in Specific Locations, Through Sponsorship of School Events, or on Student Apparel or Merchandise, or Prohibited All Tobacco Advertising, Selected U.S. Sites: School Health Profiles, Principal Surveys, 2006 (continued)

Site	In school building	On school grounds*	On school buses, or other vehicles†	In school publications	Through sponsorship of school events	On tobacco brand-name apparel or merchandise‡	All tobacco advertising§
LOCAL SURVEYS							
Charlotte-Mecklenburg County	95.2	95.2	95.2	95.2	80.8	80.9	68.9
Chicago	92.6	89.5	89.0	90.4	88.4	90.0	73.2
Dallas	91.7	91.7	89.7	89.5	87.9	97.9	79.2
District of Columbia	90.3	90.3	90.3	87.4	84.1	71.6	52.2
Hillsborough County	90.9	87.2	85.5	87.5	90.8	90.8	74.1
Los Angeles	97.8	97.8	97.8	95.7	94.3	94.5	85.3
Memphis	93.0	93.0	93.0	91.5	87.7	98.3	82.5
Miami	95.7	94.6	93.5	95.5	94.3	92.2	79.8
Orange County	91.3	88.2	84.7	88.8	97.5	97.5	80.1
Philadelphia	98.0	93.3	98.0	94.2	90.4	88.7	72.6
San Diego	100.0	100.0	98.2	98.2	92.8	100.0	91.0
San Francisco	94.4	91.6	91.6	91.6	97.1	85.3	79.2
Local Median	93.7	92.4	92.3	91.6	90.6	91.5	79.2
Local Range	90.3 – 100.0	87.2 – 100.0	84.7 – 98.2	87.4 – 98.2	80.8 – 97.5	71.6 – 100.0	52.2 – 91.0

* Including on the outside of the school building, on playing fields, or other areas of the campus.

† Used to transport students.

‡ Prohibited students from wearing tobacco brand-name apparel or carrying merchandise with tobacco company names, logos, or cartoon characters.

§ In school buildings, on school grounds, on school buses or other vehicles, in school publications, and through sponsorship of school events; and prohibiting students from wearing tobacco brand name apparel or carrying merchandise with tobacco company names, logos, or cartoon characters.

|| Survey did not include schools from the New York City Department of Education.

TABLE 34. Percentage of All Schools That Provided Referrals to Tobacco Cessation Programs for Specific Groups and Posted Signs Marking a Tobacco-Free School Zone,* Selected U.S. Sites: School Health Profiles, Principal Surveys, 2006

Site	Provided Referrals		Posted signs marking a tobacco-free school zone
	Faculty and Staff	Students	
STATE SURVEYS			
Alabama	13.2	19.7	77.9
Alaska	11.8	36.4	68.4
Arizona	21.9	41.0	83.4
Arkansas	15.7	33.0	85.2
Connecticut	24.6	48.3	38.0
Delaware	29.8	55.8	70.9
Florida	23.4	49.8	61.3
Georgia	23.1	24.0	64.7
Hawaii	17.9	61.9	66.5
Idaho	17.4	65.2	62.3
Illinois	16.8	35.4	55.7
Iowa	17.5	47.5	59.8
Kansas	12.0	32.7	56.3
Maine	38.3	67.6	72.5
Massachusetts	36.9	53.2	61.4
Michigan	16.6	45.6	56.3
Mississippi	10.5	17.8	85.3
Missouri	14.0	24.4	56.4
Montana	13.2	44.0	93.2
Nebraska	20.5	40.4	60.9
New Hampshire	40.4	63.5	83.0
New York [†]	32.1	56.7	72.6
North Carolina	29.9	50.2	74.0
North Dakota	28.9	44.2	71.1
Oregon	29.3	63.2	79.1
Pennsylvania	22.1	62.3	62.0
Rhode Island	39.0	70.8	73.2
South Carolina	25.4	39.0	47.1
South Dakota	11.6	30.5	59.5
Tennessee	9.9	33.5	59.6
Texas	11.1	23.9	80.7
Utah	21.4	81.2	54.7
Vermont	32.1	65.8	68.9
Virginia	22.5	50.8	61.2
Washington	29.0	74.3	88.4
West Virginia	35.8	71.4	90.7
State Median	22.0	47.9	67.5
State Range	9.9 – 40.4	17.8 – 81.2	38.0 – 93.2

TABLE 34. Percentage of All Schools That Provided Referrals to Tobacco Cessation Programs for Specific Groups and Posted Signs Marking a Tobacco-Free School Zone,* Selected U.S. Sites: School Health Profiles, Principal Surveys, 2006 (continued)

Site	Provided Referrals		Posted signs marking a tobacco-free school zone
	Faculty and Staff	Students	
LOCAL SURVEYS			
Charlotte-Mecklenburg County	24.3	83.1	88.2
Chicago	12.3	19.8	41.3
Dallas	12.4	30.6	81.7
District of Columbia	16.5	39.2	66.9
Hillsborough County	11.3	49.8	47.4
Los Angeles	39.5	85.9	85.3
Memphis	20.4	60.3	54.7
Miami	27.3	44.9	83.3
Orange County	32.9	57.1	85.1
Philadelphia	15.4	38.3	28.4
San Diego	42.7	96.4	79.5
San Francisco	21.0	77.3	82.5
Local Median	20.7	53.5	80.6
Local Range	11.3 – 42.7	19.8 – 96.4	28.4 – 88.2

* Prohibited the use of all tobacco, including cigarettes, smokeless tobacco (i.e., chewing tobacco, snuff, or dip), cigars, and pipes; by students, faculty and school staff, and visitors; in school buildings, outside on school grounds (including parking lots and playing fields), on school buses or other vehicles used to transport students, and at off-campus school-sponsored events.

† Survey did not include schools from the New York City Department of Education.

TABLE 35a. Percentage of All Schools That Implemented Specific Safety and Security Measures, Selected U.S. Sites: School Health Profiles, Principal Surveys, 2006

Site	Required visitors to report to main office or reception area upon arrival	Maintained a closed campus	Used staff or adult volunteers to monitor halls during and between classes	Routinely conducted locker searches	Required school uniforms
STATE SURVEYS					
Alabama	100.0	97.5	96.5	66.6	21.7
Alaska	92.3	40.8	74.2	25.0	2.8
Arizona	99.7	86.7	90.2	20.1	30.2
Arkansas	100.0	97.1	91.6	61.8	8.6
Connecticut	99.5	94.7	84.4	17.5	7.6
Delaware	100.0	95.5	94.0	29.9	22.6
Florida	100.0	91.5	95.3	40.0	26.6
Georgia	100.0	96.8	96.0	62.2	4.6
Hawaii	98.4	94.1	88.5	2.3	32.2
Idaho	98.4	54.0	94.5	62.2	0.0
Illinois	100.0	86.1	84.1	56.6	4.9
Iowa	98.4	75.6	87.5	40.1	0.7
Kansas	99.2	84.5	93.7	45.0	1.7
Maine	97.7	83.3	74.3	20.8	0.3
Massachusetts	99.6	90.1	85.2	30.9	6.5
Michigan	99.4	88.1	87.5	53.9	14.6
Mississippi	100.0	94.9	93.8	58.9	24.9
Missouri	100.0	95.9	88.9	56.1	4.4
Montana	99.2	43.6	91.9	55.4	0.4
Nebraska	99.1	71.7	92.5	52.6	0.4
New Hampshire	99.5	82.3	81.5	16.7	0.0
New York*	99.8	83.2	87.2	42.6	2.6
North Carolina	100.0	90.4	92.9	56.3	9.1
North Dakota	95.0	52.9	90.9	42.1	1.3
Oregon	99.1	66.4	87.8	33.1	0.9
Pennsylvania	100.0	91.2	89.6	47.8	21.9
Rhode Island	98.8	93.2	89.7	33.7	2.3
South Carolina	100.0	93.9	93.0	60.2	6.8
South Dakota	95.8	59.5	91.3	56.3	0.0
Tennessee	100.0	98.4	92.3	62.1	13.9
Texas	100.0	85.5	95.9	60.7	20.0
Utah	98.3	58.9	91.4	56.9	3.3
Vermont	100.0	75.2	73.9	12.4	0.7
Virginia	100.0	95.7	93.3	49.4	3.2
Washington	99.7	72.0	87.8	26.6	3.5
West Virginia	100.0	97.9	92.2	66.5	0.4
State Median	99.7	87.4	91.1	48.6	4.0
State Range	92.3 – 100.0	40.8 – 98.4	73.9 – 96.5	2.3 – 66.6	0.0 – 32.2

TABLE 35a. Percentage of All Schools That Implemented Specific Safety and Security Measures, Selected U.S. Sites: School Health Profiles, Principal Surveys, 2006 (continued)

Site	Required visitors to report to main office or reception area upon arrival	Maintained a closed campus	Used staff or adult volunteers to monitor halls during and between classes	Routinely conducted locker searches	Required school uniforms
LOCAL SURVEYS					
Charlotte-Mecklenburg County	100.0	95.1	97.7	57.1	14.2
Chicago	100.0	97.0	92.3	40.2	77.6
Dallas	100.0	93.7	95.8	50.1	68.7
District of Columbia	100.0	93.2	87.4	67.8	34.6
Hillsborough County	100.0	100.0	94.6	35.1	22.6
Los Angeles	98.9	96.7	96.8	62.7	41.6
Memphis	100.0	100.0	96.7	71.3	100.0
Miami	100.0	97.7	94.5	22.2	69.0
Orange County	100.0	100.0	96.9	30.4	0.0
Philadelphia	99.0	98.3	93.7	35.9	97.5
San Diego	100.0	100.0	92.8	7.7	14.5
San Francisco	100.0	77.9	91.6	17.8	31.8
Local Median	100.0	97.4	94.6	38.1	38.1
Local Range	98.9 – 100.0	77.9 – 100.0	87.4 – 97.7	7.7 – 71.3	0.0 – 100.0

* Survey did not include schools from the New York City Department of Education.

TABLE 35b. Percentage of All Schools That Implemented Specific Safety and Security Measures, Selected U.S. Sites: School Health Profiles, Principal Surveys, 2006

Site	Required students to wear identification badges	Used metal detectors	Used security or surveillance cameras	Used police, school resource officers, or security guards during the regular school day
STATE SURVEYS				
Alabama	10.7	37.6	85.2	65.7
Alaska	1.3	1.9	20.5	20.2
Arizona	13.5	3.6	30.2	50.2
Arkansas	9.0	17.7	55.5	63.9
Connecticut	8.3	4.8	49.0	57.8
Delaware	3.0	3.0	65.6	67.0
Florida	18.4	22.2	55.5	85.1
Georgia	13.3	29.7	75.5	85.5
Hawaii	42.8	1.6	20.9	76.9
Idaho	2.6	0.9	58.2	64.0
Illinois	10.7	6.2	58.2	38.2
Iowa	1.5	0.8	35.0	24.1
Kansas	4.4	3.1	40.2	41.3
Maine	1.0	1.4	35.1	28.6
Massachusetts	12.4	6.9	55.6	53.6
Michigan	11.3	10.8	54.4	47.8
Mississippi	12.2	30.7	69.9	67.6
Missouri	10.4	7.7	50.9	51.2
Montana	1.7	2.4	35.5	35.4
Nebraska	1.3	1.2	42.7	26.1
New Hampshire	4.3	1.1	34.2	48.1
New York*	9.1	6.8	64.3	64.5
North Carolina	6.1	27.0	64.4	81.6
North Dakota	1.9	2.6	30.9	14.9
Oregon	1.6	0.0	36.5	43.8
Pennsylvania	9.5	19.8	74.6	53.9
Rhode Island	4.5	0.0	46.0	61.2
South Carolina	40.8	29.0	68.0	90.1
South Dakota	3.8	1.4	65.2	24.8
Tennessee	10.8	26.4	85.3	68.0
Texas	21.0	17.3	56.2	61.5
Utah	2.8	0.0	67.0	70.9
Vermont	0.7	0.8	30.1	30.5
Virginia	6.5	25.3	63.2	89.0
Washington	3.4	3.3	44.0	52.7
West Virginia	4.6	12.2	69.3	32.9
State Median	6.3	4.2	55.5	53.8
State Range	0.7 – 42.8	0.0 – 37.6	20.5 – 85.3	14.9 – 90.1

TABLE 35b. Percentage of All Schools That Implemented Specific Safety and Security Measures, Selected U.S. Sites: School Health Profiles, Principal Surveys, 2006 (continued)

Site	Required students to wear identification badges	Used metal detectors	Used security or surveillance cameras	Used police, school resource officers, or security guards during the regular school day
LOCAL SURVEYS				
Charlotte-Mecklenburg County	9.6	55.0	39.1	97.7
Chicago	23.3	82.5	79.5	97.9
Dallas	98.0	89.7	31.5	94.0
District of Columbia	66.3	93.2	100.0	100.0
Hillsborough County	17.1	7.6	6.2	100.0
Los Angeles	13.3	95.3	35.4	97.3
Memphis	22.4	95.0	96.6	85.6
Miami	45.2	50.3	74.9	97.7
Orange County	26.7	8.1	90.7	96.9
Philadelphia	34.0	55.2	67.7	100.0
San Diego	19.9	1.8	18.3	81.8
San Francisco	3.2	3.2	33.8	88.5
Local Median	22.9	55.1	53.4	97.5
Local Range	3.2 – 98.0	1.8 – 95.3	6.2 – 100.0	81.8 – 100.0

* Survey did not include schools from the New York City Department of Education.

TABLE 36. Percentage of All Schools That Had or Participated in Specific Violence Prevention Programs and That Had a Comprehensive Plan to Address Crisis Preparedness, Response, and Recovery in the Event of a Natural Disaster or Other Emergency or Crisis Situation, Selected U.S. Sites: School Health Profiles, Principal Surveys, 2006

Site	Violence prevention programs				Comprehensive plan to address crisis preparedness, response, and recovery
	Peer mediation program	Safe-passage to school program	Program to prevent gang violence	Program to prevent bullying	
STATE SURVEYS					
Alabama	38.3	10.5	36.5	68.3	99.7
Alaska	23.6	3.0	12.8	55.7	96.6
Arizona	39.0	10.4	41.0	68.6	91.0
Arkansas	29.1	5.6	26.6	69.1	99.3
Connecticut	61.4	5.6	21.6	72.0	96.1
Delaware	38.8	6.1	23.8	56.8	100.0
Florida	66.0	14.5	49.4	69.0	99.0
Georgia	60.9	9.1	41.6	66.5	98.8
Hawaii	56.4	3.4	42.0	61.9	98.4
Idaho	43.9	4.2	28.1	59.3	94.9
Illinois	40.9	5.5	25.6	63.7	98.3
Iowa	27.6	2.8	11.5	62.4	97.2
Kansas	23.9	4.0	14.8	58.0	97.9
Maine	32.0	5.1	8.1	65.3	97.1
Massachusetts	57.4	6.0	21.9	67.6	96.4
Michigan	46.5	6.0	18.8	59.6	97.5
Mississippi	21.8	8.1	21.1	30.8	97.6
Missouri	39.1	6.2	18.6	56.7	98.2
Montana	27.2	5.3	19.1	64.9	93.7
Nebraska	19.4	4.9	17.1	61.2	98.7
New Hampshire	48.6	2.2	11.1	69.4	96.5
New York*	57.3	8.0	22.2	69.1	100.0
North Carolina	48.9	8.5	36.3	67.7	99.3
North Dakota	15.8	1.9	16.1	61.7	87.1
Oregon	38.6	4.4	26.0	63.2	95.6
Pennsylvania	54.9	12.5	19.1	72.2	98.5
Rhode Island	47.5	3.5	24.0	55.8	94.1
South Carolina	43.3	8.1	37.6	54.7	98.8
South Dakota	15.8	2.5	8.5	46.5	90.3
Tennessee	42.2	8.8	33.8	76.2	97.2
Texas	37.9	11.1	38.4	64.8	97.6
Utah	40.4	20.8	55.7	72.8	98.5
Vermont	30.8	1.5	8.0	79.6	97.1
Virginia	59.3	8.9	44.9	71.1	100.0
Washington	32.0	5.0	23.9	70.6	98.7
West Virginia	78.8	7.5	27.2	83.6	96.1
State Median	39.8	5.8	23.9	65.1	97.6
State Range	15.8 – 78.8	1.5 – 20.8	8.0 – 55.7	30.8 – 83.6	87.1 – 100.0

TABLE 36. Percentage of All Schools That Had or Participated in Specific Violence Prevention Programs and That Had a Comprehensive Plan to Address Crisis Preparedness, Response, and Recovery in the Event of a Natural Disaster or Other Emergency or Crisis Situation, Selected U.S. Sites: School Health Profiles, Principal Surveys, 2006 (continued)

Site	Violence prevention programs				Comprehensive plan to address crisis preparedness, response, and recovery
	Peer mediation program	Safe-passage to school program	Program to prevent gang violence	Program to prevent bullying	
LOCAL SURVEYS					
Charlotte-Mecklenburg County	76.3	22.7	53.9	68.1	100.0
Chicago	42.1	60.0	67.4	66.7	98.2
Dallas	64.8	20.7	52.0	53.9	96.0
District of Columbia	80.6	33.7	68.0	45.3	100.0
Hillsborough County	92.5	12.6	55.1	79.6	100.0
Los Angeles	61.1	50.2	60.3	72.8	98.9
Memphis	73.6	12.0	89.7	89.9	100.0
Miami	88.6	27.0	72.3	87.3	100.0
Orange County	93.8	41.6	84.5	96.9	100.0
Philadelphia	66.4	39.9	50.8	78.6	99.0
San Diego	42.0	7.1	42.1	56.6	100.0
San Francisco	58.1	6.5	55.9	78.3	100.0
Local Median	70.0	24.9	58.1	75.6	100.0
Local Range	42.0 – 93.8	6.5 – 60.0	42.1 – 89.7	45.3 – 96.9	96.0 – 100.0

* Survey did not include schools from the New York City Department of Education.

TABLE 37. Percentage of All Schools That Had a School Nurse Who Provided Standard Health Services and the Percentage Where a Student Would Ever Be Permitted to Carry and Self-Administer Specific Medications, Selected U.S. Sites: School Health Profiles, Principal Surveys, 2006

Site	School nurse who provided standard health services to students	Medication				
		Prescription quick-relief inhaler	Epinephrine auto-injector (e.g., EpiPen®)	Insulin or other injected medications	Any other prescribed medications	Any over-the-counter medications
STATE SURVEYS						
Alabama	93.3	72.6	35.6	20.9	2.6	3.1
Alaska	38.1	67.3	36.3	31.5	21.2	27.3
Arizona	66.7	62.1	34.2	20.0	5.8	7.8
Arkansas	98.8	82.1	47.1	31.0	7.9	10.5
Connecticut	98.5	66.4	52.5	33.3	9.7	8.9
Delaware	100.0	51.4	23.9	10.5	3.0	4.4
Florida	71.4	65.7	47.4	19.3	6.5	8.4
Georgia	86.7	76.1	36.3	23.1	5.6	8.5
Hawaii	74.0	80.5	44.7	31.0	43.1	49.0
Idaho	63.1	86.7	55.4	52.7	29.1	39.8
Illinois	78.7	78.6	43.6	25.4	7.3	9.1
Iowa	85.9	80.7	45.1	27.4	12.1	24.7
Kansas	82.8	84.5	49.7	40.0	16.6	25.5
Maine	92.6	82.5	58.9	33.4	5.6	9.3
Massachusetts	98.8	66.7	52.6	23.9	6.7	9.9
Michigan	32.2	85.5	48.3	33.3	15.6	14.3
Mississippi	71.6	69.4	32.3	29.0	13.2	13.2
Missouri	98.2	67.8	29.9	21.6	8.7	14.4
Montana	59.3	87.6	45.5	46.8	21.7	33.8
Nebraska	87.3	75.9	29.6	32.3	11.4	19.7
New Hampshire	98.9	78.3	64.1	35.4	3.9	6.0
New York*	99.3	62.1	44.0	27.6	11.6	14.8
North Carolina	91.9	87.1	55.7	44.1	18.1	20.3
North Dakota	28.9	86.6	51.0	49.5	23.8	30.5
Oregon	60.9	68.6	35.5	29.9	15.4	30.7
Pennsylvania	99.7	71.4	42.1	18.8	3.3	5.2
Rhode Island	98.8	72.2	58.1	36.0	11.8	21.0
South Carolina	97.2	75.7	48.0	34.9	14.0	11.1
South Dakota	59.4	79.5	33.6	34.7	25.2	42.2
Tennessee	89.3	73.4	43.3	25.6	4.6	6.0
Texas	95.0	76.1	40.1	28.4	5.3	5.6
Utah	71.7	95.1	63.8	64.4	45.2	59.3
Vermont	99.3	69.0	53.6	43.1	10.7	13.8
Virginia	96.7	78.0	51.7	22.3	5.7	8.5
Washington	92.9	81.7	47.5	36.6	15.9	20.7
West Virginia	95.5	68.8	32.3	25.8	8.5	11.4
State Median	90.6	76.0	45.3	31.0	11.1	13.5
State Range	28.9 – 100.0	51.4 – 95.1	23.9 – 64.1	10.5 – 64.4	2.6 – 45.2	3.1 – 59.3

TABLE 37. Percentage of All Schools That Had a School Nurse Who Provided Standard Health Services and the Percentage Where a Student Would Ever Be Permitted to Carry and Self-Administer Specific Medications, Selected U.S. Sites: School Health Profiles, Principal Surveys, 2006 (continued)

Site	School nurse who provided standard health services to students	Medication				
		Prescription quick-relief inhaler	Epinephrine auto-injector (e.g., EpiPen®)	Insulin or other injected medications	Any other prescribed medications	Any over-the-counter medications
LOCAL SURVEYS						
Charlotte-Mecklenburg County	97.7	73.3	62.6	36.7	14.7	9.8
Chicago	91.0	73.6	26.5	21.8	19.8	10.6
Dallas	98.0	43.2	8.3	2.1	2.1	8.3
District of Columbia	100.0	63.3	14.7	4.0	10.8	13.7
Hillsborough County	97.9	64.5	36.9	10.8	1.7	7.3
Los Angeles	98.9	46.3	23.7	14.9	12.7	13.9
Memphis	96.6	69.0	27.6	24.1	22.3	20.3
Miami	16.0	75.6	67.7	41.6	22.8	31.4
Orange County	83.9	67.1	36.0	11.8	2.5	2.5
Philadelphia	100.0	48.8	13.0	7.2	6.2	8.5
San Diego	98.3	66.4	33.7	36.2	22.9	22.9
San Francisco	31.0	79.5	35.7	30.3	34.3	31.1
Local Median	97.8	66.8	30.7	18.4	13.7	12.2
Local Range	16.0 – 100.0	43.2 – 79.5	8.3 – 67.7	2.1 – 41.6	1.7 – 34.3	2.5 – 31.4

* Survey did not include schools from the New York City Department of Education.

TABLE 38. Percentage of All Schools That Provided Specific Health Services to Students, Selected U.S. Sites: School Health Profiles, Principal Surveys, 2006

Site	Identification or school-based management of chronic health conditions	Identification or school-based management of acute illnesses	Asthma Action Plan or Individualized Health Plan for all students with asthma	Immunizations	Assistance with enrolling in Medicaid or SCHIP*
STATE SURVEYS					
Alabama	82.8	73.4	80.9	45.8	59.9
Alaska	52.5	43.3	34.0	67.4	42.0
Arizona	70.7	60.2	50.1	51.4	47.1
Arkansas	72.5	66.1	67.0	48.5	58.9
Connecticut	80.3	73.5	76.4	41.5	48.8
Delaware	95.5	87.8	75.8	61.2	53.8
Florida	74.2	67.7	65.1	54.5	47.3
Georgia	73.0	63.4	58.3	34.7	37.0
Hawaii	73.8	66.7	75.7	41.4	47.5
Idaho	57.3	50.6	43.5	44.4	53.4
Illinois	69.3	60.3	59.4	41.3	48.0
Iowa	74.6	66.6	58.4	51.4	65.5
Kansas	62.6	55.4	47.9	51.0	55.4
Maine	77.2	69.9	63.0	42.8	49.7
Massachusetts	88.5	82.4	75.1	51.8	73.4
Michigan	54.8	43.4	41.9	39.1	42.0
Mississippi	49.5	43.5	36.0	32.1	44.8
Missouri	79.4	73.4	72.9	60.4	56.2
Montana	61.5	51.9	57.3	61.8	55.3
Nebraska	82.5	69.2	86.8	57.6	58.2
New Hampshire	87.4	79.8	68.5	53.3	75.8
New York†	79.5	73.6	56.7	49.7	51.4
North Carolina	90.4	78.8	82.7	60.1	47.9
North Dakota	51.0	39.0	42.0	80.7	41.4
Oregon	63.2	55.9	55.0	50.1	54.4
Pennsylvania	81.8	74.6	67.7	48.7	66.1
Rhode Island	65.6	59.3	53.6	59.6	51.6
South Carolina	84.5	77.0	67.2	55.3	65.9
South Dakota	38.9	34.3	22.8	44.8	45.8
Tennessee	72.4	63.4	67.5	46.1	37.5
Texas	82.8	74.6	65.6	64.2	62.7
Utah	70.7	64.8	62.6	53.6	51.8
Vermont	92.8	81.4	70.1	49.5	86.4
Virginia	85.4	75.2	75.5	39.4	55.6
Washington	83.9	77.3	81.1	48.2	78.6
West Virginia	86.0	78.0	75.0	44.0	65.4
State Median	74.4	67.2	65.4	49.9	53.6
State Range	38.9 – 95.5	34.3 – 87.8	22.8 – 86.8	32.1 – 80.7	37.0 – 86.4

TABLE 38. Percentage of All Schools That Provided Specific Health Services to Students, Selected U.S. Sites: School Health Profiles, Principal Surveys, 2006 (continued)

Site	Identification or school-based management of chronic health conditions	Identification or school-based management of acute illnesses	Asthma Action Plan or Individualized Health Plan for all students with asthma	Immunizations	Assistance with enrolling in Medicaid or SCHIP*
LOCAL SURVEYS					
Charlotte-Mecklenburg County	87.8	68.3	80.2	41.6	50.2
Chicago	81.0	70.1	74.7	66.9	71.2
Dallas	73.7	67.7	67.1	59.6	66.7
District of Columbia	89.2	63.0	51.0	76.8	33.3
Hillsborough County	79.3	68.8	64.5	69.3	52.4
Los Angeles	79.4	74.9	71.2	62.2	66.0
Memphis	74.0	65.2	71.3	48.0	44.6
Miami	38.8	32.9	19.7	47.5	38.3
Orange County	70.8	66.9	64.6	65.8	67.7
Philadelphia	92.8	87.8	86.9	51.5	85.8
San Diego	94.5	85.5	79.7	69.5	83.8
San Francisco	71.0	65.0	49.3	36.1	29.4
Local Median	79.4	68.0	69.2	60.9	59.2
Local Range	38.8 – 94.5	32.9 – 87.8	19.7 – 86.9	36.1 – 76.8	29.4 – 85.8

* State Children's Health Insurance Program.

† Survey did not include schools from the New York City Department of Education.

TABLE 39a. Percentage of All Schools with a Policy on Students and/or Staff who Have HIV* Infection or AIDS† and Among Those Schools, Percentage Whose Policy Addressed Specific Issues, Selected U.S. Sites: School Health Profiles, Principal Surveys, 2006

Site	Had a policy	Issue addressed by policy			
		Attendance at school of students with HIV infection	Procedures to protect HIV-infected students and staff from discrimination	Maintenance of confidentiality of HIV-infected students and staff	Work site safety
STATE SURVEYS					
Alabama	62.2	96.5	98.5	99.3	100.0
Alaska	40.5	90.5	94.7	93.5	93.6
Arizona	41.7	91.9	96.3	96.2	98.4
Arkansas	33.1	90.6	98.9	98.6	98.2
Connecticut	59.3	92.3	97.7	98.5	96.2
Delaware	30.3	100.0	100.0	100.0	94.8
Florida	43.9	90.6	96.7	100.0	100.0
Georgia	42.6	92.1	99.2	100.0	99.1
Hawaii	51.2	93.0	100.0	100.0	100.0
Idaho	60.4	93.0	93.3	96.0	94.0
Illinois	39.7	92.2	96.2	99.2	99.2
Iowa	42.9	95.1	94.1	95.0	96.6
Kansas	39.6	94.9	95.0	100.0	98.0
Maine	66.4	96.7	97.9	99.5	99.4
Massachusetts	58.2	93.3	97.8	98.9	97.0
Michigan	32.3	90.1	95.5	98.3	100.0
Mississippi	27.0	97.6	95.1	92.7	97.6
Missouri	52.9	91.5	96.0	98.1	96.7
Montana	48.3	94.4	97.8	99.3	94.7
Nebraska	53.5	94.2	95.9	99.2	96.9
New Hampshire	76.9	94.5	99.3	100.0	98.4
New York‡	59.0	85.2	96.5	99.0	97.5
North Carolina	36.0	91.1	95.7	99.0	100.0
North Dakota	39.7	93.6	96.8	98.5	96.8
Oregon	66.4	94.9	98.4	99.0	98.9
Pennsylvania	59.9	92.8	98.1	99.6	98.0
Rhode Island	64.8	98.2	100.0	100.0	100.0
South Carolina	57.9	92.7	96.5	100.0	99.3
South Dakota	51.9	95.8	95.1	98.0	95.6
Tennessee	58.2	93.6	98.8	99.4	98.2
Texas	30.8	87.7	95.4	98.9	95.9
Utah	52.5	98.2	99.1	99.1	100.0
Vermont	89.5	97.6	99.2	100.0	100.0
Virginia	55.2	90.2	96.6	98.6	96.5
Washington	45.4	94.8	97.9	97.9	98.9
West Virginia	27.1	93.8	100.0	100.0	97.9
State Median	51.6	93.5	97.3	99.1	98.1
State Range	27.0 – 89.5	85.2 – 100.0	93.3 – 100.0	92.7 – 100.0	93.6 – 100.0

TABLE 39a. Percentage of All Schools with a Policy on Students and/or Staff who Have HIV* Infection or AIDS† and Among Those Schools, Percentage Whose Policy Addressed Specific Issues, Selected U.S. Sites: School Health Profiles, Principal Surveys, 2006 (continued)

Site	Had a policy	Issue addressed by policy			
		Attendance at school of students with HIV infection	Procedures to protect HIV-infected students and staff from discrimination	Maintenance of confidentiality of HIV-infected students and staff	Work site safety
LOCAL SURVEYS					
Charlotte-Mecklenburg County	35.2	93.0	93.0	93.0	93.0
Chicago	48.3	90.4	100.0	100.0	99.0
Dallas	30.5	93.0	100.0	100.0	100.0
District of Columbia	28.1	65.4	100.0	100.0	75.7
Hillsborough County	45.3	82.8	100.0	100.0	100.0
Los Angeles	57.7	91.5	98.0	100.0	97.5
Memphis	51.1	89.8	100.0	100.0	100.0
Miami	44.6	78.3	97.3	100.0	91.8
Orange County	37.8	100.0	100.0	100.0	100.0
Philadelphia	57.7	85.2	100.0	100.0	100.0
San Diego	100.0	97.6	100.0	100.0	100.0
San Francisco	50.0	100.0	100.0	100.0	100.0
Local Median	46.8	91.0	100.0	100.0	100.0
Local Range	28.1 – 100.0	65.4 – 100.0	93.0 – 100.0	93.0 – 100.0	75.7 – 100.0

* Human immunodeficiency virus.

† Acquired immunodeficiency syndrome.

‡ Survey did not include schools from the New York City Department of Education.

TABLE 39b. Percentage of All Schools with a Policy on Students and/or Staff who Have HIV* Infection or AIDS† and Among Those Schools, Percentage Whose Policy Addressed Specific Issues, Selected U.S. Sites: School Health Profiles, Principal Surveys, 2006

Site	Issue addressed by policy			
	Confidential counseling for HIV-infected students	Communication of the policy to students, school staff, and parents	Adequate training about HIV infection for school staff	Procedure for implementing the policy
STATE SURVEYS				
Alabama	91.0	95.0	92.3	95.3
Alaska	73.7	84.7	83.7	87.6
Arizona	73.9	83.0	89.3	91.7
Arkansas	88.7	97.4	86.8	96.1
Connecticut	76.2	83.5	87.1	88.2
Delaware	89.4	78.9	89.5	94.8
Florida	87.7	89.1	87.5	94.5
Georgia	82.1	92.0	87.0	90.2
Hawaii	77.6	82.0	91.8	95.7
Idaho	73.9	77.3	67.2	84.7
Illinois	79.2	95.8	94.8	98.0
Iowa	72.7	84.7	90.9	93.1
Kansas	75.6	89.4	87.6	88.6
Maine	78.8	81.7	92.7	91.2
Massachusetts	79.1	84.8	76.0	86.0
Michigan	78.0	85.0	88.1	88.8
Mississippi	85.5	95.3	90.5	95.4
Missouri	76.5	87.9	83.5	93.6
Montana	85.6	93.4	80.2	94.5
Nebraska	75.9	90.1	83.9	92.6
New Hampshire	74.7	85.8	79.6	89.7
New York ‡	68.9	82.9	87.1	86.0
North Carolina	78.5	90.7	91.0	92.9
North Dakota	81.6	95.5	90.8	96.9
Oregon	80.1	89.4	90.4	96.0
Pennsylvania	73.3	89.5	79.7	93.3
Rhode Island	79.9	94.4	94.7	96.3
South Carolina	89.1	86.0	95.5	94.1
South Dakota	83.0	93.8	78.6	87.9
Tennessee	84.7	91.6	85.3	94.0
Texas	85.0	86.2	91.1	94.3
Utah	77.0	86.6	81.5	89.1
Vermont	73.8	86.9	91.8	91.7
Virginia	79.9	83.8	89.2	90.3
Washington	75.5	90.4	95.4	96.3
West Virginia	82.4	92.9	80.9	91.6
State Median	79.0	88.5	87.9	93.0
State Range	68.9 – 91.0	77.3 – 97.4	67.2 – 95.5	84.7 – 98.0

TABLE 39b. Percentage of All Schools with a Policy on Students and/or Staff who Have HIV* Infection or AIDS† and Among Those Schools, Percentage Whose Policy Addressed Specific Issues, Selected U.S. Sites: School Health Profiles, Principal Surveys, 2006 (continued)

Site	Issue addressed by policy			
	Confidential counseling for HIV-infected students	Communication of the policy to students, school staff, and parents	Adequate training about HIV infection for school staff	Procedure for implementing the policy
LOCAL SURVEYS				
Charlotte-Mecklenburg County	69.5	84.6	86.0	78.7
Chicago	90.4	89.7	81.9	93.4
Dallas	86.6	93.2	93.2	86.3
District of Columbia	69.2	79.5	55.1	79.5
Hillsborough County	91.8	91.8	91.8	95.9
Los Angeles	89.0	90.5	91.0	96.0
Memphis	96.2	100.0	96.8	96.8
Miami	89.3	94.7	78.2	88.8
Orange County	91.5	83.0	100.0	91.5
Philadelphia	78.4	78.3	59.2	84.5
San Diego	90.4	92.8	97.7	100.0
San Francisco	88.1	88.1	87.3	100.0
Local Median	89.2	90.1	89.2	92.5
Local Range	69.2 – 96.2	78.3 – 100.0	55.1 – 100.0	78.7 – 100.0

* Human immunodeficiency virus.

† Acquired immunodeficiency syndrome.

‡ Survey did not include schools from the New York City Department of Education.

TABLE 40. Percentage of All Schools That Had One or More School Health Councils* and Engaged Parents and Families in Specific Health Education Activities During the 2005–2006 School Year, Selected U.S. Sites: School Health Profiles, Principal and Lead Health Education Teacher Surveys, 2006

Site	School health council	Provided families with information on school health education	Met with a parents' organization to discuss school health education	Invited family members to attend health education classes
STATE SURVEYS				
Alabama	44.5	63.3	30.3	37.2
Alaska	48.7	52.8	17.8	31.6
Arizona	35.6	53.5	19.8	25.3
Arkansas	68.7	64.9	23.6	35.5
Connecticut	60.1	71.8	29.6	32.6
Delaware	63.4	80.3	31.7	37.6
Florida	49.4	62.7	26.0	29.6
Georgia	50.9	66.5	27.3	42.8
Hawaii	47.2	66.2	17.3	32.4
Idaho	49.7	58.9	14.5	42.9
Illinois	55.1	NA [†]	NA	NA
Iowa	59.5	61.8	16.7	22.3
Kansas	60.6	51.0	10.5	25.0
Maine	69.2	70.2	13.7	32.4
Massachusetts	59.0	80.0	32.8	27.2
Michigan	71.0	70.2	32.2	45.6
Mississippi	34.7	58.8	18.4	38.1
Missouri	69.4	69.6	25.0	28.5
Montana	61.1	63.2	13.1	33.2
Nebraska	54.7	63.3	13.4	24.5
New Hampshire	64.7	79.3	18.1	31.0
New York [‡]	67.4	82.7	32.5	28.2
North Carolina	57.9	66.0	22.1	41.4
North Dakota	54.1	54.3	13.6	29.7
Oregon	52.6	77.6	15.6	39.8
Pennsylvania	70.4	67.5	23.7	28.3
Rhode Island	51.8	71.5	32.5	23.0
South Carolina	56.6	67.4	22.3	45.2
South Dakota	46.7	47.6	13.6	21.0
Tennessee	43.4	59.5	26.1	32.2
Texas	63.0	59.7	23.9	31.4
Utah	45.4	64.4	23.0	41.3
Vermont	73.9	80.5	14.0	41.0
Virginia	49.3	73.8	22.2	37.8
Washington	51.4	NA	NA	NA
West Virginia	49.2	67.0	24.8	32.9
State Median	54.9	66.1	22.3	32.4
State Range	34.7 – 73.9	47.6 – 82.7	10.5 – 32.8	21.0 – 45.6

TABLE 40. Percentage of All Schools That Had One or More School Health Councils* and Engaged Parents and Families in Specific Health Education Activities During the 2005–2006 School Year, Selected U.S. Sites: School Health Profiles, Principal and Lead Health Education Teacher Surveys, 2006 (continued)

Site	School health council	Provided families with information on school health education	Met with a parents' organization to discuss school health education	Invited family members to attend health education classes
LOCAL SURVEYS				
Charlotte-Mecklenburg County	34.5	97.4	24.7	64.2
Chicago	37.9	60.0	37.3	31.9
Dallas	24.1	66.7	24.5	40.8
District of Columbia	49.6	59.9	35.5	59.3
Hillsborough County	35.5	57.2	29.7	18.8
Los Angeles	48.9	80.7	34.7	57.4
Memphis	45.4	82.1	63.3	75.4
Miami	51.0	60.5	18.8	26.3
Orange County	62.7	71.8	22.8	42.7
Philadelphia	53.7	53.7	25.4	27.7
San Diego	47.9	79.7	37.8	56.2
San Francisco	79.0	87.9	53.1	29.0
Local Median	48.4	69.3	32.2	41.8
Local Range	24.1 – 79.0	53.7 – 97.4	18.8 – 63.3	18.8 – 75.4

* A group, committee, or team that offers guidance on the development of policies or coordinates activities on health topics.

† Data not available.

‡ Survey did not include schools from the New York City Department of Education.

TABLE 41. Percentage of All Schools That Asked Students to Participate in Health-Related Community Activities as Part of a Required Health Education Course During the 2005–2006 School Year, Selected U.S. Sites: School Health Profiles, Lead Health Education Teacher Surveys, 2006

Site	Performed volunteer work*	Participated in or attended a health fair	Gathered information about health services [†]	Visited a store to compare prices of health products	Identified potential injury sites [‡]	Identified advertising designed to influence health behaviors [§]	Advocated for a health-related issue	Completed homework or projects that involved family members
STATE SURVEYS								
Alabama	21.4	24.2	47.2	24.7	56.6	57.2	39.1	59.5
Alaska	14.0	30.1	37.1	16.1	34.7	33.3	42.3	53.8
Arizona	6.2	10.5	20.1	9.7	22.4	28.2	23.1	34.1
Arkansas	11.1	19.6	49.4	30.7	59.5	69.4	42.8	72.0
Connecticut	13.4	20.7	33.2	17.3	43.3	61.8	52.3	70.2
Delaware	22.1	32.6	50.3	34.1	61.1	78.4	65.3	75.3
Florida	15.4	14.6	28.2	17.7	30.3	41.5	30.7	44.1
Georgia	17.7	25.8	49.8	28.1	59.5	69.9	44.2	75.1
Hawaii	15.9	28.8	62.3	35.4	54.0	79.5	82.2	77.7
Idaho	16.4	28.4	44.5	30.5	48.6	70.4	49.5	77.9
Iowa	11.1	9.0	29.6	13.1	36.5	46.1	33.4	58.2
Kansas	9.8	12.8	31.9	15.0	30.0	41.4	31.1	59.4
Maine	6.6	11.5	48.8	27.4	48.3	66.0	52.2	72.8
Massachusetts	10.9	16.6	37.0	21.8	40.1	59.8	44.7	68.9
Michigan	9.9	10.1	32.9	15.9	37.7	56.0	38.4	66.2
Mississippi	24.8	30.7	66.4	38.5	70.0	69.0	53.4	77.7
Missouri	10.3	28.1	39.9	22.6	52.3	61.3	41.8	71.2
Montana	10.9	28.9	39.6	21.3	53.1	61.6	45.1	71.0
Nebraska	11.4	16.7	29.2	15.2	43.4	56.1	42.4	68.6
New Hampshire	9.9	16.6	40.2	16.9	43.4	59.9	51.3	71.4
New York	23.1	28.5	62.3	31.6	57.5	79.6	71.1	90.2
North Carolina	10.2	24.5	43.5	23.6	51.3	57.7	44.5	71.1
North Dakota	9.8	35.6	42.5	15.2	45.3	61.4	45.9	70.8
Oregon	15.2	18.2	43.5	25.7	51.2	71.5	50.3	78.0
Pennsylvania	17.2	29.1	42.4	25.6	54.9	70.2	48.6	73.5
Rhode Island	18.1	28.0	46.3	28.0	53.3	70.0	63.2	79.6
South Carolina	14.1	20.0	36.0	23.8	48.1	50.9	39.6	59.5
South Dakota	6.7	23.3	21.5	5.6	28.9	33.8	27.7	54.4
Tennessee	12.6	25.0	35.1	18.8	43.6	41.6	32.2	52.1
Texas	13.8	19.1	31.6	23.3	40.5	49.3	33.8	56.5
Utah	15.7	17.1	32.9	26.9	53.5	73.2	47.8	81.8
Vermont	7.6	19.0	29.7	22.1	34.3	64.2	58.1	72.5
Virginia	11.9	16.7	40.0	29.4	48.9	58.0	40.3	67.4
West Virginia	21.9	35.0	57.8	30.1	66.7	77.2	52.0	75.0
State Median	13.0	22.0	40.0	23.5	48.5	61.4	44.6	71.1
State Range	6.2 – 24.8	9.0 – 35.6	20.1 – 66.4	5.6 – 38.5	22.4 – 70.0	28.2 – 79.6	23.1 – 82.2	34.1 – 90.2

TABLE 41. Percentage of All Schools That Asked Students to Participate in Health-Related Community Activities as Part of a Required Health Education Course During the 2005–2006 School Year, Selected U.S. Sites: School Health Profiles, Lead Health Education Teacher Surveys, 2006 (continued)

Site	Performed volunteer work [*]	Participated in or attended a health fair	Gathered information about health services [†]	Visited a store to compare prices of health products	Identified potential injury sites [‡]	Identified advertising designed to influence health behaviors [§]	Advocated for a health-related issue	Completed homework or projects that involved family members
LOCAL SURVEYS								
Charlotte-Mecklenburg County	7.9	26.9	37.9	48.8	70.0	75.9	56.5	94.5
Chicago	11.4	16.6	28.7	17.5	32.9	29.5	24.0	44.1
Dallas	16.7	20.8	39.6	22.9	37.5	52.1	33.3	45.8
District of Columbia	20.6	45.4	62.1	38.2	44.8	61.1	58.8	59.1
Hillsborough County	8.1	2.0	24.5	2.0	15.9	31.8	17.9	27.5
Los Angeles	19.5	44.3	62.0	43.0	53.4	88.2	59.7	90.7
Memphis	36.9	55.6	61.0	38.6	65.9	66.3	52.0	74.7
Miami	20.3	20.3	30.4	17.7	30.3	38.4	30.3	43.0
Orange County	13.3	13.6	27.3	21.9	30.0	38.3	38.3	41.2
Philadelphia	19.3	40.4	56.2	34.1	53.0	58.5	49.3	68.3
San Diego [¶]	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
San Francisco	14.3	40.8	35.7	17.9	28.6	50.0	42.9	50.0
Local Median	15.5	23.9	36.8	22.4	35.2	51.1	40.6	47.9
Local Range	0.0 – 36.9	0.0 – 55.6	0.0 – 62.1	0.0 – 48.8	0.0 – 70.0	0.0 – 88.2	0.0 – 59.7	0.0 – 94.5

* At a hospital, a local health department, or any other local organization that addresses health issues.

† That are available in the community.

‡ At school, home, or in the community.

§ In the community.

|| Survey did not include schools from the New York City Department of Education.

¶ San Diego does not have a required health education course, but requires that health education be taught in science and physical education classes.

PROFILES 2006

School Health Profiles Characteristics of Health Programs Among Secondary Schools

Printer:

**Please adjust spine
width to fit publication**