


Epidemiology of HIV Infection through 2011

National Center for HIV/AIDS, Viral Hepatitis, STD & TB Prevention

Division of HIV/AIDS Prevention




Diagnoses of HIV Infection among Adults and Adolescents, by Sex, 2008–2011—United States and 6 Dependent Areas



Note. Data include persons with a diagnosis of HIV infection regardless of stage of disease at diagnosis. All displayed data have been statistically adjusted to account for reporting delays, but not for incomplete reporting.



Diagnoses of HIV Infection among Adults and Adolescents, by Transmission Category, 2008–2011—United States and 6 Dependent Areas



Note. Data include persons with a diagnosis of HIV infection regardless of stage of disease at diagnosis. All displayed data have been statistically adjusted to account for reporting delays and missing transmission category, but not for incomplete reporting.


^a Heterosexual contact with a person known to have, or to be at high risk for, HIV infection.

^b Includes hemophilia, blood transfusion, perinatal exposure, and risk factor not reported or not identified.



Diagnoses of HIV Infection among Adults and Adolescents, by Transmission Category, 2011—United States and 6 Dependent Areas

N = 50,007





Note. Data include persons with a diagnosis of HIV infection regardless of stage of disease at diagnosis. All displayed data have been statistically adjusted to account for reporting delays and missing transmission category, but not for incomplete reporting.

^a Heterosexual contact with a person known to have, or to be at high risk for, HIV infection.

^b Includes hemophilia, blood transfusion, perinatal exposure, and risk factor not reported or not identified.



Diagnoses of HIV Infection among Adults and Adolescents, by Sex and Transmission Category, 2011—United States and 6 Dependent Areas



Male-to-male sexual contact

Injection drug use (IDU)

Male-to-male sexual contact and IDU

Heterosexual contact^a

Other^b


Note. Data include persons with a diagnosis of HIV infection regardless of stage of disease at diagnosis. All displayed data have been statistically adjusted to account for reporting delays and missing transmission category, but not for incomplete reporting.

^a Heterosexual contact with a person known to have, or to be at high risk for, HIV infection.

^b Includes hemophilia, blood transfusion, perinatal exposure, and risk factor not reported or not identified.





Diagnoses of HIV Infection among Adults and Adolescents, by Race/Ethnicity, 2008–2011—United States and 6 Dependent Areas



Note. Data include persons with a diagnosis of HIV infection regardless of stage of disease at diagnosis. All displayed data have been statistically adjusted to account for reporting delays, but not for incomplete reporting.
^a Hispanics/Latinos can be of any race.



Diagnoses of HIV Infection among Adults and Adolescents, by Sex and Race/Ethnicity, 2011—United States and 6 Dependent Areas



- American Indian/Alaska Native
- Asian
- Black/African American
- Hispanic/Latino^a
- Native Hawaiian/other Pacific Islander
- White
- Multiple races

Note. Data include persons with a diagnosis of HIV infection regardless of stage of disease at diagnosis. All displayed data have been statistically adjusted to account for reporting delays, but not for incomplete reporting.

^a Hispanics/Latinos can be of any race.



Diagnoses of HIV Infection among Adults and Adolescents, by Transmission Category, 2011—United States and 6 Dependent Areas

Transmission Category	No.	%
Male-to-male sexual contact	30,896	61.8
Injection drug use (IDU)	3,836	7.7
Male-to-male sexual contact and IDU	1,423	2.9
Heterosexual contact ^a	13,801	27.6
Other ^b	51	0.1
Total	50,007	100.0

Note. Data include persons with a diagnosis of HIV infection regardless of stage of disease at diagnosis. All displayed data have been statistically adjusted to account for reporting delays and missing transmission category, but not for incomplete reporting.

^a Heterosexual contact with a person known to have, or to be at high risk for, HIV infection.

^b Includes hemophilia, blood transfusion, perinatal exposure, and risk factor not reported or not identified.



Diagnosed HIV Infections Attributed to Male-to-Male Sexual Contact, by Race/Ethnicity, 2011—United States and 6 Dependent Areas

Race/ethnicity	No.	%
American Indian/Alaska Native	120	0.4
Asian	705	2.3
Black/African American	11,810	38.2
Hispanic/Latino ^a	7,266	23.5
Native Hawaiian/other Pacific Islander	61	0.2
White	10,375	33.6
Multiple races	558	1.8
Total	30,896	100.0

Note. Data include persons with a diagnosis of HIV infection regardless of stage of disease at diagnosis. All displayed data have been statistically adjusted to account for reporting delays and missing transmission category, but not for incomplete reporting.

^a Hispanics/Latinos can be of any race.



Diagnosed HIV Infections Attributed to Heterosexual Contact^a, by Sex and Race/Ethnicity, 2011—United States and 6 Dependent Areas

Race/ethnicity	Males		Females		Total	
	No.	%	No.	%	No.	%
American Indian/Alaska Native	12	0.3	32	0.4	44	0.3
Asian	65	1.4	141	1.6	206	1.5
Black/African American	3,124	65.4	5,882	65.2	9,006	65.3
Hispanic/Latino ^b	979	20.5	1,522	16.9	2,501	18.1
Native Hawaiian/other Pacific Islander	3	0.1	8	0.1	11	0.1
White	537	11.3	1,325	14.7	1,862	13.5
Multiple races	54	1.1	116	1.3	170	1.2
Total	4,775	100.0	9,026	100.0	13,801	100.0

Note. Data include persons with a diagnosis of HIV infection regardless of stage of disease at diagnosis. All displayed data have been statistically adjusted to account for reporting delays and missing transmission category, but not for incomplete reporting.

^a Heterosexual contact with a person known to have, or to be at high risk for, HIV infection.

^b Hispanics/Latinos can be of any race.



Diagnosed HIV Infections Attributed to Male-to-Male Sexual Contact and Injection Drug Use, by Race/Ethnicity, 2011— United States and 6 Dependent Areas

Race/ethnicity	No.	%
American Indian/Alaska Native	12	0.9
Asian	24	1.7
Black/African American	434	30.5
Hispanic/Latino ^a	297	20.9
Native Hawaiian/other Pacific Islander	0	0.0
White	621	43.7
Multiple races	34	2.4
Total	1,423	100.0

Note. Data include persons with a diagnosis of HIV infection regardless of stage of disease at diagnosis. All displayed data have been statistically adjusted to account for reporting delays and missing transmission category, but not for incomplete reporting.

^a Hispanics/Latinos can be of any race.



Diagnosed HIV Infections Attributed to Injection Drug Use, by Sex and Race/Ethnicity, 2011—United States and 6 Dependent Areas

Race/ethnicity	Males		Females		Total	
	No.	%	No.	%	No.	%
American Indian/Alaska Native	17	0.7	19	1.3	36	0.9
Asian	23	1.0	10	0.7	32	0.8
Black/African American	1,087	46.0	716	48.7	1,802	47.0
Hispanic/Latino ^a	707	29.9	252	17.1	959	25.0
Native Hawaiian/other Pacific Islander	4	0.2	1	0.1	6	0.2
White	495	20.9	445	30.3	940	24.5
Multiple races	33	1.4	28	1.9	60	1.6
Total	2,365	100.0	1,471	100.0	3,836	100.0

Note. Data include persons with a diagnosis of HIV infection regardless of stage of disease at diagnosis. All displayed data have been statistically adjusted to account for reporting delays and missing transmission category, but not for incomplete reporting.


^a Hispanics/Latinos can be of any race.



Rates of Diagnoses of HIV Infection among Adults and Adolescents, 2011—United States and 6 Dependent Areas

N = 50,007


Total Rate = 19.1



Note. Data include persons with a diagnosis of HIV infection regardless of stage of disease at diagnosis. All displayed data have been statistically adjusted to account for reporting delays, but not for incomplete reporting.



Rates of Diagnoses of HIV Infection among Adults and Adolescents, by Sex and Race/Ethnicity, 2011—United States



Note. Data include persons with a diagnosis of HIV infection regardless of stage of disease at diagnosis. All displayed data have been statistically adjusted to account for reporting delays, but not for incomplete reporting. Rates are per 100,000 population.

^a Hispanics/Latinos can be of any race.



Diagnoses of HIV Infection among Adult and Adolescent Males, by Race/Ethnicity, 2011—United States

Race/ethnicity	No.	Rate
American Indian/Alaska Native	161	18.0
Asian	821	13.8
Black/African American	16,447	112.8
Hispanic/Latino ^a	8,605	43.4
Native Hawaiian/other Pacific Islander	70	34.2
White	12,041	14.5
Multiple races	679	38.5
Total	38,825	30.8

Note. Data include persons with a diagnosis of HIV infection regardless of stage of disease at diagnosis. All displayed data have been statistically adjusted to account for reporting delays, but not for incomplete reporting. Rates are per 100,000 population.

^a Hispanics/Latinos can be of any race.



Diagnoses of HIV Infection among Adult and Adolescent Females, by Race/Ethnicity, 2011—United States

Race/ethnicity	No.	Rate
American Indian/Alaska Native	51	5.5
Asian	153	2.3
Black/African American	6,595	40.0
Hispanic/Latino ^a	1,530	7.9
Native Hawaiian/other Pacific Islander	8	3.9
White	1,776	2.0
Multiple races	144	7.5
Total	10,257	7.7

Note. Data include persons with a diagnosis of HIV infection regardless of stage of disease at diagnosis. All displayed data have been statistically adjusted to account for reporting delays, but not for incomplete reporting. Rates are per 100,000 population.

^a Hispanics/Latinos can be of any race.



Deaths of Persons with Diagnosed HIV Infection, by Race/Ethnicity, 2010—United States



Race/ethnicity	No.	Rate
American Indian/Alaska Native	90	3.9
Asian ^a	91	0.6
Black/African American	9,517	25.0
Hispanic/Latino ^b	2,846	5.6
Native Hawaiian/other Pacific Islander	10	2.1
White	5,981	3.0
Multiple races	803	14.2
Total^c	19,343	6.3

Note. Data include persons with a diagnosis of HIV infection regardless of stage of disease at diagnosis. Deaths of persons with a diagnosed HIV infection may be due to any cause. All displayed data have been statistically adjusted to account for reporting delays, but not for incomplete reporting. Rates are per 100,000 population.

- ^a Includes Asian/Pacific Islander legacy cases.
- ^b Hispanics/Latinos can be of any race.
- ^c Includes persons of unknown race/ethnicity.



Adults and Adolescents Living with Diagnosed HIV Infection, by Sex and Race/Ethnicity, Year-end 2010—United States and 6 Dependent Areas



- American Indian/Alaska Native
- Asian^a
- Black/African American
- Hispanic/Latino^b
- Native Hawaiian/Other Pacific Islander
- White
- Multiple races

Note. Data include persons with a diagnosis of HIV infection regardless of stage of disease at diagnosis. All displayed data have been statistically adjusted to account for reporting delays, but not for incomplete reporting.

^a Includes Asian/Pacific Islander legacy cases.

^b Hispanics/Latinos can be of any race.


* Total males include 582 persons and total females include 180 persons with unknown race/ethnicity.



Adults and Adolescents Living with Diagnosed HIV Infection, by Sex and Transmission Category, Year-end 2010— United States and 6 Dependent Areas

Males


N = 665,872



- Male-to-male sexual contact
- Injection drug use (IDU)
- Male-to-male sexual contact and IDU

Females

N = 223,045



- Heterosexual contact^a
- Perinatal exposure
- Other^b

Note. Data include persons with a diagnosis of HIV infection regardless of stage of disease at diagnosis. All displayed data have been statistically adjusted to account for reporting delays and missing transmission category, but not for incomplete reporting.


^a Heterosexual contact with a person known to have, or to be at high risk for, HIV infection.

^b Includes hemophilia, blood transfusion, and risk factor not reported or not identified.



Rates of Adults and Adolescents Living with Diagnosed HIV Infection, Year-end 2010—United States and 6 Dependent Areas

N = 888,921 Total Rate = 342.2




Note. Data include persons with a diagnosis of HIV infection regardless of stage of disease at diagnosis. All displayed data have been statistically adjusted to account for reporting delays, but not for incomplete reporting.



Rates of Children Aged <13 Years Living with Diagnosed HIV Infection, Year-end 2010—United States and 6 Dependent Areas

N = 2,936


Total Rate = 5.5



Note. Data include persons with a diagnosis of HIV infection regardless of stage of disease at diagnosis. All displayed data have been statistically adjusted to account for reporting delays, but not for incomplete reporting.




Stage 3 (AIDS) Classifications and Deaths of Persons with HIV Infection Ever Classified as Stage 3 (AIDS), among Adults and Adolescents, 1985–2010—United States and 6 Dependent Areas



Note. All displayed data have been statistically adjusted to account for reporting delays, but not for incomplete reporting. Deaths of persons with HIV infection, stage 3 (AIDS) may be due to any cause.



Stage 3 (AIDS) Classifications among Adults and Adolescents with HIV Infection, by Race/Ethnicity and Year of Diagnosis, 1985–2011—United States and 6 Dependent Areas




Note. All displayed data have been statistically adjusted to account for reporting delays, but not for incomplete reporting.

^a Hispanics/Latinos can be of any race.

^b Includes Asian/Pacific Islander legacy cases.



Percentages of Stage 3 (AIDS) Classifications among Adults and Adolescents with HIV Infection, by Race/Ethnicity and Year of Diagnosis, 1985–2011—United States and 6 Dependent Areas




Note. All displayed data have been statistically adjusted to account for reporting delays, but not for incomplete reporting.

^a Hispanics/Latinos can be of any race.

^b Includes Asian/Pacific Islander legacy cases.



Stage 3 (AIDS) Classifications among Adults and Adolescents with HIV Infection, by Transmission Category and Year of Diagnosis, 1985–2011—United States and 6 Dependent Areas




Note. All displayed data have been statistically adjusted to account for reporting delays and missing transmission category, but not for incomplete reporting.

^a Heterosexual contact with a person known to have, or to be at high risk for, HIV infection.

^b Includes hemophilia, blood transfusion, perinatal exposure, and risk factor not reported or not identified.



Percentages of Stage 3 (AIDS) Classifications among Adults and Adolescents with HIV Infection, by Transmission Category and Year of Diagnosis, 1985–2011—United States and 6 Dependent Areas





Note. All displayed data have been statistically adjusted to account for reporting delays and missing transmission category, but not for incomplete reporting.

^a Heterosexual contact with a person known to have, or to be at high risk for, HIV infection.

^b Includes hemophilia, blood transfusion, perinatal exposure, and risk factor not reported or not identified.



Stage 3 (AIDS) Classifications among Adults and Adolescents with HIV Infection, by Sex and Transmission Category, 2011—United States and 6 Dependent Areas



Male-to-male sexual contact

Injection drug use (IDU)

Male-to-male sexual contact and IDU

Heterosexual contact^a

Other^b

Note. All displayed data have been statistically adjusted to account for reporting delays and missing transmission category, but not for incomplete reporting.

^a Heterosexual contact with a person known to have, or to be at high risk for, HIV infection.

^b Includes hemophilia, blood transfusion, perinatal exposure, and risk factor not reported or not identified.



Stage 3 (AIDS) Classifications among Adults and Adolescents with HIV Infection, by Race/Ethnicity, 2011—United States

Race/ethnicity	No.	Rate
American Indian/Alaska Native	146	8.0
Asian ^a	492	3.9
Black/African American	15,946	51.3
Hispanic/Latino ^b	6,353	16.2
Native Hawaiian/other Pacific Islander	47	11.6
White	8,303	4.9
Multiple races	750	20.0
Total	32,037	12.4

Note. All displayed data have been statistically adjusted to account for reporting delays, but not for incomplete reporting.

Rates are per 100,000 population.

^a Includes Asian/Pacific Islander legacy cases.


^b Hispanics/Latinos can be of any race.



Rates of Stage 3 (AIDS) Classifications among Persons with HIV Infection, 2011—United States and 6 Dependent Areas

N = 32,561


Total Rate = 10.3



Note. All displayed data have been statistically adjusted to account for reporting delays, but not for incomplete reporting. q




Adults and Adolescents Living with Diagnosed HIV Infection Ever Classified as Stage 3 (AIDS), by Sex, 1993–2010—United States and 6 Dependent Areas



Note. All displayed data have been statistically adjusted to account for reporting delays, but not for incomplete reporting.



Persons Living with Diagnosed HIV Infection Ever Classified as Stage 3 (AIDS), by Race/Ethnicity, 1993–2010—United States and 6 Dependent Areas



Note. All displayed data have been statistically adjusted to account for reporting delays, but not for incomplete reporting.


^a Hispanics/Latinos can be of any race.

^b Includes Asian/Pacific Islander legacy cases.



Rates of Adults and Adolescents Living with Diagnosed HIV Infection Ever Classified as Stage 3 (AIDS), Year-end 2010—United States and 6 Dependent Areas

N = 498,704 Total Rate = 192.0




Note. All displayed data have been statistically adjusted to account for reporting delays, but not for incomplete reporting.



Rates of Children Aged <13 Years Living with Diagnosed HIV Infection Ever Classified as Stage 3 (AIDS), Year-end 2010—United States and 6 Dependent Areas

N = 463 Total Rate = 0.9



Note. All displayed data have been statistically adjusted to account for reporting delays, but not for incomplete reporting.

