

Table 18b. Persons living with an AIDS diagnosis, by race/ethnicity and selected characteristics, year-end 2008—United States and 5 U.S. dependent areas

	American Indian/ Alaska Native		Asian ^a		Black/African American		Hispanic/Latino ^b		Native Hawaiian/ Other Pacific Islander		White		Multiple races		Total ^c	
	No.	Est. No. ^d	No.	Est. No. ^d	No.	Est. No. ^d	No.	Est. No. ^d	No.	Est. No. ^d	No.	Est. No. ^d	No.	Est. No. ^d	No.	Est. No. ^{d,e}
Age at end of year																
<13	2	2	6	6	469	481	131	134	2	2	79	82	12	13	701	720
13–14	1	1	3	3	406	413	123	125	1	1	68	70	9	9	612	624
15–19	7	7	9	9	1,843	1,899	567	587	2	2	315	322	40	42	2,783	2,868
20–24	20	21	42	45	3,829	4,037	1,448	1,533	5	5	856	903	104	110	6,304	6,654
25–29	73	78	181	196	8,111	8,490	3,932	4,164	26	27	2,796	2,946	224	236	15,343	16,138
30–34	132	136	406	431	13,568	14,044	7,645	7,978	42	44	6,055	6,288	429	451	28,278	29,371
35–39	263	266	821	862	23,500	24,087	13,435	13,866	64	66	15,536	15,936	758	779	54,388	55,873
40–44	353	352	949	979	37,399	38,046	20,929	21,354	87	89	30,926	31,338	1,118	1,141	91,787	93,325
45–49	371	370	826	845	42,023	42,498	21,412	21,680	94	95	39,823	39,988	1,381	1,403	105,957	106,907
50–54	256	253	644	654	34,971	35,147	15,407	15,501	59	60	29,486	29,368	955	963	81,821	81,989
55–59	137	135	429	433	21,898	21,873	9,033	9,067	34	34	18,829	18,646	610	621	50,993	50,832
60–64	65	64	237	240	10,278	10,234	4,757	4,777	12	12	10,074	9,923	268	269	25,702	25,530
≥65	49	48	177	179	8,046	7,926	4,067	4,025	11	11	7,674	7,476	187	186	20,225	19,865
Transmission category																
Male adult or adolescent																
Male-to-male sexual contact	723	786	2,515	3,152	54,121	65,600	39,026	45,962	264	300	102,341	110,279	2,208	2,555	201,271	228,727
Injection drug use	173	187	150	231	27,487	34,234	16,646	19,795	16	18	11,034	11,914	759	888	56,274	67,287
Male-to-male sexual contact and injection drug use	231	245	165	191	9,964	11,465	5,260	5,876	21	22	12,944	13,514	456	503	29,045	31,822
Heterosexual contact ^f	76	86	318	403	19,760	26,041	7,802	9,123	20	22	4,754	5,313	400	472	33,135	41,467
Other ^g	108	12	779	72	25,279	948	11,189	562	35	2	10,879	1,582	550	45	48,857	3,224
Subtotal	1,311	1,315	3,927	4,049	136,611	138,288	79,923	81,318	356	364	141,952	142,603	4,373	4,463	368,582	372,528
Female adult or adolescent																
Injection drug use	133	158	50	80	14,918	20,313	5,453	6,925	11	14	6,701	7,851	481	630	27,754	35,981
Heterosexual contact ^f	192	237	475	658	33,991	46,851	12,188	15,210	46	60	9,762	11,753	785	1,046	57,451	75,831
Other ^g	82	12	260	78	18,278	1,172	4,444	458	21	6	3,569	550	408	35	27,070	2,311
Subtotal	407	407	785	816	67,187	68,336	22,085	22,593	78	79	20,032	20,154	1,674	1,711	112,275	114,123
Child (<13 yrs at diagnosis)																
Perinatal	10	10	14	14	2,451	2,459	826	828	5	5	478	475	42	42	3,827	3,835
Other ^h	1	1	4	4	91	92	52	52	0	0	55	54	5	5	208	208
Subtotal	11	11	18	18	2,542	2,551	878	880	5	5	533	529	47	47	4,035	4,043

Table 18b. Persons living with an AIDS diagnosis, by race/ethnicity and selected characteristics, year-end 2008—United States and 5 U.S. dependent areas (cont)

	American Indian/ Alaska Native		Asian ^a		Black/African American		Hispanic/Latino ^b		Native Hawaiian/ Other Pacific Islander		White		Multiple races		Total ^c	
	No.	Est. No. ^d	No.	Est. No. ^d	No.	Est. No. ^d	No.	Est. No. ^d	No.	Est. No. ^d	No.	Est. No. ^d	No.	Est. No. ^d	No.	Est. No. ^{d,e}
Region of residence																
Northeast	134	139	1,084	1,141	59,809	61,821	37,419	38,899	24	25	34,259	34,899	2,996	3,123	135,812	140,136
Midwest	204	202	345	357	23,213	23,341	5,075	5,192	15	16	22,159	22,126	654	666	51,668	51,902
South	445	452	655	677	107,291	107,814	23,709	23,871	64	66	57,981	58,081	1,852	1,842	192,021	192,827
West	944	938	2,625	2,687	15,835	16,007	25,756	26,284	329	334	48,071	48,133	584	583	94,182	95,003
U.S. dependent areas	2	2	21	21	193	192	10,927	10,544	7	8	47	48	9	8	11,211	10,828
Totalⁱ	1,729	1,733	4,730	4,883	206,341	209,175	102,886	104,791	439	449	162,517	163,286	6,095	6,222	484,894	490,696

^a Includes Asian/Pacific Islander legacy cases (see Technical Notes).

^b Hispanics/Latinos can be of any race.

^c Includes persons of unknown race/ethnicity.

^d Estimated numbers resulted from statistical adjustment that accounted for reporting delays and missing risk-factor information, but not for incomplete reporting.

^e Because the estimated totals were calculated independently of the corresponding values for each subpopulation, the subpopulation values may not sum to the totals shown here.

^f Heterosexual contact with a person known to have, or to be at high risk for, HIV infection.

^g Includes hemophilia, blood transfusion, perinatal exposure, and risk factor not reported or not identified.

^h Includes hemophilia, blood transfusion, and risk factor not reported or not identified.

ⁱ Because column totals for estimated numbers were calculated independently of the values for the subpopulations, the values in each column may not sum to the column total.