

**Table 4b. AIDS diagnoses, by race/ethnicity and selected characteristics, 2009—United States and 5 U.S. dependent areas**

	American Indian/ Alaska Native		Asian <sup>a</sup>		Black/African American		Hispanic/Latino <sup>b</sup>		Native Hawaiian/ Other Pacific Islander		White		Multiple races		Total	
	No.	Est. No. <sup>c</sup>	No.	Est. No. <sup>c</sup>	No.	Est. No. <sup>c</sup>	No.	Est. No. <sup>c</sup>	No.	Est. No. <sup>c</sup>	No.	Est. No. <sup>c</sup>	No.	Est. No. <sup>c</sup>	No.	Est. No. <sup>c,d</sup>
<b>Age at diagnosis (yr)</b>																
<13	0	0	0	0	6	8	3	3	0	0	1	1	0	0	10	13
13–14	0	0	0	0	24	38	16	21	0	0	0	0	0	0	40	60
15–19	2	2	4	5	257	334	75	94	0	0	32	39	9	13	379	488
20–24	8	9	14	17	1,026	1,298	340	431	2	3	262	319	27	34	1,679	2,110
25–29	17	20	40	55	1,369	1,725	715	924	6	8	601	740	47	61	2,795	3,531
30–34	18	21	57	72	1,504	1,878	896	1,135	4	5	761	948	47	61	3,287	4,120
35–39	18	23	78	96	1,754	2,220	917	1,160	6	7	1,100	1,364	93	125	3,966	4,996
40–44	22	27	55	72	2,048	2,607	1,001	1,276	5	6	1,392	1,731	92	121	4,615	5,840
45–49	19	24	39	54	2,031	2,589	787	1,009	8	10	1,433	1,809	85	106	4,402	5,602
50–54	14	19	13	17	1,590	2,029	514	660	6	8	1,005	1,258	70	90	3,212	4,080
55–59	7	8	11	14	860	1,095	288	371	1	1	558	706	37	49	1,762	2,244
60–64	1	1	10	12	406	518	141	181	2	2	259	315	8	10	827	1,039
≥65	1	1	8	14	331	420	140	177	1	1	194	242	13	17	688	872
<b>Transmission category</b>																
<b>Male adult or adolescent</b>																
Male-to-male sexual contact	64	90	145	260	3,653	6,191	2,377	3,806	26	37	4,405	6,460	221	328	10,891	17,171
Injection drug use	10	15	12	21	742	1,654	479	886	1	1	386	575	31	54	1,661	3,207
Male-to-male sexual contact and injection drug use	8	11	9	17	295	550	210	330	4	5	470	658	24	38	1,020	1,608
Heterosexual contact <sup>e</sup>	7	10	30	49	1,402	2,635	492	799	1	1	258	390	47	73	2,237	3,956
Other <sup>f</sup>	13	2	70	3	2,599	79	1,037	32	4	0	981	43	57	1	4,761	159
<b>Subtotal</b>	<b>102</b>	<b>127</b>	<b>266</b>	<b>349</b>	<b>8,691</b>	<b>11,109</b>	<b>4,595</b>	<b>5,852</b>	<b>36</b>	<b>45</b>	<b>6,500</b>	<b>8,126</b>	<b>380</b>	<b>495</b>	<b>20,570</b>	<b>26,102</b>
<b>Female adult or adolescent</b>																
Injection drug use	4	8	2	6	474	1,135	179	366	0	0	245	409	32	57	936	1,982
Heterosexual contact <sup>e</sup>	13	20	28	68	2,109	4,414	636	1,186	4	7	502	915	75	130	3,367	6,740
Other <sup>f</sup>	8	0	33	5	1,926	93	420	34	1	0	350	20	41	4	2,779	157
<b>Subtotal</b>	<b>25</b>	<b>28</b>	<b>63</b>	<b>80</b>	<b>4,509</b>	<b>5,642</b>	<b>1,235</b>	<b>1,586</b>	<b>5</b>	<b>7</b>	<b>1,097</b>	<b>1,344</b>	<b>148</b>	<b>192</b>	<b>7,082</b>	<b>8,879</b>
<b>Child (&lt;13 yrs at diagnosis)</b>																
Perinatal	0	0	0	0	6	8	2	2	0	0	1	1	0	0	9	12
Other <sup>g</sup>	0	0	0	0	0	0	1	1	0	0	0	0	0	0	1	1
<b>Subtotal</b>	<b>0</b>	<b>0</b>	<b>0</b>	<b>0</b>	<b>6</b>	<b>8</b>	<b>3</b>	<b>3</b>	<b>0</b>	<b>0</b>	<b>1</b>	<b>1</b>	<b>0</b>	<b>0</b>	<b>10</b>	<b>13</b>

**Table 4b. AIDS diagnoses, by race/ethnicity and selected characteristics, 2009—United States and 5 U.S. dependent areas (cont)**

	American Indian/ Alaska Native		Asian <sup>a</sup>		Black/African American		Hispanic/Latino <sup>b</sup>		Native Hawaiian/ Other Pacific Islander		White		Multiple races		Total	
	No.	Est. No. <sup>c</sup>	No.	Est. No. <sup>c</sup>	No.	Est. No. <sup>c</sup>	No.	Est. No. <sup>c</sup>	No.	Est. No. <sup>c</sup>	No.	Est. No. <sup>c</sup>	No.	Est. No. <sup>c</sup>	No.	Est. No. <sup>c,d</sup>
<b>Region of residence</b>																
Northeast	2	3	68	99	2,787	3,956	1,550	2,186	8	11	1,221	1,681	171	236	5,807	8,171
Midwest	16	26	30	37	1,702	2,178	344	449	1	1	1,264	1,602	78	101	3,435	4,394
South	30	34	76	95	7,862	9,525	1,881	2,171	8	10	3,108	3,693	223	278	13,188	15,806
West	79	94	155	197	845	1,082	1,510	1,913	23	28	2,002	2,490	56	71	4,670	5,875
U.S. dependent areas	0	0	0	0	10	18	548	723	1	1	3	4	0	0	562	747
<b>Total<sup>h</sup></b>	<b>127</b>	<b>155</b>	<b>329</b>	<b>429</b>	<b>13,206</b>	<b>16,759</b>	<b>5,833</b>	<b>7,442</b>	<b>41</b>	<b>52</b>	<b>7,598</b>	<b>9,471</b>	<b>528</b>	<b>686</b>	<b>27,662</b>	<b>34,993</b>

<sup>a</sup> Includes Asian/Pacific Islander legacy cases (see Technical Notes).

<sup>b</sup> Hispanics/Latinos can be of any race.

<sup>c</sup> Estimated numbers resulted from statistical adjustment that accounted for reporting delays and missing risk-factor information, but not for incomplete reporting.

<sup>d</sup> Because the estimated totals were calculated independently of the corresponding values for each subpopulation, the subpopulation values may not sum to the totals shown here.

<sup>e</sup> Heterosexual contact with a person known to have, or to be at high risk for, HIV infection.

<sup>f</sup> Includes hemophilia, blood transfusion, perinatal exposure, and risk factor not reported or not identified.

<sup>g</sup> Includes hemophilia, blood transfusion, and risk factor not reported or not identified.

<sup>h</sup> Because column totals for estimated numbers were calculated independently of the values for the subpopulations, the values in each column may not sum to the column total.