

Table 9. AIDS diagnoses among adult and adolescent Hispanics/Latinos, by transmission category and place of birth, 2009—United States and 5 U.S. dependent areas

Transmission category	Place of birth																				
	United States			Central America			South America			Cuba			Mexico			Puerto Rico			Total		
	No.	Est. No. ^a	%	No.	Est. No. ^a	%	No.	Est. No. ^a	%	No.	Est. No. ^a	%	No.	Est. No. ^a	%	No.	Est. No. ^a	%	No. ^b	Est. No. ^{a,b}	%
Male adult or adolescent																					
Male-to-male sexual contact	964	1,412	67.0	148	302	65.9	121	177	76.0	80	100	69.6	525	868	75.2	161	236	33.0	2,377	3,806	65.0
Injection drug use	176	328	15.6	8	37	8.0	1	12	5.0	8	13	9.0	33	71	6.2	194	273	38.1	479	886	15.1
Male-to-male sexual contact and injection drug use	115	171	8.1	6	15	3.3	6	8	3.4	3	6	3.9	26	43	3.7	35	50	7.0	210	330	5.6
Heterosexual contact ^c	126	181	8.6	56	102	22.3	18	36	15.5	14	23	16.0	95	170	14.7	104	155	21.7	492	799	13.7
Other ^d	282	16	0.8	147	2	0.5	29	0	0.2	19	2	1.6	235	2	0.2	53	2	0.2	1,037	32	0.5
Subtotal	1,663	2,109	100	365	459	100	175	233	100	124	144	100	914	1,155	100	547	715	100	4,595	5,852	100
Female adult or adolescent																					
Injection drug use	102	198	32.4	2	9	7.2	1	5	12.7	1	2	9.7	7	17	10.2	43	64	20.4	179	366	23.1
Heterosexual contact ^c	210	393	64.4	57	114	90.4	17	33	86.6	14	18	90.3	80	144	88.8	159	246	77.9	636	1,186	74.8
Other ^d	166	19	3.2	40	3	2.3	12	0	0.7	2	0	0.0	43	2	1.0	42	5	1.7	420	34	2.1
Subtotal	478	610	100	99	126	100	30	38	100	17	20	100	130	162	100	244	316	100	1,235	1,586	100
Total^e	2,141	2,718	100	464	585	100	205	271	100	141	164	100	1,044	1,317	100	791	1,031	100	5,830	7,438	100

^a Estimated numbers resulted from statistical adjustment that accounted for reporting delays and missing risk-factor information, but not for incomplete reporting.

^b Entries include persons whose place of birth is not among those listed and persons whose place of birth is unknown.

^c Heterosexual contact with a person known to have, or to be at high risk for, HIV infection.

^d Includes hemophilia, blood transfusion, perinatal exposure, and risk factor not reported or not identified.

^e Because column totals for estimated numbers were calculated independently of the values for the subpopulations, the values in each column may not sum to the column total.