

Table 6. HIV infection cases¹ by age group, exposure category, and sex, reported through December 2001, from areas with confidential HIV infection reporting²

Adult/adolescent exposure category	Males				Females				Totals ³			
	2001		Cumulative total		2001		Cumulative total		2001		Cumulative total	
	No.	(%)	No.	(%)	No.	(%)	No.	(%)	No.	(%)	No.	(%)
Men who have sex with men	7,674	(32)	52,139	(43)	—	—	—	—	7,674	(22)	52,139	(31)
Injecting drug use	1,844	(8)	14,904	(12)	1,097	(10)	8,609	(17)	2,941	(8)	23,514	(14)
Men who have sex with men and inject drugs	614	(3)	6,651	(6)	—	—	—	—	614	(2)	6,651	(4)
Hemophilia/coagulation disorder	24	(0)	455	(0)	6	(0)	37	(0)	30	(0)	492	(0)
Heterosexual contact:	1,466	(6)	8,597	(7)	3,071	(28)	19,157	(39)	4,537	(13)	27,754	(16)
<i>Sex with injecting drug user</i>	208		1,694		544		4,594		752		6,288	
<i>Sex with bisexual male</i>	—		—		150		1,330		150		1,330	
<i>Sex with person with hemophilia</i>	4		17		13		147		17		164	
<i>Sex with transfusion recipient with HIV infection</i>	7		82		11		120		18		202	
<i>Sex with HIV-infected person, risk not specified</i>	1,247		6,804		2,353		12,966		3,600		19,770	
Receipt of blood transfusion, blood components, or tissue	47	(0)	447	(0)	52	(0)	467	(1)	99	(0)	914	(1)
Other/risk not reported or identified ⁴	12,230	(51)	37,675	(31)	6,907	(62)	20,956	(43)	19,137	(55)	58,639	(34)
Adult/adolescent subtotal	23,899	(100)	120,868	(100)	11,133	(100)	49,226	(100)	35,032	(100)	170,103	(100)
Pediatric (<13 years old) exposure category												
Hemophilia/coagulation disorder	8	(3)	107	(6)	1	(0)	2	(0)	9	(2)	109	(3)
Mother with/at risk for HIV infection:	205	(73)	1,603	(83)	185	(71)	1,733	(87)	390	(72)	3,336	(85)
<i>Injecting drug use</i>	31		472		30		473		61		945	
<i>Sex with injecting drug user</i>	17		168		7		178		24		346	
<i>Sex with bisexual male</i>	2		19		0		15		2		34	
<i>Sex with person with hemophilia</i>	0		1		0		7		0		8	
<i>Sex with transfusion recipient with HIV infection</i>	0		8		0		5		0		13	
<i>Sex with HIV-infected person, risk not specified</i>	47		317		56		403		103		720	
<i>Receipt of blood transfusion, blood components, or tissue</i>	4		15		1		16		5		31	
<i>Has HIV infection, risk not specified</i>	104		603		91		636		195		1,239	
Receipt of blood transfusion, blood components, or tissue	2	(1)	21	(1)	2	(1)	26	(1)	4	(1)	47	(1)
Risk not reported or identified ⁴	67	(24)	202	(10)	73	(28)	229	(12)	140	(26)	431	(11)
Pediatric subtotal	282	(100)	1,933	(100)	261	(100)	1,990	(100)	543	(100)	3,923	(100)
Total	24,181		122,801		11,394		51,216		35,575		174,026	

¹Includes only persons reported with HIV infection who have not developed AIDS.

²See table 3 for areas with confidential HIV infection reporting.

³Includes 9 persons whose sex is unknown.

⁴For HIV infection cases, "risk not reported or identified" refers primarily to persons whose mode of exposure was not reported and who have not been followed up to determine their mode of exposure, and to a smaller number of persons who are not reported with one of the exposures listed above after follow-up. See Technical Notes.